

324
E.J.
55573
142

MALAYAN FERNS.

MALAYAN ARCHIVES

QR
529
A4115Z
Bot.

MALAYAN FERNS.

HANDBOOK

TO THE DETERMINATION

OF

THE FERNS

OF

THE MALAYAN ISLANDS

(INCL. THOSE OF THE MALAY PENINSULA, THE PHILIPPINES
AND NEW GUINEA).

BY

*Cornelis
J. van
Sullow
Ard*
CAPT. C. R. W. K. VAN ALDERWERELT VAN ROSENBURGH.

PUBLISHED BY

THE DEPARTMENT OF AGRICULTURE

NETHERLANDS INDIA.

BATAVIA
LANDSDRUKKERIJ
1908.

211757
C

287.5
A355
Bot

PREFACE.

Handbooks are like watches: the worst is better than none, and the best cannot be expected to go quite true.

Dr. JOHNSON. — (Variant).

When acting for some few years as an Assistant Curator at the Buitenzorg Herbarium I was directed to elaborate the non-determined fern-material of the Malay Archipelago. Malayan Ferns have much been written about in handbooks and periodicals, but seeing no work exists treating with them more particularly I resolved to collect all available descriptions in one volume. Originally my intention was to do so for my private use, but it occurred to me that publication of this compilation might prove to be of some value to those interested in the Malayan Ferns. Moreover the Director of the Agricultural Department in Netherlands India, Prof. Dr. M. TREUB, desired the work to be printed.

The descriptions are taken from various sources and have been compared with living or dried material if on hand in the Buitenzorg Botanical Gardens or in the Herbarium. Those thus controlled and more or less modified if necessary are marked with an asterisk. I used in the first place RACIBORSKI'S *Die Pteridophyten der Flora von Buitenzorg* and in the second HOOKER & BAKER'S *Synopsis Filicum*, the works my studies in pteridology are founded upon and only in cases where these books could not provide me with such information as required I had recourse to other sources, principally CHRIST'S *Die Farnkräuter der Erde*, BEDDOME'S *Handbook to the Ferns of British India* and COPELAND'S *The Polypodiaceae of the Philippine Islands*. Thence specific names and synonyms are not followed by an exhaustive and somewhat unnecessary list of old sources, but in most cases only by one or at the utmost two of the newer sources consulted. Likewise enumeration of synonyms has been limited as much as possible. Those desirous to have further particulars on the subject should consult CHRISTENSEN'S *Index Filicum*.

A definite time being fixed for the completion of the work, I saw myself

compelled to discontinue the proper compilation-work just when the last part of the *Index Filicum* appeared, the remainder of my term of office to be spared for revising, comparing and classifying the various genera and species. Therefore no diagnoses published after January 1st, 1907 are to be found in this work, except those which can be inserted while printing, but as a matter of fact I have not been able to do much more than enumerating as synonyms the new names given to well-known species in consequence of changed conceptions.

I regret that owing to the want of several books in the Buitenzorg Library I am not in a position to give diagnoses of some few species of ferns, a lack however which may be considered of but little importance seeing the ferns in question belong to species which are rather rare. Nevertheless I intend to procure their descriptions and publish them later on in an appendix or other supplementary work, together with the other new species that have still been discovered since the date above quoted.

In giving the diagnoses of the various species I have endeavoured to stick to facts only and to avoid any speculative opinion even when my conception or errors in other authors' work may have tempted me to alter specific names. Of course I do not claim completeness or entire correctness and shall be pleased to have the no doubt many errors and omissions pointed out to me in order to correct them in a supplement.

As to the limitation and the nomenclature of the genera and species I have chiefly followed the *Index Filicum*, which with a few exceptions agrees with **DIELS'** fern-system in **ENGLER & PRANTL'S** *Die natürlichen Pflanzenfamilien*, Vol. I, part 4. In both works however a *natural system* has been adopted, but as the object of this book is *determination* only, I have deemed it more practical to adopt for my purpose an *artificial arrangement*. Consequently some clashing between system and arrangement could not be avoided, the same as some few modifications in **CHRISTENSEN'S** limitation of the genera. So, for instance, I have kept the truly indusiate ferns separate from those which have no indusium or only a spurious one. Accordingly my arrangement is of no systematic value, but only a guide to those studying Malayan Pteridology. Therefore I have quoted under each species the name given in the *Index Filicum* wherever I followed my own conception. Where I, for convenience' sake, refer to **DIELS**, in **ENGL. & PRANTL**, Nat. Pfl.Fam., I⁴, one should bear in mind that **DIELS'** *fern-system* is meant, for the articles on the Hymenophyllaceae are written by **SADEBECK** and those on the Marattiaceae and Ophioglossaceae by **BITTER**.

The reason why the ferns found in Malacca (the Malay Peninsula), the Philippines and New Guinea are inserted in this work on the ferns of the Malay Archipelago is that in botany no strict limits may be expected to exist, and fern-species growing closely outside geographical boundaries may reasonably be expected to be found inside these boundaries as well.

In giving habitats I have confined myself to regions and not merely to the places where the species are reported to have been found. By *Malaya* is understood the entire territory this work is dealing with. The smaller islands, if not treated of separately, are included in the nearest larger territory to which they geographically belong. It will however not be advisable to adhere too closely to the habitats given, seeing it is not at all impossible that species, up till now found in one place only, may also be discovered elsewhere far away from the present known place of origin.

For this work the English language was chosen only in behalf of the foreigners who visit these regions, seeing a book written in Dutch would be useless to them.

In conclusion I beg to tender my best thanks to all who assisted me to render the completion of this work possible and especially to Mrs. H. GUDE—*Murray*, who kindly undertook the grammatical correction of the work.

BUITENZORG, January 1908.

v. A. v. R.

ABBREVIATIONS.

Abh. Nat. Ver. Brem.	= Naturwissenschaftlicher Verein zu Bremen; Abhandlungen.
A. Br.	= A. Braun.
A. Cunn.	= Allan Cunningham.
Adans.	= M. Adanson.
Agardh. (Ag).	= J. G. Agardh.
Ait.	= W. Aiton.
Amann	= J. Amann (? Kurz).
Anal.	= <i>Analecta pteridographica</i> , etc. (Kze, 1857).
Ann. Botz.	= <i>Annales du Jardin botanique de Buitenzorg</i> .
Ann. Mus. Bot. L. B.	= <i>Annales Musei Botanici Lugduno-Batavi</i> . (Miq., 1867—'69).
Ann. Nat. Hist.	= <i>Annals of Natural History</i> .
Ann. of Bot.	= <i>Annals of Botany</i> .
Bail.	= F. M. Bailey.
Balf.	= J. Baylay Balfour.
Bau. & Hk.	= F. Bauer & W. J. Hooker.
Becc.	= O. Beccari.
Bedd.	= R. H. Beddome.
Beech. Voy.	= <i>The botany of Captain Beechey's Voyage</i> . (Hk. & Arn., 1830—'41).
Bél.	= Bélanger.
Berk.	= Berkeley.
Berl. Mag.	= <i>Der Gesellschaft Naturforschender Freunde zu Berlin Magazin etc.</i> (1807—'18).
Bernh.	= J. J. Bernhardt.
Bitt.	= G. Bitter.
Bk. (Bak.)	= J. G. Baker.
Bl.	= C. L. Blume.
Bory	= Bory de Saint-Vincent.
Bot. Centralbl.	= <i>Botanisches Centralblatt</i> .

Bot. Jahrb.	= Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie. (Engl.).
Bot. Misc.	= Botanical Miscellany.
Bot. Voy. Herald.	= Botany of the Voyage of H. M. S. Herald. (Seemann, 1854—'57).
Bot. Zeit.	= Botanische Zeitung. (Mohl & Schlechtendal).
Brack.	= Brackenridge.
Br. Ferns	= The British Ferns, or coloured figures and descriptions, etc. (Hk., 1861).
Brong.	= A. Brongniart.
Bull. Ac. Crac.	= Bulletin international de l'Académie des Sciences de Cracovie.
Bull. Ac. Int. Bot.	= Bulletin de l'Académie internationale de géographie botanique.
Bull. Dép. Agr. I. N.	= Bulletin du Département de l'Agriculture aux Indes Néerlandaises.
Bull. Herb. Boiss.	= Bulletin de l'Herbier Boissier.
Bull. Soc. Bot. Belg.	= Bulletin de la Société botanique de Belgique.
Bull. Soc. Bot. Fr.	= Bulletin de la Société botanique de France.
Bull. Torr. Cl.	= Bulletin of the Torrey Botanical Club.
Burck	= W. Burck.
Burm.	= N. L. Burmann.
Carr.	= W. Carruthers.
Cav.	= A. J. Cavanilles.
2 nd Cent. of Ferns.	= A second Century of Ferns. (Hk., 1861).
C. Chr.	= C. Christensen.
Ces.	= V. Cesati.
Christ	= H. Christ.
Clarke	= C. B. Clarke.
Colen.	= W. Colenso.
Copel.	= E. B. Copeland.
Dec.	= M. J. Decaisne.
Descr.	= Descripción de las Plantas que D. Antonio Josef Cavanilles demostró en las lecciones publicas del año 1801. (Cav., 1802).
Desv.	= A. N. Desvaux.

- de Vr. = W. H. de Vriese.
 de Vr. & Teysm. = de Vriese & Teysmann.
 Dict. d' Hort. = Dictionnaire d' Horticulture et de Jardinage.
 (Nich.).
 Diels = L. Diels.
 Dissert. = Dissertatio inauguralis medico-botanica de
 duabus novis Trichomanarum speciebus de
 earum nec non aliarum huius generis plan-
 tarum structura. (Taschn., 1845).
 Don = D. Don.
 Dry. = J. Dryander.
 Eat. = D. C. Eaton.
 Endl. = S. L. Endlicher.
 Engl. = A. Engler.
 Engl. Jahrb. = Botanische Jahrbücher für Systematik,
 Pflanzengeschichte und Pflanzengeographie,
 ed. A. Engler.
 Engl. & Prantl = A. Engler & K. Prantl.
 Enum. = Enumeratio Plantarum Javae et Insularum
 adjacentium. (Bl., 1828).
 Enum. Fil. = Enumeratio Filicum, etc. (Klf., 1824).
 Epim. = Epimeliae botanicae. (Pr., 1849).
 Exot. Flor. = Exotic Flora. (Hk., 1823—'27).
 Expl. Exp. = U.S. Exploration Expedition, XVI. (Wilk.,
 1854).
 Farng. = Über einige Farngattungen. (Mett.); I, Poly-
 podium (1857); II, Plagiogyria (1858); III,
 Pteris (1858); IV, Phegopteris & Aspidium
 (1858); V, Cheilanthes (1859); VI, Asple-
 nium (1859).
 Farnkr. = Die Farnkräuter in kolorirten Abbildungen,
 etc. (Kze, 1840—'51).
 Farnkr. d. Erde = Die Farnkräuter der Erde. (Christ, 1897).
 Fed. = F. Fedde.
 Fée = A. L. A. Fée.
 Fel. Becc. Polin. = Prospetto delle Felci raccolte dal Signor O.
 Beccari nella Polinesia, etc. — Rend. Acad.
 Napoli, XVI. — (Ges., 1877).

- Fel. di Bor. = Felci e specie nei gruppi affini raccolte a Borneo dal Signor O. Beccari. — Atti Acad. Sci. fis. e mat. Napoli. — (Ces., 1876).
- Ferns Br.I. = The Ferns of British India. (Bedd., 1865—'70).
- Ferns S.I. = The Ferns of Southern India (Bedd., 1863—'64).
- Ferns S.Br.I., Suppl. = Supplement to the Ferns of Southern India and British India. (Bedd., 1876).
- Fil. Exot. = Filices Exoticae or Figures and Descriptions of exotic Ferns, etc. (Hk., 1857—'59).
- Fil. Gen. & Spec. = Filicum Genera et Species recentiori Methodo accomodatae analytice descriptae. (Hedw., 1799—1803).
- Fil. Lechl. = Filices Lechlerianae. (Mett., 1856—'59).
- Fil. Saras. = Filices Sarasinianae, in »Verhandlungen der Naturforschenden Gesellschaft in Basel, XI.» — (Christ, 1895).
- Fisch. = Fischer.
- Flor. Btz. = Die Pteridophyten der Flora von Buitenzorg (Band I der Flora von Buitenzorg; Rac., 1898).
- Flor. deut. Sch.geb. = Die Flora der deutschen Schutzgebiete in der Südsee. (Schum. & Laut., 1901). — Suppl. (1905).
- Flor. Filip. = Flora de Filipinas, etc. (Blanco, 1837).
- Flor. Jav., II. = Flora Javae nec non Insularum adjacentium. Filices. (Bl. 1828—'29).
- Flor. Kais. Wilhl. = Die Flora von Kaiser Wilhelmsland. (Schum. & Hollr., 1889).
- Flor. Trist. d'Ac. = Esquisse de la Flore de Tristan d'Acugna. (Thouars, 1804?).
- Forb. = H. O. Forbes.
- Forschr. Gaz., IV, Farne. = Farne und bärlappartige Gewächse in „Die Forschungsreise S. M. S. Gazelle.“ (1874—'76).
- Forsk. = P. Forskål.
- Forst. = G. Forster.
- Fragm. = Fragmenta Phytographiae Australiae. (Muell., 1864—'81).

Fragm. Philipp.	= Fragmenta Florae Philippinae. (Perk., 1905).
Freyc.	= Freycinet.
Gard. Chron.	= The Gardener's Chronicle.
» » , N. S.	= » » » , New Series.
Gard. Ferns	= Garden Ferns. (Hk., 1862).
Gardn.	= G. Gardner.
Gaud.	= C. Gaudichaud Beaupré.
Gen.	= Genera Filicum. (5 ^e Mémoire sur la famille des Fougères; Fée, 1850—'52).
Gen. Fil.	= Genera Filicum. (Schott, 1854).
Gen. Fil.	= Genera Filicum. (Bau. & Hk., 1842).
Ges. Nat. Fr.	= Gesellschaft Naturforschender Freunde, Berlin.
Gies.	= K. Giesenhagen.
Gled.	= J. G. Gleditsch.
Grev.	= R. K. Greville.
Grev. & Hk.	= R. K. Greville & W. J. Hooker.
Griff.	= W. Griffith.
Gris.	= A. Grisebach.
Hance	= H. F. Hance.
Handb. Ind. Ferns.	= Handbook to the Ferns of British India, Ceylon and the Malay Peninsula. (Bedd., 1885 .
» » » , Suppl.	= Handbook to the Ferns of British India, Ceylon and the Malay Peninsula, Supplement. (Bedd., 1892).
Harr.	= M. V. Harrington.
Hassk.	= J. K. Hasskarl.
Hedw.	= J. Hedwig.
Herb. Berol.	= Herbarium Beroliense (Berlin).
Herb. Bog.	= Herbarium Bogoriense (Buitenzorg).
Hier.	= G. Hieronymus.
Hist. Fil.	= Historia Filicum. (J. Sm., 1875).
Hk. (Hook.)	= W. J. Hooker.
Hk. & Arn.	= W. J. Hooker & G. A. Walker—Arnott.
Hk. Bk.	= W. J. Hooker & J. G. Baker.
Hk. & Grev.	= W. J. Hooker & R. K. Greville.
Hoffm.	= G. F. Hoffmann.
hort.	= hortus.
Hort. Bog.	= Hortus Bogoriensis.

Houtt.	= M. Houttuyn.
Huds.	= W. Hudson
Humb. & Bonpl.	= A. von Humboldt & A. Bonpland.
Hym.	= Hymenophyllaceae. (Pr., 1843).
Hymen. Jav.	= Hymenophyllaceae Javanicae etc. (V. D. B., 1861).
Ic. Fil.	= Icones Filicum. (Hk. & Grev., 1827—'31)
Icon. Pl.	= Icones plantarum. (Hk).
Ind.	= Index Filicum. (Moore, 1857—'62).
Ind. Ferns	= Handb. Ind. Ferns.
Ind. Fil.	= Index Filicum. (C. Chr., 1905—'06).
Journ. of Bot.	= The Journal of Botany, British and Foreign
Journ. of Linn. Soc., Bot.	= Journal of the Linnaean Society; Botany.
J. Sm.	= John Smith.
Karst.	= G. Karsten.
Kell.	= A. Kellogg.
Kew Bull.	= Royal Gardens Kew Bulletin of miscellaneous information
Keys.	= A. Keyserling.
Kl.	= J. F. Klotzsch.
Klf. (Kaulf.)	= G. F. Kaulfuss.
K. Müll.	= K. Müller.
Kön.	= D. König.
Kuhn	= Max. Kuhn.
Kurz	= S. Kurz.
Kze	= G. Kunze.
L. (Linné)	= Linnaeus (C. von Linné).
Lab.	= de Labillardière.
Lam.	= J. B. A. P. Monet de Lamarck
l.c.	= loco citato.
Leafl.	= Leaflets on Philippine Botany.
L. f.	= C. von Linné, fil.
L. & F.	= G. H. Langsdorff & F. Fischer.
L' Hérit.	= L' Héritier de Brutelle.
Link.	= H. F. Link.
Linn.	= Linnaea, Ein Journal für die Botanik, u.s.w.
List.	= A numerical List of dried specimens of plants, etc. (Wall., 1828—'29).

- Lond. Journ. of Bot. = The London Journal of Botany.
 Luer. = C. Luer.
 Mal. = Malesia. (Becc., 1886).
 Mann = H. Mann.
 Masters = Maxwell Tylden Masters.
 Mém. = Mémoires sur la famille des Fougères. (Fée, 1844—'66).
 Mett. = G. Mettenius.
 Mchx = A. Michaux.
 Meyen = F. J. F. Meyen.
 Milde = J. Milde.
 Miq. = F. A. V. Miquel.
 Mon. Maratt. = Monographie des Marattiacées. (de Vr., 1855).
 Mons. = Monsunia. (Warb., 1901).
 Moore = T. Moore.
 Morph. d. Gefässcr. = Untersuchungen zur Morphologie der Gefässcryptogamen. (Prantl); I, Hymen. (= Die Hymenophyllaceen, 1875); II, Schiz. (= Die Schizaeaceen, 1881).
 MS. = Manuscript.
 Muell. = F. von Mueller.
 Muell. & Bk. = F. von Mueller & J. G. Baker.
 Nat. Geneesk. Arch. = Natuur- en Geneeskundig Archief voor Nederlandsch-Indië.
 Nat. Pfl.Fam., I⁴. = Die natürlichen Pflanzenfamilien, Teil I, Abteilung 4. Pteridophyta. (Engl. & Prantl, 1902).
 Nat. Tydschr. Ned. Ind. = Natuurkundig Tijdschrift voor Nederlandsch-Indië.
 Ned. Kruidk. Arch. = Nederlandsch kruidkundig Archief.
 Nees = Nees ab Esenbeck.
 Nees & Bl. = Nees ab Esenbeck & Blume.
 Nich. = G. Nicholson.
 Niph. = Die Farngattung Niphobolus. (Gies., 1901).
 Nouv. Ann. Mus. = Nouvelles annales du Musée d'Histoire naturelle.
 Nova Acta = Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae curiosorum.

Nov. Holl.	= Novae Hollandiae plantarum specimen. (Lab., 1806).
Obs. Fil. Jav.	= Observationes Botanicae, etc., in »Verhandlungen der Naturkundige Vereeniging in Nederlandsch-Indië.»
O. Ktze	= O. Kuntze.
Ophiogl.	= Beiträge zur Systematik der Ophioglosseen (Prantl), in »Jahrbuch des Kgl. botanischen Gartens, etc. zu Berlin, III.» (1884).
Pap. Pl.	= Descriptive notes on Papuan Plants. (Muell., 1875—'85).
Par.	= Parish.
Perk.	= J. Perkins.
Philipp. Journ.	= The Philippine Journal of Science. (I + Suppl.; II ^c , Bot.; III ^c , Bot.).
Pl. escul.	= De plantis esculentis insularum oceani australis, &c. (Forst. 1786).
Poir.	= J. L. M. Poiret.
Pol. Cyath. Bung.	= Polypodiaceae & Cyatheaceae Herbarii Bungeani. (Keys., 1873).
Polypod. Philipp.	= The Polypodiaceae of the Philippine Islands. (Copel., 1905).
p. p.	= pro parte.
Pr.	= C. B. Presl.
Pr. Amer. Ac.	= Proceedings of the American Academy of Arts and Science.
Prantl	= K. Prantl.
Prod.	= Prodrome de la famille des fougères, in »Mémoires de la Soc. Linnéenne, VI.» (Desv., 1827).
Prod. Flor. Nepal.	= Prodomus Florae Nepalensis. (Don, 1825).
Prod. N.H.	= Prodomus Florae Novae-Hollandiae et Insulae Van Diemen. (R. Br., 1803).
Queensl. Agr. Journ.	= The Queensland Agricultural Journal.
Rac.	= M. Raciborski.
Raddi	= J. Raddi.
Rb. (Roxb.)	= W. Roxburgh.
R. Br.	= R. Brown.

Rec.	= Recensio specierum generis Pteridis. (Ag., 1839).
Reinw. (Rw.)	= C. G. C. Reinwardt.
Rel. Haenk.	= Pteridophyta, in »Reliquiae Haenkeanae.» (Pr., 1825).
Rep.	= Repertorium novarum specierum regni vegetabilis. (Fed.).
Rich.	= L. Richard.
Ridl.	= H. N. Ridley.
Rev. Gen. Pl.	= Revisio generum plantarum, etc. (O. Ktze, 1891).
Roth	= A. W. Roth.
Rst. (Rosenst.)	= E. Rosenstock.
Rud.	= Rudolphi.
Sadeb.	= R. Sadebeck.
Schk.	= C. Schkuhr.
Schl.	= D. F. L. von Schlechtendal.
Schott	= H. Schott.
Schrad.	= H. A. Schrader.
Schroet.	= C. Schroeter.
Schum. & Hollr.	= K. Schumann & M. Hollrung.
Schum. & Laut.	= K. Schumann & K. Lauterbach.
Scort.	= Revd Father Scortechini.
Scott	= J. Scott.
Sieb.	= C. von Siebold.
Sm.	= J. E. Smith.
Spec.	= Caroli à Liné. Species plantarum. (Willd., 1810).
Spec. Fil.	= Species Filicum. (Hk., 1846—'64).
Spr.	= C. Sprengl.
Sw.	= O. Swartz.
St. Hil.	= St. Hilaire.
Syn.	= Synopsis Filicum. (Sw., 1806).
Syn. Fil.	= Synopsis Filicum. (Hk. Bk., 1874, 1885).
Syst.	= Carolini Linnaei. Systema Vegetabilium ed. XVI curante Curtio Sprengl. (1827).
Syst. Verz.	= Systematisches Verzeichnis, etc. (Zoll., 1854).
Taschn.	= C. F. Taschner.

Tasm. Journ.	= Tasmanian Journal of Natural Science.
T. & B.	= Teysmann & Binnendijk.
Tent. Pterid.	= Tentamen Pteridographiae, etc. (Pr., 1836).
Thb. (Thunb.)	= K. P. Thunberg.
Thouars	= Aubert du Petit-Thouars.
Thw.	= G. H. K. Thwaites.
Trans. Linn. Soc., Bot.	= The Transaction of the Linnaean Society of London; Botany.
Trev.	= V. Trevisan.
Tijdschr. N. Gesch. Phys.	= Tijdschrift voor Nat. Geschiedenis en Physiologie.
Und.	= L. Underwood.
Urb.	= J. Urban.
V. D. B.	= R. van den Bosch.
Verh. Zool. Bot. Ges.	= Verhandlungen der K.K. Zoologischen-botanischen Gesellschaft in Wien.
Voy. Bot.	= Voyage aux Indes orientales; Botanique. (Bél., 1835).
Voy. Ur. Phys., Bot.	= Voyage autour du monde, etc. sur l'Uranie et la Physicienne; Botanique. (Freyc., 1826—'29).
Wall.	= N. Wallich.
Warb.	= O. Warburg.
Wawra	= H. Wawra.
Wilk.	= Ch. Wilkes.
Willd.	= K. L. Willdenow.
Zeill.	= Zeiller.
Zenk.	= J. C. Zenker.
Zipp.	= Zippelius.
Zoll.	= H. Zollinger.
Zoll. & Mor.	= H. Zollinger & A. Moritz.

COMPENDIUM

OF THE FAMILIES, TRIBES AND GENERA AS
FOLLOWED IN THIS WORK.

ORDER FILICALES

Green plants with 2 distinct generations, a pro-embryonal and an embryonal, the pro-embryonal consisting of a minute, tuberous, flat or filiform, green or white, simple, entire, incised or branched thalloid plant (prothallium) with hair-like roots and bearing one or more, ♂ and (or) ♀, tuberculiform protuberances (antheridia, archegonia). The embryonal generation, i. e. the normal, cormophytical plant with a solid, short or elongate, rarely tuberous stem (rhizome or caudex) and true roots, stalks (stipes) and leaves (fronds), traversed by one or more fibro-vascular bundles and producing capsules (sporangia) with spores, develops when a ripe archegonium (♀) is fecundated by the spiral-shaped spermatozoids of a ripe antheridium (♂). Fronds seriate or circinate, rarely placed in whorls but then not connected, commonly more strongly developed than the stems, the fertile ones whether or not contracted, rarely metamorphosed. (*)

SUBORDER I. FILICES (Isospore Filicales, True Ferns).
Spores uniform, producing independent prothallia on which are developed

(*) The other fern-allies being also green plants with 2 generations and fibro-vascular bundles are readily distinguished from the **Filicales** in having the fronds commonly less strongly developed than the stems and by the following principal characters:

Order Equisetales. Fronds very small, in whorls, connected so as to form distinct, toothed, longitudinally ribbed, tubular, ochrea-like organs enclosing the bases of the internodations of the stems. Stems elongate, hollow. Sporangia placed on the under side of metamorphosed, scale-like, peltate, stalked sporophylla forming terminal spikes. Spores uniform. (1 Family).

Order Lycopodiales. Fronds various, seriate, circinate or in whorls, not connected. Stems tuberous or elongate, solid. Sporangia solitary at the base of the fronds on the upper side, superficial or imbedded. Sporophylla forming close or lax terminal spikes on an elongated stem, or placed with the barren fronds in a dense rosette surrounding a tuberous stem. Spores uniform or dimorphous. (4 Families).

antheridia and archegonia. Sporangia dorsal (on the under surface) or marginal on leafy fronds, rarely in spikes or panicles. (10 Families).

SUBORDER II. HYDROPTERIDES (Heterospore Filicales, Water-ferns). Spores dimorphous, macrospores and microspores, both developing more or less rudimentary prothallia, those of the macrospores bearing archegonia and those of the microspores antheridia. Sporangia enclosed in metamorphosed fronds forming 1—4-celled, globose or ellipsoidal sporocarpia. (2 Families; not treated with in this work).

FILICES.

Capsules (sporangia) 1-celled, sessile or stalked, usually membranaceous, dorsal or marginal, encircled by a complete or incomplete, \pm vertical or horizontal, jointed, elastic ring (annulus) and opening by a stomium at right angles to the ring, collected in clusters (sori) of a definite but varied form with or without a covering (indusium), or crowded densely over the parenchyma, sometimes scattered, spicate or paniculate; or the capsules have the ring obsolete, or none, or the ring is confined to the apex of the capsule, where it forms a longitudinally striated crown bursting vertically; or the capsules are sunk in many-celled, fleshy or corky receptacles of varied forms, opening by pores or clefts on the upper or inner side. Spores minute, various in form, mostly ellipsoidal, bilateral, tetrahedral or globose, and in intermediate forms.

CLASS I. PLANITHALLOSÆ. Prothallium flat or filiform, green, entire or variously incised or branched.

Series I. Annulatae. Sporangia with a complete or incomplete ring consisting of a close or open row or a group of thickened cells.

FAMILY I. CYATHEACEÆ. Ring complete, subvertical. Indusium wanting or present, but then neither umbrella-shaped nor linear. Caudex arborescent in the full-grown plants.

Tribe I. Cyatheae. Ring not depressed at the place of the stomium. Sori globose, dorsal on the veins.

GENUS 1. CYATHEA. Indusium globose, at least when young, at length cup-shaped, entire, lacinate or evanished.

GENUS 2. HEMITELIA. Indusium consisting of a *rather large*, convex, inferior scale attached at the inner side of the sorus.

GENUS 3. ALSOPHILA. Indusium wanting, or consisting of a *minute*, inferior scale or a whorl of hairs or fibres.

Tribe II. Dicksonieae. Ring depressed at the place of the stomium. Sori roundish or suboblong, terminal on the veins.

GENUS 4. DICKSONIA. Indusium cucullate, 2-valved, the outer valve consisting of a small lobe of the frond.

GENUS 5. CIBOTIUM. Indusium subcucullate, 2-valved, the outer valve differentiated from the margin of the frond.

FAMILY II. MATONIACEAE. Ring complete, subvertical. Sori round. Indusium umbrella-shaped. Caudex not arborescent.

GENUS 6. MATONIA. Characters of the family.

FAMILY III. GLEICHENIACEAE. Ring complete, horizontal, surrounding the sporangium like a belt. Sori roundish, dorsal on the veins. Indusium wanting.

GENUS 7. GLEICHENIA. Characters of the family.

FAMILY IV. HYMENOPHYLLACEAE. Ring complete, \pm horizontal, surrounding the compressed sporangium like a belt. Sori terminal on the veins. Indusium present.

GENUS 8. HYMENOPHYLLUM. Indusium more or less deeply 2-valved. Surface of the sporangium divided into many, uniform cells.

GENUS 9. TRICHOMANES. Indusium tubular or funnel-shaped, the mouth entire or 2-lipped. Surface of the sporangium divided into few, unequal cells.

FAMILY V. SCHIZAEACEAE. Ring complete, horizontal, crowning the apex of the sporangium like an operculum.

GENUS 10. LYGODIUM. Sori consisting of 1—2 sporangia placed in the axils of 2 rows of clasping, imbricating indusia. Fronds long and twining.

GENUS 11. SCHIZAEA. Sori linear, consisting of 2—4 rows of sporangia. Indusium wanting. Fronds small and erect.

FAMILY VI. POLYPODIACEAE. Ring incomplete, vertical, the extremities sometimes separated by the stalk of the sporangium only. Sori definite or indefinite, but then closely crowded over the parenchyma. (Ring sometimes as in Family I, but then the sori and indusia linear and the caudex not arborescent).

Indusiatae. Indusium present.

Tribe I. Woodsieae. Sori roundish or linear-oblong. Indusium globose, cupuliform or sausage-shaped, fastened all round the receptacle or apparently so in genus 13.

GENUS 12. DIACALPE. Indusium globose, bursting down irregularly. Veins free.

GENUS 13. DIPLAZIOPSIS. Indusium sausage-shaped, bursting down irregularly. Veins anastomosing.

GENUS 14. DENNSTAEDTIA. Indusium cup-shaped, open at the top.

Tribe II. Aspidieae. Sori roundish or oblong. Indusium superior, reniform, round or oblong, if reniform attached at the top of a more or less deep sinus, if round peltate, if oblong attached by its longitudinal axis, all opening all round the edge or nearly so.

GENUS 15. OLEANDRA. Indusium reniform. Fronds simple, articulate to the rhizome; veins free.

GENUS 16. ARTHROPTERIS. Indusium reniform. Fronds pinnate, articulate to the rhizome; pinnae articulate to the rachis; veins free.

GENUS 17. NEPHROLEPIS. Like genus 16, but the fronds not articulate to the rhizome.

GENUS 18. CYCLOPELTIS. Like genus 17, but the indusium round.

GENUS 19. POLYSTICHUM. Indusium round or reniform. Fronds divided, not articulate to the rhizome. Leaflets subrhomboidal, not articulate to the rachis; upper base broadly, lower narrowly cuneate; veins free.

GENUS 20. DIDYMOCHLAENA. Like genus 19, but the indusium oblong. Leaflets deciduous.

GENUS 21. PLEOCNEMIA. Indusium reniform, rarely round. Fronds divided, not articulate to the rhizome; veins anastomosing in costal and costular arches.

GENUS 22. DRYOPTERIS. Indusium reniform, rarely round. Fronds divided, not articulate to the rhizome; veins free, but then the leaflets not as in genus 19, or the lower veins anastomosing and forming costal rows of regular, \pm deltoid areolae.

GENUS 23. MESOCHLAENA. Like genus 22, but the veins not free. Indusium oblong.

GENUS 24. LUERSSENIA. Indusium oblong, $\pm 3 \times$ as long as broad. Fronds simple, not articulate to the rhizome; veins anastomosing copiously.

GENUS 25. ASPIDIUM. Indusium round or reniform, rarely nearly $2 \times$ as long as broad. Fronds simple or divided, not articulate to the rhizome; veins anastomosing copiously; included free veinlets wanting or present, but then not as in genus 26, if excurrent then solitary in the areolae.

GENUS 26. CYRTOMIUM. Indusium round. Fronds as in genus 25, but only simply pinnate; included free veinlets excurrent, 2—3 to each areola.

Tribe III. Davallieae. Sori roundish, oblong or linear, if roundish or longitudinally elongated then solitary and terminal on the veins (dorsal in genus 27), if transversally oblong or linear then uniting the apices of 2 or more veinlets. Indusium \pm of the same shape, attached at the inner side of the sorus, opening outwardly.

GENUS 27. CYSTOPTERIS. Sori roundish, dorsal on the veins.

GENUS 28. ODONTOSORIA. Sori roundish or transversally oblong, terminal on the cuneate, ultimate segments.

GENUS 29. LINDSAYA. Sori roundish, transversally oblong or linear, submarginal. Fronds divided; leaflets dimidiate, i.e. the half on the lower side of the midrib cut away.

GENUS 30. SCHIZOLOMA. Sori transversally oblong or linear, submarginal. Fronds simple or divided; leaflets not dimidiate.

GENUS 31a. SACCOLOMA. Sori roundish, submarginal, immersed in conspicuously cucullate lobes, covered

by a roundish indusium fixed by a broad base, or placed in submarginal depressions covered by a cuneate indusium. Indusium with the outer edge \pm entire. Fronds divided, not articulate to the rhizome.

GENUS 31*b*. LEPTOLEPIA. Sori roundish, submarginal or on marginal, recurved, lobe-like outgrowths. Indusium roundish or transversally oblong, fixed by the apex of the veinlet only, for the rest free, the outer edge lacerato-lobate. Fronds divided, not articulate to the rhizome.

GENUS 32. HUMATA. Sori roundish, submarginal, superficial. Indusium fixed by a broad base, the sides and apex free, \pm entire. Fronds simple or divided, scattered or subapproximate, articulate to the rhizome.

GENUS 33. ACROPHORUS. Like genus 32, but the fronds copiously divided, tufted, not articulate to the rhizome.

GENUS 34. DAVALLIA. Sori roundish or longitudinally elongated, submarginal. Indusium attached at the base and sides, the apex free. Fronds divided, scattered, articulate to the rhizome. Rhizome scaly, the scales not minute, not fibril-like.

GENUS 35. MICROLEPIA. Sori roundish, submarginal. Indusium as in genus 34. Fronds divided, rather scattered, not articulate to the rhizome. Rhizome hairy or fibrillose. Ultimate segments unequal-sided at the base. Receptacle distinct.

GENUS 36. TAPEINIDIUM. Like genus 35, but the rhizome fibrillose or minutely scaly and the fronds rather approximate. Ultimate segments rather equal-sided at the base. Receptacle indistinct.

Tribe IV. Pterideae. Sori roundish, oblong or linear, if roundish or longitudinally elongated then solitary and terminal on the veins, but not rarely laterally confluent into a submarginal line, if transversally oblong or linear then uniting the apices of several veinlets. Indusium \pm of the same shape, marginal, i.e. consisting of the reflexed, modified margin of the frond, opening inwardly.

GENUS 37. ADIANTUM. Differs from the following genera by the indusium bearing the capsules on its inner surface.

GENUS 38. HYPOLEPIS. Sori roundish, distant, confined to the tops (angles) of the ultimate sinuses, 1 to each sinus. Indusium roundish, small, hardly covering the sorus at full maturity.

GENUS 39. CHEILANTHES. Sori roundish or oblong, solitary and terminal on the veins, soon confluent into a submarginal line. Indusium roundish, transversally oblong or linear. Vein tips thickened. Stipes not as in genus 40. Fronds uniform.

GENUS 40. PLAGIOGYRIA Sori as in genus 39. Indusium linear. Vein tips not or hardly thickened. Stipes abruptly thickened at the base and generally purveyed there with \pm 2—12 spongy glands or tubercles on the outer side. Fronds dimorphous, the fertile segments contracted, narrowly linear.

GENUS 41. PELLAEA. Sori linear, or round to oblong, but then soon confluent into a submarginal line. Indusium linear. Fronds divided, the leaflets \pm stalked, articulate at the base.

GENUS 42a. DORYOPTERIS. Sori and indusia as in

genus 41. Fronds simple or incised, the segments adnate, confluent, not articulate at the base.

GENUS 42*b*. SCHIZOSTEGE. Sori and indusia terminal on the veins, marginal, solitary on T-shaped expansions or uniting the fork branches of the veins. Fronds divided; segments not articulate.

GENUS 43. HISTIOPTERIS. Sori and indusia oblong or linear. Fronds compoundly divided; lowest pinnulae short, close to the main rachis, stipule-like; leaflets not articulate at the base.

GENUS 44. ONYCHIUM. Sori and indusia as in genus 43. Fronds finely dissected; ultimate segments narrow, with the sori and indusia (which are occupying both sides of the segments) reaching nearly or quite to the midrib; leaflets not articulate at the base. Spores globose to tetrahedral.

GENUS 45*a*. PTERIS. Like genus 44, but the fronds not finely dissected, the ultimate segments broader with the sori and indusia not reaching the midrib. Stipes with a single V- or U-shaped vascular strand.

GENUS 45*b*. HEMIPTERIS. Like genus 45*a*, but the sori occupying only the posterior edge of the ultimate segments, the anterior side destitute of sori.

GENUS 46. PAESIA. Like genus 45*a*, but the indusia 2-valved and the spores 2-lateral.

GENUS 47. PTERIDIUM. Like genus 46, but the inner valve of the indusium sometimes obsolete, the spores as in genus 44. Stipes with many, irregularly placed vascular strands.

GENUS 48. LOMARIA. Sori and indusia linear, uniting the vein tips or running them transversely. Fronds simple or pinnate, dimorphous, the fertile fronds or segments narrowly linear. Stipes with 3 to many vascular strands placed in a circle.

Tribe V. Blechneae. Sori transversally oblong or linear, distant from the margin, parallel to the midribs of the fronds or their segments. Indusium of the same shape as the sorus, not marginal, attached at the outer side of the sorus, opening inwardly.

GENUS 49. BLECHNUM. Sori long, linear, sometimes interrupted, \pm close to the midribs. Veins of fertile fronds united near the base by subcostal strands bearing the sori.

GENUS 50. SADLERIA. Like genus 49, but the veins of both barren and fertile fronds united near the base.

GENUS 51. DOODIA. Sori oblong, superficial, in one or more rows on each side, not \pm close to the midribs.

GENUS 52. WOODWARDIA. Like genus 51, but the sori in 1 row, immersed in cavities.

Tribe VI. Asplenieae. Sori oblong or linear, attached to the sides of the veins which unite the midribs or main veins with the margin, therefore mostly oblique to the midribs, rarely placed between the veins. Indusia of the same shape, springing from the veins on the same side as the sori.

GENUS 53. DIPLAZIUM. Stipes with 2 \pm flattened vascular strands mostly uniting towards the apex of the stipe (or higher) into a single diarchous one ($\backslash /$, \vee , \surd , \sqcup , \cup). Sori linear or linear-oblong, the lower of each group of veins mostly double.

GENUS 54. ATHYRIUM. Like genus 53, but the sori oblong, the higher never, the lower rarely double, but not rarely hippocrepiform, hamato-recurvate or subreniform.

GENUS 55. ASPLENIUM. Stipes with 1, or 2 \pm concave vascular strands mostly uniting towards the apex of the stipe (or higher) into a single tetrarchous one [) (,)(,)(, ><, X, \underline{V}, \underline{V}]. Sori and indusia not in pairs; indusia opening towards the midribs, occasionally towards the margin.

GENUS 56. PHYLLITIS. Like genus 55, but the sori and indusia in opposite pairs, the indusia of each pair opening towards each other.

GENUS 57. TRIPHLEBIA. Like genus 56, but the sori medial between the veins, i. e. 1 sorus to each pair of indusia.

Exindusiatae. Indusium wanting, but the sori sometimes covered by the unaltered, recurved margin or placed in dorsal or \pm marginal cavities or indusium-like depressions.

Tribe VII. Phegopterideae. Sori punctiform, round or oblong, rarely more than 2 X as long as broad, medial or terminal on the veins and then sometimes running into a submarginal line, if medial on the veins then sometimes confluent in pairs. Stipes not articulate to the rhizome.

GENUS 58. NOTHOLAENA. Veins free. Sori round or oblong, terminal, not rarely running into a submarginal line. Fronds hairy or fibrillose. Spores globoso-tetrahedral.

GENUS 59. MONACHOSORUM. Veins free. Sori oblong,

terminal, always distant. Fronds naked. Spores 5-lobato-tetrahedral.

GENUS 60a. PHEGOPTERIS. Veins anastomosing as in *Nephrodium* (*Dryopteris* §), or free but then the spores 2-lateral. Fronds naked or hairy. Sori round or oblong, rarely linear-oblong, medial or terminal.

GENUS 60b. DICTYOPTERIS. Veins anastomosing more or less copiously, but not as in genus 60a. Fronds simple or divided, but not as in genus 61. Sori \pm round.

GENUS 61. DIPTERIS. Veins anastomosing copiously and irregularly. Fronds flabellato-dichotomous. Sori round or punctiform.

Tribe VIII. Gymnogrammeae. Sori linear or linear-oblong, more than 2 \times as long as broad, several on each side of the costae, mostly oblique, rarely subparallel to the latter and the margin, free or reticulate. Fronds not articulate to the rhizome.

GENUS 62. CONIOGRAMME. Veins free quite to the margin. Under surface of the fronds not powdery.

GENUS 63. CEROPTERIS. Veins free quite to the margin. Under surface of the fronds white- or yellow-powdery.

GENUS 64. ANTROPHYUM. Veins anastomosing copiously. Sori longitudinal, running along the veins, irregular. Costa wanting, or present in the lower half of the fronds only.

GENUS 65. LOXOGRAMME. Veins anastomosing copiously. Sori parallel, \pm oblique to the costa and margin. Costa distinct; veins invisible.

GENUS 66. SYNGRAMMA. Veins mostly uniting 1—3 \times at or near the margin, sometimes anastomosing copiously, but then the veins distinctly visible like the costa. Sori with paraphyses.

GENUS 67. HEMIONITIS. Veins anastomosing copiously, distinctly visible like the costa. Sori without paraphyses.

Tribe IX. Grammitideae. Sori linear, short or long, 1 on each side of the costa, parallel to the latter or the margin. Fronds not articulate to the rhizome.

GENUS 68. MONOGRAMMA. Sori costal, often protected by a lateral, raised outgrowth of leaf-tissue. Fronds simple, grass-like; lateral veins mostly wanting.

GENUS 69. VITTARIA. Sori deeply immersed in marginal or submarginal grooves. Fronds simple or forked, mostly grass-like; lateral veins present, free.

GENUS 70. TAENITIS. Sori not deeply immersed, medial or submarginal. Fronds pinnate or simple, but then not grass-like; lateral veins present, anastomosing.

Tribe X. Polypodieae. Sori round, oblong or linear. Fronds articulate to the rhizome.

GENUS 71. POLYPODIUM. Sori round, oblong or linear, several to numerous on each side, the oblong ones parallel or oblique to the costa, the linear ones oblique. Fronds uniform or dimorphous, simple or divided, the barren ones never oakleaf-like; surfaces naked, scaly or hairy, if hairy the hairs not stellately branched.

GENUS 72. CYCLOPHORUS. Sori round or oblong, several to numerous. Fronds as in genus 71,

but simple; at least the under surface always more or less hairy, the hairs stellately branched.

GENUS 73. DRYNARIA. Sori round, many or numerous. Fronds dimorphous. Barren fronds (cup-leaves) oakleaf-like, sessile, humus-collecting, fertile ones pinnatifid or pinnate, stipitate.

GENUS 74. DRYOSTACHYUM. Sori oblong or sub-quadrangular, many. Fronds pinnatifid, the higher segments fertile, much contracted. Barren fronds not oakleaf-like.

GENUS 75. DRYMOGLOSSUM. Sori linear, 1 on each side, parallel to the costa or the margin. Barren fronds roundish or oblong, fertile ones linear or linear-oblong.

GENUS 76. CHRISTOPTERIS. Sori linear, 1 on each side, parallel to the costa. Barren fronds sagittate, fertile ones 3-partite.

Tribe XI. Acrosticheæ. Sori indefinite, i.e. quite covering portions of the under surface or occasionally both surfaces of the fertile segments at full maturity, but apparently linear in the forms with much contracted fertile segments.

GENUS 77. PLATYCERIUM. Distinguished from the following genera by the dichotomously branched fertile fronds. Barren (basal) fronds adpressed to the substratum, humus-collecting.

GENUS 78. ELAPHOGLOSSUM. Veins free, the apices not rarely united by a \pm marginal, cartilagineous or decoloured line. Fronds simple.

GENUS 79. STENOCHLAENA. Veins free, those of the fertile fronds united at the base by a \pm

costal strand. Fronds pinnate, Lomaria-like. Rhizome high-scandent.

GENUS 80. POLYBOTRYA. Veins free, not united at the base in the fertile fronds. Fronds pinnately divided, not Lomaria-like. Rhizome erect or wide-creeping, not high-scandent. Sori often occupying the apices of the veins only when young.

GENUS 81. STENOSEMIA. Veins anastomosing, forming costal and costular arches, otherwise free. Barren fronds ternate or pinnate, fertile ones conspicuously contracted, branched.

GENUS 82. HYMENOLEPIS. Veins anastomosing copiously. Fronds simple, the soriferous apex narrowly linear or spoon-shaped.

GENUS 83. ACROSTICHUM. Veins anastomosing copiously. Fronds pinnate; pinnae not articulate to the rachis, the higher ones fertile, rarely much contracted.

GENUS 84. PHOTINOPTERIS. Veins anastomosing copiously. Fronds pinnatifid or pinnate; segments articulate to the rachis, the higher ones fertile, always much contracted.

GENUS 85. CHEIROPLEURIA. Veins anastomosing copiously. Fronds dimorphous, the barren ones simple or more commonly 2-fid, with the segments sometimes 2-fid again, each segment with 2 or more stout, longitudinal ribs. Fertile fronds simple, narrowly linear.

GENUS 86a. LEPTOCHILUS. Veins anastomosing copiously. Fronds dimorphous, simple, 3-fid or pinnate, each segments with a single, central rib. Spores oblong, ovate or bilateral.

GENUS 86*b*. PLATYTAENIA. Like genus 86*a*, but the fronds pinnate and the spores globoso-tetrahedral.

FAMILY VII. PARKERIACEAE. Ring incomplete, vertical. Sori indefinite, consisting of numerous sporangia scattered irregularly along the few, longitudinal veins, covered by the unaltered, recurved margin of the frond when young.

GENUS 87. CERATOPTERIS. Characters of the family.

FAMILY VIII. OSMUNDACEAE. Ring incomplete, short and horizontal or consisting of a lateral group of thickened cells only. Indusium wanting.

GENUS 88. OSMUNDA. Sori globose, placed in opposite, marginal clusters. Segments articulate to the rachis, the fertile ones more or less contracted.

GENUS 89. LEPTOPTERIS. Sori flat, dorsal. Segments not articulate to the rachis, the fertile ones not contracted.

Series II. Exannulatae. Sporangia without a ring.

FAMILY IX. MARATTIACEAE. Fronds mostly large, the stipes flanked by 2 stipulaeform appendages at the base. Indusium wanting or inconspicuous.

GENUS 90. ANGIOPTERIS. Sporangia in 2 opposite, close rows, not concrete. Fronds large, compound.

GENUS 91. MARATTIA. Sporangia concrete in 2-valved synangia, the valves opposite, boat-shaped. Fronds large, compound.

GENUS 92. CHRISTENSENIA. Sporangia concrete in

circular synangia. Fronds small, ternate or quinate.

CLASS II. TUBERITHALLOSAE. Prothallium tuberous, white.

FAMILY X. OPHIOGLOSSACEAE. Sporangia without a ring, sessile, placed on much contracted, spicate or paniculate fertile segments opposite to and springing from the base, the centre or the margin of the normal, frondose, barren segment. Indusium wanting.

GENUS 93. OPHIOGLOSSUM. Fertile segment spicate, with the sporangia in 2 opposite, lateral rows. Barren segment simple; veins anastomosing.

GENUS 94. HELMINTOSTACHYS. Fertile segment spicate, with the sporangia in numerous crested clusters spreading in all directions. Barren segment divided; veins free.

GENUS 95. BOTRYCHIUM. Fertile segment paniculate, with the sporangia placed in 2 rows on the inner face of the branches. Barren segment divided; veins free.

**Abbreviated natural arrangement
of the
FAMILIES, TRIBES AND GENERA
as construed in this work.**

I. CYATHEACEAE. (Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 115).

1. **Cyathea.** (Diels, l.c., 123).

- 1 *Cyathea*,
- 2 *Hemitelia*,
- 3 *Alsophila*.

- II. **Dicksonieae**. (Diels, l. c., 119).
 4 Dicksonia,
 5 Cibotium.
- II. **MATONIACEAE**. (Diels, l. c., 343).
 6 Matonia.
- III. **GLEICHENIACEAE**. (Diels, l. c., 350).
 7 Gleichenia.
- IV. **HYMENOPHYLLACEAE**. (Sadeb., in Engl. & Prantl, Nat. Pfl. Fam., 1⁴, 91).
 8 Hymenophyllum,
 9 Trichomanes.
- V. **SCHIZAEACEAE**. (Diels, l. c., 356).
 10 Lygodium,
 11 Schizaea.
- VI. **POLYPODIACEAE**. (Diels, l. c., 139).
 I. **Woodsieae**. (Diels, l. c., 159).
 I. **WOODSIINAE**.
 12 Diacalpe,
 27 Cystopteris,
 33 Acrophorus.
 II. **ONOCLEINAE**. — .
- II. **Aspidieae**. (Diels, l. c., 166).
 I. **ASPIDIINAE**.
 1. **INDUSIATAE**.
 22 *Dryopteris* (*Nephrodium*), *excl. Phegopteris*,
 24 *Luerssenia*,
 23 *Mesochlaena*,
 20 *Didymochlaena*,
 18 *Cyclopeltis*,
 25 *Aspidium*, *excl. Pleocnemia & Dictyopteris*.
 21 *Pleocnemia*,
 19 *Polystichum*, *excl. Cyrtomium*,
 26 *Cyrtomium*.
 2. **EXINDUSIATAE**.
 60a *Phegopteris*,

- 60*b* Dictyopteris,
- 80 Polybotrya,
- 81 Stenosemia,
- 86*a* Leptochilus (*Gymnopteris*).

II. DIPTERIDINAE.

- 61 Dipteris.

II*a*. Oleandreae. (Diels, l. c., 203).

- 15 Oleandra.

III. Davallieae. (Diels, l. c., 204).

1. INDUSIATAE.

- 16 Arthropteris,
- 17 Nephrolepis,
- 32 Humata,
- 31*a* Saccoloma,
- 31*b* Leptolepia,
- 34 Davallia, *excl. Prosaptia*,
- 35 Microlepia,
- 28 Odontosoria,
- 36 Tapeinidium (*Wibelia*),
- 14 Dennstaedtia,
- 30 Schizoloma,
- 29 Lindsaya.

2. EXINDUSIATAE.

- 59 Monachosorum.

IV. Pterideae. (Diels, l. c., 254).

I. GYMNOGRAMMINAE.

- 66 Syngramma,
- 62 Coniogramme,
- 67 Hemionitis,
- 63 Ceropteris (*Neurogramme*).

II. CHEILANTHINAE.

1. INDUSIATAE.

- 41 Pellaea,
- 42*a* Doryopteris,
- 39 Cheilanthes,
- 38 Hypolepis,
- 44 Onychium (*Cryptogramme* §),
- 40 Plagiogyria.

2. EXINDUSIATAE.

58 Notholaena (*Nothochlaena*).

III. ADIANTINAE.

37 Adiantum.

IV. PTERIDINAE.

45a Pteris, *excl. Schizostege*,

42b Schizostege,

45b Hemipteris,

43 Histiopteris,

47 Pteridium,

46 Paesia.

V. Blechnaeae. ——— .

VI. Asplenieae. (Diels, l. c., 222).

I ASPLENIINAE.

54 Athyrium,

53 Diplazium,

13 Diplaziopsis (*Allantodia*),

57 Triphlebia,

56 Phyllitis (*Scolopendrium*),

55 Asplenium.

II BLECHNINAE.

1. INDUSIATAE.

49 Blechnum, *excl. Lomaria*,

48 Lomaria,

50 Sadleria,

52 Woodwardia,

51 Doodia.

2. EXINDUSIATAE.

79 Stenochlaena.

VII. Phegopterideae. ——— .

VIII. Gymnogrammeae. ——— .

IX. Grammitideae (*Vittarieae*, Diels, l. c., 297).

I. VITTARIINAE.

68 Monogramma,

69 Vittaria.

II. ANTROPHIINAE.

64 Antrophyum.

X. Polypodieae. (Diels, l. c., 302).

I. TAENITIDINAE.

1. ARTICULATAE.

75 *Drymoglossum*,76 *Christopteris*,86*b* *Platytaenia*,82 *Hymenolepis*.

2. INARTICULATAE.

65 *Loxogramme*,70 *Taenitis*.

II. POLYPODIINAE.

71 *Polypodium*, *excl. Loxogramme, incl. Prosaptia & Lecanopteris*,72 *Cyclophorus (Niphobolus)*,74 *Dryostachyum*,84 *Photinopteris*,75 *Drynaria*.

XI. Acrosticheae. (Diels, l.c., 330).

I. ACROSTICHINAE.

78 *Elaphoglossum*,83 *Acrostichum*.

II. PLATYCERIINAE.

85 *Cheiropleuria*,77 *Platyterium*.

VII. PARKERIACEAE. (Diels, l. c. 339).

87 *Ceratopteris*.

VIII. OSMUNDACEAE. (Diels, l. c., 372).

89 *Leptopteris*,88 *Osmunda*.IX. MARATTIACEAE. (Bitt., in Engl. & Prantl., Nat. Pfl.Fam., I⁴, 422).90 *Angiopteris*,91 *Marattia*,92 *Christensenia (Kaulfussia)*.

X. OPHIOGLOSSACEAE. (Bitt., l. c. 449).

93 *Ophioglossum*; 95 *Botrychium*; 94 *Helmintostachys*.

Following pages contain besides descriptions keys to the determination of the families, tribes, genera and species.

If one supposes to have found in the keys the name sought after, he should not omit reading the diagnosis itself, seeing the plants are determined by more characteristics than those few admitted in the keys. These are mere expedients, no infallible guides; they chiefly had to be extracted from the available diagnoses since I did not set eyes on a great part of the plants. Furthermore it would have been like looking for a needle in a haystack to insert all deviations from the normal forms. Consequently the use of the keys may be sometimes the cause of going wrong.

In comparing the collected plants with the descriptions given in this and other works one should bear in mind that different authors are not always unanimous in using botanical terms. So, for instance, the word *ovate* is not rarely used for *oval* or even *deltoid* or another term, *spreading* for *horizontal*, *erecto-patent* for *more or less oblique*, *conspicuous* and *inconspicuous* for *distinct* and *indistinct*, *costa* for *rachis*, *costula* for *main vein*, *vein* for *veinlet*, *pinna* for *lobe* or *lacinia*, *medial* for *medial on the veins* or *medial between the costae* (*costulae*) and *edge*, *excurrent* for *directed towards the margin*, *forking* for *fork branch*, &c.

FILICES.

FILICES (Ferns).

Arrangement of the families.

a. CLASS I. **PLANITHALLOSÆ.** Prothallium green, flat or filiform, entire, incised or branched, epigeous or epiphytcal, addressed to the substratum. Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 11.

b. SERIES I. **ANNULATÆ.** Sporangia with a ring consisting of a close or broken row or a group of conspicuously prominent, thickened cells.

FAMILY I. **Cyatheaceæ.** Ring complete, oblique, subvertical, sometimes interrupted by a group of depressed, transversally oblong cells, which are however distinctly differentiated from the other cells of the sporangium. Sori globose or oblong, consisting of numerous, densely crowded sporangia; indusium wanting or present, but then neither umbrella-shaped nor linear. Arborescent ferns. Diels l. c., fig. 82, D; Hk. Bk., Syn. Fil., tab. I, fig. 4—6.

FAMILY II. **Matoniaceæ.** Ring as in Cyatheaceæ but never interrupted. Sori consisting of 4—8 sporangia placed in a whorl; indusium umbrella-shaped. No arborescent ferns. Diels, l. c., fig. 181 & 182, B; Hk. Bk., l. c., fig. 8.

FAMILY III. **Gleicheniaceæ.** Ring complete, \pm horizontal, surrounding the sporangium like a belt. Sori dorsal, consisting of 2—12 sporangia, mostly placed in a whorl; indusium wanting; receptacle dot-like. Diels, l. c., fig. 186; Hk. Bk., l. c., fig. 2.

FAMILY IV. **Hymenophyllaceæ.** Ring as in Gleicheniaceæ. Sori marginal, consisting of numerous sporangia crowded around an erected, oblong, cuneate, capitate, malleiform or columnar-filiform receptacle; indusium present. Diels, l. c., fig. 72 & 75, D; Hk. Bk., l. c., tab. II, fig. 16—17.

FAMILY V. **Schizaeaceae**. Ring operculiform, crowning the apex of the sporangium. Sporangia *either* 1—2, placed in the axils of imbricate, clasping, scale-like indusia *or* numerous, but then placed in 2—4 regular, naked rows. Diels, l. c., fig. 192; Hk. Bk., l. c., tab. VIII, fig. 64, tab. IX, fig. 68.

FAMILY VI. **Polypodiaceae**. Ring incomplete, vertical, the extremities sometimes separated by the stalk of the sporangium only. Sori definite or indefinite and then the sporangia densely crowded over the parenchyma; indusium wanting or present. (Ring sometimes as in Cyatheaceae, but then the indusium linear and the ferns not arborescent). Diels, l. c., fig. 57 & 87, D & H; Hk. Bk., l. c., tab. I, fig. 7, d.

FAMILY VII. **Parkeriaceae**. Ring as in Polypodiaceae. Sori indefinite, consisting of many sporangia scattered irregularly along the few, longitudinal veins, protected by the much recurved margin of the frond when young. Diels, l. c., fig. 178—179; Hk. Bk., l. c., tab. III, fig. 52.

FAMILY VIII. **Osmundaceae**. Ring incomplete, short and horizontal or consisting of a roundish, lateral group of thickened cells. Indusium wanting. Diels, l. c., fig. 205; Hk. Bk., l. c., tab. VIII, fig. 62, c, & 65, b & d.

b.b. SERIES II. **EXANNULATAE**. Sporangia without ring.

FAMILY IX. **Marattiaceae**. Sori consisting of 8—24 sporangia. Sporangia free, placed in 2 close, short rows or joined into 2-valved or circular synangia. Diels, l. c., fig. 259; Hk. Bk., l. c., tab. IX, fig. 69, 70 & 72.

a.a. CLASS II. **TUBERITHALLOSAE**. Prothallium white, tuberous, hypogeous or immersed in the substratum.

FAMILY X. **Ophioglossaceae**. Sporangia without ring, rather large, not gathered into sori but placed on much contracted, spicate or paniculate segments, originating from the base, the centre or the margin of the barren, normal-shaped segment of the frond. Diels, l. c., fig. 259, A—B & 260; Hk. Bk., l. c., fig. 73—75.

Key to the families.

- a.** Sporangia with a ring.
- b.** Ring complete.
- c.** Ring oblique, subvertical.
- d.** Ferns arborescent. Indusium wanting or present, but then neither linear nor umbrella-shaped.
- I. **Cyatheaceae.**
- d.d.** Ferns not arborescent. Indusium linear or umbrella-shaped.
- e.** Indusium umbrella-shaped. II. **Matoniaceae.**
- e.e.** Indusium linear.
- VI. **Polypodiaceae, No. 40, Plagiogyria.**
- c.c.** Ring \pm horizontal, surrounding the sporangium like a belt.
- i.** Receptacle inconspicuous, dot-like. Indusium wanting.
- III. **Gleicheniaceae.**
- i.i.** Receptacle conspicuous, erect, oblong to columnar-filiform. Indusium present. IV. **Hymenophyllaceae.**
- c.c.c.** Ring operculiform, crowning the apex of the sporangium.
- V. **Schizaeaceae.**
- b.b.** Ring incomplete.
- x.** Ring vertical, long, the extremities sometimes separated by the stalk of the sporangium only.
- !.** Sori indefinite. Sporangia scattered irregularly on the few, longitudinal veins. VII. **Parkeriaceae.**
- !?.** Sori definite or indefinite, but then the sporangia densely crowded over the parenchyma.
- VI. **Polypodiaceae.**
- x.x.** Ring horizontal, short or consisting of a lateral group of thickened cells only. VIII. **Osmundaceae.**
- a.a.** Sporangia without ring.
- *** Sori on the under surface of normal fronds.
- IX. **Marattiaceae.**
- *.*.** Sporangia not gathered into sori, placed on much contracted, spicate or paniculate segments originating from the base, the centre or the margin of the leafy, barren segment.
- X. **Ophioglossaceae.**
-

FAMILY I. CYATHEACEAE.



I. CYATHEACEAE (*).

Sori globose or ellipsoid, dorsal or terminal on the veins; sporangia numerous, sessile or stalked, furnished with a complete, oblique (subvertical) ring, which is sometimes interrupted at the place of the stomium by a row of depressed, transversally oblong cells; stomium opening at right angles to the ring; receptacle often elevated. Indusium wanting or more generally present and then scale-like, 2-valved, subcupuliform or globose, but then afterwards evanishing or becoming cup-shaped, sometimes consisting of a whorl of hairs or fibres, always fastened beneath the sorus.

Caudex arborescent, sometimes subarborescent. Fronds often very large, mostly compoundly divided; veins free or the costal ones sometimes anastomosing in *Hemitelia*.

Tropical and subtropical.

Arrangement of the tribes.

- § 1. **CYATHEAE.** Sori dorsal, placed on the back or at the forking of a vein. Annulus not interrupted by a row of depressed cells.
- § 2. **DICKSONIEAE.** Sori terminal on the veins. Annulus interrupted by a row of depressed cells at the place of the stomium.

Key to the tribes and genera.

- ‡. Sori on the back or at the forking of a vein. **CYATHEAE.**
- a.* Sori indusiate.
- b.* Indusium distinct, globose, quite enclosing the sorus when young, bursting down irregularly, afterwards evanishing or becoming ± cup-shaped. Receptacle central. **No. 1. Cyathea.**
- b.b.* Indusium distinct, squamiform, convex, placed at the inner side of the receptacle. **No. 2. Hemitelia.**

(*) The species of this family are difficult of determination unless the whole stipe and the lowest pinnae besides the central pinna and a portion of the main rachis are present.

b.b.b. Indusium spurious, consisting of a minute scale placed at the inner side of the receptacle or of a circular, entire or lobed scale or a whorl of fibres with the receptacle in the centre.

No. 3. **Alsophila.**

a.a. Sori exindusiate.

No. 5. **Alsophila.**

‡.‡. Sori terminal on the veins.

DICKSONIEAE.

*. Indusium 2-valved, the outer valve consisting of an unmodified, cucullate lobe of the frond, forming with the inner valve a saccate involucre to the sorus.

No. 4. **Dicksonia.**

,. Indusium 2-valved, the outer valve like the inner one, distinctly differentiated from the margin of the frond, both forming a saccate involucre to the sorus.

No. 5. **Cibotium.**

TRIBE I. CYATHEAE.

1. CYATHEA, *Smith.*

Sori globose, dorsal on a vein or veinlet or in the axil of a veinlet; receptacle elevated, globose or shortly columnar. Indusium inferior, membranaceous or somewhat horny, globose, at first quite enclosing the sorus, afterwards evanishing or breaking down and becoming cupshaped, with an entire or more or less irregularly lobed or lacinated margin.

Caudex arborescent, often beautifully marked with the scars of fallen fronds. Fronds mostly large and divided, rarely simple (not in malayan material); veins free, simple or forked. Hk. Bk., Syn. Fil., tab. I, fig. 4.

Tropical and extra-tropical in the southern hemisphere.

-
- a.** *Fronds simply pinnate.* (1) *C. moluccana.*
- a.a.** *Fronds 2-pinnatifid to 2-pinnate.*
- b.** *Main rachis densely scaly.* (2) *C. philippinensis.*
- b.b.** *Main rachis naked or nearly so.*
- c.** *Pinnae pinnatifid throughout.* (5) *C. aneitensis.*
- c.c.** *Pinnae pinnate, at least at the base.*
- d.** *Pinnae pinnate near the base.*
- e.** *Pinnae $\pm 2\frac{1}{2}$ c.M. broad. Secondary segments ± 5 m.M. broad.* (4) *C. dulitensis.*
- e.e.** *Pinnae 3—8 c.M. broad. Segments ± 10 m.M. broad.* (5) *C. alternans.*
- d.d.** *Pinnae pinnate nearly throughout.* (6) *C. sarawakensis.*
- a.a.a.** *Fronds 3-pinnatifid to 3-pinnate.*
- 1.** *Tertiary segments very small, roundish, not more than $\pm 2\frac{1}{2}$ m.M. long.* (7) *C. Macgregorii.*
- 1.1.** *Tertiary segments oblong to linear, much more than $2\frac{1}{2}$ m.M. long.*
- 2.** *Pinnulae pinnatifid, sometimes pinnate at the base.*
- 3.** *Main rachis not prickly, naked.*
- 4.** *Tertiary segments entire or slightly crenate.*
- ***. *Veins 7—8 on each side of the costulae, deeply forked.* (8) *C. polypoda.*
- *.*.** *Veins ± 12 on each side of the costulae, 2—3 \times forked.* (9) *C. leucotricha.*

4.4. *Tertiary segments ± toothed.*

5. *Veins 5—6 on each side of the costulae.*

(10) *C. suluensis.*

5.5. *Veins 9—12 on each side.*

6. *Texture papyraceous. Sori in ± 4—5 pairs to a lobe.*

(11) *C. saccata.*

6.6. *Texture coriaceous. Sori in ± 8 pairs to a lobe.*

(12) *C. cyclodonta.*

3.3. *Main rachis not prickly, but scaly, hairy or tomentose, at least on the upper side.*

α. *Texture membranaceous to papyraceous.*

β. *Main rachis hairy or tomentose.*

γ. *Ultimate segments obscurely crenate, slightly scaly beneath.*

(13) *C. fusca.*

γ.γ. *Ultimate segments serrate, naked beneath.*

(14) *C. integrā.*

β.β. *Main rachis squamose.*

(15) *C. assimilis.*

α.α. *Texture chartaceous to coriaceous.*

β'. *Main rachis purveyed with large glands on the upper side near the base of the pinnae.*

(16) *C. Zollingeriana.*

β'.β'. *Main rachis without glands.*

γ'. *Main rachis hairy or tomentose, not scaly.*

δ'. *Sori costular.*

ε'. *Lobes entire or crenate, indusium rupturing irregularly.*

***. *Secondary rachises hirsute above, slightly scaly beneath.*

(17) *C. Korthalsii.*

**.*.* *Secondary rachises tomentose above.*

(18) *C. crenulata.*

ε'.ε'. *Lobes serrate. Indusium at length regularly cup-shaped.*

(19) *C. orientalis.*

δ'.δ'. *Sori medial on the veins.*

(20) *C. runensis.*

γ'.γ'. *Main rachis scaly, or scaly and hairy.*

§. *Pinnae under 20 c.M. long; veins ± 4—5 on each side of the costulae.*

(21) *C. Havilandii.*

§.§. *Pinnae more than 20 c.M. long; veins ± 7—10 on each side.*

*. *Lower sori divaricating. Indusium at length regularly cup-shaped.*

(22) *C. javanica.*

*. *. *Lower sori not divaricating. Indusium deciduous or breaking down irregularly.*

†. *Sori costular. (25) C. sumatrana.*

†. †. *Sori rather medial.*

(24) *C. inquinans.*

3.3.3. *Main rachis ± prickly.*

4. *Upper surface whitish when dry. (25) C. leucophaës.*

4.4. *Upper surface not whitish when dry.*

5. *Sori distant from the costulae. (26) C. strigosa.*

5.5. *Sori costular.*

6. *Rachises tomentose above. Pinnulae 6—7½ c.M. long. Sori few to a lobe. (27) C. hymenodes.*

6.6. *Rachises not tomentose. Pinnulae 8—12 c.M. long. Sori many to a lobe. (28) C. spinulosa.*

2.2. *Pinnulae pinnate ± throughout.*

§. *Main rachis not prickly, naked. (29) C. Muelleri.*

§.§. *Main rachis not prickly, but scaly, hairy or tomentose.*

*. *Edges crenate; each crenation corresponding with a bullate vault, prominent on the upper surface. (30) C. oinops.*

*. *. *Edges entire or crenulate; prominent, bullate vaults wanting. (31) C. schizochlamys.*

§.§.§. *Main rachis prickly and tomentose or ± furfuraceous.*

†. *Tertiary segments adnate to the rachis. Under surface cobwebby. (32) C. arachnoidea.*

†. †. *Lower tertiary segments slightly stalked. Under surface naked or inconspicuously scaly. (33) C. celebica.*

* (1) **C. moluccana**, *R. Br.*, *Desv.*, *Prod.*, 522; *C. brunonis*, *Wall.*, *Hk. Bk.*, *Syn. Fil.*, 16, *Schizocaena brunonis*, *J. Sm.*, *Bau. & Hk.*, *Gen. Fil.*, tab. II.

Stipes 50—100 c.M. long, the base slightly rough and furnished with brown scales. Fronds large, ± 100 c.M. long, simply pinnate. Pinnae nu-

merous, distant, lanceolate, \pm 15—50 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, the edge repando-crenate, or slightly and irregularly undulate, the base subcuneate or rounded, the apex acuminate, crenate-dentate. Texture thinly coriaceous; surfaces naked or the upper one hairy on the costa near the base like the rachis; costae and veins distinct; veins parallel, several times branched. Sori in \pm 2, often very irregular rows on each side of the costa; indusium thin, breaking down irregularly.

Malaya.

(2) **C. philippinensis**, *Bk.*, Ann. of Bot., V, 186.

Trunk short, slender; the scales large, linear, pale-brown. Fronds bipinnate, oblong-rhomboid, \pm 90 c.M. long, \pm 30 c.M. broad, narrowed from the middle to the base; rachises of frond and pinnae densely scaly. Pinnae lanceolate, central largest, 15—18 c.M. long, \pm $3\frac{1}{2}$ —4 c.M. broad. Pinnules ligulate, crowded, sessile, 6—7 m.M. broad, entire or crenate, with forked veins and a single row of sori between the midrib and the edge; indusium moderately firm, but breaking up into pieces.

Philippines.

(3) **C. aneitensis**, *Hk.*, Hk. Bk., Syn. Fil., 26.

Pinnae oblong, 30—45 c.M. long, 10—15 c.M. broad, sessile or nearly so, short-acuminate, pinnatifid nearly to the costa. Lobes rather distant, oblong, acute, subserrate. Texture coriaceo-membranaceous; rachis smooth, purple or ebeneous; costae and costulae scaly with hair-like scales; veins forked. Sori copious, intermediate between the costulae and the margin; indusium thin-membranaceous, persistent, soon breaking down and generally opening at the back so as to form a large, lax, almost transparent, 2-lobed and ragged cup, often again irregularly lacerated.

Ternate; Aneitum.

(4) **C. dulitensis**, *Bk.*, Kew Bull., 1896, 40.

Caudex short. Stipes densely tufted, \pm 15 c.M. long, brown, naked above, the base covered with large, spreading, linear-subulate, membranaceous scales. Fronds oblong-lanceolate, \pm 50 c.M. long, 12—15 c.M. broad, 2-pinnate. Pinnae lanceolate, \pm $2\frac{1}{2}$ c.M. broad, pinnate at the base, the lower ones short-stalked, the lowest reduced, deflexed. Segments linear-oblong, \pm 5 m.M. broad, nearly entire, the margin revolute. Texture thick, rigid, coriaceous; surfaces naked, rachis nearly so, brown; veins obscure, immersed,

erecto-patent, forked. Sori in 1 row on each side of the costa, medial; indusium campanulate, fragile, breaking down irregularly, persistent.

Borneo.

* (5) **C. alternans**, *Pr.* (not *Hk.*), *Christ*, *Farnkr. d. Erde*, 327; *Alsophila alternans*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 41; *Bedd.*, *Ferns Br.I.*, II, tab. CCXXXVI; *Amphicosmia alternans*, *Moore*, *Bedd.*, *Handb. Ind. Ferns*, 10; *Hemitelia alternans*, *Hk.*, *Icon. Pl.*, VII, tab. DCXXII; — var. *Lobbiana*; *C. Lobbiana*, *Hk.*, *Hk. Bk.*, l. c., 24.

Fronds large, sub-2-pinnate. Pinnae short-stalked, remote, 20—35 c.M. long, 4—8 c.M. broad, linear-oblong, acuminate, deeply pinnatifid, but pinnate at the base. Segments linear-oblong, acute, subfalcate, spreading, remote or slightly so, \pm alternate, \pm 1 c.M. broad, the edge entire or serrate, the lowest ones sessile with a broad base. The extremity of the frond suddenly pinnate with oblong, undivided, sessile pinnulae, serrate at the margin. Texture subcoriaceous; rachis and surfaces naked or the rachis tomentose above; veins approximate, 1—2 \times forked. Sori 2-seriate on each pinnula or lobe, close, mixed with rather copious, jointed hairs; indusium at length a hemispherical cup, completely surrounding the base of the sorus.

Var. Lobbiana: Pinnae strongly acuminate nearly to the costa; segments obtuse. Indusium very thin, breaking down irregularly.

Penang; — var. *Lobbiana*: *Borneo*.

(6) **C. sarawakensis**, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 23; (? *C. alternans*, *Pr.*, *C. Chr.*, *Ind. Fil.*).

Fronds large, 2-pinnate. Pinnae distant, stalked, to 50 c.M. long or longer, \pm 15 c.M. broad, oblong, pinnatifid only at the acuminate apex, pinnate towards the base. Pinnulae numerous, approximate, horizontal, sessile, 10—15 m.M. broad, linear, acuminate, crenato-serrate, obliquely cuneate at the base. Texture subcoriaceous; surfaces and rachises naked; veins approximate, 2—3-furcate. Sori dorsal, rarely in the axils of the veins; indusium thin, fragile, breaking down into irregular lobes.

Borneo.

(7) **C. Macgregorii**, *Muell.*, *Journ. of Bot.*, XXVIII, 104.

Caudex short, under 60 c.M. long. Fronds 3-pinnatifid; rachises more or less clothed with an intertangled coat of brown scales. Pinnae \pm 10 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad. Pinnulae lanceolate, 4—5 m.M. broad. Segments

round or oblong, bullate, rigid, with the edges much recurved. Texture rigid; surfaces copiously scaly when young. Sori 1—2 to each ultimate segment, quite covering its under surface; indusium globose, fragile.

New Guinea.

(8) **C. polypoda**, *Bk.*, Trans. Linn. Soc., Bot., 2d Series, IV, 250.

Fronds large, 5-pinnatifid; main rachis naked. Pinnae lanceolate-oblong, \pm 45 c.M. long, 10—15 c.M. broad. Pinnulae lanceolate, the lower stalked, \pm 15 m.M. broad, pinnatifid nearly to the costa. Lobes oblong, obtuse, entire, $3\frac{1}{2}$ —5 m.M. broad. Texture moderately firm; under surface covered with numerous white, ovate scales, especially on the costae; veins 7—8 on each side, deeply forked. Sori medial between the costulae and the margin; indusium cupuliform, small, naked.

Borneo.

* (9) **C. leucotricha**, *Christ*, Ann. Btz., XX, 135.

Fronds large, tripinnatifid, subtripinnate; main rachis naked, glossy, reddish-brown. Pinnae stalked, approximate, to 50 c.M. long, \pm 18 c.M. broad, acuminate, \pm 30 on each side below the pinnatifid apex. Pinnulae \pm sessile, \pm 9 c.M. long, \pm 18 m.M. broad, acuminate, the base slightly narrowed, the edge cut down nearly to the costa; lowest pinnulae not the largest. Ultimate segments pectinate, bluntish or subacute, falcate, \pm 25 on each side below the pinnatifid apex, \pm $2\frac{1}{2}$ m.M. broad, linear-oblong, entire or slightly crenate towards the apex, the margin hardly reflexed. Texture coriaceous; under surface not glaucous; costae, costulae and secondary rachises covered beneath with pale, \pm adpressed, firm hairs like the costae above; veins \pm 12 on each side, oblique, 2—3 \times forked. Sori medial on the lower veins, 4—5 on each side of the costulae; indusium thin, hyaline, globose.

Borneo.

(10) **C. suluensis**, *Bk.*, Journ. of Bot., XVII, 65.

Fronds large, 5-pinnatifid or subtripinnate; main rachis pale-brown, naked. Pinnae sessile, oblong-lanceolate, 50 c.M. or more long, 12—15 c.M. broad. Pinnulae lanceolate, sessile, 1—2 c.M. broad, cut down nearly (or occasionally at the base quite) to the costa into ligulate-oblong, toothed segments \pm 4—5 m.M. broad. Texture moderately firm; rachises naked, pale-brown; under surface covered with many small, white, bullate scales on the midribs of the pinnulae and ultimate segments; veins 5—6-jugate, erecto-patent, dis-

tinct, deeply forked. Sori rather small, placed at the forkings of the lower veins; indusium persistent, semiglobose, entire or slightly broken up as it matures.

Sulu Islands.

(11) **C. saccata**, *Christ.*, Ann. Btz., XIX, 42.

Dwarf tree. Stipes glossy, sulcate below, convex above, yellow-brown, covered with minute warts and prickles, ± 25 c.M. long. Fronds ± 115 c.M. long, ± 80 c.M. broad, ovate, acuminate, narrowed towards the base, 3-pinnatifid. Pinnae distant, stalked, ± 12 c.M. broad, ovate, acuminate, caudate, narrowed at the base. Pinnulae distant, $2-2\frac{1}{2}$ c.M. apart, $\pm 1\frac{1}{2}$ c.M. broad, the lower ones shorter, stalked, lanceolate, the apex serrate, the edge pinnatifid. Segments 13—14 on each side, ligulate, $\pm 2\frac{1}{2}$ m.M. broad, obtuse, toothed. Texture papyraceous; upper surface dark-green, under one paler; rachis reddish, those of the pinnae brown or ebeneous; upper surface furfuraceous on the costae, the scales brown; veins distinct, 9—12 on each side, forked. Sori 4—5 on each side, medial; indusium membranaceous, afterwards saccate with the margin entire.

Celebes.

* (12) **C. cyclodonta**, *v. A. v. R.*; *Alsophila cyclodonta*, *Christ.*, Ann. Btz., XX, 137.

Main rachis naked, glossy, brownish-green. Pinnae ± 60 c.M. long, 20—25 c.M. broad, acuminate, the base not or hardly narrowed, on stalks 1—6 c.M. long, the rachis hairy above. Pinnulae subsessile, spreading, close, $2-2\frac{1}{2}$ c.M. broad, numerous, the base not narrowed, the apex acuminate, subcaudate, toothed, the costa covered above with adpressed, pale-coloured hairs, clothed beneath with longer, spreading, subulate setae. Tertiary segments linear, falcate, obtuse, $\pm 2\frac{1}{2}$ m.M. broad, the higher ones confluent, the others sessile, 2—3 m.M. apart, united by a narrow wing, decurrent with a rounded, adnate lobe, the lowest subsessile, with a truncate base, the margin dentate-crenate, the crenations semiorbicular, ± 12 on each side. Texture coriaceous; under surface hairy on the costulae; veins fine, ± 12 on each side. Sori ± 8 on each side, contiguous, \pm medial, not occupying the higher veins, 1 to each crenation; indusium at length a shallow, deeply and irregularly lobed, membranaceous, persistent, brown cup.

Borneo.

(15) **C. fusca**, *Bk.*, Becc., Mal., III, 31.

Fronds large, 3-pinnatifid; main rachis brownish-pubescent. Pinnae oblong-

lanceolate. Pinnulae sessile, lanceolate. Ultimate segments oblong, obscurely crenate. Texture moderately firm; under surface covered with small scales on the veins; veins distinct, erecto-patent, 5—6 on each side, mostly forked. Sori medial on the veins; indusium firm, persistent, naked, the opening broad, entire or cut down to the receptacle.

New Guinea.

(14) **C. integra**, *J. Sm.*, Hk. Bk., Syn. Fil., 23; Hk., Icon., Pl., VII, tab. DCXXXVIII; — var. *petiolata*; *C. petiolata*, *J. Sm.*, Hk., l. c., fig. 2.

Fronds 3-pinnatifid; main rachis \pm short-hairy or tomentose above (conf. Hooker's drawing). Pinnae large, \pm 45—50 c.M. long. Pinnulae sessile, \pm 15 on each side below the pinnatifid apex, \pm 7 $\frac{1}{2}$ —12 $\frac{1}{2}$ c.M. long, the apex acuminate, toothed, the base \pm truncate, \pm 1 $\frac{1}{2}$ c.M. broad, the edge cut $\frac{1}{2}$ — $\frac{2}{3}$ -way down to the costa into broad, oblong, subfalcate, serrate lobes. Texture firm-membranaceous; upper surface brownish-green, under one paler, both surfaces naked. Sori in 1 row on each side, medial between the costulae and the margin, occupying the lower veins; indusium pale, membranaceous, breaking down into a persistent, lobed cup.

Var. petiolata: Pinnulae often stalked. ✓

Amboyna, Philippines.

(15) **C. assimilis**, *Hk.*, Hk. Bk., Syn. Fil., 24; *C. Beccariana*, *Ces.*, Fel. di Bor., 3.

Stipes bright-chestnut-brown, slightly muricate and fringed on one side with long, ferruginous scales. Fronds large, 3-pinnatifid; superior rachis \pm squamose like the underside of the costae. Pinnae nearly 50 c.M. long, 7 $\frac{1}{2}$ —10 c.M. broad, acuminate, petiolate. Pinnulae and superior pinnae deeply pinnatifid almost to the costa. Lobes oblong, obtuse, subfalcate, coarsely toothed, the lowest inferior lobe always the smallest. Texture thinly herbaceous to rigidly membranaceous; veins simple or 1—2 \times forked. Sori dorsal on the veins or in the axils of the forks; indusium very thin and filmy, breaking down into irregular lobes.

Borneo, Celebes.

(16) **C. Zollingeriana**, *Mett.*, Ann. Mus. Bot. L. B., I, 57.

Stipe and rachis blackish-purpureous; stipes rough, nodose below, furfuraeous; rachis rough in the lower part, slightly scaly, densely hirsute above, purveyed with prominent, tuberculiform glands at the base of the pinnae. Fronds

3-pinnatifid. Pinnae lanceolate, short-stalked. Pinnulae sessile, \pm 7 c.M. long, \pm 17 m.M. broad, linear-oblong, the apex narrowed, obscurely serrate. Lobes close, linear-oblong, slightly falcate, entire, serrate at the obtuse apex. Texture coriaceous; upper surface hirsute or furfuraceous on the costae, under one scaly on the costulae; veins fine, forked. Sori close to the costae, 1 on each side of the costulae; indusium firm, membranaceous, cupuliform, the mouth constricted, \pm repand, afterwards cut down irregularly.

Java.

(17) **C. Korthalsii**, *Mett.*, Hk. Bk., Syn. Fil., 25; *Mett.*, Ann. Mus. Bot. L. B., I, 57.

Fronds 3-pinnatifid, main rachis hardly nodulose, short-ferrugineo-tomentose above. Pinnae 30—45 c.M. long, subsessile, elongate-oblong. Pinnulae 5—8 c.M. long, 1—2 c.M. broad, sessile, oblong, acuminate, the point bluntish, the edge deeply pinnatifid. Lobes oblong, approximate, subfalcate, subentire, obtuse, narrow. Texture firm, chartaceous; colour deep-green above, paler beneath; rachises slightly scaly beneath, \pm hirsute above; upper surface ferrugineo-hirsute on the costae, under one sparingly bullato-paleaceous; veins visible, slender, oblique, the lower forked. Sori more or less copious, close to the costulae, chiefly in the lower part of the lobes; indusium firm-membranaceous, falling away in fragments.

Sumatra, Java, Celebes.

* (18) **C. crenulata**, *Bl.*, Rac., Flor. Btz., I, 36; *C. excelsa*, *Kze* (not *Sw.*), Bot. Zeit., 1848, 284.

Fronds 3-pinnatifid; main rachis ferrugineo-tomentose above like the secondary ones and the costae. Pinnae subsessile, to 50 c.M. long, to 20 c.M. broad, oblong, acuminate. Pinnulae oblong-lanceolate, short-acuminate, 5—10 c.M. long, to 2 c.M. broad. Lobes approximate, oblong, subacute, subfalcate, \pm 3 m.M. broad, slightly crenate, subdistinct. Texture subcoriaceous; colour dark-green, paler beneath; veins to \pm 10 on each side of the costulae, forked. Sori close to the costulae, occupying the lower 1—3 veins; indusium membranaceous, firm, soon breaking into very irregular lobes. Under surface sparingly scaly on the costulae.

Java.

* (19) **C. orientalis**, *Moore*, Rac., Flor. Btz., I, 37; *C. arborea*, *Sm.*, var. *pallida*, Hassk., Obs. Fil. Jav., I, 15.

Fronds large, ovate-lanceolate, to 2½ M. long, to 1½ M. broad, 3-pinnatifid;

main rachis yellowish, punctato-muricate, tomentose above. Pinnae to 60 c.M. long, on stalks 2—4 c.M. long. Pinnulae $7\frac{1}{2}$ —10 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, sessile, oblong-lanceolate, caudato-acuminate, the point serrate, the edge deeply pinnatifid nearly to the costa. Lobes linear-oblong, approximate, subfalcate, the margin serrate and rather recurved. Texture coriaceous; colour black-green, paler beneath; rachises and costae \pm tomentose above; costae and costulae furfuraceous beneath, the hairs and scales brownish; veins 8—12 on each side of the costulae, mostly forked. Sori 4—9 on each side, close to the costulae which they conceal; indusium firm, persistent, becoming subglobose, the mouth a little contracted, entire.

Java, Celebes.

* (20) **C. runensis**, *v. A. v. B.*

Fronds large, 3-pinnatifid; main rachis setose above like the secondary ones and the costae; the setae subulate, \pm adpressed, reddish-brown. Pinnae \pm 55 c.M. long, 22—25 c.M. broad. Pinnulae sessile, spreading, 2— $2\frac{1}{2}$ c.M. broad at the base, the apex acuminate, toothed or crenate, the edge cut \pm $\frac{4}{5}$ -way down to the costa into numerous (\pm 15), oblong, bluntish, subfalcate lobes with a narrow space between them, the margin crenate, recurved, the lowest anterior lobes spreading, 5 m.M. broad, the lowest posterior \pm erecto-patent; lowest pinnulae not the longest. Texture coriaceous; under surface paler than the upper one; veins distinct, \pm 8 on each side of the costulae, the lower 1—3 forked. Sori medial on the veins, occupying nearly all the veins, except the highest ones, filling up the whole space between the costulae and the margin; indusium thin, hyaline, breaking down irregularly, the lower part persistent.

Pulu Run, (Teysmann).

(21) **C. Havilandii**, *Bl.*, Trans. Linn. Soc., Bot., 2d Series, IV, 249.

Stipes strong; the scales copious, lanceolate, brown, crisped. Fronds oblong-lanceolate, 3-pinnatifid, \pm 60 c.M. long, 20—50 c.M. broad; main rachis paleaceous. Pinnae sessile, oblong-lanceolate, \pm 5 c.M. broad, the lowest not reduced. Pinnulae numerous, close, lanceolate, sessile, the lower ones deeply pinnatifid. Lobes linear-oblong, erecto-patent. Texture subcoriaceous; under surface densely scaly on the costae; veins simple, 4—5 on each side of the costulae, erecto-patent. Sori in 1 row on each side of the costae; indusium firm, cupuliform, naked, persistent.

Borneo.

* (22) **C. javanica**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 55.

Fronds large, 5-pinnatifid; main rachis unarmed but not rarely rough, paleaceo-tomentose like the secondary ones. Pinnae to 60 c.M. long, on stalks to 2 c.M. long, linear-lanceolate, acuminate. Pinnulae lanceolate, to $1\frac{1}{2}$ c.M. broad, deeply pinnatifid. Lobes linear, subfalcate, entire or obtuso-crenulate. Texture subcoriaceous; under surface bullato-squamulose on the costae; veins 8—10 on each side of the costulae, the lower forked. Sori 2—6 on each side of the costulae, the lower divaricating; indusium thin, breaking down irregularly, the base persistent, cupuliform, firmer.

Java, Sumatra.

* (23) **C. sumatrana**, *Bk.*, *Journ. of Bot.*, XVIII, 209.

Fronds large, 5-pinnatifid to subtripinnate; rachises tomentose, especially on the upper side and clothed with long, linear, acuminate, scattered, brown scales, which make the rachises rough when they fall. Pinnae \pm 50 c.M. long, with the opposite pinnulae of contiguous pinnae overlapping. Pinnulae numerous, spreading, \pm 6—8 c.M. long, \pm $1\frac{1}{2}$ —2 c.M. broad, oblong, suddenly acuminate at the narrow, mostly linear, broadly crenate apex, the point blunt, the edges parallel, cut down nearly to the costa, pinnate at the very base. Segments linear-oblong, \pm 3 m.M. broad, falcate, blunt, entire or crenate, not conspicuously recurved at the edge, the lowest 1—2 free, mostly bluntly toothed; lowest pinnulae not the largest. Texture coriaceous; costae rough-hairy above; upper surface dark, the under one paler, covered with pale, bullate scales on the costae and the lower part of the costules; veins 8—10 on each side, forked, the higher simple. Sori close to the costules, occupying the lower 1—4 veins; indusium membranaceous, thin, breaking down irregularly.

Sumatra, Java.

(24) **C. inquinans**, *Christ*, *Ann. Btz.*, XV, 85, tab. XIII, fig. 5, a—g.

Fronds 5-pinnatifid. Stipes and rachises more or less densely scaly and hairy, the scales to $1\frac{1}{2}$ c.M. long, pale-ferruginous, the hairs crisped, ferruginous, joined into groups. Pinnae 27—50 c.M. long, imbricate, sessile, oblong, bluntish, slightly narrowed at the base. Pinnulae sessile, close, to 6 c.M. long, to $1\frac{1}{2}$ c.M. broad, bluntish, cut down nearly to the costa into short-acuminate, subfalcate, crenulate to toothed lobes. Texture coriaceous, under surface hairy between the sori; colour dark-green, paler beneath; veins inconspicuous, forked at the base. Sori large, 6—8 to each lobe; indusium large, flaccid, yellowish, soon deciduous.

Celebes.

* (25) **C. leucophaës**, *Hassk.*, Journ. of Bot., VII, 323.

Stipe and rachis muricate. Pinnae linear-lanceolate, \pm 10—12 c.M. broad at the base, short-stalked, the apex acuminate. Pinnulae linear-lanceolate, acuminate, the lowest longest, \pm 1—1½ c.M. broad at the base, the edge deeply pinnatifid. Lobes linear-oblong, acute, falcate, \pm 1½—2 m.M. broad at the base, the lowest \pm free, the edge recurved, crenato-serrate towards the point. Texture coriaceous; surfaces naked, the upper one nearly white when dry; veins forked. Sori globose; indusium membranaceous, lucid, soon breaking down irregularly, at length evanished.

Java.

(26) **C. strigosa**, *Christ*, Ann. Btz., XV, 84, tab. XIII, fig. 6, a—c.

Stipes \pm 45 c.M. long, angular, orange-brown, covered with small, furfureous scales and strong, cylindric, glossy, blackish-red prickles. Fronds \pm 65 c.M. long, \pm 40 c.M. broad, deltoid; main rachis prickly. Pinnae 10—12, \pm 20 c.M. long, \pm 8 c.M. broad, ovate, short-acuminate, sessile, the apex pinnatifid, the lowest not shorter. Pinnulae 15—18 on each side, sessile, linear-lanceolate, acute, pinnatifid nearly to the costa. Lobes falcate, deltoid, the edge entire, revolute. Texture coriaceous, firm but flexible; upper surface naked or sparingly hairy, under surface opaque-glaucous; veins covered with subulate hairs and white, ovate-bullate scales. Sori small, nearer the edge than the costulae; indusium afterwards flat and orbicular.

Celebes.

(27) **C. hymenodes**, *Mett.*, Hk. Bk., Syn. Fil., 24.

Stipes \pm 17 c.M. long, dark-brown, densely short-aculeolate, with few, black-brown, lanceolate scales. Fronds large, 5-pinnatifid; rachis tomentose above, the base subasperous or muricate. Central pinnae elongate-lanceolate to \pm 60 c.M. long, 12—15 c.M. broad; rachis densely tomentose above. Pinnulae 2—2½ c.M. broad, elongate-oblong, acuminate, slightly narrowed towards the base, sessile, the apex crenate, the edge deeply pinnatifid nearly to the costa. Lobes compact, 4—5 m.M. broad, oblong, falcate, obtuse, entire or rarely crenate. Texture subcoriaceous; costa hirsute above, sparingly covered beneath with minute, pale, bullate scales; veins \pm 9 on each side, oblique, forked. Sori few, close to the costulae, occupying the lower 1—2 veins; indusium thin, membranaceous, soon breaking down and falling away. Mettenius describes the texture as membranaceous.

Sumatra.

* (28) **C. spinulosa**, *Wall.*, Hk. Bk., Syn. Fil., 23; Hk., Spec. Fil., I, tab. XII, C; — var. *muriculata*, Hassk., Journ. of Bot., VII, 322.

Stipe and rachis strongly prickly, often dark-purple. Fronds large, to 3 M. long, 3-pinnatifid. Pinnulae oblong, moderately long-acuminate, \pm 8—12 c.M. long, \pm 2 c.M. broad, pinnatifid nearly to the costa. Lobes numerous, very close, lanceolate-oblong, acute, subfalcate, serrulate. Texture rather flaccid; surfaces naked, except that the costulae are covered beneath with small, bullate scales; veins forked. Sori small, numerous, close to the costulae, not occupying the higher veins; indusium thin, membranaceous, fragile, soon breaking down irregularly.

Var. muriculata: Stipe and rachis muriculate, not spinulose.

Malaya; Formosa, Japan, South. China, North. India.

(29) **C. Muelleri**, *Bk.*, Journ. of Bot., XXVIII, 104.

Fronds large, 3-pinnate; main rachis naked. Lowest pinnae oblong, \pm 30 c.M. long. Pinnulae lanceolate, to $2\frac{1}{2}$ c.M. broad. Tertiary divisions linear-oblong, \pm 3—4 m.M. broad, adnate to the rachis, the edge not recurved, deeply crenate. Texture moderately firm; surfaces naked, except the midribs of the pinnules above; colour dark-green, paler beneath; veins forked in the lower laciniae of the ultimate divisions. Sori 8—10 to the largest leaflets; indusium cup-shaped, entire, persistent.

New Guinea.

* (30) **C. oinops**, *Hassk.*, Journ. of Bot., VII, 322; *C. sinops*, *Rac.* (not *Hassk.*), Flor. Btz., I, 36.

Stipes to 30 c.M. long, clothed with very deciduous scales. Fronds oblong, to 150 c.M. long, to 70 c.M. broad, 3-pinnate; main rachis tomentose above. Pinnae not or hardly narrowed at the base, to 24 c.M. broad, on stalks to 2—3 c.M. long. Pinnulae sessile, linear, to 22 m.M. broad, acuminate, pinnate. Tertiary divisions, \pm close, bluntish, subfalcate, to $2\frac{1}{2}$ m.M. broad, crenate, each crenation corresponding with a small vault. Texture subcoriaceous; colour dark-green, paler beneath; costae densely scaly, with 8—12 forked veins on each side. Sori occupying the lower 3—7 veins; indusium thin, breaking down irregularly.

Java.

(31) **C. schizochlamys**, *Bk.*, Journ. of Bot., XVIII, 209.

Fronds large, 3-pinnate; main rachis thinly tomentose and clothed with

small, acuminate scales. Pinnae oblong-lanceolate, \pm 45 c.M. long, 12—15 c.M. broad. Pinnulae lanceolate, $1\frac{1}{2}$ —2 c.M. broad, narrowed from the middle to the apex, cut down to the rachis into linear, entire or rarely crenulated divisions \pm $2\frac{1}{2}$ m.M. broad, with a distinct space between them. Texture firm, subrigid; both surfaces green; rachises and under surface densely covered with small, ovate-lanceolate, cuspidate scales on the midribs; veins 9—10 on each side, distinct, forked from the base. Sori close to the midribs; indusium fragile, membranaceous, soon breaking down to the base.

Sumatra.

* (52) **C. arachnoidea**, *Hk.*, Hk. Bk., Syn. Fil., 24.

Fronds 3-pinnate; rachises all dark-coloured, opaque, clothed with rusty tomentum and muricate with short, black, sharp spines. Pinnulae 10—12 c.M. long, 1—2 c.M. broad, oblong, pinnate, pinnatifid only near the narrow, acuminate point. Tertiary divisions scarcely $7\frac{1}{2}$ m.M. long, linear from a broad base, acute, subfalcate. Texture coriaceous; under surface cobwebby; veins immersed, inconspicuous. Sori near the middle of the pinnulae, occupying the whole space between the midribs of the ultimate divisions and the recurved margin; indusium thin-membranaceous, whitish, breaking down into variously lobed and lacinated segments. The javan plant (Leg. Scheffer) has the ultimate segments crenate, \pm 1 c.M. long.

Ternate, Java; Queensland.

* (53) **C. celebica**, *v. A. v. R.*

Fronds large, 3-pinnate; main rachis prickly and furfuraceous like the secondary ones; the prickles blackish, strong, sharp; the tomentum ferrugineous. Pinnae 50—60 c.M. long, \pm 20 c.M. broad, on stalks \pm 5 c.M. long. Pinnulae sessile, the lower ones stalked, \pm 2 c.M. broad, the apex short-acuminate, crenate; lowest pinnulae distant, \pm 5 c.M. apart. Tertiary divisions linear, the highest sessile, subfalcate, crenate, blunt, the lowest \pm longest, \pm 2 m.M. broad, spreading, subpetiolate, lobed towards the base, with the lowest segments free, the lowest anterior segments often larger than the other ones. Texture coriaceous but not firm; tertiary rachises tomentose above; surfaces naked or the under surface sparingly and minutely scaly on the veins; veins distinct, \pm 8 on each side, mostly forked. Sori filling up the space between the costae and the edge, occupying all the veins except the simple highest and the 2—3 \times forked lowest ones; indusium thin, gradually evanishing.

Celebes, (Teysmann, No. 15681).

2. HEMITELIA, R. Brown.

Characters of *Cyathea*, but the indusium consisting of a lateral, mostly semiorbicular, convex scale, attached at the inner side of the receptacle, too small to cover the sorus permanently. Fronds never simple and the veins sometimes anastomosing near the costae. Hk. Bk., Syn. Fil., tab. I, fig. 5; Diels, in Eng. & Prantl., Nat. Pfl.Fam., I⁴, fig. 82, A—C.

Tropical and subtropical.

Arrangement of the sections.

§ 1. **EUHEMITELIA.** Costal veins anastomosing. (Not in Malaya).

§ 2. **AMPHICOSMIA.** Veins all free.

§ 2. AMPHICOSMIA.

§. *Stipes not prickly.*

*. *Ultimate segments entire or crenulate.* (1) *H. crenulata.*

.. *Ultimate segments strongly serrate.* (2) *H. capensis.*

§.§. *Stipes prickly, at least at the base.*

‡. *Main rachis smooth.* (3) *H. Junghuhniana.*

‡.‡. *Main rachis prickly.* (4) *H. sumatrana.*

* (1) ***H. crenulata*, Mett.**, Rac., Flor. Btz., I, 58; *Alsophila crenulata*, Mett., Hk. Bk., Syn. Fil., 44.

Stipes different in length, often bearing at the base some contracted or reduced pinnae. Fronds to 150 c.M. long, to 70 c.M. broad, oblong-elliptic, narrowed towards both ends, 3-pinnate. Pinnae lanceolate, to 10 c.M. broad, on stalks to 1 c.M. long. Pinnulae sessile, to 15 m.M. broad. Ultimate segments linear, bluntish, the edge entire or slightly crenate. Texture herbaceous; rachises villose above or beneath; under surface scaly on the costae. Sori 2—6 on each side of the costae, near the base.

Sumatra, Java, Philippines.

(2) ***H. capensis*, R. Br.**, Hk. Bk., Syn. Fil., 29; Bau. & Hk., Gen. Fil., tab. XLII, A; *Cyathea capensis*, Sm., Christ, Farnkr. d. Erde, 522, fig. 1023; *Amphicosmia riparia*, Gardn., Lond. Journ. of Bot. I, tab. XII.

Stipes scaly at the base, often bearing anomalous pinnae, which are 5—10 c.M. long, compoundly and pinnately divided with a rigid rachis and

linear-filiform segments. Fronds ample, 2—5-pinnate. Pinnae petiolate, ovate-oblong. Pinnulae subsessile, 5—7½ c.M. long, 1½—2 c.M. broad, oblong, acuminate, deeply pinnatifid or pinnate again. Ultimate segments linear-oblong, acute, strongly serrate. Texture membranaceo-papyraceous; rachises and costae clothed beneath with scattered, deciduous, bullate scales; veins simple, 1 to each serrature. Sori frequently solitary at the base of the ultimate segments, rarely 3—4; receptacle large, prominent; indusium brown, fimbriate.

Java, South. Africa, Brasilia.

* (3) **H. Junghuhniana**, *Mett.*, *Rac.*, *Flor. Btz.*, I, 38; *Allophila Junghuhniana*, *Kze*, *Bot. Zeit.*, 1848, 284; — var *dissoluta*, *Rac.*, l. c.

Stipes aculeate at the base. Fronds 3-pinnatifid or 3-pinnate; main rachis deciduously tomentose. Pinnae lanceolate-oblong, to ± 60 c.M. long. Pinnulae sessile or subsessile, 5—10 c.M. long, ± 1½ c.M. broad, linear-lanceolate, acuminate. Ultimate segments free or slightly joined at the base, 2—5 m.M. broad, linear-oblong, straight or subfalcate, bluntish, the edge crenate, toothed or lobed. Texture membranaceous to chartaceous; secondary rachises tomentose; colour deep-green above, paler beneath; under surface ± hairy or scaly on the costae and costulae, the scales mostly deciduous; upper surface naked or hairy on the costae; veins 6—9 on each side, forked, the highest simple. Sori 1—4 on each side, occupying the lower veins, ± costular.

Var. *dissoluta*: Ultimate segments 1—5 m.M. apart.

Java, Sumatra.

* (4) **H. sumatrana**, *v. A. v. B.*

Fronds 3-pinnatifid, sub-3-pinnate; main and secondary rachises muricate with numerous, short, sharp, dark-tipped warts, densely but rather deciduously furfuraceo-tomentose and purveyed with many, deciduous, linear, spreading scales, the main one rough between the warts. Pinnae ± 55 c.M. long, stalked. Pinnulae linear-lanceolate, short-acuminate, 8—12 c.M. long, ± 2 c.M. broad, deeply pinnatifid, pinnate at the base. Ultimate segments linear-oblong, subacute, ± 2½ m.M. broad, the lower ones free, the edge ± crenato-dentate. Texture chartaceous, fragile when dry; colour dark above, paler beneath; upper surface piloso-tomentose on the costae (tertiary rachises), under one densely scaly on the costae and costulae; veins 1—2 × forked. Sori 1—4 on each side, subcostular, occupying the lower veins, densely mixed with the fimbriated scales; indusium brown.

Sumatra, (Teysmann, No. 2436).

3. ALSOPHILA, R. Brown.

Sori globose, dorsal on a vein or veinlet or in the forking of a vein; receptacle elevated, rarely punctiform. Indusium wanting or spurious and minute and then consisting of a semiorbicular, inferior, lateral scale placed at the inner side or of a circular, entire or lobed scale or a whorl of fibres with the receptacle in the centre.

Caudex arborescent. Fronds mostly large, divided; veins simple or forked, free or very rarely joining towards the margin. Hk. Bk., Syn. Fil., tab. I, fig. 6; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 85.

Tropical, and extra-tropical in the southern hemisphere.

a. Pinnulae entire, crenate or lobed to $\frac{1}{3}$ -way down to the costa. (1) *A. dubia*.

a.a. Pinnulae lobed or pinnatifid $\pm \frac{1}{3}$ — $\frac{2}{3}$ -way down to the costa.

b. Main rachis naked, but sometimes rough.

§. Sori divaricating, the higher ones nearer the main veins than the lower ones, so as to form a Λ . (2) *A. glabra*.

§.§. Sori not divaricating, medial. (3) *A. Ridleyi*.

b.b. Main rachis more or less scaly, hairy or tomentose.

c. Texture \pm herbaceous.

*. Main rachis very densely scaly and long-hairy. (4) *A. Burbidgei*.

.. Main rachis not densely scaly or hairy.

✧. Main rachis short-hairy above, otherwise naked or paleaceous. (5) *A. squamulata*.

✧.✧. Main rachis moderately long-hairy above, the hairs articulate, mixed with long, linear scales. (6) *A. Margarethae*.

c.c. Texture chartaceous to coriaceous.

d. Stipes bearing 3—5 long, branched spines at the base. (7) *A. ramispina*.

d.d. Stipes without branched spines at the base.

e. Lobes entire. Capsules intermixed with fine hairs. (8) *A. obscura*.

e.e. Lobes crenate. Capsules intermixed with paraphyses, which are clavate at the apex. (9) *A. commutata*.

a.a.a. Pinnulae deeply pinnatifid, often nearly to the costa, sometimes pinnate at the base.

1. Stipes unarmed, but sometimes scabrous with minute, more or less acute warts.

- 2.** *Stipe or rachis scabrous.*
 *. *Veins 7—8 on each side of the costulae. Sori medial.*
 (10) *A. trichodesma.*
 .. *Veins 12—14 on each side. Sori subcostular.* (11) *A. scaberula.*
- 2.2.** *Stipe and rachis not scabrous.*
3. *Pinnulae or tertiary segments ± dimorphous.*
 §. *Barren pinnulae slightly lobed, fertile ones deeply pinnatifid.* (12) *A. dimorpha.*
 §.§. *Both barren and fertile pinnulae deeply pinnatifid but the fertile lobes much smaller than the barren ones.*
 (13) *A. Kingi.*
- 3.3.** *Pinnulae and tertiary segments uniform.*
4. *Under surface densely hairy.* (14) *A. Wallacei.*
4.4. *Under surface naked or ± scaly.*
5. *Main rachis at best pubescent.*
 †. *Lobes entire. Pinnulae not caudate.*
 (15) *A. modesta.*
 †.†. *Lobes serrate. Pinnulae caudate.*
 (16) *A. caudata.*
5.5. *Main rachis pubescent and scaly.*
6. *Surfaces naked.* (17) *A. sangirensis.*
6.6. *Surfaces scaly on the costulae.*
 (18) *A. lepidoclada.*
- 1.1.** *Stipes aculeate or muricate.*
2. *Main rachis smooth, rough or slightly prickly near the base.*
3. *Receptacle with a minute, spurious indusium at the inner side.*
 (19) *A. latebrosa.*
3.3. *Spurious indusium wanting or consisting of a minute, orbicular scale with the receptacle in the centre.*
4. *Main rachis furfuraceous, hairy or scaly, at least on the upper side.*
5. *Spurious indusium present.* (20) *A. saparuensis.*
5.5. *Spurious indusium (most probably) wanting.*
6. *Main rachis furfuraceous.* (21) *A. fuliginosa.*
6.6. *Main rachis hairy and slightly scaly.*
 (22) *A. lepifera.*
4.4. *Main rachis naked or nearly so; spurious indusium (probably) wanting.*

- !. *Sori medial.* (23) *A. batjanensis.*
- !!. *Sori ± costular*
- *. *Lobes obscurely crenato-serrate.* (24) *A. melanopus.*
- *. *. *Lobes distinctly serrate.* (25) *A. ornata.*
- 2'.2'. *Main rachis conspicuously muricate.*
- ‡. *Spurious indusium wanting.* (26) *A. crinita.*
- ‡.‡. *Spurious indusium present, consisting of a minute, inferior, orbicular, entire or lobed scale or a wool of fibres, with the receptacle in the centre. Under surface glaucous.* (27) *A. glauca.*
- a.a.a.a. *Pinnulae pinnate; the segments free or sometimes slightly joined at the base.*
- b. *Texture membranaceous to papyraceous.*
- c. *Pinnae under 15 c.M. broad.*
- d. *Main rachis rough and strigulose. Pinnae ± 8 c.M. broad. Spurious indusium wanting.* (28) *A. Christii.*
- d.d. *Main rachis muricate and furfuraceo-tomentose. Pinnae ± 10—12 c.M. broad. Spurious indusium present.* (29) *A. truncata.*
- c.c. *Pinnae 15—20 c.M. broad.*
- *. *Tertiary segments ± 2½ m.M. broad, crenulate at the apex.* (30) *A. celebica.*
- *. *. *Tertiary segments ± 5 m.M. broad, distinctly crenate.* (31) *A. polyphlebia.*
- b.b. *Texture ± coriaceous.*
- c'. *Main rachis prickly.*
- d'. *Costae densely paleaceous beneath with ragged, tomentose scales.* (32) *A. tomentosa.*
- d'.d'. *Costae naked or not densely scaly beneath.* (33) *A. extensa.*
- c'.c'. *Main rachis not prickly.*
- *. *Pinnulae ± 1—1½ c.M. broad, the costae densely scaly.* (34) *A. lurida.*
- *. *. *Pinnulae ± 4 c.M. broad, the costae fibrillose.* (35) *A. mindanensis.*

* (1) **A. dubia**, *Bedd.*, Journ. of Bot., XXVI, 1, tab. CCLXXIX, A; *A. Teysmanniana*, *Rac.*, MS. in Herb. Bog.

Fronds large, 2-pinnate; rachises brown, naked and glossy beneath, to-

mentose above, the tomentum dark-reddish-brown. Pinnae \pm crecto-patent, stalked, 40—60 c.M. long, with a terminal segment, which is pinnatifid at the base, toothed towards the acuminate point. Pinnulae spreading, short-stalked, \pm 25 on each side, distant, the lowest longest or sometimes somewhat shorter than the next higher ones, \pm 7—10 c.M. long, 10—15 m.M. broad, with the apex acuminate, toothed, the base truncate, the edge broadly crenate or lobed to $\frac{1}{3}$ -way down, the lobes or crenations entire, crenate or denticulate. Texture coriaceous; colour dark above, paler beneath; surfaces naked; costae and veins distinct, dark-coloured; main veins pinnate, veins simple. Sori large in 1 row medial between the costa and the margin or in 2 rows rather near the costa, on the lower 1—2 veins, sometimes in 3—4 rows, not divaricating.

Malacca, Banca, Java, Borneo.

* (2) **A. glabra**, *Hk.*, *Rac.*, *Flor. Btz.*, I, 34; *Bedd.*, *Ferns S.I.*, tab. LX; *A. gigantea*, *Wall.*, (not *Hk.*), *Hk.*, *Spec. Fil.*, I, 53; *A. vexans*, *Ces.*, *Fel. di Bor.*, 4.

Stipe more or less rough, scaly at the base, and as well as the main rachis reddish-black, polished. Fronds oblong-lanceolate, hardly narrowed at the base, 3-pinnatifid, to 3 M. long. Pinnae stalked, 45—60 c.M. long or longer, the apex acuminate and pinnatifid. Pinnulae linear, acuminate, (upper ones sessile, lower ones petiolate), 7—15 c.M. long, 12—24 m.M. broad, the base truncate, the edge pinnatifid \pm $\frac{1}{3}$ — $\frac{1}{2}$ -way down to the costa. Lobes triangular or rounded, crenate or serrate. Texture firm, but not coriaceous; rachises often scaly or tomentose above; surfaces naked; veins simple, 5—6 on each side of the costulae. Sori dorsal on the veins, mostly arranged in the shape of a Λ , not extending to the apices of the lobes.

Trop. Asia.

(3) **A. Ridleyi**, *Bk.*, *Ann. of Bot.*, VIII, 122.

Fronds large, deltoid, 5-pinnatifid; main rachis naked, black-brown, unarmed. Pinnae oblong-lanceolate, the lowest \pm 30 c.M. long; rachis brown, hairy above. Pinnulae sessile, linear, subobtuse, $3\frac{1}{2}$ —5 c.M. long, 10—15 m.M. broad, the base truncate, the lower ones pinnatifid \pm $\frac{1}{2}$ -way down to the costa. Lobes oblong, entire. Texture rigid; surfaces naked; main veins pinnate in the lobes; veins 4—6-jugate, conspicuous, curved, ascending, simple or the lower ones forked. Sori medial.

Singapore.

* (4) **A. Burbidgei**, *Bk.*, Journ. of Bot., XVII, 58.

Fronds ample, 3-pinnatifid. Stipe and rachis densely covered with long, linear, acuminate, yellow-brown, membranaceous scales to 2 c.M. long and with very fine, long and soft, brownish hairs to 8 m.M. long. Pinnae oblong-lanceolate, 30 c.M. or more long, acuminate, ± 10 c.M. broad at the base, the rachis hairy like the main one. Pinnulae sessile or subsessile, 4—7 c.M. long, $1-1\frac{1}{2}$ c.M. broad, cut $\frac{1}{2}-\frac{2}{3}$ -way down into oblong, blunt, subentire lobes ± 3 m.M. broad. Texture herbaceous, but firm; colour dark above, paler beneath; costae hairy and the veins slightly so beneath; veins 4—5-jugate, simple or forked. Sori medial, occupying the lower veins. — The description is taken from a specimen gathered by Hallier, determined by Christ (Ann. Btz., XX, 138).

Borneo.

* (5) **A. squamulata**, *Hk.* (not *Mett.*), Rac., Flor. Btz., I, 53; *A. comosa*, *Wall.*, *Hk. Bk.*, Syn. Fil., 41; *Hk.*, Spec. Fil., I, tab. XX, A; *Bedd.*, Ferns Br.I, I, tab. LXXXIV.

Stipes to 30 c.M. long, rough, paleaceous with light-brown, linear-subulate scales $\pm 2\frac{1}{2}$ c.M. long. Fronds ovate, narrowed towards both ends, to 180 c.M. long; main rachis setose or tawny-villous above, the rest naked or paleaceous. Pinnae 35—50 c.M. long, lanceolate, stalked, slightly narrowed towards the base. Pinnulae close, sessile or short-stalked, linear, acuminate, 15—20 c.M. long, 10—15 m.M. broad, the base truncate, the edge cut $\frac{1}{2}-\frac{2}{3}$ -way down to the costa with narrow sinusses. Lobes close, \pm crenulate, to 4 m.M. broad, oblong, bluntish, slightly falcate. Texture herbaceous; surfaces glabrous; colour yellowish-green; secondary rachises like the main one; costulae pinnate in the lobes, naked or slightly scaly beneath; veins simple or forked. Sori medial, 1—3 or rarely more on each side of the costulae, occupying the lower veins.

Malaya.

* (6) **A. Margarethae**, *Schroet.*, Ann. Btz., XX, 156.

Main rachis unarmed, rough, hairy above, the hairs castaneous, not short, intermixed with long, pale-coloured scales. Pinnae short-stalked, ± 40 c.M. long, ± 15 c.M. broad, ovate-oblong, acuminate. Pinnulae sessile, ± 25 on each side below the pinnatifid apex, $\pm 10-11$ m.M. broad, close, lobed $\pm \frac{1}{2}$ -way down to the costa. Lobes pectinate, nearly imbricate, ± 4 m.M. long,

rhomboid-oblong, \pm rectangular, the apex truncate, the margin entire. Texture herbaceous; upper surface naked, the under one and the margin covered with scattered, long, white hairs and few, minute scales; veins prominent, forked. Sori medial, not confluent, mixed with hairs; receptacle small, elevated, pale-coloured.

Borneo.

* (7) **A. ramispina**, *Hk.*, Hk. Bk., Syn. Fil., 42.

Stipes subasperous, tomentose above, ebeneous or dark-brown like the rachises, slightly paleaceous at the base and bearing there 5—5 subsecund, black spines $2\frac{1}{2}$ —4 c.M. long, with many sharp, divaricating branches. Fronds 5-pinnatifid; rachises tomentose above. Pinnae \pm 50 c.M. long or longer, gradually narrowed at the pinnatifid apex. Pinnulae 4— $7\frac{1}{2}$ c.M. long, 1—2 c.M. broad, oblong-lanceolate, sessile or short-stalked, the apex acuminate, the point bluntish, the base truncate, the edge pinnatifid $\pm \frac{1}{2}$ — $\frac{2}{3}$ of the way to the costa. Lobes oblong, subfalcate, obtuse, crenate or serrate towards the apex. Texture subcoriaceous; colour dark-green above, paler beneath where the costae and costulae are \pm scaly; veins mostly simple. Sori small, of few sporangia, nearer the costulae than the edge, not reaching the apices of the lobes.

Borneo.

* (8) **A. obscura**, *Scort.*, Journ. of Bot., XXV, 521, tab. CCLXXVIII, fig. 2.

Stipes 30—60 c.M. long, densely clothed near the base with long, linear-lanceolate, acuminate, sharply serrated scales. Fronds 1— $1\frac{1}{2}$ M. long, \pm 60—80 c.M. broad, 5-pinnatifid, the rachis scaly or tomentose above, naked beneath. Central pinnae \pm 30—40 c.M. long, 7—15 c.M. broad, the base a little narrower, the apex pinnatifid, acuminate, the rachis often hairy above. Pinnulae linear-oblong, \pm 1— $1\frac{1}{2}$ c.M. broad, the base truncate, the apex bluntish, toothed, the edge cut $\pm \frac{1}{2}$ -way down into broad, obtuse, entire segments. Texture \pm coriaceous; costae hairy above; costules naked or \pm scaly beneath, pinnate in the lobes; veins simple or forked, 4—5 on each side. Sori medial on the lower veins, not occupying the lobes; capsules densely intermixed with long, filiform hairs.

Malacca.

* (9) **A. commutata**, *Mett.*, Hk. Bk., Syn. Fil., 45; *A. squamulata*,

Bedd., Ferns Br.I., II, tab. CCXXXV; *Gymnosphaera squamulata*, *J. Sm.* (not *Bl.*), *Bau. & Hk.*, Gen. Fil., tab. C.

Stipes paleaceous, the base tuberculate-asperous. Fronds 3-pinnatifid, the main rachis dark-purple-ebeneous like the stipe and slightly squamose, (tomentose above?). Pinnae \pm 45 c.M. long, 15—20 c.M. broad, stalked. Pinnulae rather distant, stalked, linear, acuminate, the point serrulate, the base truncate, the edge cut \pm $\frac{1}{2}$ -way down to the costa. Segments oblong, obtuse, rotundato-serrate; the margin slightly thickened or recurved. Texture coriaceous; costulae with a few deciduous, bullate scales; veins simple or forked. Sori mostly confined to the lower part of the lobes, the higher ones close to the costulae, the lower slightly divaricating; capsules mixed with paraphyses, which are clavate at the apex.

Malacca, Batjan.

(10) **A. trichodesma**, *Scort.*, Journ. of Bot., XXV, 321.

Stipes scaly at the base. Fronds 120—180 c.M. long, 3-pinnatifid, the main rachis scabrid above, clothed with adpressed, spreading, semi-viscous, copious, jointed hairs. Pinnae short-stalked, narrow-lanceolate, 60—75 c.M. long, 15—20 c.M. broad, the rachis stramineous, hairy like the main one. Pinnulae linear-lanceolate, \pm 2 c.M. broad, cut down nearly to the costa into linear, obtuse, crenulate, rather distant segments. Texture thinly herbaceous; surfaces hairy like the main rachis, principally on the veins; costulae pinnate; veins forked, 7—8 on each side. Sori medial, 5—6 on each side of the costulae.

Malacca.

(11) **A. scaberula**, *Christ*, Schum. & Laut., Flor. deut. Sch.geb., 110.

Stipes yellowish-brown, covered with short, often glanduligerous hairs and with numerous minute, acute warts. Fronds large, \pm 2 M. long, 3-pinnatifid. Pinnae sessile, \pm 45 c.M. long, lanceolate; rachis clothed with narrow, short scales. Pinnulae \pm 8 c.M. long, broadly lanceolate, short-acuminate, not caudate, alternate, distant, \pm 2 c.M. broad, sessile, cut down nearly to the costa into \pm close, falcate, broadly ligulate, obtuse lobes, which are regularly serrato-crenate. Texture thin, herbaceous, flaccid; veins conspicuous, forked, 12—14 on each side; surfaces naked, but the costae bullato-squamose beneath. Sori occupying the lower portions of the lobes, nearly close to the costulae; receptacle globose.

New Guinea.

(12) **A. dimorpha**, *Christ*, Ann. Btz., XIX, 41.

Stipe and rachis unarmed, castaneous, angular, covered with dark, furfureous scales. Fronds large, ± 2 M. long, ± 40 c.M. broad, 2-pinnate in the barren, 5-pinnatifid in the fertile portions. Lower and higher pinnae sterile and shorter, central fertile and longer. Barren pinnae ± 20 c.M. long, ± 7 c.M. broad, distant, acuminate, stalked; the pinnulae stalked, with the base cordate, hardly 1 c.M. broad, narrowed gradually to the point, the apex acuminate, the edge slightly lobed into close, not deep, roundish or acute, crenulate lobes. Texture coriaceous; upper surface dark-green, under one paler, scaly at or near the costae. Fertile pinnulae $\pm \frac{1}{2}$ c.M. broad, pinnatifid nearly to the costa, the lobes not close, ± 5 m.M. broad, obtuse, deltoid. Sori large, 5—6 to a lobe, occupying its whole under surface.

Celebes.

* (13) **A. Kingi**, *Clarke*, Bedd., Handb. Ind. Ferns, 475.

Fronds 5-pinnatifid; rachises tomentose above, naked, brown, somewhat glaucous beneath. Pinnae 50—60 c.M. long, the apex pinnatifid, toothed at the point. Pinnulae distant, short-stalked, the lowest largest or a little shorter, ± 6 —8 c.M. long, ± 1 — $1\frac{1}{2}$ c.M. broad, the apex acuminate and bluntly crenate, the edge cut down nearly to the costa into not very close, oblong, subfalcate, blunt, slightly crenate lobes, the lowest lobes \pm largest. Texture coriaceous; surfaces naked, of a bluish-glaucous tint; costae \pm scaly beneath, hairy above, very prominent especially in the fertile pinnulae, which have the soriferous segments mostly much smaller than the barren segments; veins forked or sometimes simple. Sori in 1 row on each side of the costulae, filling up the whole under surface of the fertile segments, which have the edges recurved.

Malacca.

(14) **A. Wallacei**, *Mett.*, Hk. Bk., Syn. Fil., 459.

Fronds large, 5-pinnatifid; rachises stramineous, densely pubescent. Pinnae oblong-lanceolate, 50—40 c.M. long. Pinnulae ligulate, distinctly stalked, 3—5 c.M. long, ± 1 c.M. broad, cut down to a narrow wing. Segments close, ligulate, blunt, obscurely toothed, $\pm 2\frac{1}{2}$ m.M. broad. Texture subcoriaceous; surfaces green, the under one densely pilose, with a few bullate scales on the ribs; veins 4—5-jugate, simple. Sori costular.

Borneo.

(15) **A. modesta**, *Bk.*, Journ. of Bot., XVIII, 210.

Stipes to 30 c.M. long; the scales glossy, dark-brown, linear, to $1\frac{1}{2}$ c.M. long. Fronds 65—90 c.M. long, rhomboid, 3-pinnatifid, the rachis pubescent, unarmed. Pinnae oblong-lanceolate, the central ones the largest, 20—30 c.M. long, $6\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad, short-stalked, the lower smaller, decidedly petiolate. Pinnulae lanceolate, sessile, 1 — $1\frac{1}{2}$ c.M. broad, cut down in the lower part to the rachis and in the upper to a narrow wing into close, ligulate, entire, obtuse segments to $\pm 2\frac{1}{2}$ m.M. broad. Texture moderately firm; surfaces green and nearly naked except a few lanceolate scales on the midribs of the pinnulae beneath; veins distinct, 6—7-jugate, simple or forked. Sori costular, not reaching the margin.

Sumatra.

* (16) **A. caudata**, *J. Sm.*, Hk. Bk., Syn. Fil., 42; Hk., Spec. Fil., I, tab. XX, B.

Fronds 3-pinnatifid; rachises tomentose or pubescent above (after Hooker's drawing). Pinnulae sessile, oblong-lanceolate, ± 6 —8 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, broadest at the base, the apex suddenly contracted into a long, narrow, serrate or toothed, tail-like acumen, the edge cut down nearly to the costa into close, oblong, slightly falcate, rather obtuse, \pm serrate lobes ± 4 m.M. broad. Texture semicoriaceous, firm; surfaces naked; veins simple or forked. Sori close to the costulae, occupying the lower half of the ultimate segments.

Luzon.

(17) **A. sangirensis**, *Christ*, Warb., Mons., I, 90.

Stipe stramineous, angular, unarmed, but covered like the main rachis with minute, filiform and subulate scales. Fronds ± 70 c.M. long, broadly ovate-deltoid, 3-pinnatifid. Pinnae 10—15, distant, ovate, the rachis strigoso-pilose. Pinnulae ± 4 c.M. long, ± 1 c.M. broad, deeply incised. Lobes separated by rounded sinusses, ovate, rounded, crenulate. Surfaces naked, the under one paler than the upper one; veins simple. Sori close, ± 4 on each side of the costulae; receptacle not elevated. — Fronds with the aspect of *Dryopteris setigera*, *O. Ktze*, but firmer.

Sanggir Islands.

(18) **A. lepidoclada**, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., Suppl., 37.

Stipes unarmed, fusco-stramineous, castaneous at the base, hardly 10 c.M.

long, covered with ovate-acuminate, scariose, ochreous scales $\pm \frac{1}{2}$ c.M. long. Fronds ± 60 c.M. long, ± 28 c.M. broad, ovate-elongate, acuminate, narrowed towards the base. Pinnæ sessile, close, spreading, ± 50 on each side below the pinnatifid apex, the central ones the longest, bluntish, ± 3 c.M. broad. Pinnulae close, ± 25 on each side, sessile, ovate-elongate, obtuse, $\pm \frac{1}{2}$ c.M. broad, cut down nearly to the costa, the segments ovate, ± 7 on each side, apiculate or obtuse, $\pm 1\frac{1}{2}$ m.M. broad. Texture herbaceous; upper surface dark-green, the under one paler, not glaucous; rachis ochreo-stramineous, short-hairy, covered like the costae with scariose, whitish, lanceolate, acute scales, the costulae covered with broadly ovate scales; veins conspicuous, 3—4-jugate. Sori minute, 3—4 to each ultimate segment, placed at the bases of the veins; receptacle punctiform or hardly elevated; indusium none.

New Guinea.

* (19) **A. latebrosa**, *Wall.* (not *Hk.*), *Hk. Bk., Syn. Fil.*, 43; *Bedd., Ferns S.I.*, tab. LVIII; *Hemitelia latebrosa*, *Mett., Rac., Flor. Btz.*, I, 39.

Stipes dark-brown, aculeate towards the base, muricate upwards; main rachis muricate towards the base. Fronds 3-pinnatifid, to 2 M. long, to 1 M. broad; rachises often covered with long, linear, fine, brown, deciduous scales on the upper side. Pinnæ lanceolate-oblong, acuminate, stalked, 30—60 c.M. long, 12—20 c.M. broad, the rachis \pm tomentose above. Pinnulae linear-lanceolate, acuminate, subsessile, $\pm 1\frac{1}{2}$ —2 c.M. broad, the base subtruncate, the edge deeply pinnatifid into linear-oblong, blunt, subfalcate, crenate or slightly toothed lobes ± 2 m.M. broad. Texture firm-herbaceous; surfaces naked or the under one slightly hairy, under surface not glaucous; costae and costulae more or less scaly beneath with \pm bullate scales; veins forked. Sori on the 1—7 lower veins, nearly costular; receptacle often with a minute, spurious indusium at the inner side.

Malaya; North. India.

* (20) **A. saparuensis**, *v. A. v. B.*

Stipes aculeate; main rachis slightly so towards the base; secondary rachises somewhat rough towards the base with few, scattered, minute warts and covered like the main one with copious, very fine, \pm deciduous, crisped, yellowish or brownish, fibril-like hairs. Central pinnæ ± 45 c.M. long, 12—16 c.M. broad, lanceolate, acuminate. Pinnulae sessile or subsessile, the lower ones spreading, smaller than the central ones, which are $\pm 1\frac{1}{2}$ c.M. broad, cut down nearly to the costa. Lobes \pm close, oblong, blunt, crenate

or toothed especially towards the apex, the lowest lobes a little smaller and somewhat narrowed at the base. Texture firm but not rigid; veins distinct, forked, 8—10 on each side; costae and costulae hairy like the rachises, especially beneath. Sori occupying nearly all the veins except the higher ones, medial; receptacle hairy, with a distinct, flat, brown, subentire, orbicular, spurious indusium. — Under surface paler than the upper one, but not glaucous.

Saparua, (Teysmann), *Buru*.

(21) **A. fuliginosa**, *Christ*, Bull. Herb. Boiss., VI, 138.

Stipes blackish-brown, the base covered with aculeate, rough tubercles and long, erect, curved, rigid, dark-coloured bristles. Fronds broadly ovate, \pm 100 c.M. long, \pm 40 c.M. broad, 5-pinnatifid, narrowed towards the base; main rachis yellowish, rough, furfureaceous. Pinnae close, broadly ovate, \pm 8 c.M. broad, sessile, short-acuminate, the rachis clothed above like the costae with large scales, some of which are dark-coloured and lanceolate and the others bullate or cupuliform and greyish with a black centre. Pinnulae ligulate, short-acuminate, \pm 5 m.M. broad, \pm 22 on each side below the pinnatifid apex, the lower ones shorter, cut down nearly to the costa into erectopatent, pectinate, falcate, subacute, slightly crenate lobes, with the margin revolute; veins conspicuous, 4—5 on each side, naked or scaly. Sori minute, numerous; indusium not seen.

Philippines.

(22) **A. lepifera**, *J. Sm.*, Hk., Spec. Fil., I, 54; (? *A. tomentosa*, *Hk.*, C. Chr., Ind. Fil.); — var. *congesta*, *Christ*, Bull. Herb. Boiss., VI, 37.

Stipe furfureaceous at the base and covered like the lower portion of the main rachis with small, prominent, aculeate tubercles with dark-red or blackish points; upper portion of the rachis clothed with long, moderately firm, coarsish, brown hairs and with few, slender, yellowish, ovate, crinite scales like the stipe. Fronds 5-pinnatifid, sub-5-pinnate. Pinnulae sessile, broadest at the base, narrow, lanceolate, deeply pinnatifid, the point slender, subulate. Segments linear-oblong, distant, falcate, \pm obtuse, the lower ones free. Texture herbaceous; under surface slightly glaucous and covered with flattish, whitish scales on the costae; veins forked. Sori occupying the whole under surface of the segments.

Var. congesta: Segments close or imbricating, the lower ones overlapping the rachis.

Philippines.

(23) **A. batjanensis**, *Christ*, Warb., Mons., I, 90.

Caudex \pm 3 M. high. Stipes very aculeate, the prickles short, conical; rachis rough, less prickly, yellowish-brown. Fronds deltoid, 3-pinnatifid, 100 c.M. and more long. Pinnae 8—10, very distant, petiolate, deltoid-ovate, 8—10 c.M. long, \pm 6 c.M. broad, short-acuminate. Pinnulae distant, deeply incised; segments obtuse, ligulate-ovate, 6—8 m.M. long, \pm 3 m.M. broad, serrate. Texture coriaceous; surfaces naked, the under one not glaucous; veins simple, conspicuous. Sori small, not close, medial.

Batjan.

* (24) **A. melanopus**, *Hassk.*, Journ. of Bot., VII, 525.

Stipe very aculeate and densely scaly at the base, the rest rough like the rachis, the scales large, blackish, deciduous. Fronds 3-pinnatifid, ovate-oblong, acute. Pinnae elongate-oblong and acute, or lanceolate-oblong and acuminate. Pinnulae linear-oblong, deeply pinnatifid. Ultimate segments linear-oblong, obtuse or subacute, subfalcate, obscurely crenato-serrate. Texture membranaceous; under surface slightly glaucous; costulae clothed with minute, deciduous scales. Sori near the costulae, at length confluent.

Java.

(25) **A. ornata**, *Scott*, Hk. Bk., Syn. Fil., 460; Bedd., Ferns Br.I., II, tab. CCCXLII; — var. *sikkimensis*, Bedd., Journ. of Bot., XXXI, 225; *A. sikkimensis*, *Clarke & Bk.*, Journ. of Linn. Soc., Bot., XXIV, 409.

Stipe paleaceous at the base, slightly muricate and as well as the main rachis glossy, purplish-brown. Fronds 3-pinnatifid. Pinnae oblong-lanceolate, rather abruptly contracted into a short, deeply pinnatifid apex. Pinnulae nearly all petiolate, lanceolate-oblong, acuminate, deeply cut down nearly to the costa. Ultimate segments ovate in the sterile, narrow-oblong in the fertile pinnulae, serrate, slightly falcate. Texture herbaceo-membranaceous; costae slightly hairy above and scaly below; veins forked, or a few 3-branched. Sori subcostular, occupying the lower half of the lobes (conf. Beddome's drawing).

Var. sikkimensis: Stipe and main rachis more prominently muricate.

Sikkim; — var. *sikkimensis*: *Malacca*; *Sikkim*.

* (26) **A. crinita**, *Hk.*, Hk. Bk., Syn. Fil., 42; Hk., Icon. Pl., VII, tab. DCLXXI; Bedd., Ferns S.I., tab. LIX.

Stipe and main rachis strongly muricate, stramineous-brown, long-hispido-

paleaceous; spines short, tipped with a black gland. Fronds large, 3-pinnatifid. Pinnae \pm 60 c.M. long, \pm 25 c.M. broad. Pinnulae oblong, acuminate, pinnatifid nearly to the costa. Lobes narrow, oblong, subobtuse, subfalcate, the margin recurved, serrate. Texture subcoriaceous; rachises pilose above, densely paleaceo-crinite beneath; costae and veins often pilose, or scaly below; veins 1 \times forked. Sori subcostal (conf. Hooker's drawing), mixed with crinite scales, often covering the whole underside of the lobes.

Java; Ceylon, South. India.

* (27) **A. glauca**, *J. Sm.*, Journ. of Bot., III, 419; *A. contaminans*, *Wall.*, Rac., Flor. Btz., I, 35; *Hk.*, Spec. Fil., I, tab. XVIII; *A. Brunomiana*, *Bedd.*, Ferns Br.I., I, tab. LXXXVI; *Gnoophora glauca*, *Bl.*, (oldest name); — var. *celebica*, *Christ* in *Ann. Btz.*, XV, 79; — var. *longepaleata*, *Christ*, l. c., XIX, 42; — var. *squamulata*, *densa* & *setulosa*, *Hassk.* in *Journ. of Bot.*, VII, 524.

Stipe and main and secondary rachises more or less aculeate with short prickles and whether or not tomentose above. Fronds large, to 5 M. long, to 1½ M. broad, 3-pinnatifid to sub-3-pinnate. Pinnae \pm 12—24 c.M. broad, lanceolate-oblong, acuminate, more or less stalked. Pinnulae linear- or oblong-lanceolate, 1½—5 c.M. broad, cut down nearly to the costa or even pinnate at the base into numerous, linear-oblong, blunt, acute or acuminate, subfalcate, obscurely crenate to serrate segments, the fertile segments not rarely narrower than the barren ones, the margin subrevolute. Texture more or less coriaceous; under surface glaucous; veins \pm 8—12 on each side, mostly 1—2 \times forked. Sori commonly nearer the costulae than the margin; receptacle surrounded by a spurious indusium, consisting of a flat, orbicular, entire or lobed, minute scale or a whorl of fibres. — A variable species. — The typical form (*var. robusta*, *Hassk.*) has the rachis and the stipe reddish, often glaucous, the stipe densely clothed at the base with long, pale-coloured scales; the costae and costulae naked or covered beneath with few, small, deciduous, bullate scales.

Var. celebica: Pinnae and pinnulae short and broad; ultimate segments not numerous, 2—3 m.M. apart; veins 3-furcate.

Var. longepaleata: Rachis yellowish-ochraceous, covered with short, dark-coloured prickles and long, brown, linear-subulate, spreading scales. Texture less coriaceous, less glaucous.

Var. squamulata: Costae and costulae densely scaly beneath.

Var. densa: Sori dense, confluent, at length covering the whole under surface of the lobes.

Var. setulosa: Costae and costulae clothed beneath with moderately long, whitish, subulate bristles especially in the barren portions.

Trop. Asia. — *The varieties in Malaya, chiefly Celebes or Java.*

(28) **A. Christii**, *v. A. v. R.*; *A. comosa*, *Christ* (not *Wall.*), *Ann. Btz.*, XV, 80.

Stipes strong, ± 2 c.M. thick, pale-brown, rough with numerous, minute, acute warts, not scaly. Fronds large; main rachis slightly rough and strigulose. Pinnae sessile, ± 40 c.M. long, ± 8 c.M. broad, lanceolate. Pinnulae spreading, acuminate, $\pm \frac{3}{4}$ c.M. broad, pinnate. Ultimate divisions (leaflets) lanceolate-falcate, obtuse, ± 5 m.M. broad, the apex rounded, crenate. Texture herbaceous; costae clothed above with short, whitish hairs and covered beneath with scattered, brown, bullate scales; veins forked. Sori large, yellowish-brown, 6—8 on each side, not reaching to the apex of the ultimate segments.

Celebes.

* (29) **A. truncata**, *Brack.*, *Hk. Bk.*, *Syn. Fil.*, 59; *Hemitelia truncata*, *Christ*, *Ann. Btz.*, XV, 81, tab. XIII, fig. 4. — var. *nivea*, *Christ* in *Schum. & Laut*, *Flor. deut. Sch.geb.*, *Suppl.*, 56.

Main & second. rachises dark-purple or brownish, glossy, muricate, decidedly furfuraceo-tomentose as well as the costae and costulae beneath. Fronds 3-pinnate. Pinnae 25—45 c.M. long, 10—12 c.M. broad, oblong, acuminate. Pinnulae to $7\frac{1}{2}$ c.M. long, to $1\frac{1}{2}$ c.M. broad, sessile, lanceolate, acuminate, numerous (25—30 on each side), pinnate, the short apex pinnatifid. Ultimate segments to 12 m.M. long, to 3 m.M. broad, herbaceous, firm, the lower ones \pm petiolate, the base subauriculate or hastate, the margin roundly crenate, recurved. Costules thick and prominent, bullato-squamulose; veins simple or forked; colour bright-green above, paler beneath. Sori small, dark-orange, lax, near the costulae; indusium consisting of a narrow, flat, thin, orbicular scale.

Var. nivea: Fronds 2—2 $\frac{1}{2}$ M. long, $\pm 1\frac{1}{2}$ M. broad; under surface covered with adpressed, white tomentum.

Celebes, Batjan, New Guinea; Fiji, Samoa; — var. *nivea: New Guinea.*

(30) ? **A. celebica**, *Mett.*, *Ann. Mus. Bot. L. B.*, I, 53; ? *Cyathea celebica*, *Bl.*, *Enum.*, 245.

Fronds \pm tripinnatisect; main rachis ferrugineo-hirsute in the upper portion, subfurfuraceous, scaly and slightly aculeate or rough in the lower portion. Pinnae ± 30 c.M. long or longer. Pinnulae approximate, ± 10

c.M. long, $1\frac{1}{2}$ —2 c.M. broad, linear-oblong, acuminate, inciso-serrate at the apex. Tertiary divisions numerous, linear-oblong, subfalcate, obtuse, sessile, $\pm 2\frac{1}{2}$ m.M. broad, the apex crenulate. Texture membranaceous; veins fine, pellucid, forked at the base. Sori?

Celebes, Ternate, Batjan.

(31) **A. polyphlebia**, *Bk.*, Journ. of Linn. Soc., Bot., XII, 104.

Trunk to $7\frac{1}{2}$ M. high. Fronds ample, tripinnate. Rachis of pinnae pale-brown, slightly furfuraceous and rough with raised points. Pinnae oblong-lanceolate, ± 60 c.M. long, 15—20 c.M. broad. Pinnulae sessile, lanceolate-ligulate, $\pm 2\frac{1}{2}$ c.M. broad, cut down to the rachis into close, blunt, distinctly crenate, tertiary segments ± 5 m.M. broad. Texture thin, but firm; surfaces naked, the under one with a few bullate scales on the midribs of the tertiary segments; veins 12—14-jugate, close, fine, distinct, forked, the lowest sometimes bifurcate. Sori placed close to the costae, with a distinct space between each, sometimes 12—15 to a segment, absent from the upper $\frac{1}{3}$ of the segments.

Aru Islands.

* (32) **A. tomentosa**, *Hk.*, Rac., Flor. Btz., I, 52.

Stipes densely covered with light-brown scales 2—5 c.M. long, muricate like the more or less densely scaly rachises; the spines short, often black-tipped. Fronds large, to $2\frac{1}{2}$ M. long, ovate-lanceolate, 3-pinnate. Pinnae to 45 c.M. long or longer, ± 15 c.M. broad, stalked. Pinnulae sessile, $\pm 1\frac{1}{2}$ c.M. broad, linear-lanceolate, acuminate. Tertiary segments sessile, free or slightly joined at the base, $\pm 2\frac{1}{2}$ m.M. broad, entire, blunt, slightly convex. Texture coriaceous; upper surface naked or slightly hairy on the costae; under surface densely covered with pale-brown-yellowish hairs and tomentose, ragged scales on the costae and tertiary rachises; veins 8—12 on each side, mostly forked. Sori close to the central ribs of the ultimate segments, hidden between the scales.

Java.

* (33) **A. extensa**, *Spr.*, Syst., IV, 124; *A. lunulata*, *R. Br.*; *Hk. Bk.*, Syn. Fil., 41; *A. Haenkei*, *Pr.*, Rel. Haenk., I, 68; *Polypodium extensum*, *Forst.*, (oldest name); — var. *intermedia*; *A. intermedia*, *Mett.*, *Hk. Bk.*, l. c.

Fronds large, 3-pinnate; the rachises yellowish or pale-brown, \pm densely muricate, naked below, often hairy above. Pinnae oblong-lanceolate, 45—60 c.M. long. Pinnulae close or approximate, ligulate, sessile, 10—15 c.M. long,

1½—2½ c.M. broad. Segments close, free or slightly joined at the base, ligulate, falcate, blunt, ± 2½ m.M. broad, obscurely crenulate. Texture subcoriaceous; surfaces naked; midribs below naked or furfuraceous; veins crowded, ± 10—12-jugate, forked. Sori minute, subcostular; receptacle pilose.

Var. intermedia: Ultimate segments broader; veins and sori more distant.

Java & Celebes to Polynesia; — var. intermedia: New Guinea; New Caledonia.

* (34) **A. lurida**, *Hk.*, Hk. Bk., Syn. Fil., 45.

Stipes smooth; rachis slightly scabrous and scaly below, densely tomentose and scaly above. Fronds broadly lanceolate. Pinnae to ± 30 c.M. long and to 10 c.M. or more broad. Pinnulae more than 5 c.M. long, more than 1 c.M. broad, cut down to the rachis into linear-oblong, subfalcate, nearly entire leaflets. Texture subcoriaceous; upper surface dark-green, naked, under surface densely furnished with linear scales on the rachis and costae.

Java, Celebes.

(35) **A. mindanensis**, *Christ*, Warb., Mons., I, 90.

Rachis stramineous, unarmed, covered with a woolly, brown tomentum and long, reddish-brown, fine, fibrillose scales. Pinnae ± 40 c.M. long, ovate, subsessile. Pinnulae numerous, 20—30, ligulate-lanceolate, ± 8 c.M. long, short-acuminate, the lower ones pinnatifid quite to the costa. Segments of the sterile pinnae ± 2 c.M. long, close, ± 6 m.M. broad, deeply incised, with the lobes ovate, bluntish, crenate; those of the fertile pinnae, smaller, narrower, rather distant, ligulate, entire or slightly crenulate. Texture very coriaceous; costae fibrillose like the rachis; veins numerous, forked. Sori minute, close to the central veins of the segments, not reaching the apex.

Philippines.

Species of an indefinite systematical place.

(36) ? **A. xantholepis**, *Christ*, Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 138.

Fronds somewhat larger than in *A. tristis*, *Bl.* (which is *Cystopteris tristis*, *Mett.*), becoming black when dried.

Batjan.

TRIBE II. DICKSONIEAE.

4. DICKSONIA, *l'Héritier*.

Sori roundish or transversally suboblong, terminal on the veins, submarginal. Indusium 2-valved, the outer valve consisting of a conspicuously cucullate, small lobe of the frond, forming with the inner valve a saccate involucre to the sorus, the inner valve being of another texture than the outer one.

Caudex arborescent. Fronds compound, \pm elongate as are the segments; veins free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 78, B—D.

Tropical and subtropical.

* **D. Blumei**, *Moore*, C. Chr., Ind. Fil.; *D. chrysotricha*, *Moore*, Rac., Flor. Btz., I, 121; *Balantium Blumei*, *Kze*, (oldest name).

Caudex arborescent. Stipes to 80 c.M. long, densely pilose like the rachises and costae, the base covered with long, yellowish or brownish hairs to $3\frac{1}{2}$ c.M. long. Fronds 5-pinnatifid to 3-pinnate, to 3 M. long, narrowed towards both ends. Pinnae 50—75 c.M. long, 15—25 c.M. broad, erecto-patent, on stalks to $1\frac{1}{2}$ c.M. long, oblong-lanceolate, acuminate. Pinnulae \pm sessile, approximate, linear-lanceolate, acuminate, the lowest not the longest, 15—20 m.M. broad at the base. Tertiary segments erecto-patent, linear-oblong, obtuse, 3—4 m.M. broad, free or slightly joined at the base; the barren ones decurrent, toothed, the teeth acute, entire or slightly denticulate; the fertile ones deeply incised, the lobes rounded, cucullate, forming the outer valve of the indusium. Texture coriaceous; surfaces naked, except on the costae or costulae beneath; veins simple or forked. Sori numerous, 1 to each lobe, transversally suboblong; inner valve of the indusium \pm cucullate, brown, firm-membranaceous, at length reflexed.

Java, Celebes.

5. CIBOTIUM, *Kaulfuss*.

Sori roundish or oblong, terminal on the veins, submarginal. Indusium 2-valved, the outer valve similar to the inner one, distinctly differentiated from the margin of the frond, forming with the inner one a saccate involucre to the sorus.

Caudex often, but not invariably arborescent. Fronds compound; segments linear or oblong; veins free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 78, E—F; Christ, Farnkr. d. Erde, fig. 1001.


Trop. Asia, China, Sandwich Islands, Centr. America.

* **C. barometz, J. Sm.** (not *Link*), Christ, Farnkr. d. Erde, 315; *C. assamicum*, *Hk.*, Spec. Fil., I, tab. XXIX, B; *C. glaucescens*, *Kze*, Farnkr., I, tab. XXXI; *Dicksonia Barometz*, *Link*, Hk. Bk., Syn. Fil., 49.

Caudex short, strong. Stipes firm, the base densely covered with very long, yellow, hair-like scales. Fronds ovate-oblong, to 2 M. or more long. Lower pinnae ovate-lanceolate, \pm 50—60 c.M. long. Pinnulae linear-lanceolate, acuminate, cut nearly and sometimes quite down (in the lower part) to the costa into oblong, narrow, subfalcate, approximate segments. Texture subcoriaceous; surfaces \pm naked, shining, the under one glaucous, sometimes furfuraceous; veins commonly simple. Sori 2—12, rarely 1 to a segment; indusium brownish, transversely oblong.

Malaya; Assam, South. China.

FAMILY II. MATONIACEAE.



II. MATONIACEAE.

Sori round, consisting of few (4—10) decumbent capsules placed in a whorl. Capsules as in Cyathea. Receptacle columnar, bearing a superior, umbrella-shaped, obscurely lobed indusium.

Rhizome creeping. Fronds *either* erect and pedato-flabellate *or* pendulous and pinnate, the pinnae simple or 1—2 × forked; veins free or those around the sori reticulate.

Malayan.

No. 6. **Matonia.**

6. MATONIA, R. Brown.

Characters of the family.

- *. *Fronds pendulous, pinnate, not pedato-flabellate.* (1) *M. sarmentosa.*
.. *Fronds erect, pedato-flabellate, not pinnate.* (2) *M. pectinata.*
-

(1) **M. sarmentosa**, *Bk.*, Christ, Farnkr. d. Erde, 556; Diels, in Engl. & Prantl, Nat. Pfl.Fam., fig. 183; Ann. of Bot., V, tab. XIV.

Rhizome short, creeping, scaly. Stipes slender. Fronds pendulous, to ± 1 M. long, not fan-shaped, pinnate or irregularly branched and the branches pinnate again. Leaflets simple or once or more times forked, the ultimate segments linear, 5—15 c.M. long, 4—10 m.M. broad, the apex blunt, the edge entire. Texture flaccido-coriaceous; surfaces naked; colour pale; costae stout; veins immersed, simple or forked. Sori terminal on short veinlets, placed ± midway between the costae and the margin, consisting of ± 8—10 sporangia.

Borneo.


* (2) **M. pectinata**, *R. Br.*, Christ, Farnkr. d. Erde, 555, fig. 1064; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 182; Bedd., Ferns Br.I., II, tab. CLXXXVI; Bau. & Hk., Gen. Fil., tab. XLIII.

Rhizome creeping, hairy. Stipes stout, erect, to ± 2 M. long. Fronds fan-shaped, conjugato-pedatisect, 40—60 c.M. broad, forked. Forkings spreading or divaricating, each pinnate with many pinnulae on the upper side below

the pinnatifid apex, the lower side destitute of pinnulae. Pinnulae linear, the lowest largest, to 45 c.M. long, cut down nearly to the costa into linear or linear-oblong, erecto-patent, subfalcate lobes. Lobes pectinate, linear-lanceolate, acute, to 2 c.M. long and 5 m.M. broad, dilated at the base. Texture very coriaceous; costae distinct; veins simple or forked, free or those around the sori reticulate. Sori solitary at the base of the lobes or in 1 row on each side of the costulae.

Malacca, Lingga Islands, Borneo, Amboyna.

FAMILY III. GLEICHENIACEAE.



III. GLEICHENIACEAE.

Sori dorsal, consisting of few (2—12) capsules. Capsules commonly decumbent and placed in a whorl, with the receptacle in the centre, sessile or short-stalked, surrounded by a horizontal or subhorizontal, complete ring, bursting vertically. Receptacle punctiform. Indusium wanting.

Rhizome mostly creeping. Fronds rarely unbranched, generally dichotomously forked or pinnate and if pinnate the pinnae or pinnulae dichotomous, frequently bearing axillary buds; the leafy portion of the branches commonly pectinate; veins always free.

Tropical and extra-tropical in the southern hemisphere.

No. 7. **Gleichenia.**

7. GLEICHENIA, *Smith.*

Sori consisting of few, commonly 2—5, rarely 6—12 sessile sporangia, placed mostly in a whorl. No indusium.

Rhizome creeping, rarely erect. Stipes mostly scattered, not articulate to the rhizome. Fronds mostly forked, rarely pinnate, the primary branchings pinnate or once or more times forked again, *or* pinnate and forked. The dichotomous forms are often purveyed with axillary buds, which not rarely grow out to fully developed branches, which give the fronds the appearance of pinnate or 2-pinnate fronds with dichotomous branches of a higher order. The thus formed spuriously pinnate fronds are often to be recognized by the presence of mostly divided stipulae or a mass of densely crowded scales placed in the axils of the principal branchings and wanting in the truly pinnate forms. Ultimate segments (lobes or leaflets) semiorbicular and contiguous or linear and pectinate; veins free, simple or forked, often immersed or obsolete.

Tropical and extra-tropical in the southern hemisphere.

Arrangement of the sections.

§ 1. **PLATYZOMA.** Fronds unbranched, simply pinnate; leaflets roundish or ovate, cucullate, to 2 m.M. long. Hk. Bk., Syn. Fil., tab. I, fig. 1, a—c; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 189, C. (Not in Malaya).

§ 2. **EUGLEICHENIA.** Fronds branched; ultimate segments roundish, 1—3 m.M. long. Diels, l.c., fig. 189, B; Christ, Farnkr. d. Erde, fig. 1068.

§ 3. **MERTENSIA.** Fronds branched; ultimate segments oblong to linear, 1—10 c.M. long. Christ, l.c., fig. 1074, 1077, 1081, 1082.

- a. **Dicranopteris.** Fronds forked; forkings variously branched again.
- b. **Protogleichenia.** Fronds pinnate; pinnae variously branched again.

§ 2. EUGLEICHENIA.

- *. *Leaflets slightly recurved at the edge. Rachises naked or clothed with chaffy pubescence.* (1). *G. circinata.*
- *.*. *Leaflets subhemispherical. Rachises scaly or woolly.* (2). *G. vulcanica.*

* (1) ***G. circinata*, Sw., Hk. Bk., Syn. Fil., 11; *G. semivestita*, Lab., Hk., Spec. Fil., I, tab. II, A; *G. microphylla*, R. Br., Hk., l.c., 3.**

Stipes to \pm 50 c.M. long, slender, wiry, flexuose, naked or \pm villose. Fronds to \pm 1 M. broad, dichotomously branched, the branches and internodations with numerous spreading pinnulae \pm 5 c.M. long. Leaflets roundish, \pm 1 m.M. each way, the edges slightly recurved. Texture firm; under surface glaucous; branches and rachises naked or more or less clothed with chaffy pubescence. Capsules 3—4, superficial.

Malaya; Australia, Tasmania, New Zealand, New Caledonia.

* (2) ***G. vulcanica*, Bl., Rac., Flor. Btz., I, 10; *G. dicarpa*, R. Br., var. *alpina*, Hk. Bk., Syn. Fil., 12; *G. hecistophylla*, A. Cunn., Hk., Spec. Fil., I, tab. II, B.**

Fronds 20—80 c.M. long, dichotomously branched, the branches and internodations with spreading, linear pinnulae \pm 3 c.M. long, $1\frac{1}{2}$ —3 m.M. broad. Leaflets roundish, subhemispherical, \pm as long as broad, the edges much recurved. Texture coriaceous; rachises covered with brown scales or ferruginous, furfuraceous tomentum. Capsules 2—3, concealed within the almost slipper-shaped leaflets.

Malaya.

§ 3. MERTENSIA. a. **Dicranopteris.**

- a.** *Fronds forked; branches pinnate; secondary branches (pinnulae) whether or not branched again, (Diplopterygium, Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 188, A).*
- b.** *Pinnulae not branched.*
- c.** *Pinnulae distant.* (3) *G. Norrisii.*
- c.c.** *Pinnulae close.*
- d.** *Under surface naked or sparingly scaly.* (4) *G. glauca.*
- d.d.** *Under surface densely tomentose with ciliate or fimbriate scales.* (5) *G. arachnoidea*
- b.b.** *Pinnulae dichotomously branched.* (6) *G. Warburgii.*
- a.a.** *Fronds forked; branches dichotomously forked again.*
- 1.** *The ultimate branches only leafy. The forkings whether or not purveyed with a pair of deflexed, opposite, leafy, accessory branches, (Acropterygium & Heteropterygium, Diels, l. c., fig. 188, C—D).*
- (7) *G. linearis.*
- 1.1.** *All the branchings leafy like the ultimate ones, but the 1—3 lower ones not rarely nudate, (Holopterygium, Diels, l. c., fig. 188, B).*
- 2.** *Ultimate segments (lobes or leaflets) horizontally spreading.*
- (8) *G. laevigata.*
- 2.2.** *Ultimate segments (lobes or leaflets) more or less erecto-patent.*
- 3.** *Lobes or leaflets (not the rachises) naked beneath.*
- ***. *Texture coriaceous; under surface glaucous.*
- (9) *G. oceanica.*
- **.** *Texture membranaceous; under surface not glaucous.*
- (10) *G. flabellata.*
- 3.3.** *Lobes or leaflets hairy, scaly or tomentose beneath, at least on the central ribs.*
- 4.** *Rachises hairy, scaly and white-powdery.*
- (11) *G. Koordersii.*
- 4.4.** *Rachises hairy or scaly, but not white-powdery.*
- 5.** *Lobes or leaflets toothed at the apex.* (12) *G. hirta.*
- 5.5.** *Lobes or leaflets not toothed at the apex.*
- 6.** *Rachises long-hairy beneath.*
- (13) *G. hispida.*
- 6.6.** *Rachises densely scaly beneath.* (14) *G. vestita.*
-

* (3) **G. Norrisii**, *Mett.*, Hk. Bk., Syn. Fil., 449; Bedd., Ferns S.Br.I., Suppl., tab. CCCXLVI.

Fronds forked. Branches oblong-lanceolate, 60—100 c.M. long, pinnate. Pinnulae lanceolate, distant, alternate, the lowest stalked, 15—25 c.M. long, cut down to a narrow wing into \pm close, ligulate, spreading, blunt segments, the higher ones growing gradually smaller. Texture coriaceous; surfaces naked, the under one green or glaucous; veins once forked. Sori medial, 12—20 to a segment.

Malacca.

* (4) **G. glauca**, *Hk.*, Christ, Farnkr. d. Erde, 340; Hk., Spec. Fil., I, tab. III, B; *G. longissima*, *Bl.*, Rac., Flor. Btz., I, 10; *G. gigantea*, *Wall.*, Bedd., Ferns Br.I., I, tab. XXX; Hk., l. c., tab. III, A; *Polypodium glaucum*, *Thb.*, (oldest name).

Fronds large, forked. Branches to 6 M. or more long, bearing numerous closely placed pinnulae. Pinnulae as long, 10—25 c.M. long, 2—5 c.M. broad, cut down nearly or quite to the costa into numerous segments. Segments linear-lanceolate or oblong, acute or rounded, pale or more or less glaucous beneath, to 5 m.M. broad. Under surface naked or sparingly covered with small, ciliate scales. Capsules 3—5, often mixed with hairs.

Malaya; Japan, China, Trop. Asia, Australia and Polynesia.

* (5) **G. arachnoidea**, *Mett.*, Rac., Flor. Btz., I, 11; *G. longissima*, *Bl.*, var. *rigida*, *Bl.*, Enum., 251; *G. longissima*, *Bl.*, var. *arachnoides*, *Hk.* Bk., Syn. Fil., 12; *G. glauca*, *Hk.*, var., C. Chr., Ind. Fil.

Like *G. glauca*, *Hk.*, but the stipes and rachises densely tomentose with ferruginous, membranaceous, crisped, ciliate or fimbriate scales leaving the rachises \pm rough when they fall. Under surface densely tomentose with the same scales. Capsules immersed between the tomentum.

Malayan Islands.

* (6) **G. Warburgii**, *Christ*, Ann. Btz., XV, 78.

Fronds forked. Branches flexuose, pinnate, bifurcate at the apex, \pm 50 c.M. long. Pinnulae distant, alternate, spreading, dichotomously forked. The 1—3 lowest branchings nudate, the other ones leafy, with the rachises naked or covered with deciduous, reddish tomentum. Ultimate branches 5—15 c.M. long, 2—4 c.M. broad, acuminate. Lobes or leaflets spreading, 15—30 on each side, linear, 2—5 m.M. broad, free or joined at the base, the apex blunt; higher lobes abbreviated, reduced. Texture \pm herbaceous to semi-

coriaceous; surfaces naked or the under one tomentose on the costulae like the rachises. — The description is taken from specimens gathered in Amboyna by Boerlage, determined by Christ.

Borneo, Celebes, Batjan, Amboyna.

* (7) **G. linearis**, *Clarke*, Trans. Linn. Soc., Bot., 2^d Series, I, 428; *G. dichotoma*, *Hk.*, Rac., Flor. Btz., I, 13; *Bedd.*, Ferns S.I., tab. LXXIV; *G. Hermannii*, *R. Br.*, Bl., Enum., 248; *Polypodium lineare*, *Burm.*, (oldest name); — var. *rigida*, *Rac.*, l. c.; — var. *normalis*, *Rac.*, l. c.; — var. *ferruginea*, *Rac.*, l. c.; *G. ferruginea*, *Bl.*, Enum., 249; — var. *emarginata*, *Christ*, Ann. Btz., XV, 77; — var. *malayana*, *Christ*, l. c.

Fronds large, \pm 3—4 \times dichotomous, each forking bearing a pair of deflexed, unbranched branches at the base on the outer side, except the terminal forkings, which are commonly without accessory lateral branches. Ultimate branches mostly with numerous \pm spreading segments on each side. —

Forma typica: Texture subcoriaceous; surfaces naked, not glaucous. Ultimate segments under 4 c.M. long.

Var. rigida: Texture rigidly coriaceous.

Var. normalis: Under surface glaucous.

Var. ferruginea: Under surface clothed with ferrugineous, furfuraeous tomentum, at least on the costae.

Var. emarginata: Texture very rigid; under surface clothed throughout with ferrugineous hairs.

Var. malayana: Very large; branches much elongated, with the ultimate segments 4—6 c.M. long or longer.

Tropical and extra-tropical.

* (8) **G. laevigata**, *Hk.*, Rac., Flor. Btz., I, 11; *G. bifurcata*, *Bl.*, Enum., 250; ? *G. flagellaris*, *Spr.*, *Christ*, *Farnkr. d. Erde*, 342.

Stipes terete or nearly so, often narrowly marginate. Fronds large, repeatedly dichotomous; the lowest forkings divaricating or rectangularly spreading, not rarely nudate. Ultimate segments linear, to 5 c.M. long, to 2½ m.M. broad, horizontally spreading, obtuse or retuse. Texture \pm herbaceous; upper surface bright-green, under one paler, naked, or the costae cobwebby or tomentose with ciliate, ferrugineous scales. Sporangia \pm 3—4. — *G. flagellaris*, *Spr.*, which has the under surface very glaucous, is probably another species very near our plant, but not malayan.

Malaya.

(9) **G. oceanica**, *Kuhn*, Hk. Bk., Syn. Fil., 449.

Fronds repeatedly dichotomous, above 30 c.M. long. Terminal branches 10—12½ c.M. long, acuminate. Ultimate segments close, to 2½ m.M. broad, acuminate, erecto-patent, subfalcate. Texture subcoriaceous; under surface glaucous; rachises densely clothed with ferrugineous, lanceolate, fimbriate scales.

Celebes; Polynesia.

(10) **G. flabellata**, *R. Br.*, Hk. Bk., Syn. Fil., 12; Hk., Fil. Exot., tab. LXXI.

Stipes firm, polished, dark-brown. Fronds repeatedly dichotomous, flabelliform, to 1 M long, more broad than long. Terminal branches 10—20 c.M. long, 2½—5 c.M. broad, acuminate. Ultimate segments close, linear, acute. Texture membranaceous; surfaces naked, not glaucous beneath. Sporangia 1—6, mostly 4.

New Guinea; Australia, Tasmania, New Zealand, New Caledonia.

(11) **G. Koordersii**, *Christ*, Ann. Bot., XV, 76, tab. XIII, fig., 1, a—b.

Ultimate segments numerous (to \pm 70 on each side of the terminal branches of the fronds), \pm 1½ m.M. broad, linear, falcate, the margin revolute. Texture coriaceous, under surface hairy; rachises covered with deciduous, fibrillose, woolly, reddish-brown hairs besides linear, flaccid, tortuose, ciliate scales and a white powder. Sori consisting of 4—5 sporangia.

Celebes, Luzon.

* (12) **G. hirta**, *Bl.*, Rac., Flor. Botz., I, 12.

Fronds dichotomous, to \pm 1 M. long, the rachises, costae and small axillary buds furnished with copious, ferrugineous, lanceolate to subulate, ciliato-dentate, deciduous scales. Primary branches 3—4 \times forked. Lowest internodations 1½—2½ c.M. long, nudate, the secondary ones 2—4 c.M. long, subnudate. Terminal branches 7½—25 c.M. long, linear, narrowed gradually towards the apex. Ultimate segments ¾—2 c.M. long, 2—4 m.M. broad, linear, obtuse, subfalcate, the margin entire, revolute, the apex denticulate, those of the lower forkings gradually becoming shorter, the lowest segments semidecurrent. Texture membranaceo-chartaceous; under surface caeruleo-pruinose; rachises and costae glaucous.

Malaya.

* (13) **G. hispida**, *Mett.*, *Rac.*, *Flor. Btz.*, I, 12.

Fronds to \pm 50 c.M. long, forked. Forkings 1—5 \times forked again. Ultimate segments erecto-patent, linear, entire, close or \pm distant, to 2 c.M. long, \pm 2 m.M. broad, the edges recurved. Texture coriaceous; stipes, rachises and under surface more or less densely covered with a tomentum of long, ferruginous hairs, the younger portions of the fronds very densely tomentose. — Specimens in the Buitenzorg herbarium have the fronds with the axillary buds copiously developed and without stipulae, as in *Protogleichenia*.

Java; Celebes, Ternate.

* (14) **G. vestita**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 11; — var. *paleacea*, *Bk.* in *Journ. of Bot.*, XVII, 38.

Frons forked, $\frac{1}{2}$ —1 M. long. Forkings 1—3 \times forked again, the lowest internodations 2—5 c.M. long, leafy. Terminal branches 10—25 c.M. long. Ultimate segments approximate, oblong to triangular-oblong, $7\frac{1}{2}$ — $12\frac{1}{2}$ m.M. long, $2\frac{1}{2}$ — $6\frac{1}{2}$ m.M. broad, entire, obtuse, \pm erecto-patent. Texture chartaceous to coriaceous; under surface glaucous; rachises and costae densely covered with yellow or ferruginous, fimbriate scales. Capsules 3—5.

Var. paleacea: Scales of the rachises linear-subulate

Malaya.

§ 3. MERTENSIA. b. **Protogleichenia** (*).

a. *Lowest pinnae forked, the forkings unbranched or once or more times forked again. Texture herbaceous.* (15) *G. Hallieri*.

a.a. *Lowest pinnae pinnate, the pinnulae variously branched. Texture \pm coriaceous.*

b. *Pinnulae dichotomously forked like the apex of the frond.*

(16) *G. amboinensis*.

b.b. *Pinnulae pinnate, the tertiary divisions pseudo-dichotomous like the apex of the frond.* (17) *G. dolosa*.

* (15) **G. Hallieri**, *Christ*, *Ann. Btz.*, XX, 138.

Fronds pinnate, the rachis castaneous, nearly glossy, covered with scattered, subulate, brown hairs. Pinnae flabellate, repeatedly dichotomous, \pm 20 c.M. long, \pm 30 c.M. broad, the internodations \pm 3 c.M. long. Ultimate branches

(*) See also **G. hispida**, *Mett.*

± 15 c.M. long, ± 2 c.M. broad, linear-lanceolate, acuminate. Segments of the last internodes but one, abbreviate, deciduous, those of the ultimate branches pectinate, erecto-patent, not imbricate, linear-lanceolate, ± 1 c.M. long, ± 2 m.M. broad, bluntish, entire. Texture thinly herbaceous, weak; veins inconspicuous, forked; surfaces covered with few, scattered hairs, nearly naked, Sori fugacious.

Borneo.

* (16) **G. amboinensis**, v. A. v. B.

Fronds pinnate, ± 50—100 c.M. long, the main and secondary rachises deciduously tomentose or furfuraceous, at length naked, the apex of the frond 3—4 × dichotomous. Pinnae 2—5-jugate. Lowest pinnae erecto-patent, once pinnate again, the apex dichotomously forked as are the pinnulae. Highest pinnae 2—3 × dichotomous. All the forkings (not the *pinniform* primary branchings) leafy and clothed beneath with 2 opposite rows of spreading, triangular, elongate, ciliate, pale-yellow-brown scales, which are roundish or cordate at the base. Ultimate segments subspreading, linear-oblong, 1—2 c.M. long, ± 2 m.M. broad, the apex bluntish and crenate, in ± 20—50 pairs on the ultimate forkings which are gradually narrowed towards the acuminate, broadly toothed apex. Leaflets of the lower forkings often reduced and ± orbicular. Texture subcoriaceous; under surface of the leaflets deciduously tomentose especially on the midribs, not rarely glaucous.


Amboyna (Teysmann, Boerlage No. 488), *Buru* (Binnendijk No. 153).

(17) **G. dolosa**, C. Chr., Ind. Fil.; *Dicranopteris dolosa*, Copel., Perk., *Fragm. Philipp.*, 193, tab. IV, fig. C.

Rhizome creeping; the scales lanceolate-subulate, squarrose, dirty-brown. Stipes erect, ± 1—2 M. long, terete, firm, black-brown, scaly near the base. Fronds ± 1 M. long, deltoid, pinnate, subdichotomous at the apex. Pinnae 5—5-jugate, the lower ones 2-pinnate again. Highest pinnae and tertiary branches in the lowest part of the frond repeatedly pseudo-dichotomous, the forkings leafy, cut down nearly or quite to the rachis. Ultimate segments 10—12 m.M. long, ± 2 m.M. broad, close, entire, obtuse, 50—80 on each side of the ultimate forkings. Texture coriaceous, surfaces naked or nearly so; under surface bluish-glaucous; rachises clothed with distant, spreading, brown or whitish scales. Capsules fugacious, 5—5 to a sorus.

Philippines.

FAMILY IV. HYMENOPHYLLACEAE.



IV. HYMENOPHYLLACEAE.

Sori marginal, terminal on the veins, exserted or immersed in the substance of the frond. Receptacle erect, oblong, capitate, malleiform, clavate or columnar-filiform, included or exserted. Sporangia dense, imbricate, compressed, surrounded by a broad, horizontal or subhorizontal, complete ring, opening vertically. Indusium inferior, 2-valved or \pm tubular to obconical with the mouth truncate, dilated or \pm 2-lipped, mostly of the same texture as the frond, opening outwardly.

Rhizome frequently creeping. Fronds often small and membranaceous, rarely subcoriaceous; veins free.

Tropical and subtropical, rarely inhabiting cooler climates.

Key to the genera.

*. Indusium more or less deeply 2-valved; receptacle mostly included; surface of the sporangia divided into many, uniform cells.

No. 8. **Hymenophyllum.**

.. Indusium tubular or funnel-shaped, the mouth sometimes 2-lipped; receptacle mostly exserted; surface of the sporangia divided into few, unequal cells.

No. 9. **Trichomanes.**

8. HYMENOPHYLLUM, *Smith.*

Sori marginal, terminal on the veins. Indusium more or less deeply 2-valved, of the same texture as the frond, exserted or more or less immersed in the margin. Receptacle mostly short and included, rarely elongated and prominent beyond the sporangia. Surface of the sporangia divided into many, uniform cells.

Rhizome mostly creeping, often filiform. Fronds generally small, delicately membranaceous, more or less compoundly divided, with mostly linear or linear-oblong ultimate segments. Hk. Bk., Syn. Fil., tab. II, fig. 16; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 74.

Tropical, and extra-tropical in the southern, rarely in the northern hemisphere.

Arrangement of the sections.

§ 1. **EUHYMENOPHYLLUM.** Ultimate segments of the fronds entire.

§ 2. **LEPTOCIONIUM.** Ultimate segments spinuloso-denticulate.

§ 1. **EUHYMENOPHYLLUM.**

a. *Fronds naked.*

b. *Main rachis not winged, or winged towards the apex only.*

c. *Fronds under 5 c.M. long.* (1) *H. discosum.*

c.c. *Fronds more than 5 c.M. long.*

d. *Ultimate segments crisped or complicate.* (2) *H. ooides.*

d.d. *Ultimate segments flattened.*

e. *Pinnae rather distant.* (3) *H. eximium.*

e.e. *Pinnae rather imbricating.*

f. *Ultimate segments emarginate at the apex. Valves of the indusium slightly repand.* (4) *H. modestum.*

f.f. *Ultimate segments entire as are the valves of the indusium.* (5) *H. flabellatum.*

b.b. *Main rachis winged throughout.*

1. *Valves of the indusium more or less toothed at the outer edge.*

§. *Edges crisped.* (6) *H. australe.*

§.§. *Edges not crisped.*

***.** *Main rachis slightly winged.* (7) *H. demissum.*

(8) *H. subdemissum.*

***.*.** *Main rachis broadly winged.* (9) *H. badium.*

1.1. *Valves of the indusium entire, or rarely crenate at the apex.*

2. *Fronds mostly 2—5 c.M. long.*

‡. *Sori numerous on the higher pinnae. Receptacle included.*

(10) *H. paniculiflorum.*

‡.‡. *Sori solitary on the higher pinnae. Receptacle exerted.*

(11) *H. ringens.*

2.2. *Fully developed fronds mostly 5—15 c.M. long.*

3. *Edges crisped.* (12) *H. thuidium.*

3.3. *Edges not crisped.*

4. *Stipes winged ± throughout.* (15) *H. salakense.*

4.4. *Stipes not winged, or winged towards the apex only.*

5. *Indusium obovate, narrowed at the apex.*

(14) A. Blumeanum.

5.5. *Indusium oblong, the apex bluntly rounded or subtruncate.*

6. *Pinnae ± 4—5 on each side. Indusium exerted, deeply 2-valved.* (15) H. Treubii.

6.6. *Pinnae ± 6—8 on each side. Indusium immersed, cut $\frac{1}{2}$ -way down.* (16) H. rarum.

2.2.2. *Fully developed fronds more than 15 c.M. long.*

‡. *Ultimate segments 1—1 $\frac{1}{2}$ m.M. broad. Receptacle globosocapitate.* (17) H. formosum.

‡.‡. *Ultimate segments 1 $\frac{1}{2}$ —2 m.M. broad. Receptacle malleiform.* (18) H. Junghuhnii.

a.a. *Fronds hairy or ciliate, at least on the central veins.*

b. *Central ribs of the segments purveyed with 2 wing-like, crisped crests.*

‡. *Ribs slightly crested. Fronds rather long.* (19) H. Zollingerianum.

‡.‡. *Ribs broadly crested. Fronds rather short.* (20) H. fuscum.

b'.b'. *Central ribs not crested.*

c'. *Fronds under 5 c.M. long, ± 1 c.M. broad.* (21) H. borneense.

c'.c'. *Fully developed fronds mostly 5—10 c.M. long, 1—5 c.M. broad.*

d'. *Fronds under 2 c.M. broad. Indusium free, cleft nearly to the base.* (22) H. pedicularifolium.

d'.d'. *Fronds more than 2 c.M. broad. Indusium rather slightly immersed, cleft ± $\frac{1}{2}$ -way down.* (23) H. obtusum.

c'.c'.c'. *Fully developed fronds more than 10 c.M. long, 10—15 c.M. broad.* (24) H. pachydermicum.

(1) **H. discosum**, *Christ*, Bull. Herb. Boiss., VI, 140.

Stipes filiform. Fronds ± 2 $\frac{1}{2}$ c.M. long, ± 1 $\frac{1}{2}$ c.M. broad, broadly ovate; rachis not winged. Pinnae alternate, nearly flabellately cut down to the rachis into oblong segments 4 m.M. long, which are deeply forked. Surfaces naked; colour fulvous. Sori terminal on the segments, numerous, large, round; indusium with the valves entire; receptacle short, not exerted.

Philippines.

(2) **H. ooides**, *Muell. & Bk.*, Journ. of Bot., XXVIII, 105.

Stipes short, filiform, naked. Fronds lanceolate, 2—5-pinnate, pendulous,

naked, 10—15 c.M. long, to $2\frac{1}{2}$ c.M. broad; rachis filiform. Pinnae very numerous, lanceolate, erecto-patent, the central ones the largest, usually simply pinnate, rarely 2-pinnate. Ultimate divisions obovate, obtuse, 1-nerved, emarginate, crowded, ± 2 m.M. long, more or less crisped or complicate. Sori small, terminal on the ultimate divisions; indusium immersed at the base, the valves cuneate, with the outer edge rounded.

New Guinea.

* (5) **H. eximium**, *Hæ*, V. D. B., Hymen. Jav., 57, tab. XLVI; *H. dilatatum*, *Sw.*, Rac., Flor. Btz., I, 17, p.p.; C. Chr., Ind. Fil.

Rhizome creeping, naked. Stipes 7—10 c.M. long, firm, terete, not winged. Fronds ± 10 —15 c.M. long, 5—8 c.M. broad, ovate or deltoid-oblong; main rachis narrowly winged towards the apex, terete, angular or marginate in the lower part. Pinnae rather distant, ovate, acuminate to caudate, cut down to a winged rachis, short-stalked. Pinnulae erecto-patent, cut down to a winged rachis into linear, simple, forked or branched segments 2—10 m.M. long, ± 1 — $1\frac{1}{2}$ m.M. broad. Sori small, terminal on short segments, 2—12 to a pinna; receptacle clavate-oblong; indusium deeply 2-valved, the valves obovate, with the outer edge minutely and obscurely eroso-denticulate.

Java, Sumatra.

(4) **H. modestum**, *V. D. B.*, Ned. Kruidk. Arch., V³, 165.

Rhizome creeping. Stipes terete, flexuose, 5—7 c.M. long. Fronds ± 10 c.M. long, 4—5 c.M. broad, oblong, narrowed at the base, pinnate, the apex pinnatifid. Pinnae oblong, broadly cuneate at the base, spreading or divaricating, the lowest remote, the rest contiguous, imbricate, slightly decurrent on the rachis, sub-2-pinnatifid. Segments obtuse, contiguous, erecto-patent, the lower ones pinnatifid, the other ones 1—3-furcate; lacinulae short, nearly straight, slightly undulate, emarginate at the apex. Texture membranaceous; colour olivaceo-fuscous, subopaque; main rachis flexuose; veins straight, slender. Sori terminal on the lacinulae; indusium oblong, conical at the base, 2-valved $\pm \frac{3}{4}$ -way down, the valves narrowly rounded, slightly repand; receptacle cylindrical.

Philippines.

* (5) **H. flabellatum**, *Lab.*, Hk. Bk., Syn. Fil., 61; *H. nitens*, *R. Br.*, Hk. & Grev., Ic. Fil., tab. CXCVII.

Rhizome creeping. Stipes 5—15 c.M. long, firm, erect. Fronds 10—30

c.M. long, $2\frac{1}{2}$ —10 c.M. broad, ovate, acuminate; main rachis winged above. Pinnae rather imbricating; the lower ones distant, broadly rhomboidal-acuminate, subflabellate, erecto-patent or subspreading, divided down to a broadly winged rachis into several cuneate-based pinnulae which are forked or pinnatifid with several simple or forked to pinnatifid segments on each side. Texture flaccid; surfaces somewhat glossy. Sori 6—20 to a pinna, terminal on the segments; indusium divided more than $\frac{1}{2}$ -way down, the valves entire, rounded, the base cuneate; receptacle columnar-oblong. — Ultimate segments not emarginate at the apex, entire.

? *Sumatra*, ? *New Guinea*, *Batjan*; *Australia*, *New Zealand*, *Samoa*.

* (6) **H. australe**, *Willd.*, *Christ*, *Farnkr. d. Erde*, 17; *Diels*, in *Engl. & Prantl*, *Nat. Pfl.Fam.*, I⁴, fig. 74, C; *H. javanicum*, *Spr.*, *Rac.*, *Flor. Btz.*, I, 17; *V. D. B.*, *Hymen. Jav.*, tab. XL; *H. micranthum*, *V. D. B.*, *H. erosum*, *Bl.*, *H. fimbriatum*, *J. Sm.*, *V. D. B.*, l. c., tab. XLI, XLIII & XLIV; — var. *atrovirens*; *H. atrovirens*, *Colen.*, *Tasm. Journ.*, II, 186; *H. tasmanicum*, *V. D. B.*, *Ned. Kruidk. Arch.*, IV, 599.

Rhizome wide-creeping, slender, naked or nearly so. Stipes 5—7 c.M. long, erect, margined with a broad, crisped wing, except at the base. Fronds 5—20 c.M. long, 5—10 c.M. broad, ovate to deltoid, 3—4-pinnatifid, main & secondary rachises provided with a \pm broad wing. Lower pinnae oblong or deltoid-obovate, erecto-patent. Pinnulae oblong, cut down to a crisped wing into simple or bifid segments, which are naked, entire, crisped, \pm 1 m.M. broad. Sori \pm 6—20 to a pinna, terminal on the ultimate segments; indusium to 1 m.M. broad, divided down nearly to the base, the valves roundish or oblong, with the outer edge denticulate; receptacle \pm filiform, included.

Var. atrovirens: Segments nearly plane.

Malaya; *North. India*, *Australia*, *New Zealand*, *Antarct. Isles*.

* (7) **H. demissum**, *Sw.*, *Hk. Bk.*, *Syn. Fil.*, 61; *H. productum*, *Kze.*, *Rac.*, *Flor. Btz.*, I, 17; *V. D. B.*, *Hymen. Jav.*, tab. XLV.

Rhizome filiform, creeping, naked. Stipes 5—15 c.M. long, erect, firm, wingless or narrowly winged above. Fronds ovate-triangular to elongate, 6—30 c.M. long, $5\frac{1}{2}$ —10 c.M. broad, 3- or 4-pinnatifid; main rachis slightly winged. Lower pinnae $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, triangular-rhomboidal, divided down very nearly to the rachis into numerous pinnulae on each side, which are again divided down nearly to the rachis into pinnatifid segments. Ultimate lobes $2\frac{1}{2}$ —5 m.M. long, \pm $1\frac{1}{2}$ m.M. broad. Sori very numerous,

20—50 to a pinna, on both sides and terminal on the segments; indusium divided down nearly to the base, the valves ovate to lanceolate, the outer edge denticulate (in malayan material).

Malaya; Polynesia, New Zealand.

(8) **H. subdemissum**, *Christ*, Bull. Herb. Boiss., VI, 140.

Like *H. demissum*, *Sw.* but the dimensions $\frac{1}{2} \times$ as long; stipes and rachises tender, flexuose, filiform; the segments short, narrowed, less numerous, distant: the valves of the indusium minute.

Philippines.

(9) **H. badium**, *Hk. & Grev.*, Hk. Bk., Syn Fil., 60; Hk. & Grev., Ic. Fil., tab. LXXVI; Bedd., Ferns Br.I., II, tab. CCLXXXII; *H. javanicum*, *Spr.*, var. *badium.*, Bedd., Handb. Ind. Ferns, 53.

Stipes 5—7 $\frac{1}{2}$ c.M. long, slender, winged above. Fronds 10—30 c.M. long, 5—7 $\frac{1}{2}$ c.M. broad, oblong or ovate-lanceolate, acuminate, 3-pinnatifid; main rachis broadly winged. Lower pinnae lanceolate-rhomboidal, divided down to a broadly winged rachis into several pinnules on each side, the lowest of which are a few times deeply incised. Ultimate divisions broadly linear. Sori 2—12 to a pinna, on both sides and terminal on the segments; indusium nearly 2 $\frac{1}{2}$ m.M. broad, divided down nearly to the base, the valves rounded, nearly 2 \times as broad as long, the outer edge toothed; receptacle very short.

Malacca; Himalaya.

* (10) **H. paniculiflorum**, *Pr.*, Rac., Flor. Btz., I, 16; V. D. B., Hymen. Jav., tab. XXXIX.

Rhizome filiform, sparingly hairy. Stipes naked, slender, 1—5 c.M. long, naked, winged above. Fronds ovate to lanceolate-oblong, 2—5 c.M. long, 1—2 $\frac{1}{2}$ c.M. broad, cut down to a narrowly winged rachis. Pinnae spreading or erecto-patent, ovate or suborbicular, deeply pinnatifid. Pinnulae simple, forked or pinnatifid again. Ultimate segments to 5 m.M. long, $\pm \frac{1}{2}$ m.M. broad. Colour reddish or brownish. Sori numerous, confined to the uppermost pinnae, very close as to form a panicle; indusium large, reddish, elliptical or broadly obovate, ± 2 m.M. long, ± 1 m.M. broad, joined in the lower half, the valves entire; receptacle short-columnar, included.

Malaya; Japan.

(11) **H. ringens** *Christ*, Ann. Btz., XIX, 54.

Rhizome filiform. Stipes slender, $\pm 1\frac{1}{2}$ c.M. long. Fronds ovate-elongate,

\pm 3 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, cut down to a winged rachis into 4—7 segments on each side, the base narrowed and decurrent. Segments ovate, obtuse, deeply incised. Lobes linear, obtuse, entire, 2—4 on each side, \pm 1 m.M. broad. Texture firm; colour dark-brown; both surfaces nearly naked. Sori solitary on the superior primary segments, exserted, often short-stalked; indusium campanulate, deeply 2-valved; valves triangular, entire, acute, at length reflexed; receptacle 3 m.M. long, broad, exserted.

Celebes.

(12) **H. thuidium**, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 25.

Stipes slender, $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, winged. Fronds 5— $12\frac{1}{2}$ c.M. long, $2\frac{1}{2}$ —5 c.M. broad, ovate to oblong, 3—4-pinnate, erect; rachis and branches winged like the stipe. Pinnae to \pm $2\frac{1}{2}$ c.M. long, deltoid; wings and ultimate divisions finely crisped throughout. Surfaces naked. Sori few to many in the upper part of the frond, on the ends of the ultimate divisions, large; indusium divided nearly to the base; valves large, ovate, crisped.

Philippines.

* (15) **H. salakense**, *Rac.*, Flor. Btz., I, 18.

Rhizome wide-creeping, slender, sparingly pilose. Stipes distant, 4—8 c.M. long, winged throughout or nearly so. Fronds lanceolate or obovate, 5—15 c.M. long, 3—10 c.M. broad, sometimes subdeltoid; main rachis broadly winged. Pinnae erecto-patent, cut down to a broadly winged rachis into simple, forked or 2 \times forked pinnulae. Ultimate segments 5—15 m.M. long, 2—3 m.M. broad, linear or oblong, bluntly rounded or emarginate at the apex. Surfaces naked; a single central rib in each segment. Sori terminal on the segments; indusium deeply 2-valved; valves roundish or transversely oblong, entire, broader than the segments, to $5\frac{1}{2}$ m.M. broad; receptacle malleiform.

Java.

* (14) **H. Blumeanum**, *Spr.*, *Rac.*, Flor. Btz., I, 16; V. D. B., Hymen. Jav., tab. XXXVI; Bedd., Ferns S.I., tab. CCLXVI; *H. pycnocarpum* & *integrum*, V. D. B., l. c., tab. XXXVII—XXXVIII; *H. polyanthos*, *Sw.*, Hk. Bk., Syn Fil., 60, p. p.

Rhizome filiform, creeping. Stipes slender, 1—4 c.M. long, not rarely marginate, not winged, or winged in the upper half only. Fronds obovate to lanceolate or even linear, 5—15 c.M. long, \pm 1—7 c.M. broad; main rachis narrowly winged. Central pinnae the longest, oblong or lanceolate, deltoid or rhom-

boidal, cut down to a winged rachis into simple or once or more times divided segments on each side. Ultimate lobes \pm 2—4 m.M. long, $\frac{1}{2}$ —1 m.M. broad, entire, each with a single central rib. Surfaces naked. Sori on the higher pinnae, terminal on narrowed segments; indusium obovate, the base rounded, the valves entire, narrowed towards the apex; receptacle linear, not exserted.

Trop. Asia.

* (15) **H. Treubii**, *Rac.*, Flor. Btz., I, 15; Nat. Tijdschr. Ned. Ind., LIX, tab. III.

Rhizome slender, wide-creeping, sparingly hairy. Stipes filiform, distant, 2—3 c.M. long, slightly winged towards the frond. Fronds \pm ovate in general outline, 5—8 c.M. long, 5—5 c.M. broad; main rachis flexuose, narrowly winged. Pinnae erecto-patent, to 4—5 on each side, cut down to a winged rachis into \pm 5—5 simple, forked or pinnatifid pinnulae. Ultimate segments oblong or linear, 2—15 m.M. long, \pm 2 m.M. broad, the apex often emarginate. Surfaces naked; a single central rib in each segment. Sori terminal on short segments; indusium deeply 2-valved; valves suboblong, the apex rounded or subtruncate, entire, or slightly and irregularly erose; receptacle erect-oblong.

Java.

(16) **H. rarum**, *R. Br.*, Hk. Bk., Syn. Fil., 58; Kze, Farnkr., I, tab. XCVI, fig. 1; *H. semibivalve*, *Hk. & Grev.*, Ic. Fil., tab. LXXXIII.

Stipes very slender, $2\frac{1}{2}$ —8 c.M. long. Fronds flaccid, pendent, 5—15 c.M. long, $2\frac{1}{2}$ —5 c.M. broad, linear-oblong, 2-pinnatifid, the main rachis winged throughout. Pinnae simple, linear, or forked to pinnatifid with 2—3 deeply cut segments on each side, which are $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long, \pm $2\frac{1}{2}$ m.M. broad. Sori large, terminal on the segments of the upper pinnae, often $2\frac{1}{2}$ m.M. broad; indusium divided $\frac{1}{2}$ -way down or more, the base cuneate, the valves rounded; receptacle erect-oblong.

? *Malacca; Tasmania, New Zealand.*

* (17) **H. formosum**, *Brack.*, V. D. B., Hymen. Jav., 59, tab. XLVII—XLVIII; *H. dilatatum*, *Sw.*, *Rac.*, Flor. Btz., I, 17, p. p.; C. Chr., Ind. Fil.

Rhizome creeping, hairy. Stipes 5—10 c.M. long, mostly narrowly winged. Fronds 15—30 c.M. long, or to 40 c.M. long, \pm 5—10 c.M. broad; main rachis winged throughout. Pinnae erecto-patent, cut down to a winged rachis, short-stalked. Pinnulae erecto-patent, deeply pinnatifid again into linear, simple,

forked or branched ultimate segments 2—10 m.M. long, $\pm 1-1\frac{1}{2}$ m.M. broad. Sori terminal on short lobes, 2—12 to a pinna; receptacle globosocapitate; indusium deeply 2-valved, the valves orbicular or oblong-obovate, entire or slightly emarginate at the apex.

Malaya; Polynesia, Trop. Australia, New Zealand.

* (18) **H. Junghuhnii**, *V. D. B.*, Rac., Flor. Btz., I, 18; V. D. B., Hymen. Jav., tab. XLIX; *H. dilatatum*, *Bl.*, Enum., 221.

Rhizome wide-creeping, sparingly hairy. Stipes 5—12 c.M. long, furnished at the base with few, short, brown, spreading, deciduous hairs, narrowly and shortly winged above. Fronds 3—4-pinnatifid, linear-obovate, 15—25 c.M. long, sometimes shorter, 4—10 c.M. broad; rachises more or less broadly winged. Pinnae erecto-patent, to ± 10 c.M. long and to ± 2 c.M. broad. Pinnulae erecto-patent, to $\pm 1\frac{1}{2}$ c.M. long, to ± 5 m.M. broad, cut down into 1—3 simple or bifid, erecto-patent ultimate segments 3—12 m.M. long, $1\frac{1}{2}$ —2 m.M. broad. Sori terminal on lateral segments; indusium roundish, transversely oblong or reniform, $1\frac{1}{2}$ — $2\frac{1}{2}$ m.M. long, 2—5 m.M. broad, the outer edge rounded, entire; receptacle malleiform.

Sumatra, Java to Polynesia.

* (19) **H. Zollingerianum**, *Kze*, Hk. Bk., Syn. Fil., 62; V. D. B., Hym. Jav., tab. L.; (*H. fuscum*, *V. D. B.*, Rac., Flor. Btz., I, 19, p. p.).

Rhizome creeping, \pm hairy. Stipes 5—10 c.M. long, deciduously hairy. Fronds lanceolate and 10—20 c.M. long or linear and 20—40 c.M. long, acuminate. Pinnae rather distant, $1-2\frac{1}{2}$ c.M. long, 7—15 m.M. broad, the lower ones triangular or oblong, acute or rounded at the apex, divided more than $\frac{1}{2}$ -way down into oblong, erecto-patent, deeply pinnatifid segments. Ultimate lobes linear-oblong, ± 1 m.M. broad. Rachis narrowly winged, often slightly hairy; a single central vein in each ultimate segment; veins \pm hairy and slightly crested like the rachises. Sori 2—8 to a pinna, large, terminal on lateral segments; indusium obovate, divided $\pm \frac{1}{2}$ -way down, the mouth rounded, entire; receptacle mostly short.— An abnormal form has the upper pinnae much elongated.

Java.

* (20) **H. fuscum**, *V. D. B.*, Rac., Flor. Btz., I, 19, p. p.; V. D. B., Hymen. Jav., 62, tab. LI—LII.

Rhizome filiform, creeping, hairy. Stipes 3—7 c.M. long, hairy.

Fronds deltoid-obovate or triangular-lanceolate, with a broadly winged rachis, 6—16 c.M. long, 4—5 c.M. broad, 3-pinnatifid. Pinnae moderately close, 1—2 c.M. broad, lowest largest, with a winged rachis, ovate-oblong in outline. Pinnulae close or imbricating, pinnatifid. Segments simple or forked, with a single central vein in each ultimate segment. Veins hairy and purveyed with long and broad, undulate and crisped, entire crests like the rachises. Sori terminal on lateral segments, 1 or more to a pinna; indusium obovate, divided about $\frac{1}{2}$ -way down; valves entire; receptacle exerted, filiform.

Malaya.

(21) **H. borneense**, *Hk.*, Hk. Bk., Syn. Fil., 62.

Stipes $\pm 2\frac{1}{2}$ c.M. long, very slender. Fronds square or fan-shaped in general outline, ± 1 —2 c.M. long, ± 1 c.M. broad, divided down like a fan nearly to the base, the divisions linear, dichotomously branched. Ultimate segments $2\frac{1}{2}$ — $7\frac{1}{2}$ m.M. long, ± 1 m.M. broad; the surface hairy and the margin ciliate with branched, rufous hairs. Sori terminal, solitary, orbicular, the cuneate base sunk in the frond; valves ciliate.

Borneo.

(22) **H. pedicularifolium**, *Ces.*, Fel. Becc. Polin., 2 & 6.

Fronds sessile, 5—10 c.M. long, 1 — $1\frac{1}{2}$ c.M. broad, pendulous, oblong in general outline, pinnatifid to within a short distance of the costa. Lower lobes rhomboidal-cuneate, cleft on both sides, or on one side only; upper lobes simple, linear-oblong, the margin undulate or crisped, the costulae beneath covered with reddish hairs. Sori terminal on the segments of the lobes; indusium free, cleft nearly to the base.

New Guinea.

* (23) **H. obtusum**, *Hk. & Arn.*, Hk. Bk., Syn. Fil., 65; Hk., Spec. Fil., I, tab. XXXIII, D.

Stipes filiform, to 5 c.M. long, ciliate. Fronds 5—10 c.M. long, $2\frac{1}{2}$ —5 c.M. broad, oblong or lanceolate-oblong; main rachis winged troughout, but sometimes nudate in the lower part. Pinnae lanceolate, oblong or cuneate-oblong, deeply pinnatifid into few or several simple, or forked segments on each side. Ultimate lobes linear or oblong, bluntly rounded. Texture very thin; margin and central ribs as well as the rachises densely clothed with stalked, stellately branched hairs. Sori 2—6 to a pinna, terminal on the

lateral segments; indusium as broad as the segments, divided \pm $\frac{1}{2}$ -way down; valves rounded, strongly ciliated, the base \pm cuneate, immersed.

Borneo, Philippines, New Guinea; Hawaii.

(24) **H. pachydermicum**, *Ces.*, Fel. di Bor., 8.

Rhizome slender, intricate, covered with brown bristles. Stipes slender, \pm setulose, 1—5 c.M. long. Barren fronds short, subflabellate, as broad as long; fertile ones oblong or lanceolate, \pm 20 c.M. long, 10—15 c.M. broad. Primary segments obtuse or truncate, obovato-linear. Lobes 3—5 to a segment, densely barbate at the apex in the barren fronds. Under surface covered with short, brown hairs, especially on the veins. Sori immersed; valves of the indusium ovate, acute; receptacle filiform, exserted.

Borneo.

§ 2. LEPTOCIONIUM.

a. *Fronds 3-partite, subflabellate.* (25) **H. subflabellatum.**

a.a. *Fronds pinnately divided.*

b. *Edges not crisped.*

c. *Main rachis narrowly marginate.* (26) **H. edulentum.**

c.c. *Main rachis distinctly winged throughout or nearly so.*

d. *Sori terminal on the main or higher secondary rachises.*

e. *Valves of the indusium hardly toothed.* (27) **H. Kurzii.**

e.e. *Valves of the indusium subulate-dentate.* (28) **H. Lobbii.**

d.d. *Sori axillary on the anterior side of the pinnae.*

*. *Indusium without crests at the base.*

(29) **H. tunbridgense.**

.. *Indusium surrounded at the base by a row of longitudinal crests.*

(30) **H. holochilum.**

c.c.c. *Main rachis not winged, or winged in the upper half only.*

d'.d'. *Valves of the indusium toothed.* (31) **H. blandum.**

d'.d'.d'. *Valves of the indusium entire.*

e. *Fronds under 5 c.M. long.* (32) **H. praetervisum.**

e'.e'. *Fully developed fronds 5—20 c.M. long.*

f'.f'. *Veins paleaceo-hirsute.* (33) **H. violaceum.**

f'.f'.f'. *Veins naked.*

g'.g'. *Indusium short-2-lipped, the lips rounded.*

(34) **H. multifidum.**

g.g. *Indusium divided $\frac{1}{2}$ -way down, the lips narrowed at the apex.* (35) *H. serrulatum.*

b.b. *Edges crisped.*

1. *Indusium quite naked on the back.*

2. *Fronds under 8 c.M. long; valves of the indusium elliptical.*
(36) *H. fraternum.*

2.2. *Fronds more than 8 c.M. long; valves of the indusium \pm orbicular.*

3. *Edges bluntly dentate, the teeth not deep.*
(37) *H. Reinwardtii.*

3.3. *Edges aculeate-dentate, the teeth deep.*
(38) *H. physocarpum.*

1.1. *Indusium spinulose on the back.*

*. *Edges of the segments deeply toothed nearly to the midribs.*
(39) *H. sabinifolium.*

.. *Edges not deeply toothed.* (40) *H. Neesii.*

1.1.1. *Indusium surrounded by irregularly spinoso-dentate, transversal crests at the base.*

2. *Valves of the indusium serrate or dentate.*

3. *Fronds 5—8 c.M. long. Indusium divided $\pm \frac{2}{3}$ -way down.* (41) *H. denticulatum.*

3.3. *Fronds $2\frac{1}{2}$ —5 c.M. long. Indusium divided $\pm \frac{1}{3}$ -way down.* (42) *H. aculeatum.*

2.2. *Valves of the indusium entire.* (45) *H. bismarckianum.*

1.1.1.1. *Indusium surrounded by a row of rather close, longitudinal crests at the base.* (44) *H. brachyglossum.*

(25) **H. subflabellatum**, *Ces.*, Fel. di Bor., 8.

Rhizome creeping, branched, hairy. Stipes slender, 5—10 m.M. long, not winged. Fronds 5—10 m.M. long, 3-partite, nearly flabellate. Primary segments \pm pinnatifid. Lobes linear, obtuse, distant, spinuloso-denticulate. Sori terminal on the lobes; indusium ovate, with the edges of the valves spinuloso-ciliate; receptacle exserted.

Borneo.

(26) **H. edulentum**, *C. Chr.*, Ind. Fil.; *Leptocionium edulentum*, *V. D. B.*, Ned. Kruidk. Arch., V³, 148.

Rhizome creeping, nearly naked, filiform. Stipes filiform, terete, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long. Fronds 4—5 c.M. long, 2 — $2\frac{1}{2}$ c.M. broad, ovate or ovate-oblong, broadly rounded at the base, narrowed towards the apex, the main rachis narrowly marginate, subterete below. Pinnæ spreading, (the lower divaricating), subcontiguous, suboblong, the apex adscendent. Pinnulae spreading, subremote, simple or dichotomous, the lowest pinnatifid. Lacinulae linear, elongate, approximate, flat, the apex entire, the margin subentire or denticulate, the teeth small, remote. Texture membranaceous, diaphanous, firm; colour olivaceo-fuscous; main rachis flexuose; veins straight, blackish-brown. Sori on the lacinulae; indusium moderately immersed, elliptical, turgid, 2-valved $\pm \frac{1}{2}$ -way down, the valves entire or obscurely denticulate; receptacle setiform.

Borneo; Assam.

* (27) **H. Kurzii**, *Prantl*, *Morph. d. Gefässcr.*, I, *Hymenoph.*, 54.

Main rachis narrowly winged, the wing entire, not toothed. Ultimate segments toothed. Sori terminal on the main and higher secondary rachises, the valves of the indusium hardly toothed. — If this plant is the same as lately found by the Ladies von Grävenitz and Stein, the description will be: Rhizome creeping, filiform, sparingly hairy. Stipes distant, more than ± 1 c.M. apart, sparingly hairy, 1—2 c.M. long, narrowly winged (rather marginate) towards the frond. Fronds lanceolate-oblong to subdeltoid-obovate, 4—5 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad; main rachis narrowly winged, the wing entire, not toothed. Pinnæ cut down to a winged, not toothed rachis into linear, simple or forked segments, which are blunt at the apex, toothed at the margin. Texture thin, hyaline; under surface covered with scattered, spine-like hairs on the costae. Sori terminal on the main and higher secondary rachises and some anterior segments of the higher pinnulae; indusium oblong, the base conical, covered with spine-like hairs on the under side, the valves subacute, slightly toothed at the point; receptacle linear, slightly exserted.

Java.

(28) **H. Lobbii**, *Moore*, *V. D. B., Ned. Kruidk. Arch.*, V³, 176.

Rhizome filiform, branched, hirsute with deciduous, yellowish hairs. Stipes filiform, terete, flexuose, 10—15 m.M. long. Fronds 3—4 c.M. long, 6—8 m.M. broad; main rachis narrowly winged by the decurrent pinnæ, filiform, flexuose. Pinnæ narrow, spreading, slightly incurved, contiguous, obtuse, either 1—2-furcate, or pinnatifid with a single lacinula on each

side. Lacinulae subfastigate, narrow, linear, flat, the margin serrate, the teeth remote, subulate, elongate. Texture membranaceous, rather firm; colour reddish-brown, subopaque. Sori small, terminal on the frond; indusium tubular, compressed, 2-fid \pm to the base, the valves truncate-rotundate and unequally subulate-dentate at the outer edge; receptacle setiform, thick.

Malaya; Assam.

(29) **H. tunbridgense**, *Sm.*, Hk. Bk., Syn. Fil., 67; Hk., Br. Ferns, tab. XLIII; Bedd., Ferns S.I., tab. CCLXV; Bau. & Hk., Gen. Fil., tab. XXXII.

Stipes 1—4 c.M. long, winged in the upper part. Fronds oblong-lanceolate, $2\frac{1}{2}$ —8 c.M. long, 1— $2\frac{1}{2}$ c.M. broad; main rachis winged throughout. Pinnæ distichous, subflabellato-pinnatifid, the lobes $2\frac{1}{2}$ — $7\frac{1}{2}$ m.M. long, and (as well as the usually, solitary, axillary, suborbicular indusia) spinuloso-serrate.

In the temperate climates and the cool, alpine regions of the tropics.

* (50) **H. holochilum**, *C. Chr.*, Ind. Fil.; *H. affine*, *Rac.* (not *V. D. B.*), Flor. Btz., I, 20; *H. Smithii*, *Hk.*, Hk. Bk., Syn. Fil., 69, p. p.; *Leptocionium holochilum & affine*, *V. D. B.*, Hymen. Jav., tab. XXXIV—XXXV.

Rhizome slender, filiform, sparingly hairy. Stipes filiform, 2—5 c.M. long. Fronds oblong to lanceolate-oblong, 4—10 c.M. long, $1\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad; main rachis winged throughout or in the upper half only. Pinnæ cut down to a winged rachis into 1—4 simple or forked pinnulae. Ultimate segments \pm 2 m.M. broad, the margin toothed. Sori 1—4 to a pinna, terminal on short lobes, occupying the anterior side of the higher pinnæ; indusium obovate-oblong, the base \pm obconical, surrounded by a row of longitudinal crests, the valves blunt, more or less distinctly denticulate, or quite entire.

Java.

* (31) **H. blandum**, *Rac.*, Flor. Btz., I, 20.

Rhizome filiform, sparingly hairy. Stipes $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, filiform, very slender. Fronds irregularly obovate, 2—4 c.M. long, 1—2 c.M. broad, simply pinnate or 2-pinnatifid, the rachis at best winged above. Lower pinnæ short-stalked, often forked or pinnatifid, the higher ones commonly undivided, sessile with a narrowed base. Ultimate segments from 1—10 m.M. long. Sori terminal on the segments; indusium oblong, the valves with the apex blunt and toothed.

Java.

(52) **H. praetervisum**, *Christ*, Farnkr. d. Erde, 21, fig. 26; *Trichomanes denticulatum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 82.

Rhizome slender, creeping. Stipes slender, naked, \pm 5—12 m.M. long. Fronds $2\frac{1}{2}$ —4 c.M. long, to $1\frac{1}{2}$ c.M. broad, ovate or oblong, 2-pinnatifid; main rachis winged above, nudate below. Pinnae pinnatifid down to a narrowly winged rachis. Ultimate segments linear, serrate, about 5 m.M. long, 1—2 m.M. broad; a single costa only in each segment. Sori 1 to a pinna, terminal on the lowest segments on the upperside; tube exerted, the mouth deeply 2-valved, the valves entire.

Borneo.

(53) **H. violaceum**, *Meyen*, MS. in *Herb. Berol.*; *Leptocionium violaceum*, *V. D. B.*, *Ned. Kruidk. Arch.*, V³, 147.

Stipes flexuose, terete, hirsute, 6—8 c.M. long. Fronds 8—10 c.M. long, $4-4\frac{1}{2}$ c.M. broad, oblong, acuminate. Pinnae spreading, oblong. Pinnulae erecto-patent, broad-oblong or obcuneate, rather remote or subdistant. Lacinulae straight, parallel, elongate, the margin minutely and irregularly denticulate. Texture firm; colour red-brownish, opaque; main rachis broadly winged above, the wing entire, flat; veins flexuose, all paleaceo-hirsute. Sori axillary, at length turned back on the frond (resupinate); indusium immersed, 2-valved $\frac{2}{3}$ — $\frac{3}{4}$ -way down, the tube cuneate, the valves broadly oblong, entire.

Philippines.

* (54) **H. multifidum**, *Sw.*, *Hk. Bk.*, *Syn. Fil.*, 69; *Hk. & Grev.*, *Ic. Fil.*, tab. CLXVII; *H. klabatense*, *Christ*, *Fil. Saras.*, I, 4; *H. polyanthos*, *Sw.*, *var. dentatum*, *Ces.*, *Fel. Becc. Polin.*, 2.

Stipes 5—10 c.M. long, wiry, naked. Fronds broadly ovate-lanceolate, 5—20 c.M. long, $2\frac{1}{2}$ — $12\frac{1}{2}$ c.M. broad; main rachis naked or scaly, winged above. Pinnae and pinnulae rhomboidal-lanceolate, the rachises winged, \pm wavy. Ultimate segments linear, to $7\frac{1}{2}$ m.M. long, to 1 m.M. broad, conspicuously dentate with short or less short teeth. Sori 2—12 to a pinna, terminal on the ultimate segments; indusium oblong, tubular below, the valves short, entire; receptacle sometimes exerted.

Philippines and Celebes to Polynesia; New Zealand, New South Wales.

(55) **H. serrulatum**, *C. Chr.*, *Ind. Fil.*; *H. Smithii*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 69, p. p.; *Hk.*, *Spec. Fil.*, I, tab. XXXV, B; *Bedd.*, *Ferns Br.I.*, II, tab. CCLXXVII.

Stipes $2\frac{1}{2}$ —5 c.M. long, wiry, naked or slightly tomentose. Fronds $7\frac{1}{2}$ —20 c.M. long, 5— $7\frac{1}{2}$ c.M. broad, oblong, acuminate, winged in the upper half, hairy below. Pinnae lanceolate-oblong, acuminate, cut down to a winged rachis into \pm 8—10 pinnulae. Highest pinnulae simple, the lower ones to 4 \times forked. Ultimate segments linear or oblong, \pm 2 m.M. broad, the edges toothed. Sori 1—6 to a pinna, occupying the anterior side of the pinnulae, terminal on short lobes; indusium oblong, divided \pm $\frac{1}{2}$ -way down, the valves entire.

New Guinea, Philippines, Malacca.

(36) **H. fraternum**, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 26; *H. Steerei*, *C. Chr.*, Ind. Fil.

Stipes slender, \pm 5 c.M. long, winged. Fronds \pm $6\frac{1}{2}$ c.M. long, \pm $3\frac{1}{2}$ —4 c.M. broad, oblong, erect, 5-pinnatifid; rachis and branches winged. Pinnae ovate, \pm 2 c.M. long or less; ultimate divisions linear. Edges of wings and segments crisped, denticulate. Sori terminal, sessile, scattered over the frond; indusium the width of the ultimate divisions, cleft to the base; valves elliptical, lacinate at the apex, elsewhere entire and crisped.

Philippines.

(37) **H. Reinwardtii**, *V. D. B.*, Hymen. Jav., 52, tab. XLII.

Rhizome creeping, branched, covered with long, reddish-brown hairs. Stipes 5—8 c.M. long, firm, broadly winged, nudate at the base. Fronds 8—12 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, membranaceous, thin, lanceolate or long-acuminate, 5-pinnatifid. Pinnae contiguous, often erecto-patent, the higher fertile ones more distant, obovate or lanceolate, cut down to a crisped wing into several, slightly crisped pinnulae on each side. Pinnulae deeply pinnatifid again. Lobes linear or oblong, simple or 2-fid. Margins of the lobes and wings irregularly and minutely sinuato-denticulate; the apices of the lobes rounded or truncate. Sori terminal on the lobes, 2—12 or more to a pinna; indusium broad, deeply 2-valved; valves orbicular, with the edge minutely and obtusely denticulate.

Java, Sumatra.

(38) **H. physocarpum**, *Christ*, Schum. & Laut., Flor., deut. Sch. geb., Suppl., 55, tab. I, fig. A, a—b.

Near *H. sabinifolium*, *Bk.*, but longer. Fronds lanceolate, not triangular, \pm 12 c.M. long, \pm 5 c.M. broad. Pinnae short, ovate, rather distant, nu-

merous. Pinnulae short. Segments and rachises aculeato-dentate, crisped, filiform. Sori large, terminal, roundish, $\pm 1\frac{1}{2}$ m.M. broad; indusium bullate, globose, scariose, the valves finely denticulate, not spinose on the back.

New Guinea

(39) **H. sabinifolium**, *Bk.*, Hk. Bk., Syn. Fil., 71; *H. aculeatum*, *Bedd.* (not *V. D. B.*), Handb. Ind. Ferns, Suppl., 8, p. p.

Stipes $2\frac{1}{2}$ —5 c.M. long, naked or slightly winged. Fronds broadly ovate, triangular, deeply 3-pinnatifid, $2\frac{1}{2}$ —5 c.M. long, more than $2\frac{1}{2}$ c.M. broad below; rachis margined throughout with a wing which is thickly beset with aristate teeth. Pinnae crisped, the lower ones flabellato-pinnatifid nearly down to the midrib. Ultimate segments very narrow, 5—8 m.M. long, deeply cut up nearly to the midrib by numerous, strong, aristate teeth. Sori solitary, supra-axillary; indusium spinose on the back, divided $\frac{1}{2}$ -way down with ovate, spinoso-serrate valves.

Java, Sumatra.

* (40) **H. Neesii**, *Hk.*, Hk. Bk., Syn. Fil., 71; *Bedd.*, Fern Br.I., II, tab. CCLXXIX; (*H. denticulatum*, *Sw.*, Rac., Flor. Btz., I, 21, p. p.; *C. Chr.* Ind. Fil.; *H. aculeatum*, *Rac.*, l. c., p. p.; *C. Chr.*, l. c.); *Leptocionium Neesii* & *acanthoides*, *V. D. B.*, Hymen. Jav., tab. XXX & XXXII.

Rhizome creeping, filiform, nearly naked. Stipes 2—5 c.M. long, winged throughout or towards the apex only. Fronds deltoid or oblong $2\frac{1}{2}$ —5 c.M. long, 1—3 c.M. broad, the main rachis winged throughout. Pinnae cut down to a winged rachis into simple, pinnatifid to 2-pinnatifid or once to more times forked pinnulae. Ultimate segments ± 1 —5 m.M. long or longer, $\pm \frac{1}{2}$ —1 m.M. broad. Sori 1—5 to a pinna; indusium divided $\pm \frac{1}{2}$ -way down or deeper, the valves toothed, the outer surface covered with scattered, sharp prickles.

Malaya; Ceylon.

* (41) **H. denticulatum**, *Sw.*, Hk. Bk., Syn. Fil., 71; *Rac.*, Flor. Btz., I, 21, p. p.; *Bedd.*, Ferns Br.I., II, tab. CCLXXVIII; *Leptocionium denticulatum*, *V. D. B.*, Hymen. Jav., tab. XXIX.

Rhizome filiform, sparingly hairy. Stipes 2—5 c.M. long, nudate, or narrowly winged towards the apex. Fronds ovate-triangular, 5—8 c.M. long, $2\frac{1}{2}$ —5 c.M. broad at the base; main rachis margined throughout with a narrow, slightly undulated wing. Pinnae slightly crisped, the lower ones

FILICES.

6

oblong, deeply cleft with linear, flabellate or pinnatifid pinnulae. Ultimate segments toothed and crisped. Sori \pm 1—2 to a pinna, occupying the anterior pinnulae, terminal on short lobes; indusium ovate, divided about $\frac{2}{3}$ of the way down, surrounded at the base by broad, deeply and irregularly spinoso-dentate transversal crests, the valves toothed or serrate towards the apex.

Trop. Asia.

* (42) **H. aculeatum**, *Rac.*, (not *Sm.*), Flor. Btz., I, 21, p. p., *H. aculeatum*, *Bedd.* (not *V. D. B.*), Handb. Ind. Ferns, Suppl., 8, p. p. *Leptocionium aculeatum*, *V. D. B.*, Hymen. Jav., tab. XXXI.

Like *H. denticulatum*, *Sw.*, but smaller. Fronds $2\frac{1}{2}$ —5 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, the indusium not so deeply divided (\pm $\frac{1}{3}$ of the way down), the lips with the outer edge sharply toothed.

Malaya; Ceylon.

(43) **H. bismarckianum**, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., Suppl., 54.

Near *H. denticulatum*, *Sw.*, but 3 \times larger. Sori not axillary but terminal on the segments, roundish, minute, blackish-brown; valves of the indusium entire, not spinose.

New Guinea.

(44) **H. brachyglossum**, *A. Br.*, Kze, Bot. Zeit., 1847, 227; *Leptocionium Braunii*, *V. D. B.*, Hymen. Jav., 43, tab. XXXIII.

Rhizome creeping, filiform, nearly naked. Stipes scattered, filiform, naked, 2—4 c.M. long. Fronds lanceolate or lanceolate-oblong, 5—6 c.M. long, 1—2 c.M. broad; main rachis winged throughout. Pinnae slightly crisped, cut down to a winged rachis into simple or sometimes forked pinnulae. Ultimate segments linear, toothed and \pm crisped. Sori 1—2 to a pinna, occupying mostly the anterior side of the higher pinnae; indusium oblong, divided \pm $\frac{1}{3}$ -way down, the base surrounded by a row of rather close, repand, longitudinal crests, the valves \pm truncate or rounded and toothed at the apex.

Java, Borneo.

9. TRICHOMANES, *Linné.*

Sori marginal, terminal on the veins. Indusium tubular, cupuliform, obconical or turbinate, of the same texture as the frond, exerted or more or

less immersed in the margin, the mouth truncate, dilate or 2-lipped. Receptacle mostly long and exserted, prominent beyond the sporangia. Surfaces of the sporangia divided into few, unequal cells.

Rhizome creeping or erect. Fronds simple or divided, varying from delicately membranaceous to (rarely) \pm coriaceous; ultimate segments mostly linear or oblong, rarely ovate or rhomboidal, sometimes capillary. Hk. Bk., Syn. Fil., tab. II, fig. 17; Diels, in Engl. & Prantl, Nat. Pfl.Fam., 1⁴, fig. 73.

Tropical, rarely extra-tropical.

Arrangement of the sections and groups.

- § 1. **FÉEA**. Fronds dimorphous, the fertile ones spicate. Diels, in Engl. & Prantl, Nat. Pfl.Fam., 1⁴, fig. 73, D. (Not in Malaya).
- § 2. **EUTRICHOMANES**. Fronds uniform, the fertile ones not spicate. Ultimate segments flattened, often narrowly linear, but not capillary. Diels, l. c., fig. 73, A—C; Christ, Farnkr. d. Erde, fig. 54—54, 75.
- a. **Cardiomanes**. Fronds simple, reniform; veins flabellate; spurious veins wanting. Sori pantotactous ⁽¹⁾. (Not in Malaya).
- b. **Hemiphlebium**. Fronds simple or slightly lobed, not reniform; spurious veins present. Sori epitactous ⁽²⁾.
- c. **Gonocormus**. Fronds digitato-subdichotomous ⁽⁴⁾; sori paratactous ⁽³⁾.
- d. **Ptilophyllum**. Fronds pinnately divided, the main rachis winged throughout or towards the base only. Cells of the texture roundish, oblong, quadrangular or subhexagonal, not placed as in *e*. Sori epitactous or paratactous.
- e. **Abrodictyum**. Fronds pinnately divided, the main rachis whether or not winged. Cells of the texture transversally oblong, placed in obliquely longitudinal rows. Sori paratactous. (V. D. B., Hymen. Jav., tab. XII, *Habrodictyon*).
- f. **Lacostea**. Fronds pinnately divided, the main rachis not winged or winged towards the apex only, but often narrowly marginate. Cells of the texture as in *d*. Sori paratactous.

⁽¹⁾ Prantl, Morph. d. Gefässcr., I, Hymen., 7.

⁽²⁾ Prantl, l. c., 11.

⁽³⁾ Prantl, l. c., 8.

⁽⁴⁾ Fronds digitate, the segments simple or once or more times forked, the central segment not rarely elongate and pinnatifid with simple or divided secondary segments.

*. **Cephalomanes.** Fronds simply pinnate; pinnae \pm ovate, rather broad, the base cuneate or rounded.

.. **Trichomanopsis.** Fronds at least 2-pinnatifid; ultimate segments linear or oblong, narrow, the base neither cuneate nor rounded.

§ 3. **LEPTOMANES.** Fronds uniform, the fertile ones not spicate. Ultimate segments capillary, whether or not narrowly marginate (not visible except under magnifying power). Christ, l. c., fig. 68.

Key to the malayan sections and groups.

(Fronds uniform).

1. Ultimate segments flattened, often narrowly linear but not capillary.

§ 2. **EUTRICHOMANES.**

2. Fronds simple or slightly lobed. § 2, b. **Hemiphlebium.**

2.2. Fronds divided.

3. Fronds flabellato-digitate; segments simple or once or more times forked, the central segment not rarely elongate and pinnatifid. § 2, c. **Gonocormus.**

3.3. Fronds pinnately divided.

4. Cells of the texture transversally oblong, placed in obliquely longitudinal (ladder-shaped) rows. § 2, e. **Abrodictyum.**

4.4. Cells of the texture roundish, oblong, quadrangular, sub-hexagonal, &c, but not as in 4.

5. Main rachis winged throughout or towards the base only. § 2, d. **Ptilophyllum.**

5.5. Main rachis not winged, or winged towards the apex only. § 2, f. **Lacostea.**

6. Fronds simply pinnate; pinnae \pm ovate, rather broad, the base cuneate or rounded.

§ 2, f*. **Cephalomanes.**

6.6. Fronds at least 2-pinnatifid; ultimate segments linear or oblong, narrow, the base neither cuneate nor rounded. § 2, f**. **Trichomanopsis.**

1.1. Ultimate segments capillary, whether or not narrowly marginate.

§ 3. **LEPTOMANES.**

§ 2. EUTRICHOMANES. b. **Hemiphlebium.**

- a.** Veins flabellate or radiate. Fronds small, 5—15 m.M. long.
- b.** Fronds peltate; veins radiate. (1) *T. peltatum.*
- b.b.** Fronds not peltate; veins flabellate.
- c.** Fronds cordate-orbicular.
- d.** Indusium funnel-shaped, the mouth entire. (2) *T. Motleyi.*
- d.d.** Indusium subcylindrical, the mouth 2-lipped. (3) *T. pannosum.*
- c.c.** Fronds oblong or linear-oblong. (4) *T. Beccarianum.*
- a.a.** Veins pinniform. Fronds 1—7½ c.M. long.
- *. Fronds small, ± 1—1½ c.M. long, ± 2½—5 m.M. broad. Indusium with the mouth conspicuously 2-lipped. (5) *T. bimarginatum.*
- *.*. Fronds larger, ± 1½—7½ c.M. long, ± 1—2 c.M. broad. Indusium with the mouth not or obscurely 2-lipped. (6) *T. sublimbatum.*

* (1) **T. peltatum**, *Bk.*, Hk. Bk., Syn. Fil., 73; Journ. of Linn. Soc., Bot., IX, tab. VIII, C; *T. omphalodes*, *C. Chr.*, Ind. Fil.

Rhizome wiry, slender, wide-creeping, tomentose. Fronds quite sessile, attached to the rhizome near its centre or towards the base, suborbicular in general outline, 1—3 c.M. each way, the margin undulate, not thickened, sometimes irregularly, more or less deeply cleft. Texture delicately membranaceous; veins close, radiant, regularly dichotomous, with numerous interrupted spurious venules between them near the margin and towards the centre, Sori 1—3, scattered; indusium more or less exerted, with a much dilated, spreading, slightly toothed, 2-lipped mouth.

Trop. Asia, Polynesia & Australia, Liu-Kiu.

* (2) **T. Motleyi**, *V. D. B.*, Hk. Bk., Syn. Fil., 73; *T. Henzaiense*, *Bedd.*, Ferns Br.I., II, tab. CLXXXIII; *Microgonium Motleyi*, *V. D. B.*, Hymen. Jav., tab. I.

Rhizome filiform. Fronds copious, sessile, ± 6 m.M. long, ± 4 m.M. broad, cordate-orbicular, nearly entire. Texture very thin; barren fronds without costa; veins flabellate, with spurious veins; fertile fronds with a distinct costa terminating in a ± deep sinus, the edge with a marginal vein bordering the frond, slightly denticulate or entire. Sorus terminal on the costa, stalked; indusium funnel-shaped, the mouth entire, much dilated.

Trop. Asia, Queensland, Melanesia.

(3) **T. pannosum**, *Ces.*, Fel. Becc. Polin., 2 & 6; *T. Motleyi*, *V. D. B.*, var., C. Chr., Ind. Fil.

Near *T. Motleyi*, *V. D. B.* Fronds 10—13 m.M. each way, lobed, nigropannose beneath. Sorus not exserted; indusium subcylindrical, the mouth distinctly 2-lipped.

New Guinea.

(4) **T. Beccarianum**, *Ces.*, Fel. di Bor., 8, tab. I, fig. 2; (? *T. Motleyi*, *V. D. B.*, C. Chr. Ind. Fil.).

Like *T. Motleyi*, *V. D. B.*, but the fronds ovate, oblong or linear-oblong, not cordate-orbicular, less pellucid; veins close, divaricating. Sorus solitary, terminal, short; indusium short, the mouth dilated, spreading.

Borneo.

(5) **T. bimarginatum**, *V. D. B.*, Ned. Kruidk. Arch., V², 143; *T. neilgherrense*, *Bedd.*, Hk. Bk., Syn. Fil., 74; *Bedd.*, Ferns S.I., tab. VI; *Microgonium bimarginatum*, *V. D. B.* (oldest name).

Stipes 2½—5 c.M. long. Fronds 1—1½ c.M. long, 2½—5 m.M. broad, oblong-lanceolate, not lobed, sometimes bifid at the apex; spurious venules numerous. Sori 1—2, terminal; indusium more or less sunk in the frond, the mouth exserted, with 2 large, rounded lips.

Malaya to Polynesia, Formosa, South. India, Ceylon, Australia.

* (6) **T. sublimbatum**, *H. Müll.*, Rac., Flor. Btz., I, 23; *T. muscoides*, *Bk.* (not *Sw.*), Hk. Bk., Syn. Fil., 75; *Microgonium sublimbatum*, *V. D. B.*, Hymen. Jav., tab. II.

Rhizome creeping, filiform, hairy at the apex. Stipes, filiform, naked, 1—10 m.M. long, 1—3 c.M. apart. Fronds 1½—7½ c.M. long, ± 1—2 c.M. broad, varying in shape from linear to oblong or suborbicular, entire to bluntly lobed, especially above. Texture very thin; both surfaces naked; costa pinnate; veins simple or forked; spurious venules faint but copious. Sori 1—8, terminal on the lobes; indusium sunk in the frond, the mouth free, dilated, often obscurely 2-lipped.

Malaya; Khasia.

§ 2. EUTRICHOMANES. c. **Gonocormus.**

- 1.** *Fronds not proliferous.*
- 2.** *Edges not ciliate.*
- 3.** *Margin not bounded by a thickened line, nor purveyed with a submarginal spurious vein.*
- 4.** *Spurious veins wanting throughout.*
- 5.** *Fronds very small, suborbicular-flabellate, 5—7½ m.M. each way. Segments rather numerous. (7) T. saxifragoides.*
- 5.5.** *Fronds somewhat larger, 10—20 m.M. long. Segments not numerous.*
- 6.** *Stipes short, 1—4 m.M. long. Fronds cuneate-flabellate, sometimes simple and oblong. (8) T. Aswijkii.*
- 6.6.** *Stipes longer, 10—30 m.M. long. Fronds roundish in general outline. (9) T. nitidulum.*
- 4.4.** *Spurious veins present. (10) T. parvulum.*
- 3.3.** *Margin bounded by a thickened line or purveyed with a submarginal spurious vein.*
- ***. *Marginal line (spurious vein) continuous. (11) T. nanum.*
- *.*.** *Submarginal spurious vein interrupted. (12) T. rupicolum.*
- 2.2.** *Edges ciliate.*
- §.** *Ciliae tooth-like, distant. Veins naked. (13) T. digitatum.*
- §.§.** *Ciliae capillary, approximate. Veins hairy. (14) T. palmatifidum.*
- 1.1.** *Fronds proliferous, i. e. the stipe, rachis or costa bearing 1 or more sessile or stalked secondary frondiculae, which gives the whole frond the appearance of a very irregularly and pinnately divided single frond, besides which there often are also not-proliferous ones.*
- ***. *Frondiculae few.*
- §.** *Fronds and frondiculae ternate or quinate. (15) T. diffusum.*
- §.§.** *Fronds and frondiculae with more numerous segments. (16) T. minutum.*
- *.*.** *Frondiculae many.*
- †.** *Rhizome hairy. Ultimate segments complicate. (17) T. proliferum.*
- †.†.** *Rhizome nearly naked. Ultimate segments not complicate. (18) T. Teysmanni.*

(7) **T. saxifragoides, Pr.**, Hk. Bk., Syn. Fil., 75.

Stipes \pm 6 m.M. long, slender, tomentose below. Fronds 5—7 $\frac{1}{2}$ m.M. each way, flabellato-suborbicular in general outline, cut from the margin in the direction of the apex of the stipe into numerous linear, acute segments, the deepest divisions reaching scarcely $\frac{1}{2}$ -way down. Veins close, prominent, irregular. Sori 2—4, terminal; indusium sunk, the mouth dilated, spreading, scarcely 2-lipped.

Java, Philippines.

* (8) **T. Aswijkii (Asnykii), Rac.**, Natuurk. Tijdschr. N. I., LIX, 258, tab. II, fig. 6.

Rhizome creeping, short-hairy, much branched. Stipes 5—12 m.M. apart, filiform, 1—4 m.M. long. Fronds simple or more commonly 1—3 \times forked, naked, entire. Undivided fronds linear 1 $\frac{1}{2}$ —2 c.M. long, 2 $\frac{1}{2}$ —5 m.M. broad, the apex blunt, emarginate when barren, the base narrowed. Texture thin; a single central vein only carried into each segment. Sori terminal on the segments; indusium cylindrical-infundibuliform, sunk in the margin, the mouth dilated, round or slightly 2-valved; receptacle filiform, exserted.

Java.

* (9) **T. nitidulum, V. D. B., Rac.**, Flor. Btz., I, 25; V. D. B., Hymen. Jav., tab. XV.

Rhizome filiform, creeping, branched, sparingly hirsute. Stipes capillary, 1 $\frac{1}{2}$ —3 c.M. long, sparingly covered with scattered, short hairs. Fronds roundish in general outline, \pm 1—2 c.M. each way, the base broadly rotundato-cuneate, digitato-dichotomous (digitate, the segments once or more times forked); ultimate segments linear, different in length, to 1 $\frac{1}{2}$ c.M. long, 1 $\frac{1}{2}$ —2 $\frac{1}{2}$ m.M. broad, the edge entire, the apex blunt, emarginate. Texture thin, hyaline; surfaces naked, slightly hairy when young; veins 1 in each ultimate segment. Sori terminal on the segments; indusium sunk, funnel-shaped, the mouth dilated, naked.

Java, Riouw, Borneo.

(10) **T. parvulum, Poir.**, Hk. Bk., Syn. Fil., 75; Hk., Spec. Fil., I, tab. XXXIX, A; Bedd., Ferns Br.I., II, tab. CLXXIX.

Rhizome wide-creeping, interlaced. Stipes \pm 2 $\frac{1}{2}$ c.M. long, wiry, slender, tomentose below. Fronds $\frac{1}{2}$ —1 $\frac{1}{2}$ c.M. each way, orbicular in general outline, cuneate or truncate at the base, flabellately cut about half-way down

from the outer edge in the direction of the base into narrow, irregular segments. Veins close, prominent (so that the frond when dry appears channeled), dichotomous at a very small angle; spurious venules numerous. Sori 4—6, terminal on the central segments; indusium sunk, the mouth dilated at the sides.

Trop. & temp. Asia, Australia, Polynesia, East African Islands.

(11) **T. nanum**, *V. D. B.*, Hk. Bk., Syn. Fil., 77; *T. Kurzii*, *Bedd.*, Ferns. Br.I., II, tab. CCLXXXVI; C. Chr., Ind. Fil.

Rhizome slender, wide-creeping. Stipes very short. Fronds not more than $1\frac{1}{2}$ c.M. each way, cuneate below, the outer margin rounded; central portion not more than $1\frac{1}{2}$ c.M. broad, divisions simple or once branched. Ultimate segments $3\frac{1}{2}$ —5 m.M. long and $\pm \frac{1}{2}$ m.M. broad. Texture thick but not rigid; surface somewhat undulate, margin bounded by a thickened line; a central vein only in each segment. Sori 1—2 to a frond, subterminal; indusium turbinate, not at all exserted, the mouth dilated, undulated in irregular plications.

Trop. Asia.

* (12) **T. rupicolum**, *Rac.*, Flor. Btz., I, 24.

Rhizome creeping, densely covered with spreading hairs. Stipes filiform, 1—3 c.M. long, nearly naked. Fronds commonly cuneate-flabellate, ± 1 c.M. each way, the base cuneate or slightly rounded, the outer edge palmatifid into 3—5 segments; central segments 1—2 \times forked, the lateral ones simple or bifid; ultimate lobes ± 2 m.M. broad, the apex blunt, the edge entire and with a submarginal, interrupted spurious vein. Texture thin, hyaline; veins 1 in each ultimate lobe, sparingly covered with short, acute scales. Sori terminal and solitary on the lobes; indusium funnel-shaped, sunk, the mouth dilated, spreading.

Java.

(13) **T. digitatum**, *Sw.*, Hk. Bk., Syn. Fil., 76; *T. flabellatum*, *V. D. B.*, Hymen. Jav., 19, tab. XIII; *T. corticola*, *Bedd.*, Ferns S. I., tab. CCLXIV.

Rhizome slender, creeping, nearly naked, intricate. Stipes scattered, $\frac{1}{2}$ —2 c.M. long, capillary, naked. Fronds 1—5 c.M. long, 1 — $2\frac{1}{2}$ c.M. broad, irregular in general outline, 3-partite, the central portion produced, 3-fid or 1—3-furcate, the lateral segments shorter, simple or 2-fid. Ultimate segments linear, to 1 m.M. broad or hardly broader, the apex retuse or forked, the margin ciliate. Texture membranaceous, opaque, brownish; surfaces naked; veins 1 in each ultimate segment. Sori 2—6, terminal and solitary on the

ultimate segments; indusium sunk, cup-shaped, the mouth dilated, slightly 2-lipped, naked.

Trop. Asia, New South Wales, Mascarenes.

* (14) **T. palmatifidum**, *H. Müll.*, V. D. B., Hymen. Jav., 20, tab. XIV; *T. digitatum*, *Sw.*, var. *palmatifidum*, *Hk. Bk.*, Syn. Fil., 76.

Rhizome slender, creeping, branched, intricate. Stipes scattered, capillary, pubescent, 1—5 c.M. long. Fronds 1—1½ c.M. long, 1—2 c.M. broad, digitato-dichotomous, the base cuneate or rounded. Ultimate segments linear, very unequal in length, 1½—2 m.M. broad, the apex rounded, obtuse, retuse, the margin minutely denticulate, each projection purveyed with a group of short, flexuose ciliae. Texture thin, hyaline; veins 1 in each segment, ciliate, nearly reaching the apex of the segments. Sori terminal on the segments; indusium sunk, obconical or cupuliform, the surface and the margin ciliate.

Java.

* (15) **T. diffusum**, *Bl.*, Enum., 225; *Gonocormus diffusus*, *V. D. B.*, Hymen. Jav., 9, tab. IV.

Near *T. minutum*, *Bl.*, but the fronds and frondiculae mostly ternate or rarely quinate, oblong or obovate, the base ± cuneate, the edges conduplicate, undulate, the indusium short-funnel-shaped with the mouth suddenly much dilated, spreading and undulate.

Java.

* (16) **T. minutum**, *Bl.*, Rac., Flor. Btz., I, 25; *Gonocormus minutus*, *V. D. B.*, Hymen. Jav., tab. III.

Rhizome creeping, branched, intricate, hairy. Stipes filiform, villose at least at the base, 5—20 m.M. long. Fronds and frondiculae conform, roundish in general outline, to ± 1 c.M. each way, subtruncate to cordate at the base, palmatifid into many linear segments ± ½ m.M. broad. Segments simple or 1—2 × forked, the ultimate lobes to 4 m.M. long, the apex blunt, rounded or slightly emarginate; the medial segments often elongate and subpinnatifid. Texture membranaceous, diaphanous. Sori terminal and solitary on the lobes; indusium sunk, funnel-shaped, but the mouth not very much dilated.

Java, Borneo, New Guinea.

* (17) **T. proliferum**, *Bl.*, *Hk. Bk.*, Syn. Fil., 76; *Gonocormus palmatus*, *V. D. B.*, Hymen. Jav., 11, tab. VI.

Rhizome wide-creeping, branched, hairy, densely interlaced. Stipes $2\frac{1}{2}$ —5 c.M. long, slender, naked, flexuose, repeatedly proliferous. Fronds and frondiculae varying from ovate-lanceolate to roundish, to $2\frac{1}{2}$ c.M. each way, digitately divided. Segments narrow, complicate or concavo-convolute, divided down nearly to the base into forked or dichotomous lobes, the central lobes often elongate and subpinnatifid. Texture membranaceous, subopaque. Sori terminal; indusium funnel-shaped, marginate (or winged in the lower part), the mouth dilated, undulate, obscurely 2-lipped.

Trop. Asia.

* (18) **T. Teysmanni**, *V. D. B.*, Ned. Kruidk. Arch., V², 142; *Gonocormus Teysmanni*, *V. D. B.*, Hymen. Jav., 10, tab. V.

Near *T. proliferum*, *Bl.*, but the rhizome nearly naked, the frondiculae ovate-lanceolate to linear-lanceolate, the segments deeply pinnatifid, the lobes erecto-patent, 2—3-fid, not complicate, the edges slightly undulate.

Sumatra, Java.

§ 2. EUTRICHOMANES. d. **Ptilophyllum**.

- a.** *Main rachis winged towards the base only.* (19) *T. microchilum*.
- a.a.** *Main rachis winged throughout.*
- b.** *Fronds pruinose. Central ribs flanked on each side by a broad stripe consisting of numerous darker coloured cells.* (20) *T. pallidum*.
- b.b.** *Fronds not pruinose. Central ribs not flanked by darker coloured cells.*
- c.** *Marginal or submarginal spurious veins present.*
- d.** *Spurious veins quite marginal or apparently so.*
 - *. *Fronds $\pm 1-1\frac{1}{2}$ c.M. long, simply pinnatifid.* (21) *T. Nymani*.
 - *.*. *Fronds $\pm 1\frac{1}{2}-5$ c.M. long, 2-3-pinnatifid.* (22) *T. humile*.
- d.d.** *Spurious veins not quite marginal.*
- e.** *Accessory spurious veins (between the submarginal one and the midrib) many.* (23) *T. bilabiatum*.
- e.e.** *Accessory spurious veins few or wanting.*
- f.** *Submarginal spurious vein not or slightly interrupted.*
Lips of the indusium triangular.

- g.** Fronds $2\frac{1}{2}$ —10 c.M. long. Stipes $1\frac{1}{2}$ —5 c.M. long. (24) T. *hipunctatum*.
- g.g.** Fronds smaller. Stipes short or nearly wanting. (25) T. *filiculoides*.
- f.f.** Submarginal spurious vein copiously interrupted. Lips of the indusium semiorbicular. (26) T. *capillatum*.
- c.c.** Submarginal spurious vein wanting.
- 1.** Segments with the margin serrate or dentate.
- *. Fronds small, under $1\frac{1}{2}$ c.M. long, membranaceous. (27) T. *serrulatum*.
- *.*. Fronds middle-sized, 15—30 c.M. long, semicoriaceous. (28) T. *auriculatum*.
- 1.1.** Segments with the margin entire.
- 2.** Surfaces or edges hairy or ciliate.
- §. Edges ciliate with short, distant bristles. (29) T. *dichotomum*.
- §.§. Surfaces hairy, especially on the costae. (30) T. *vestitum*.
- 2.2.** Surfaces and edges naked.
- 3.** Stipes \pm tufted.
- ‡. Fronds at best 2-pinnatifid.
- *. Fronds to \pm 15 c.M. long. (31) T. *Warburgii*.
- *.*. Fronds to \pm 30 c.M. long. (32) T. *ignobile*.
- ‡.‡. Fronds 4-pinnatifid. (33) T. *hispidulum*.
- 3.3.** Stipes scattered.
- 4.** Fronds small, under 15 c.M. long.
- 5.** Fronds 2-pinnatifid. (34) T. *Lauterbachii*.
- 5.5.** Fronds 3-pinnatifid, at least at the base.
- 6.** Indusium with the mouth 2-valved. (35) T. *Hosei*.
- 6.6.** Indusium with the mouth not or scarcely 2-valved.
7. Fronds small, \pm 3 c.M. long, 7—10 m.M. broad. (36) T. *gracile*.

7.7. Fronds larger, \pm 3—15 c.M.
long, 25—45 m.M. broad.

(37) *T. pyxidiferum*.

4.4. Fronds larger, more than 15 c.M. long when
fully developed.

*. Rhizome tomentose; fronds membranaceous,
15—30 c.M. long. (38) *T. speciosum*.

.. Rhizome \pm naked; fronds subcoriaceous,
30—75 c.M. long. (39) *T. maximum*.

(19) **T. microchilum**, *Bk.*, Trans. Linn. Soc., Bot., 2^d Series, IV, 250.

Rhizome wide-creeping, filiform. Stipes slender, naked, erect, $3\frac{1}{2}$ —5 c.M. long, narrowly winged towards the apex. Fronds oblong-lanceolate, naked, decompose, 7—10 c.M. long; rachis narrowly winged at the base. Pinnae sessile, the lower lanceolate, \pm $2\frac{1}{2}$ c.M. long, the lower base cuneate. Lower pinnulae not deltoid; ultimate segments erecto-patent, narrow, linear, entire, 1-nerved. Sori few, terminal, sessile; indusium oblong, the mouth 2-lipped, the lips small, ovate, entire, erect.

Borneo.

* (20) **T. pallidum**, *Bl.*, Rac., Flor. Btz., I, 29; *T. glauco-fuscum*, *Hk.*, Spec. Fil., I, tab. XL, A; *Bedd.*, Ferns S.I., tab. CCLXIII; *Craspedoneuron album & pallidum*, *V. D. B.*, Hymen. Jav., tab. VII—VIII; — Var. *Braunii*; *C. Braunii*, *V. D. B.*, l. c., tab. IX.

Rhizome creeping, slender, filiform, covered with scattered hairs. Stipes rather distant, 5—7 c.M. long, hairy or naked. Fronds deltoid, obovate or lanceolate, 2—13 c.M. long, 1—4 c.M. broad; main rachis winged throughout. Pinnae erecto-patent, oblong or linear-oblong to rhomboidal, cut down to a winged rachis into simple, linear, bifid or more compoundly divided pinnulae. Texture thin, hyaline; surfaces glaucous or ceraceo-pruinose and covered with long, jointed, or short, not-jointed brownish hairs; the edge bounded by a \pm marginal, spurious vein; veins 1—2 in each segment, flanked on each side by a very broad, distinctly visible stripe of darker coloured cells. Sori 1—4 to a pinna, terminal on short lobes between the barren, longer ones; indusium subconical, the mouth \pm dilated or truncate.

Var. Braunii: Surfaces not hairy.

Trop. Asia, Polynesia

(21) **T. Nymani**, *Christ*, Schum. & Laut., Flor. deut. Sch.geb., Suppl., 36.

Rhizome creeping, rather woody, slender but not filiform, sparingly covered with short, brown hairs. Fronds subsessile, subtufted, erect, $1-1\frac{1}{2}$ c.M. long, $\pm \frac{1}{2}$ c.M. broad, ovate, subcuneate at the base, simply pinnatifid, the rachis narrowly winged. Segments 3—4 on each side, linear, 2—3 m.M. long, $\pm \frac{2}{3}$ m.M. broad, acute, \pm mucronulate, the higher ones shorter, rarely produced, \pm radiating. Texture thin, diaphanous; rachis and veins blackish; the margin of the pinnae bounded by a blackish, thickened line; veins 1 in each segment. Sori minute, 1—2, terminal on the higher segments; indusium campanulate, winged, the mouth 2-lipped; receptacle short but exerted.

New Guinea.

* (22) **T. humile**, *Forst.*, Rac., Flor. Btz., I, 28; Hk. & Grev., Ic. Fil., tab. LXXXV; *Crepidomanes humile*, V. D. B., Hymen. Jav., tab. XI.

Rhizome creeping, tomentose. Stipes 5—15 m.M. long, slender, winged above. Fronds lanceolate-obovate in general outline, $1\frac{1}{2}-5$ c.M. long, to $1\frac{1}{2}$ c.M. broad, 2-pinnatifid; main rachis narrowly winged. Ultimate segments linear, simple or forked. Texture membranaceous; surfaces naked but the veins often covered with distant, short scales; a central vein in each segment; no spurious veinlets, but the frond bounded by an indistinct, \pm marginal line uniting the apices of the veins. Sori solitary on the upper side of the segments, at the base; tube slightly winged, the mouth spreading, 2-lipped.

Java, New Guinea, Ternate; Formosa, Polynesia, Australia, New Zealand.

* (23) **T. bilabiatum**, *Nees & Bl.*, Nova Acta, XI, 123; *T. brevipes*, *Bk.*, Hk. Bk., Syn. Fil., 84 p. p.; *T. Filicula*, *Bory*, Rac., Flor. Btz., I, 28, p. p.; *Didymoglossum laxum*, V. D. B., Hymen. Jav., tab. XXVII.

Rhizome creeping, densely tomentose. Stipes short, 5—10 m.M. long, narrowly winged towards the apex. Fronds deltoid-oblong, 5— $7\frac{1}{2}$ c.M. long, $2\frac{1}{2}-4\frac{1}{2}$ c.M. broad; main rachis narrowly winged. Pinnae spreading, oblong, cut down to a narrowly winged rachis. Lowest pinnulae again deeply pinnatifid with narrow, linear, simple or 2-fid lobes to $2\frac{1}{2}$ m.M. long, $\pm \frac{1}{2}$ m.M. broad, yellow-green, purveyed with a submarginal, flexuose, slightly interrupted spurious vein and many short, flexuose, longitudinal ones between the latter and the single costa. Sori 2—8 to a pinna, the tube winged, with the mouth 2-lipped, the lips bluntish or rounded.

Malaya; Melanesia.

* (24) **T. bipunctatum**, *Poir.*, Christ, Farnkr. d. Erde, 52 (oldest name); *T. Filicula*, *Bory*, Rac., Flor. Btz., I, 28, p. p.; Bedd., Ferns Br.I., II, tab. CCLXXXIII; *Didymoglossum Filicula*, *Desv.*, V. D. B., Hymen. Jav., tab. XXVI; — var. *late-alatum*; *D. late-alatum*, *V. D. B.*

Rhizome creeping, filiform, densely covered with black, hair-like scales. Stipes rather distant, $1\frac{1}{2}$ —5 c.M. long, naked, slightly winged above or throughout. Fronds deltoid to lanceolate-obovate, $\pm 2\frac{1}{2}$ —10 c.M. long, 2—5 c.M. broad, 3-pinnatifid; main rachis winged. Pinnae ovate-lanceolate to rhomboidal, pinnatifid down to a winged rachis, the lower pinnulae again deeply pinnatifid. Ultimate segments linear, blunt. Texture membranaceous, hyaline, bright-green or olivaceous; a central vein in each ultimate segment, besides this a continuous or slightly interrupted, flexuose, submarginal spurious one. Sori 1—4 to a pinna; indusium winged, the mouth 2-lipped, the lips triangular, rather acute.

Var. late-alatum: Ultimate segments acute and very narrow; wing of the rachis somewhat crisped; spurious veins more distinct. Fronds sessile.

Trop. Asia & Australia, Polynesia, Japan, West. & South. Africa.

* (25) **T. filiculoides**, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., 108; *T. bipunctatum*, *Poir.*, var. *nana-subflabellata*, Christ, MS. in Herb. Bog.

Like *T. bipunctatum*, *Poir.*, but the fronds smaller, under $2\frac{1}{4}$ c.M. each way, subflabellate, subdigitato-pinnatifid. Rhizome wide-creeping, filiform, at length naked; stipes distant, short or nearly wanting.

New Guinea.

(26) **T. capillatum**, *Taschn.*, Dissert., 34, tab. I, fig. 2; *T. Filicula*, *Bory*, Rac., Flor. Btz., I, 28, p. p.; *T. bipunctatum*, *Poir.*, var., C. Chr., Ind. Fil.; *Didymoglossum capillatum*, *Pr.*, V. D. B., Hymen. Jav., tab. XXVIII.

Like *T. bipunctatum*, *Poir.*, but the colour dark-green, the submarginal spurious vein copiously interrupted, the indusium with the lips semiorbicular.

Java; North. India.

(27) **T. serrulatum**, *Bk.*, Hk. Bk., Syn. Fil., 80.

Rhizome slender, wide-creeping. Stipes slender, naked, about $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long. Fronds to 13 m.M. long, about 6 m.M. broad, oblong or sub-rhomboidal in general outline, pinnatifid down to a narrowly winged rachis. Segments erecto-patent, in 2—6 pairs, linear, simple or forked, ± 6 m.M. long and to $1\frac{1}{2}$ m.M. broad, toothed at the margin. Texture membranaceous;

a central vein only in each segment; spurious veinlets none. Sori 1—2, terminal on the segments; indusium exserted or stipitate, the mouth slightly 2-lipped, deeply ciliated with sharp, linear teeth.

Borneo.

* (28) **T. auriculatum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 26; *Cephalomanes auriculatum*, *V. D. B.*, *Hym. Jav.*, tab. XXV; *T. dissectum*, *J. Sm.*, *Bedd.*, *Ferns Br.I.*, II, tab. CLXXXII.

Rhizome strong, wide-creeping or climbing, the apex clothed with brown hairs. Stipes 5—10 m.M. long. Fronds linear-lanceolate, 15—30 c.M. long, 2—6 c.M. broad, pinnate, acuminate; rachis narrowly winged, not hairy. Pinnae short-stalked, the upper ones sessile with a narrowed base, the lower ones 2 × as long as broad, the apex rounded, the barren ones toothed, incisedentate or -crenate or irregularly pinnatifid 1/2-way down or more, the stalks winged, the base obliquely cuneate, the lowest anterior lobe often prolonged; central ribs pinnate. Fertile pinnae more deeply pinnatifid, sometimes contracted. Texture semicoriaceous; venation subdichotomous. Sori 2—12 to a pinna; indusium nearly or quite exserted, the mouth truncate.

Trop. Asia, China & Japan.

* (29) **T. dichotomum**, *Kze.*, *Rac.*, *Flor. Btz.*, I, 26; *V. D. B.*, *Hymen. Jav.*, tab. XVI.

Rhizome filiform, sparingly clothed with spreading hairs. Stipes filiform, brown, 5—6 c.M. long, covered with short hairs. Fronds 6—15 c.M. long, 1—2 1/2 c.M. broad, obovate-oblong, cut down to a winged rachis. Segments erecto-patent, ± 5—10 c.M. long, to ± 2 c.M. broad, simple, forked or pinnatifid. Lobes erecto-patent, to ± 3 c.M. long, 1 1/2—2 m.M. broad, simple or divided again, the apices of the ultimate segments bluntish or rounded. Texture thin; surfaces naked, but the margin of the segments sparingly ciliate with short bristles. Sori terminal; indusium obconical, sunk, the mouth furnished with deciduous hairs.

Java; Society Islands.

(30) **T. vestitum**, *Bl.*, *Kew Bull.*, 1894, 7.

Rhizome filiform, wide-creeping. Stipes short, distant, covered with few, subulate scales. Fronds ovate, 15—30 m.M. long, 10—15 m.M. broad, deeply pinnatifid, the base cuneate; both surfaces (especially the costa) covered with firm hairs. Lobes paucijugate, central ones 3—5-lobed. Sori few, sessile,

placed on the upper edge of the lobes; indusium subcylindrical, the mouth spreading, hardly lobed; receptacle included.

Borneo.

(51) **T. Warburgii**, *Christ*, Warb., Mons., I, 55.

Rhizome short, strong. Stipes densely tufted, filiform, not winged, brownish. Fronds to \pm 15 c.M. long, the rachis broadly winged. Segments caudato-elongate, acute and long-acuminate, undulate, simple or with few lobes. Texture hyaline, pale-coloured. Sori commonly solitary or with 2 on a pinna in the lower sinusses of the segments; indusium stalked, cylindrico-campanulate, the valves dilated; receptacle included.

Mindanao.

(52) **T. ignobile**, *Ces.*, Fel. di Bor., 9.

Rhizome fasciculate, fibrillose, short-strigose. Stipes caespitose, to 20 c.M. long, complanate, squamulose, at length naked, passing into the fronds. Fronds \pm 30 c.M. long, nearly as broad, ovate, deeply 2-pinnatifid. Segments subflabellate, the lobes fastigiate, linear, obtuse, crisped when dry. Surfaces naked; colour dark-greenish. — The description being not very complete, I don't know whether this plant has been classified here in the proper way.

Borneo.

(53) **T. hispidulum**, *Mett.*, Hk. Bk., Syn. Fil., 466.

Rhizome woody, suberect, the scales minute, subulate, brown. Stipes 7—15 c.M. long, winged, and slightly crinite upwards. Fronds 15—30 c.M. long, deltoid, 4-pinnatifid, the rachis narrowly winged throughout. Pinnae close, deltoid, lowest much the largest, 4—8 c.M. broad, produced on the lower side. Pinnulae close, lower deltoid, cuneate-truncate on lower side at base. Ultimate lobes distant, \pm 1 m.M. broad, 5—6 m.M. long. Texture rather thick; colour dark-olive-green; surfaces naked. Sori copious, placed on upper side of ultimate lobes; indusium minute, free, funnel-shaped, with a ciliated, entire mouth and a long receptacle.

Borneo, Malacca.

(54) **T. Lauterbachii**, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., 108.

Rhizome creeping, tomentose. Stipes winged, varying from very short to 2 c.M. long. Fronds \pm 2 c.M. long, \pm 1½ c.M. broad, broadly obovate, the rachis broadly winged. Segments obovate, \pm 4 on each side, cut down

to a broadly winged rachis, the lobes incised again. Ultimate segments obtuse, ± 2 m.M. long, ± 1 m.M. broad, with an acute sinus. Texture thin; veins very conspicuous, 1 in each ultimate segment. Sori 6—8 to each frond, placed on the lower lobes of the higher pinnae, 1—2 to a pinna, deeply immersed; indusium lanceolato-cylindrical; receptacle included.

New Guinea.

(55) **T. Hosei**, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 225, tab. XII.

Rhizome filiform, wide-creeping. Stipes 1—1½ c.M. long, erect, distant, naked, not winged. Fronds ovate-rhomboidal, $\pm 3\frac{1}{2}$ —4 c.M. long, 5-pinnatifid. Lowest pinnae largest, deltoid, with the lower base cuneate-truncate. Ultimate divisions linear, 3—5 m.M. long, ± 1 m.M. broad, 1-nerved, entire, erecto-patent. Sori terminal, solitary; indusium infundibuliform, narrowly winged on both sides nearly or quite to the top of the tube, the mouth 2-lipped, the lips small, semi-orbicular; receptacle shortly exerted.

Borneo.

(56) **T. gracile**, *V. D. B.*, Ned. Kruidk. Arch., V², 157.

Stipes filiform, 6—8 m.M. long. Fronds ± 5 c.M. long, 7—10 m.M. broad, oblong or obovate, narrowed at the base, pinnate. Pinnae erect, distant, the lower ones approximate, subopposite, oblong, subpinnatisect, plicate; the segments recurvo-patulous, short, simply forked, with a blunt sinus, the lobes narrow, emarginate at the apex. Wing of the rachis narrow, marginate. Sori terminal on the ultimate segments, small, immersed; indusium short, ventricose, the base narrowed, the mouth dilated, spreading, undulate. — Rhizome unknown, perhaps creeping as in many of the species with small fronds.

Java.

* (57) **T. pyxidiferum**, *L.*, Hk. Bk., Syn. Fil., 81; Hk. & Grev., Ic. Fil., tab. CCVI; *T. Filicula*, *Bedd.*, Ferns S.I., tab. VII.

Rhizome wide-creeping, rather filiform, tomentose. Stipes 2½—5 c.M. long, naked, or winged above. Fronds 2½—15 c.M. long, 2½—4½ c.M. broad, ovate-oblong or subflabellate, 5-pinnatifid; main rachis narrowly winged. Pinnae pinnatifid down to a narrowly winged rachis, the lower pinnulae again deeply pinnatifid. Ultimate segments more or less numerous, linear, often emarginate at the apex, whether or not purveyed with more or less distinct spurious veins. Texture membranaceous; a central vein only in each ultimate segment. Sori 1—4 to a pinna, the tube more or less winged, the

mouth broadly dilated, scarcely 2-lipped; receptacle filiform, more or less exerted. — A very variable species.

Pantropical.

* (58) **T. speciosum**, *Willd.*, Christ, Farnkr. d. Erde, 50; *T. radicans*, *Sw.*, Hk. Bk., Syn. Fil., 81, p. p.; Bedd., Ferns Br.I., II, tab. CLXXXI; C. Chr. Ind. Fil.

Rhizome wide-creeping, firm, tomentose. Stipes scattered, 5—15 c.M. long, narrowly winged towards the apex. Fronds lanceolate or deltoid-oblong, 15—30 c.M. long, 5—15 c.M. broad; main rachis narrowly winged. Pinnæ 2½—10 c.M. long, mostly erecto-patent, ovate-rhomboidal or lanceolate-triangular, the rachis narrowly winged. Pinnulae rhomboid-oblong, cut down to a winged rachis into linear or oblong, simple, bifid or pinnatifid segments. Texture firm-membranaceous; colour blackish-green; ultimate lobes 1-nerved. Sori lateral, 1—4 to a pinnula; indusium narrow-campanulate, the mouth slightly 2-lipped.

Scattered throughout warm-temperate regions of both hemispheres.

* (59) **T. maximum**, *Bl.*, Rac., Flor. Btz., I, 29; V. D. B., Hym. Jav., tab. XVIII; *T. anceps*, *Hk.*, var. β , Hk., Spec. Fil., I, tab. XL, fig. C, 3; *T. millefolium*, *Pr.*, V. D. B., l. c., tab. XX.

Rhizome stout, creeping, nearly naked. Stipes strong, erect, 1—3 c.M. apart, naked or slightly hairy at the base, \pm succulent, winged upwards, 7½—40 c.M. long. Fronds 50—75 c.M. long, 15—30 c.M. broad, ovate-lanceolate, 4—5-pinnatifid, \pm succulent, naked. Pinnæ erecto-patent, the lowest not much shorter than the next in order, or as long, ovate-lanceolate, acuminate. Pinnulae lanceolate, cut down into segments which are pinnate again with 1—4 divisions on each side. Ultimate divisions linear, \pm 5—5 m.M. long, \pm 1 m.M. broad, simple and truncate, emarginate or bifid, straight, falcate or curved. Texture subcoriaceous; rachises more or less narrowly winged. Sori 2—8 to a pinnula; indusium cylindrical, the mouth dilated, not 2-lipped.

Malaya; Polynesia, Queensland.

§ 2. EUTRICHOMANES. e. **Abrodictyum.**

(40) **T. Cumingii**, *C. Chr.*, Ind. Fil.; *T. Smithii*, *Hk.*, Hk. Bk., Syn. Fil., 84; Hk., Icon. Pl., VIII, tab. DCCIV; *Habrodictyon Cumingii*, *V. D. B.*, Hymen. Jav., tab. XII; *Abrodictyum Cumingii*, *Pr.*, (oldest name), Hym., tab. VII.

Rhizome short-creeping or ascending, densely hirsute. Stipes approximate or subtufed, $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, slender, erect, naked, sparingly hairy at the base. Fronds 5—20 c.M. long, $1\frac{1}{2}$ —4 c.M. broad, lanceolate or oblong; main rachis whether or not winged. Pinnae oblong, erecto-patent, the central ones the largest, pinnatifid to a narrowly winged rachis. Pinnulae linear, usually simple, occasionally forked. Ultimate segments $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long, $\pm \frac{1}{2}$ —1 m.M. broad. Texture rather flaccid, the cells transversally elongated, arranged in obliquely longitudinal (ladder-shaped) rows. Sori 1—4 to a pinna, terminal on short lobes; indusium winged, the mouth dilated. — Quite different from all the other species of Trichomanes, which have all \pm roundish, quadrangular, hexagonal or longitudinally oblong cells.

Philippines, Moluccas.

§ 2. EUTRICHOMANES. f. **Lacostea**. * **Cephalomanes**.

*. Sori on the upper edge of the pinnae. Receptacle much exerted.

(41) T. javanicum.

.. Sori mostly on the outer edge of the pinnae. Receptacle not exerted.

(42) T. asplenioides.

* (41) **T. javanicum**, *Bl.*, Rac., Flor. Btz., I, 27; Hk., Gard. Ferns, tab. XXXVII; Bedd., Ferns Br.I., II, tab. CLXXX; *Cephalomanes javanicum*, *Pr.*, V. D. B., Hymen. Jav., tab. XXII; *C. Zollingeri & rhomboideum*, V. D. B., l. c., tab. XXIII & XXIV.

Rhizome erect. Stipes tufted, $1\frac{1}{2}$ —15 c.M. long, erect, naked or villose. Fronds 5—25 c.M. long, $1\frac{1}{2}$ —5 c.M. broad, lanceolate-oblong; main rachis hairy. Pinnae short-stalked or sessile with a narrowed base, oblong, unequal-sided, the apex blunt or rounded, the margin finely toothed or ciliate (¹), the base obliquely cuneate. Texture firm; veins numerous, close, parallel. Sori 1—4 to a pinna, placed on the upper edge; indusium free, the mouth truncate or dilate; receptacle much exerted.

Trop. Asia, Polynesia & Australia, Liu-Kiu.

(¹) If ciliate, the ciliae consisting of the often very much produced veins of the pinnae.

(42) **T. asplenioides**, *Pr.*, Kze, Farnkr., I, 218, tab. LXXXIX; *T. javanicum*, *Bl.*, var., C. Chr., Ind. Fil.; *Cephalomanes asplenioides*, *Pr.*, Hym., 15 & 37.

Rhizome covered with small, subulate, black scales. Stipes tufted, flexuose, scaly. Fronds linear-lanceolate, \pm 10—14 c.M. long, $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, narrowed gradually towards both ends. Pinnae \pm 20 on each side, the central ones the largest, linear-oblong, to 18 m.M. long, \pm 5—8 m.M. broad, the apex rounded, the margin finely toothed, the base obliquely cuneate, with the upper side more or less auricled; lower pinnae \pm deflexed, the higher ones erecto-patent, confluent, soriferous. Texture rigid; surfaces naked; rachis villose and scaly, marginate by the decurrent pinnae. Sori \pm 3—7 to a pinna, often only terminal; tube not quite free, the mouth obscurely dilated; receptacle not exerted.

Luzon.

§ 2. EUTRICHOMANES. f. **Lacostea**. **. **Trichomanopsis**.

a. *Stipes naked, or hairy below.*

b. *Stipes \pm fasciculate.*

c. *Fronds small, membranaceous, 7—9 c.M. long.* (43) *T. racemulosum*.

c.c. *Fronds larger, rather rigid, 10—20 c.M. long.* (44) *T. rigidum*.

b.b. *Stipes scattered.*

*. *Rhizome densely tomentose. Fronds \pm $7\frac{1}{2}$ c.M. long. Indusium 2-lipped.* (45) *T. bilingue*.

*. *. *Rhizome naked, bristly at the extremity. Fronds \pm 40 c.M. long. Indusium with the mouth much dilated.* (46) *T. aphlebioides*.

a.a. *Stipes hairy throughout.*

1. *Rhizome wide-creeping.* (47) *T. giganteum*.

1.1. *Rhizome short-creeping or erect.*

2. *Fronds small, to 20 c.M. long.*

3. *Stipes tufted. Ultimate segments short and broad.*

(48) *T. saxatile*.

3.3. *Stipes approximate. Ultimate segments short, often tooth-like.*

(49) *T. cupressoides*.

2.2. *Fully developed fronds more than 20 c.M. long. Segments narrow-linear.*

(50) *T. Bauerianum*.

(43) **T. racemulosum**, *V. D. B.*, Ned. Kruidk. Arch., V³, 210.

Rhizome short, adscendent. Stipes fasciculate, terete, the base hirsute, thickened, 2—3 c.M. long. Fronds 7—9 c.M. long, 2—3 c.M. broad, ovate-lanceolate, acuminate; main rachis straight, compressed, firm, narrowly winged, marginate, below. Pinnae erect or erecto-patent, (the lowest largest, divaricating), \pm distant, linear-lanceolate, pinnatifid, broadly cuneate at the base. Pinnulae erect, distant, simple or forked, (the lowest rarely pinnatifid). Lacinulae straight, short, the apex entire, circinate-convolute when dry. Texture membranaceous, thin, opaque; veins flexuose. Sori seriate on the pinnae, immersed in abbreviated pinnulae or lacinulae, small, afterwards curved back; indusium short, marginate, conical-cupuliform, the mouth straight or subspreading; receptacle filiform.

Borneo.

(44) **T. rigidum**, *Sw.*, Christ, Farnkr. d. Erde, 50; Hk. Bk., Syn. Fil., 86, p. p.

Rhizome short, firm. Stipes firm, naked, tufted, 10—20 c.M. long. Fronds deltoid-ovate, 10—20 c.M. long, 5—4-pinnatifid; main rachis slightly winged towards the apex. Pinnulae deeply pinnatifid into deeply and irregularly forked, lanceolate segments. Ultimate lobes linear, acute. Texture rather firm and rigid; surfaces naked. Sori small, between the segments, hardly immersed, often free; indusium slightly dilated at the mouth.

Pantropical.

° (45) **T. bilingue**, *Hk.*, Journ. of Bot., III, 417; *T. melanorhizon*, *Hk.*, Icon. Pl., VIII, tab. DCCV; *T. brevipes*, *Bk.*, Hk. Bk., Syn. Fil., 84, p. p.; *T. bilabiatum*, *Nees & Bl.*, var., C. Chr., Ind. Fil.

Rhizome creeping, densely tomentose. Stipes short, under 1½ c.M. long. Fronds \pm 7½ c.M. long, oblong; main rachis terete. Pinnae rather distant, 2-pinnatifid, slightly decurrent. Ultimate segments narrow. Sori confined to 5—4 on the ultimate segments; indusium long, conical-tubular, marginate, the mouth slightly dilated, 2-lipped, the lips broadly rounded; receptacle filiform.

Luzon.

* (46) **T. aplebioides**, *Christ*, C. Chr., Ind. Fil.; *T. tenuissimum* *Christ*, Schum. & Laut., Flor. deut. Sch.ggeb., 106.

Rhizome wide-creeping, scandent, flexuose, naked, the extremity sparingly clothed with blackish bristles. Stipes scattered, 15—20 c.M. long, firm, erect,

blackish-green, sparingly pubescent below. Fronds \pm 40 c.M. long, \pm 20 c.M. broad, broadly deltoid-ovate, 4—5-pinnate. Pinnae distant, the lower ones shorter, stalked, narrowing from a broad base to an acuminate apex. Pinnulae regularly ovate-deltoid. Tertiary segments cut down into regular segments, the lowest of which are divided again. Ultimate segments \pm 5 m.M. long, \pm $\frac{1}{3}$ m.M. broad, with only a central vein. Sori at the base of the superior tertiary segments; indusium hispid, stalked, hypocrateriform, often incurved, the margin conspicuously dilated; receptacle moderately exserted. Texture thin; segments covered with scattered, adpressed, minute bristles. — The plant has often abnormal fronds, which are \pm 6 c.M. long, oblong-lanceolate, short-stalked, irregularly 2-pinnate with capillary divisions.

New Guinea.

(47) **T. giganteum**, *Bory*, Hk. Bk., Syn. Fil., 87; Hk., Icon. Pl., VIII, tab. DCCII.

Rhizome wide-creeping, tomentose. Stipes 10—15 c.M. long, strong, erect, hairy. Fronds 30—45 c.M. long, 15—25 c.M. broad, ovate, 4-pinnatifid; main rachis hardly winged above. Lower pinnae ovate-lanceolate, 15—20 c.M. long, cut down to the rachis. Pinnulae $2\frac{1}{2}$ c.M. long, cut down very nearly to the rachis, the divisions again deeply pinnatifid. Ult. segments linear, flat, $2\frac{1}{2}$ m.M. long, $\frac{1}{2}$ —1 m.M. broad. Colour dark-green. Surfaces naked, shining; texture semicoriaceous; a central costa only in each segment. Sori 2—20 to a pinnula; tube more or less exserted, the mouth \pm dilated, not 2-lipped.

Malaya; Fiji, Comores, Mascarenes.

(48) **T. saxatile**, *Moore*, Nich., Dict. d'Hort., V, 324.

Stipes erect, hairy, tufted. Fronds triangular-ovate. Higher pinnae decurrent, lower ones bipinnatifid. Pinnulae ovate, more or less decurved, \pm undulate. Ultimate segments broad, short, obtuse. Indusium tubuliform, slightly 2-lipped. — Plants small.

Borneo.

* (49) **T. cupressoides**, *Desv.*, (oldest name), C. Chr., Ind. Fil.; *T. obscurum*, *Bl.*, Rac., Flor. Btz., I, 30; V. D. B., Hymen. Jav., tab. XVII; *T. rigidum*, *Sw.*, Hk. Bk., Syn. Fil., 86, p. p.; *Bedd.*, Ferns S.I., tab. VIII.

Rhizome short-creeping, scaly throughout or at the extremity only. Stipes approximate, 5—20 c.M. long, slightly covered with brownish hairs. Fronds

5—20 c.M. long, $2\frac{1}{2}$ —15 c.M. broad, deltoid, obovate or lanceolate, acuminate, 3—4-pinnatifid, the main rachis mostly slightly winged towards the apex and submarginate below. Pinnae erecto-patent, $1\frac{1}{2}$ —2 c.M. broad, lanceolate-oblong, cut down nearly or quite to the rachis. Pinnulae deeply pinnatifid, with linear, simple or pinnatifid lobes. Ultimate segments short and often tooth-like. Texture \pm coriaceous; margin often slightly crisped. Sori 2—12 to a pinnula, small; indusium more or less exserted, the mouth slightly dilated.

Trop. Asia, Madagascar, Mascarenes, Comores, Seychelles.

* (50) **T. Bauerianum**, *Endl.*, C. Chr., Ind. Fil.; *T. apiifolium*, *Pr.*, *Rac.*, *Flor. Btz.*, I, 31; V. D. B., *Hymen. Jav.*, tab. XIX; *T. meifolium*, *Bl.*, *Enum.*, 227; — var. *polyanthos*; *T. polyanthos* *Hk.*, *Spec. Fil.*, I, 138; *Icon. Pl.*, VIII, tab. DCCIII.

Rhizome erect, clothed with long, brownish hairs. Stipes tufted, 10—20 c.M. long, strong, erect, densely clothed with long, spreading, reddish hairs. Fronds 20—50 c.M. long, 10—20 c.M. broad, ovate, acuminate, 3—4-pinnate; rachis hairy, slightly and narrowly winged towards the apex. Longest pinnae 5—15 c.M. long, $1\frac{1}{2}$ —4 c.M. broad, spreading or erecto-patent, lanceolate-acuminate. Pinnulae several, cut down into flattened, linear, simple, bifid or pinnate segments, the ultimate ones to 4 m.M. long, \pm $\frac{1}{2}$ m.M. broad. Texture subcoriaceous, surfaces naked; a single costa in each ultimate segment. Sori 2—12 to a pinnula; indusium small, turbinate, the mouth \pm dilated, entire or slightly 2-lipped.

Var. polyanthos: Sori larger; indusium campanulate, the mouth distinctly 2-lipped. Stipe and rachis \pm hispid.

Malaya; Polynesia; — var. polyanthos: East. Malaya; Polynesia.

§ 3. LEPTOMANES.

a. *Rhizome wide-creeping.*

‡. *Ultimate segments not spreading into various directions.* (51) *T. gemmatum.*

‡.‡. *Ultimate segments spreading into various directions.* (52) *T. meifolium.*

a.a. *Rhizome short-creeping or erect.*

b. *Ultimate segments as in a, ‡.* (53) *T. parviflorum.*

b.b. *Ultimate segments as in a, ‡.‡.*

c. *Pinnae 12—15-jugate. Sori numerous.* (54) *T. trichophyllum.*

c.c. *Pinnae 20—30-jugate. Sori few.* (55) *T. pluma.*

(51) **T. gemmatum**, *J. Sm.*, Hk. Bk., Syn. Fil., 87; *T. foeniculaceum*, *Bory*, Hk., Spec. Fil., I, 135, p. p.; *T. cellulolum*, *Kl.*, Hk., 2d Cent. of Ferns, tab. LXIII.

Rhizome strong, wiry, tomentose, beset with numerous, black wiry fibres. Stipes $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, naked, wiry, winged above. Fronds 5—15 c.M. long, $2\frac{1}{2}$ —5 c.M. broad, erect, ovate-oblong; main rachis narrowly winged. Pinnae erecto-patent, cut down to a narrowly winged rachis. Lower pinnulae deeply forked with linear-filiform segments 4—5 m.M. long. Texture subrigid; cellules large. Sori 1—8 to a pinna, minute, axillary; the tube turbinate, stalked, the mouth nearly truncate.

Malacca, Java, Philippines, New Guinea; Trop. South. America.

* (52) **T. meifolium**, *Bory*, Willd., Spec., V, 509; *T. ericoides*, *Hedw.*, Rac., Flor.; *Btz.*, I, 30; *T. longisetum*, *Bory*, V. D. B., Hymen. Jav., tab. XXI; — var. *alatum*.

Rhizome strong, wiry, wide-creeping, densely covered with brown, hair-like scales. Stipes scattered, $2\frac{1}{2}$ —10 c.M. long, erect, wiry, more or less covered with brownish, hair-like scales, more densely towards the base, whether or not marginate. Fronds 5—20 c.M. long, 2—6 c.M. broad, erect, deltoid to lanceolate; main rachis nudate, but not rarely marginate. Pinnae spreading or erecto-patent, 1—5 c.M. long. Pinnulae more or less copiously divided into capillary segments 2—5 m.M. long, spreading into various directions. Texture firm; surfaces naked; colour dark-brown or blackish when dry; all the divisions narrowly marginate. Sori mostly 1—6 to a pinna, minute, stalked; indusium \pm truncate.

Var. alatum: Stipe and rachis rather broadly winged.

Malaya; Polynesia, Réunion.

* (53) **T. parviflorum**, *Poir.*, C. Chr., Ind. Fil.; *T. foeniculaceum*, *Bory*, Hk. Bk., Syn. Fil., 88; Hk., Spec. Fil., I, 135, p. p.

Stipes nearly turfted, erect, wiry, 5—10 c.M. long, naked or tomentose. Fronds 10—20 c.M. long, 2— $7\frac{1}{2}$ c.M. broad, erect, rigid, ovate-lanceolate; main rachis naked, slightly winged above. Lower pinnae spreading or erecto-patent, $2\frac{1}{2}$ —4 c.M. long, cut down quite or very nearly to the rachis. Pinnulae regularly pinnatifid, with simple or forked, linear-filiform, bifarious segments 2—4 m.M. long. Texture subcoriaceous, colour dark-green. Sori 2—12 to a pinna, minute, the mouth rather spreading, but not 2-lipped.

Borneo, Malacca; Australia, Madagascar, Mascarenes.

(54) **T. trichophyllum**, *Moore*, Hk. Bk., Syn. Fil., 466.

Stipes densely tufted, wiry, terete, $2\frac{1}{2}$ —5 c.M. long. Fronds 5—10 c.M. long, oblong, decomposed, $2\frac{1}{2}$ —4 c.M. broad. Rachis stiff, terete. Pinnae much imbricated, 12—15-jugate, many deltoid, $1-1\frac{1}{2}$ c.M. broad. Lower pinnulae copiously pinnatifid. Ultimate divisions bristle-like, 3—7 m.M. long, spreading in all directions. Sori copious, free, ± 1 m.M. deep, placed near the base of the pinnulae on the upper side; indusium with a funnel-shaped tube and truncate mouth.

Borneo, New Guinea; New Caledonia.

(55) **T. pluma**, *Hk.*, Hk. Bk., Syn. Fil., 466; Hk., Icon. Pl., X, tab. CMXCVII; Christ, Farnkr. d. Erde, fig. 68.

Rhizome short-creeping, the scales dense, minute, fibrillose, brown. Stipes 5—10 c.M. long, terete, wiry. Fronds lanceolate, 10—15 c.M. long, $2\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, decomposed; rachis stiff, terete. Pinnae crowded, 20—30-jugate, spreading, under $1\frac{1}{2}$ c.M. broad. Lower pinnulae sparsely pinnatifid, cut into bristle-like, dichotomously forked lobes 3—7 m.M. long, spreading in all directions. Sori few, placed near the base of the upper pinnae; indusium ± 1 m.M. long, stalked; tube funnel-shaped, the mouth truncate; receptacle very long.

Malaya; Samoa, New Caledonia.

FAMILY V. SCHIZAEACEAE.

V. SCHIZAEACEAE.

Capsules *either* few (1—2), placed in the axils of imbricating, clasping indusia, which form spikes along the margin of the fertile segments *or* numerous, ranged in 2—4 regular, close rows, which cover one side of close, distichous spikes at the apex of the fertile fronds or segments (¹). Ring complete, operculiform, crowning the apex of the capsule. Stomium vertical.

Rhizome creeping or erect. Fronds small and erect or long and twining. Tropical and extra-tropical.

Key to the genera.

*. Capsules 1—2; indusium present. Fronds very long and twining.

No. 10. **Lygodium.**

.. Capsules numerous, in 2—4 rows; indusium wanting. Fronds small and erect.

No. 11. **Schizaea.**

10. **LYGODIUM**, Swartz.

Sporangia 1—2, placed in the axils of imbricating, clasping, scale-like indusia, which form spikes either on separate, contracted segments or in lax rows along the edge of normal segments.

Rhizome creeping. Fronds very long, twining, pinnate. Pinnae (²) stalked, opposite, the pairs placed on very short, alternate, primary petioles which are often inconspicuous and sometimes quite wanting; veins free or anastomosing. Hk. Bk., Syn. Fil., tab. IX, fig. 68; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 195.

Tropical and extra-tropical.

Arrangement of the sections.

§ 1. **EULYGODIUM.** Veins free.

§ 2. **HYDROGLOSSUM.** Veins anastomosing. (Not in Malaya).

(¹) The characters of the african and american genera are not mentioned here.

(²) The segments indicated here as pinnae are properly the branches of forked pinnae, the very short stalks of which are forming the so-called *primary petioles*.

§ 1. EULYGODIUM.

- a.** *Barren pinnae forked or more commonly digitate or dichotomous.*
- 1.** *Fertile segments not or hardly narrowed.*
- ‡. *Pinnae once forked.* (1) *L. semihastatum.*
- ‡.‡. *Pinnae at least 2 × forked.* (2) *L. Teysmannii.*
- 1.1.** *Fertile segments conspicuously narrowed or contracted.*
- 2.** *Spikes ± close, in continuous, short or long rows along the margin of the segments.*
- 3.** *Ultimate segments bordered by a marginal strand; the gemmiform apex of the rachis clothed with brownish hairs. Spores verruculose.* (5) *L. circinatum.*
- 3.3.** *Ultimate segments without marginal strand; the hairs of the gemmiform apex blackish. Spores coarsely tuberculate.* (4) *L. digitatum.*
- 2.2.** *Spikes in groups of 3—6 on the lobes of the segments.* (5) *L. trifurcatum.*
- a.a.** *Barren pinnae pinnate, the pinnulae simple or variously divided.*
- b.** *All the pinnulae regularly pectinato-pinnatifid throughout.* (6) *L. polystachyum.*
- b.b.** *Pinnulae simple, or the lower ones forked, hastate, subpalmate or pinnate.*
- c.** *Pinnulae articulate at the base, or the ultimate petioles distinctly nodose at the apex.*
- ‡. *Pinnulae oblong, simple, 2—3½ c.M. long. Spores reticulate.* (7) *L. scandens.*
- ‡.‡. *Pinnulae linear-oblong or lanceolate, 3½—11 c.M. long, the lower fertile ones simple or pinnate. Spores verruculose.* (8) *L. salicifolium.*
- c.c.** *Pinnulae not articulate, nor the ultimate petioles distinctly nodose at the apex.*
- d.** *Pinnulae 1—2-jugate, the lower ones simple.* (9) *L. altum.*
- d.d.** *Pinnulae 2—7-jugate, the lower ones mostly hastate, subpalmate or pinnate.*
- e.** *Barren pinnae rather long, the segments serrulate. Fertile segments often narrower, but not contracted.* (10) *L. flexuosum.*
- e.e.** *Barren pinnae rather short, the segments coarsely crenate. Fertile segments contracted and deeply incised.* (11) *L. japonicum.*

* (1) **L. semihastatum**, *Desv.*, Hk. Bk., Syn. Fil., 437; (? *L. flexuosum*, *Sw.*, C. Chr., Ind. Fil.; Prantl, Morph. d. Gefässer., II, Schiz., 73).

Primary petioles nearly or quite obsolete; secondary ones 2 c.M. or more long. Pinnae once forked; each division short-stalked, not articulate, simple, ligulate or lanceolate, 15—20 c.M. long, nearly $2\frac{1}{2}$ c.M. broad, the bases \pm unequal, sometimes very cordate at one side. Texture papyraceous to coriaceous; surfaces naked. Spikes 5—8 m.M. long, in close rows along the edge of the leaflets. Spores verruculose (in the Borneo-plant which I examined).

Borneo, Philippines; Marianne Islands.

* (2) **L. Teysmannii**, *v. A. v. R.*

Primary petioles very short; secondary ones 2—3 c.M. long. Pinnae forked, subpinnate. Pinnulae stalked, not articulate, the lower one spreading, $1-2 \times$ forked again, the stalk \pm 5—10 m.M. long; the higher one $2 \times$ forked again, the stalk $1-2\frac{1}{2}$ c.M. long, nearly straight with the secondary petiole. Ultimate segments lanceolate, 8—12 c.M. long, $1-1\frac{1}{2}$ c.M. broad, free or joined into pairs, the apex narrowed, \pm acute, the edge serrate, the base subcordate or truncate or often cuneate at the inner side when the segments are free. Texture subcoriaceous; surfaces naked or slightly covered with scattered hairs on the costae; upper surface \pm glossy, under one opaque, rachis pale-brown (when dry), slender, slightly hairy and narrowly marginate as are the stalks; costae not or slightly nodulose at the base. Fertile pinnulae similar to the barren, not or slightly narrowed. Spikes 1—4 m.M. long, contiguous. — (The capsules of the 2 specimens were all unripe, and only 2 reticulated spores are seen among the numerous examined undeveloped ones).

Pulu Pisang (Teymann Nos. 1597 & 2304).

* (3) **L. circinatum**, *Sw.*, Christ, Farnkr. d. Erde, 353; Prantl, Morph. d. Gefässer., II, Schiz., 64; *L. dichotomum*, *Sw.*, Rac., Flor. Btz., I, 8, p. p.; Bedd., Ferns S.I., tab. LXII; — var. *monstruosum*, Hort. Bog.; — var. *denticulatum*, Prantl, l. c., 65; — var. *cristatum*.

Primary petioles very short or wanting, secondary ones 2—3 c.M. long, semiterete, marginate. Pinnae digitate with 5—6 lobes reaching down nearly to the cuneate or truncate base or 2-partite with the forkings digitate. Ultimate segments 10—50 c.M. long, 1—4 c.M. broad, not articulate, lanceolate or linear, subacuminate, the margin bordered by a hypodermical strand (marginal vein). Texture firm-membranaceous to coriaceous; surfaces naked or \pm hairy on the costae, upper surface glossy, under one opaque; main rachis stramineous, the gemmiform apex clothed with pale-fuscous hairs. Fertile segments

often numerous, not rarely subpinnate, conspicuously narrowed or contracted, 1—10 m.M. broad. Spikes 2—10 m.M. long. Spores verruculose. —

Forma typica: Spores without crests. Laciniae entire, not dichotomous at the apex.

Var. monstruosum: Spores without crests. Laciniae dichotomous at the apex.

Var. denticulatum: Spores without crests. Laciniae denticulate, not dichotomous at the apex.

Var. cristatum: Like var. denticulatum, but the spores purveyed with 6 joining crests. Laciniae $\pm \frac{1}{2}$ —1 c.M. broad. *Sumatra* (Burck).

Trop. Asia, Queensland.

(4) **L. digitatum, Pr.**, Prantl, Morph. d. Gefässcr., II, Schiz., 66; *L. dichotomum, Sw.*, Rac., Flor. Btz., I, 8, p. p.

Primary petioles nearly wanting; secondary ones $1\frac{1}{2}$ —2 c.M. long, semi-terete, marginate. Pinnae 2-partite or subpinnate. Pinnulae short-stalked, not articulate, digitate with 5—6 laciniae 7—10 c.M. long, ± 1 c.M. broad, the apex acute, the margin serrulate, not bordered by a hypodermical strand. Texture subcoriaceous; surfaces naked, upper surface glossy, the under one opaque; rachis reddish-stramineous, the hairs of the gemmiform apex blackish. Fertile pinnae 2-partite, each segment with ± 3 laciniae to 55 c.M. long, ± 5 m.M. broad. Spikes to 6 m.M. long. Spores rather large, coarsely tuberculate.

Trop. Asia.

* (5) **L. trifurcatum, Bl.**, Hk. Bk., Syn. Fil., 457; Prantl, Morph. d. Gefässcr., II, Schiz., 67; *L. circinatum, Sw.*, var. *trifurcatum*, Christ, in Warb., Mons., I, 93.

Primary petioles very short or obsolete, secondary ones $2\frac{1}{2}$ —4 c.M. long. Pinnae forked, 3-furcate or 2 \times forked and each division 2-partite nearly to the base. Ultimate barren segments 10—15 c.M. long, ± 2 — $2\frac{1}{2}$ c.M. broad, not articulate, the point acute or bluntish, the edge crenulate. Texture subcoriaceous; rachis and surfaces naked. Fertile pinnae varying from dichotomous to 5-pinnate, the divisions contracted and often cut down nearly or quite to the costa with the short spikes in groups of 5—6 upon the lobes. Spores coarsely verrucose. — The Banca plant (leg. Berkhout) has the spikes rather long, in groups of 1—3, and the spores verruculose and crested. Probably an intermediate form between this species and *L. circinatum, Sw.*

Banca, Celebes, Amboyna, New Guinea; Polynesia, Melanesia.

(6) **L. polystachyum**, *Wall.*, Hk. Bk., Syn. Fil., 438; Hk., 2^d Cent. of Ferns, tab. LXXVI; Bedd., Ferns Br.I., II tab. CCC; *L. pinnatifidum*, *Prantl* (not *Sw.*), Morph. d. Gefässcr., II, Schiz., 83.

Primary petioles short or obsolete, secondary ones $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long. Pinnae 20—50 c.M. long, 10—15 c.M. broad. Pinnulae uniform, 9—12 on each side, 5— $7\frac{1}{2}$ c.M. long, to $2\frac{1}{2}$ c.M. broad, articulate at the base, pinnatifid throughout more than half-way down to the costa into close, entire, blunt lobes \pm 3 m.M. broad; upper pinnulae sessile, lower ones short-stalked. Texture subcoriaceous; rachis and surfaces pubescent. The fruit in the fertile pinnulae occupying the back of the scarcely modified lobes. Spores verruculose.

Malacca; Tenasserim.

* (7) **L. scandens**, *Sw.*, Rac., Flor. Btz., I, 8; *Prantl*, Morph. d. Gefässcr., II, Schiz., 81; Bedd., Ferns S.I., tab. LXI; *L. microphyllum*, *R. Br.*, Bedd.; Handb. Ind. Ferns, 455.

Primary petioles very short, secondary ones 1— $2\frac{1}{2}$ c.M. long. Pinnae $7\frac{1}{2}$ —20 c.M. long, with a terminal, simple or bifid pinnula and 4—6 lateral ones on each side. Pinnulae alternate, stalked, articulate at the base, 2— $5\frac{1}{2}$ c.M. long, \pm $1\frac{1}{2}$ c.M. broad, subtriangular or oblong, blunt, simple, entire or rarely slightly lobed, the base rounded, truncate, cordate or bluntly hastate. Texture thin but firm. Fertile pinnulae commonly shorter, subdeltoid, with a mostly rounded apex. Spikes $2\frac{1}{2}$ — $7\frac{1}{2}$ m.M. long, in close rows. Spores reticulate.

Trop. Africa, Asia, Australia, Polynesia.

* (8) **L. salicifolium**, *Pr.*, *Prantl*, Morph. d. Gefässcr., II, Schiz., 79; *L. pinnatifidum*, *Sw.*, Rac., Flor. Btz., I, 7, p. p.

Primary petioles short or wanting. Pinnae pinnate, 12—30 c.M. long, 7—22 c.M. broad, the rachis \pm straight and narrowly marginate, with 4—7 pinnulae on each side below the terminal one. Pinnulae stalked, linear-lanceolate, acute or bluntish, the edge serrulate, the base cordate, truncate or rounded, the stalks nodose & spuriously articulate at the apex. Lowest pinnulae sometimes pinnate with 1—3 leaflets on each side below the terminal one; lateral leaflets small, subrhomboidal, the base cordate on the lower and cuneate on the upper side; terminal leaflets with the base cuneate on both sides. Texture \pm coriaceous; upper surface glossy, naked or slightly hairy on the costae; under surface opaque; rachises rather slightly hairy. Fertile pinnulae similar to the barren ones. Spikes to \pm 4 m.M. long. Spores verruculose.

Trop. Asia.

(9) **L. altum**, v. **A. v. R.**; *L. flexuosum*, Sw., var. *alta*, Clarke, Journ. of Linn. Soc., Bot., XXV, 101, tab. XXIV; Bedd., Ind. Ferns, Suppl., 108.

Scandent, the branches pendant, to 9 M. long, pinnate. Pinnae pinnate again. Pinnulae (both sterile and fertile) \pm 20 c.M. long, \pm 2 c.M. broad, the lateral ones 1—2 on each side, simple, on stalks 1—1½ c.M. long, the edges parallel, entire, the base truncate or auriculate, the terminal ones simple or forked at the base. — Baker esteems this a species.

Malacca; Munipore.

* (10) **L. flexuosum**, Sw., Prantl, Morph. d. Gefässcr., II, Schiz., 72; Bedd., Ferns S.I., tab. LXIII; *L. pinnatifidum*, Sw., Rac., Flor. Btz., I, 7, p. p.

Primary petioles short or wanting, secondary ones to 3 c.M. long. Pinnae 15—30 c.M. long, 15—30 c.M. broad, the rachis \pm flexuose, narrowly marginate, with 2—7 pinnulae on each side below the terminal one. Pinnulae stalked, the terminal and higher ones free or joined, the base cordate, often hastato-auriculate or palmatifid, the edge serrulate, the apex acuminate, acute or blunt; central pinnulae subpalmatifid at the base, the lateral lobes similar to the central one but much shorter; lowest pinnulae similar to the central ones but the basal segments mostly free. Texture herbaceous; surfaces more or less densely hairy, rarely subglabrous; rachises hairy or nearly naked, costae rarely subnodose at the base. Fertile pinnulae similar to the barren ones but shorter, 3—5 c.M. long, the lowest nearly always pinnate with \pm 1—3 leaflets on each side similar to the terminal one but smaller. Spikes 2—10 m.M. long. Spores verruculose.

Trop. & temp. Asia, Australia.

* (11) **L. japonicum**, Sw., Rac., Flor. Btz., I, 8; Prantl, Morph. d. Gefässcr., II, Schiz., 68, tab. I, fig. 10; Bedd., Ferns S.I., tab. LXIV; Christ, Farnkr. d. Erde, fig. 1122.

Primary petioles short, to 1 c.M. long, secondary ones 1—2½ c.M. long. Pinnae 10—20 c.M. long, nearly as broad or narrower, \pm deltoid in outline, pinnate, the rachis \pm flexuose, marginate (narrowly winged) especially towards the apex. Pinnulae \pm 2—5 on each side below the terminal one, not articulate, the higher and terminal ones simple, free or confluent, elongate, linear-lanceolate, the base cuneate or subcordate, the apex bluntish, the edge coarsely crenate or toothed and crenate, or lobed in the lower part; central pinnulae hastate or subpalmatifid, the lowest ones pinnate or 2-pinnate, the rachis marginate, the segments similar to the higher or central pinnulae but

smaller. Texture herbaceous; rachises and surfaces hairy; costae subflexuose, not nodose at the base. Fertile pinnulae like the barren ones but smaller, more deeply incised and commonly so contracted that there is a little or no lamina present. Spikes to 5 or rarely to 10 m.M. long. Spores verruculose.

Trop. and temp. Asia.

11. SCHIZAEA, *Smith.*

Sori linear, consisting of 2 or 4 regular or nearly regular rows of sessile sporangia, covering the inner side of the \pm close branches of the pinnate or digitate, fertile apices of the fronds (or segments of the fronds). Indusium wanting, but the young sori \pm covered by the incurved edges of the fertile segments.

Rhizome creeping or erect. Fronds simple or dichotomous, erect. Small plants, very distinct in habit. Hk. Bk., Syn. Fil., tab. VIII, fig. 64; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 195.

Tropical and extra-tropical.

Arrangement of the sections.

- § 1. **EUSCHIZAEA.** Fertile portion pinnate, barren simple, linear. Sporangia 2-serial.
- § 2. **LOPHIDIUM.** Fertile portion pinnate, barren dichotomous. Sporangia 2-serial.
- § 3. **ACTINOSTACHYS.** Fertile portion \pm digitate, barren simple, linear. Sporangia appearantly 4-serial.

§ 1. **EUSCHIZAEA.**

*. *Fertile spikes 3—6-jugate, the lowest ones 5—7 m.M. long.* (1) *S. malaccana.*

.. *Fertile spikes 10—20-jugate, the lowest ones \pm 3 m.M. long.*

(2) *S. fistulosa.*

(1) **S. malaccana**, *Bk.*, Hk. Bk., Syn. Fil., 428; Bedd., Ferns Br. I., II, tab. CCLV; *S. fistulosa*, *Lab.*, var. *malaccana*, Prantl, Morph. d. Gefässcr., II, Schiz., 136.

Stipes dense, not distinguishable from the fronds, which are 10—20 c.M. long, weak, flexuose, subterete, simply channelled in front, not more than $\frac{1}{2}$ m.M. thick, the barren and fertile ones similar. Fertile portion erect, 8—10 m.M. long, with 5—6 slender, spreading pinnae (spikes) on each side, the lowest 5—7 m.M. long.

Malaya; Burmah.

(2) **S. fistulosa**, *Lab.*, Hk. Bk., Syn. Fil., 429; *Lab.*, Nov. Holl., tab. CCL, fig. 3.

Stipes dense, chestnut-brown, passing gradually into the fronds, which are 10—50 c.M. long, rigid, rush-like, subterete, simply channelled in front. Fertile portion suberect, unilateral, 1— $2\frac{1}{2}$ c.M. long, with 10—20 close, slender, erecto-patent spikes on each side, the lowest \pm 5 m.M. long.

Borneo; Australia, Tasmania, New Zealand, New Caledonia, Fiji, Chili, Madagascar.

§ 2. LOPHIDIUM.

* (3) **S. dichotoma**, *Sm.*, Rac., Flor. Btz., I, 6; Bedd. Ferns S.I., tab. LXV; Hk. & Grev., Ic. Fil., tab. XVII; *S. Forsteri*, *Spr.*, Muell., Pap. Pl., IV, 75.

Rhizome erect, densely covered with lanceolate scales. Stipes dense, 15—45 c.M. long, firm, erect, glossy, channelled on the face above. Fronds fan-like in general outline, $7\frac{1}{2}$ — $22\frac{1}{2}$ c.M. each way, many times dichotomously forked; ultimate divisions flattened, $1\frac{1}{2}$ — $7\frac{1}{2}$ m.M. broad, with the fertile portion pinnate, 4—12 m.M. long, the rachis often curved, with 5—10 close, spreading, hairy pinnae (spikes) on each side.

Trop. Asia, Australia & Polynesia, Madagascar, Mascarenes.

§ 3. ACTINOSTACHYS.

* (4) **S. digitata**, *Sw.*, Rac., Flor. Btz., I, 7; Bedd., Ferns S.I., tab. CCLXVIII; Hk., Gard. Ferns, tab. II.

Rhizome creeping or erect. Stipes dense, $2\frac{1}{2}$ —5 c.M. long, brownish,

subterete, passing gradually into the fronds, which are 20—35 c.M. or more long, 2—5 m.M. broad, flattened, coriaceous, entire, narrowed towards the apex, the midrib prominent beneath, crowned at the apex with 5—15 sub-triquetrous, digitate, linear spikes, which are 2—5 c.M. long, naked beneath, with the edges inflexed.

Trop. Asia & Polynesia.

January 1, 1974

FAMILY VI. POLYPODIACEAE.

VI. POLYPODIACEAE.

Sori definite, marginal, submarginal or dorsal, various in shape, punctiform, round, oblong, reniform or linear, often indefinite and consisting of numerous capsules crowded densely over the under surface or portions of the under surface, occasionally extending \pm on both surfaces of the fronds. Capsules usually stalked, provided with a longitudinal, incomplete annulus, interrupted by the stalk (except in *Plagiogyria*, which has the capsules of *Cyathea*), opening by a transversal stomium. Indusium wanting or present, various in shape and size.

Rhizome creeping or erect, rarely arborescent. Fronds various in shape, size and appearance; veins free or anastomosing variously.

A most extensive family inhabiting almost every part of the world.

Arrangement of the tribes.

A. **Indusiatae.** Indusium present.

§ 1. **WOODSIEAE.** Sori globose, oblong or linear, situated on the back or apex of a vein or veinlet. Indusium inferior, cupuliform, globose or sausage-shaped, fastened all round the receptacle, open at the top or rupturing down irregularly. Stipes not articulate to the rhizome.

§ 2. **ASPIDIEAE.** Sori oblong, round or reniform, medial or terminal on the veins. Indusium superior, of the same shape as the sorus, fastened by a longitudinal axis, a central point or at the top of a \pm deep sinus, opening all round the edge or nearly so. Stipes in most of the genera not articulate to the rhizome.

§ 3. **DAVALLIEAE.** Sori roundish, oblong or linear, rarely medial on the veins but more generally submarginal and then solitary and terminal on the vein tips or uniting the apices of 2 or more veins or veinlets. Indusium \pm of the same shape, attached at the inner side of the sorus, opening outwardly. Stipes whether or not articulate to the rhizome.

§ 4. **PTERIDEAE.** Sori roundish, oblong or linear, submarginal, solitary and terminal on the vein tips, often confluent into a submarginal line

or more commonly uniting the apices of several veins. Indusium marginal, i. e. consisting of the recurved or reflexed, modified edge of the frond or portions of the edge, opening inwardly. Stipes not articulate to the rhizome.

§ 5. **BLECHNEAE.** Sori oblong or linear, distant from the margin, parallel to the central ribs of the free or joined ultimate segments. Indusium of the same shape, not marginal, attached at the outer side of the sorus, opening inwardly. Stipes not articulate to the rhizome.

§ 6. **ASPLENIEAE.** Sori oblong or linear, straight or more or less recurved, rarely horseshoe-shaped, springing from the sides of the veins like the indusia, mostly oblique to the costa, rarely placed between the veins or occupying the single central vein of narrow, ultimate segments. Indusium of the same shape, opening towards the costa or sometimes towards the margin, if double opening towards opposite directions. Stipes not articulate to the rhizome.

B. **Exindusiatae.** Indusium wanting.

§ 7. **PHEGopterideae.** Sori punctiform, round or oblong, rarely more than 2 × as long as broad, medial or terminal on the veins, if medial the opposite pairs sometimes confluent, if terminal sometimes running into a submarginal line or more or less covered by recurved, not modified margin of the frond. Stipes not articulate to the rhizome

§ 8. **Gymnogrammeae.** Sori linear or linear-oblong, more than 2 × as long as broad, several on each side of the costa, mostly oblique to the latter, or running longitudinally and irregularly along the veins, free, joining or ± reticulate. Stipes not articulate to the rhizome.

§ 9. **Grammitideae.** Sori linear, one on each side of the costa, parallel to the latter or the margin, continuous or interrupted. Stipes not articulate to the rhizome.

§ 10. **Polypodieae.** Sori punctiform, round, oblong or linear and then either oblique or parallel to the costa or the margin. Stipes articulate to the rhizome.

§ 11. **Acrosticheae.** Sori indefinite, the sporangia densely crowded over the under surface of the fronds or portions of the under surface or occasionally extending ± on both sides of the fronds or segments, not rarely becoming narrowly linear when the fertile segments are much contracted. Stipes whether or not articulate to the rhizome.

Keys to the tribes and genera.

- 1.** Indusium present, free from the margin or consisting of the reflexed or recurved, *modified* margin or portions of the margin.
- 2.** Indusium cupuliform, globose or sausage-shaped, fastened beneath the sorus on all sides, open at the top or bursting down irregularly.
 - § 1. **Woodsieae.**
 - 2.2.** Indusium oblong, round or reniform, fastened above the sorus, attached by a longitudinal axis, a central point or at the top of a \pm deep sinus, opening all round the edge or nearly so. § 2. **Aspidieae.**
 - 2.2.2.** Indusium roundish, oblong or linear, fastened mostly at 1 side of the sorus, opening at the opposite, free edge.
 - 3.** Indusium marginal, consisting of the reflexed or recurved margin or portions of the margin, opening inwardly. § 4. **Pterideae.**
 - 3.3.** Indusium not marginal, but often reaching the margin with one of the free edges.
 - 4.** Sori *either* roundish and solitary (dorsal or terminal) on the veins, *or* oblong to linear and uniting the apices of 2 or more veins. Indusium attached at least at the inner side of the sorus, opening outwardly. § 3. **Davallieae.**
 - 4.4.** Sori linear or oblong, strictly parallel to the central ribs of the ultimate segments. Indusium attached at the outer side of the sorus, opening inwardly. § 5. **Blechnaeae.**
 - 4.4.4.** Sori linear or oblong, sometimes much recurved or horseshoe-shaped, springing from the sides of the veins, rarely placed between the veins, oblique to the costa, sometimes subparallel to the latter or occupying the single central vein of narrow, ultimate segments. Indusia springing from the same side as the sori, mostly opening obliquely towards the costa. § 6. **Asplenieae.**
- 1.1.** Indusium wanting. Sori sometimes covered by the recurved, *not modified* margin or portions of the margin, often more or less deeply sunk in marginal or dorsal grooves or pit-like, saccate or indusium-shaped cavities or depressions of the frond.
 - a.** Sori indefinite, the sporangia crowded over the under surface or portions of the under surface, not rarely becoming narrowly linear or sometimes extending on both surfaces of the fronds in the often much contracted fertile segments. § 11. **Acrosticheae.**

a.a. Sori definite. Sporangia gathered into sori, which are sometimes contiguous or confluent, but always distinct when young.

b. Fronds articulate to the rhizome. Sori punctiform, round, oblong or linear. § 10. **Polypodiaceae.**

b.b. Fronds not articulate to the rhizome.

c. Sori punctiform, round or oblong, not more than 2 × as long as broad, sometimes running into a submarginal line if terminal on the veins. § 7. **Phegopterideae.**

c.c. Sori linear or linear-oblong, more than 2 × as long as broad, continuous or interrupted, free or reticulate.

d. Sori parallel to the costa or the margin, in 1 line on each side, free. § 9. **Grammitideae.**

d.d. Sori more or less oblique to the costa and the margin, many on each side, free or reticulate.

§ 8. **Gymnogrammeae.**

§ 1. WOODSIEAE.

1. Indusium globose or sausage-shaped, bursting down irregularly.

2. Indusium globose; veins free. No. 12. **Diacalpe.**

2.2. Indusium sausage-shaped; veins anastomosing.

No. 13. **Diplaziopsis.**

1.1. Indusium cup-shaped, open at the top. No. 14. **Dennstaedtia.**

§ 2. ASPIDIEAE.

a. Stipes articulate to the rhizome. Indusium reniform, attached at the top of the sinus.

‡. Fronds simple. No. 15. **Oleandra.**

‡.‡. Fronds pinnate; pinnae articulate to the rachis.

No. 16. **Arthropteris.**

a.a. Stipes not articulate to the rhizome. Indusium oblong, round or reniform; if oblong fastened by its longitudinal axis, if round peltate, fastened by a central point, if reniform attached at the top of its sinus, with the lobes sometimes meeting or overlapping and then apparently orbicular.

b. Veins free.

c. Leaflets very unequal-sided, the upper base broadly cuneate or subtruncate, the lower obliquely cut away.

‡. Indusium oblong, terminal on the veins; leaflets articulate to the rachis. No. 20. **Didymochlaena**.

‡.‡. Indusium round, rarely reniform; leaflets not articulate to the rachis. No. 19. **Polystichum**.

c.c. Leaflets \pm equal-sided and if unequal-sided the lower base not or slightly cut away.

d. Fronds pinnate; pinnae articulate to the rachis; main veins simple or forked. Indusium round or reniform.

e. Indusium round, peltate, without a sinus.

No. 18. **Cyclopeltis**.

e.e. Indusium reniform, sinus present, the lobes sometimes meeting or overlapping and then the indusium becoming round and subpeltate. No. 17. **Nephrolepis**.

d.d. Fronds generally variously divided; leaflets not articulate; main veins pinnate. Indusium reniform, rarely round.

No. 22. **Dryopteris**, § *Lastraea*.

b.b. Veins more or less anastomosing.

1. Lowest veins of contiguous groups forming a costal row of regular, \pm deltoid areolae on each side, the tops of which with an excurrent veinlet, which is either free or joining in the angles of the superior pairs of veins, which often form 1 or more rows of regular, \pm rhomboidal areolae between the costal ones and the margin.

‡. Indusium reniform, rarely round.

No. 22. **Dryopteris**, § *Nephrodium*.

‡.‡. Indusium oblong, \pm 5 \times as long as broad.

No. 23. **Mesochlaena**.

1.1. Veins not anastomosing as in **1**.

2. Lower veins forming 1—2 rows of irregular costal areolae, the rest free or slightly anastomosing near the main veins. Indusium reniform. No. 21. **Pleocnemia**.

2.2. Veins anastomosing copiously and mostly irregularly.

3. Sori alternate, in 3—4 rows on each side of the midrib. Indusium oblong, \pm 5 \times as long as broad. Fronds simple. No. 24. **Luerssenia**.

3.3. Sori scattered irregularly or placed in 2 rows between the main veins. Indusium round or reniform, and if oblong then at best 2 × as long as broad and the fronds not simple.

4. Excurrent soriferous free veinlets 2—5 to each areola. Sori not compital. Indusium round.

No. 26. **Cyrtomium.**

4.4. Excurrent soriferous free veinlets wanting or 1 to each areola, or the soriferous veinlets spreading into various directions. Sori often compital. Indusium round or reniform. No. 25. **Aspidium.**

§ 5. DAVALLIEAE.

a. Sori roundish, dorsal on the veins. No. 27. **Cystopteris.**

a.a. Sori roundish, oblong or linear, submarginal, terminal on the veins (or veinlets, which are sometimes very short and inconspicuous), or uniting the apices of 2 or more veins or veinlets.

b. Fronds divided; leaflets (not the lobes) dimidiate, i. e. the half on the lower side of the *midrib* cut away.

No. 29. **Lindsaya.**

b.b. Fronds simple or divided, if divided the leaflets not dimidiate.

c. Sori roundish or transversally oblong, occupying the truncate apex of the cuneate, ultimate segments.

No. 28. **Odontosoria.**

c.c. Sori occupying the sides, very rarely the apex of not cuneate, ultimate segments.

d. Sori transversally oblong or linear, uniting the apices of 2 or more veins or veinlets.

‡. Fronds simple or divided, if divided the leaflets not articulate to the rachis. No. 50. **Schizoloma.**

‡.‡. Fronds pinnate, leaflets articulate to the rachis. Indusium sometimes 2-valved.

(No. 17. **Nephrolepis**, § *Lindsayopsis*).

d.d. Sori roundish or oblong, solitary on the vein tips, some of them sometimes confluent.

e. Indusium fastened at the base, open at the sides and the outer edge.

1. Sori placed in conspicuously cucullate, marginal outgrowths, which are closed by the \pm flattened, transversally oblong indusia.

No. 31. **Saccoloma** (*sorbifolium*).

1.1. Sori not on marginal outgrowths.

2. Rhizome creeping; stipes scattered, articulate to the rhizome. No. 32. **Humata**.

2.2. Rhizome erect; stipes tufted, not articulate.

No. 55. **Acrophorus**.

e.e. Indusium fastened at the base and sides, open at the outer edge.

f. Rhizome scaly, the scales often hair-pointed at the apex, not minute.

g. Stipes articulate to the rhizome, scattered.

No. 54. **Davallia**.

g.g. Stipes not articulate to the rhizome.

h. Sori 1 to each ultimate segment. Indusium half-cupshaped.

(No. 55. **Asplenium**, § *Loxoscaphe*).

h.h. Sori many to each ultimate segment. Indusium cuneate.

No. 51. **Saccoloma** (*moluccanum*).

f.f. Rhizome hairy, fibrillose or minutely scaly, creeping. Stipes not articulate to the rhizome.

*****. Ultimate divisions unequal-sided, rhomboidal or linear-triangular and then gradually narrowed towards the apex. Receptacle distinct. No. 55. **Microlepia**.

.. Ultimate divisions linear-lanceolate or divided into \pm linear segments, with the broadest part in the centre. Receptacle indistinct.

No. 56. **Tapeinidium**.

§ 4. PTERIDEAE.

1. Indusia bearing and covering the sori. No. 37. **Adiantum.**

1.1. Indusia not bearing, only covering the sori.

2. Sori submarginal, roundish, distant, terminal on the veins, confined to the tops of the sinusses, 1 to each sinus. Indusium roundish, small, hardly covering the sorus at full maturity.

No. 38. **Hypolepis.**

2.2. Sori submarginal, *either* roundish or oblong, close, soon running into a submarginal line *or* linear, uniting the apices of the veins, not confined to the tops of the sinusses. Indusia transversally oblong, approximate, contiguous, confluent or linear.

3. Sori terminal on the veins, free, roundish or oblong, soon confluent into a submarginal line.

4. Base of the stipe abruptly thickened, bearing \pm 3—6 spongy projections on the outer side. Ring oblique, complete, as in Cyatheae. Fertile pinnae narrowly linear.

No. 40. **Plagiogyria.**

4.4. Base of the stipe not abruptly thickened, without projections. Ring vertical, incomplete.

5. Vein tips thickened. No. 39. **Cheilantes.**

5.5. Vein tips not or hardly thickened.

6. Fronds divided; leaflets \pm stalked, not adnate to the rachis, more or less distinctly articulate at the base. No. 41. **Pellaea**, § *Platyloma.*

6.6. Fronds simple or divided; segments adnate or joined, not stalked, not articulate.

No. 42. **Doryopteris**, § *Doryopteridastrum.*

3.3. Sori linear or linear-oblong, the receptacle uniting the tips of the veins, sometimes running them transversally.

a. Stipes with 1—2, free or joining vascular strands.

b. Fronds compoundly divided; lowest pinnulae (secondary pinnae) short, close to the rachis, stipule-like.

No. 45. **Histiopteris.**

b.b. Fronds simple or divided, if divided the lowest pinnulae (secondary pinnae) not stipule-like.

c. Fronds divided; leaflets articulate to the rachis.

No. 41. **Pellaea**, § *Pteridella.*

c.c. Fronds mostly divided; leaflets not articulate to the rachis.

d. Veins free.

e. Spores bilateral, \pm bean-shaped; indusium mostly double. No. 46. **Paesia**.

e.e. Spores globose to tetrahedral; indusium always single.

f. Fronds finely dissected. Sori and indusia \pm meeting against the midribs of the very small and narrow ultimate segments.

No. 44. **Onychium**.

f.f. Fronds not finely dissected. Sori and indusia not meeting. Ultimate segments not very small.

No. 45. **Pteris**.

d.d. Veins anastomosing more or less copiously.

*. Stipes black.

No. 42. **Doryopteris**, § *Eudoryopteris*.

.. Stipes not black. No. 45. **Pteris**.

a.a. Stipes with 3 or more, free vascular strands.

!. Stipes with 2 stout and 1 to several finer strands placed altogether in a circle. Fronds dimorphous, at utmost 2-pinnatifid. No. 48. **Lomaria**.

!!. Stipes with many (12—24) irregularly placed strands. Fronds uniform, at least 2-pinnate at the base.

No. 47. **Pteridium**.

§ 5. BLECHNEAE.

1. Sori superficial.

2. Sori long, linear, 1 on each side of and \pm close to the midribs of the ultimate segments, continuous or interrupted.

3. Fronds dimorphous; fertile pinnae much contracted and the sori filling up nearly the whole space between the margin and the costa. Indusium submarginal. (No. 48. **Lomaria**).

3.3. Fronds not dimorphous; fertile pinnae not contracted; sori distant from the margin like the indusia.

- 4.** Veins of sterile fronds free, those of the fertile ones united near the base by subcostal strands bearing the sori.

No. 49. **Blechnum.**

- 4.4.** Veins of both sterile and fertile fronds united near the base.

No. 50. **Sadleria.**

- 2.2.** Sori oblong, in 1 or more rows on each side of the midribs, not \pm close to the latter.

No. 51. **Doodia.**

- 1.1.** Sori immersed in cavities, oblong, in 1 row on each side of the midribs of the ultimate segments.

No. 52. **Woodwardia.**

§ 6. ASPLENIEAE.

- a.** Sori between the veins; indusia 2 to each sorus, springing from the inner side of the 2 contiguous veins, opening towards one another.

No. 57. **Triphlebia.**

- a.a.** Sori springing from the sides of the veins like the indusia; indusia 1 to each sorus.

- b.** Stipes with 2 \pm flattened vascular strands mostly uniting towards the apex (of the stipe) or higher into a single diarchous one ($\backslash /$, \vee , $\backslash _ /$, \square , \cup). Scales of the rhizome finely cancellate, the areolae mostly dark-coloured.

- c.** Sori and indusia mostly long and straight; the lower sori and indusia of the same group of veins often, the higher ones rarely double, i. e. 2 opposite sori and indusia to a single vein.

No. 53. **Diplazium.**

- c.c.** Sori and indusia mostly short, often curved, sometimes hamate or horseshoe-shaped, the higher ones never, the lower rarely double.

No. 54. **Athyrium.**

- b.b.** Stipes with 2 \pm concave vascular strands mostly uniting towards the apex (of the stipe) or higher into a single tetrarchous one [$)$ (\backslash), $($), $-$ ($>$ — $<$, \times , $\underline{\vee}$, $\underline{\backslash /}$]. Scales of the rhizome coarsely cancellate, the areolae mostly decoloured.

- *. Sori in opposite pairs; indusia opening towards one another.

No. 56. **Phyllitis**

- *.*. Sori not in pairs; indusia opening towards the costa or occasionally towards the margin.

No. 55. **Asplenium.**

§ 7. PHEGOPTERIDEAE.

- 1.** Fronds flabellato-dichotomous. Sori round or punctiform. Veins anastomosing copiously. No. 61. **Dipteris.**
- 1.1.** Fronds variously divided, but not flabellato-dichotomous. Sori round or oblong, if terminal on the veins sometimes running into a submarginal line.
- 2.** Veins free.
- 3.** Sori terminal on the veins.
- 4.** Fertile segments much contracted. Sori at length covering \pm the whole under surface of the segments.
(No. 80. **Polybotrya**).
- 4.4.** Fertile segments not contracted.
- 5.** Spores bilateral.
No. 60a. **Phegopteris**, §§ *Euphegopteris* & *Leptogramme*.
- 5.5.** Spores not bilateral.
- 6.** Fronds hairy, fibrillose or scaly, at utmost 5-pinnatifid. Sori not rarely contiguous and confluent into a submarginal line, often more or less covered by the \pm recurved, hardly modified margin of the frond. Spores globoso-tetrahedral.
No. 58. **Notholaena.**
- 6.6.** Fronds naked, at least 3-pinnate. Sori distant, never confluent, nor covered by the always flattened margin of the frond. Spores 3-lobato-tetrahedral.
No. 59. **Monachosorum.**
- 3.3.** Sori medial or lower on the veins.
No. 60a. **Phegopteris**, §§ *Euphegopteris* & *Leptogramme*.
- 2.2.** Veins anastomosing.
- *. Lower veins of contiguous groups uniting as in Nephrodium.
No. 60a. **Phegopteris**, §§ *Goniopteris*, *Stegnogramme* & *Meniscium*.
- *.*. Veins anastomosing more or less copiously and irregularly.
No. 60b. **Dictyopteris**, § *Arcypteris*.

§ 8. GYMNOGRAMMEAE.

- a.** Veins free.
- b.** Fronds not powdery beneath.
- c.** Pinnae or pinnulae entire or denticulate; main veins close, simple or forked. No. 62. **Coniogramme**.
- c.c.** Pinnae or pinnulae more or less deeply pinnatifid; main veins pinnate in the lobes. (No. 60a. **Phegopteris**, § *Leptogramme*).
- b.b.** At least the fertile fronds coated beneath with a white or yellow powder. No. 63. **Ceropteris**.
- a.a.** Veins uniting 1—5 × at or near the margin.
No. 66. **Syngramme**, §§ *Toxopteris* & *Callogramme*.
- a.a.a.** Main veins and the lower ones of the same group uniting as in *Pleocnemia*. (No. 60b. **Dictyopteris**, § *Digrammaria*).
- a.a.a.a.** Veins anastomosing otherwise.
- 1.** Lower veins of contiguous groups uniting as in *Nephrodium*. Sori free or joining in pairs.
(No. 60a. **Phegopteris**, §§ *Stegnoqramme* & *Meniscium*).
- 1.1.** Veins anastomosing more copiously and commonly irregularly. Sori free or ± reticulate.
- 2.** Sori irregularly longitudinal or sublabellate, running along the veins. Costa wanting, or present in the lower half of the frond only. No. 64. **Antrophyum**.
- 2.2.** Sori oblique or sometimes subparallel to the costa. Costa distinct quite to the apex.
- 3.** Fronds thick, veins all invisible. Sori free, parallel, transversing the areolae. No. 65. **Loxogramme**.
- 3.3.** Fronds thinner, at least the main veins visible. Sori free or reticulate, running along the veins.
- 4.** Sori with paraphyses.
No. 66. **Syngramme**, § *Dictyogramme*.
- 4.4.** Sori without paraphyses. No. 67. **Hemionitis**.

§ 9. GRAMMITIDEAE.

- 1.** Fronds simple, rarely forked or subdichotomous, mostly grass-like. (*Vittariaceae*).

- 2.** Sori on or close to the costa, often protected by a lateral, raised outgrowth of leaf tissue. No. 68. **Monogramme.**
- 2.2.** Sori marginal or submarginal.
- 3.** Sori \pm deeply immersed in mostly 2-lipped, marginal or submarginal grooves or with the margins rolled back over them when young. Midrib present, veins free. No. 69. **Vittaria.**
- 3.3.** Sori not protected as in **3.** Midrib wanting, veins anastomosing. (No. 64. **Antrophyum**, p. p.).
- 1.1.** Fronds pinnate, rarely simple, but then not grass-like. Veins anastomosing. (*Taenitideae*). No. 70. **Taenitis.**

§ 10. POLYPODIEAE.

- a.** Sori punctiform, roundish or oblong, not more than $\pm 3 \times$ as long as broad, many on each side of the costae, sometimes solitary and terminal on the segments.
- b.** At least the under surface covered with scattered or more or less densely crowded stellately branched hairs. No. 72. **Cyclophorus.**
- b.b.** Fronds naked or scaly, not rarely hairy, but then the hairs simple, not stellate.
- c.** Fronds dimorphous. Barren fronds sessile, oakleaf-like, afterwards brown and rigid, fertile ones stipitate, pinnatifid or pinnate. No. 73. **Drynaria.**
- c.c.** Fronds uniform or dimorphous, if dimorphous the barren fronds not as in **c.**
- d.** Fronds deeply pinnatifid. Sori restricted to the higher, much contracted segments.
- e.** Sori oblong or subquadrangular, closely placed in 1 row on each side of and parallel to the costae. No. 74. **Dryostachyum.**
- e.e.** Sori round, placed on distant, semiorbicular lobes, 1 to each lobe. No. 71. **Polypodium** (*Meyenianum*).
- d.d.** Fronds simple or divided. Sori dorsal, marginal or on marginal outgrowths, but not on conspicuously contracted upper segments of divided fronds. No. 71. **Polypodium.**

a.a. Sori linear or linear-oblong, more than 3 × as long as broad, continuous or interrupted.

*. Sori more or less oblique to the costa and the margin, many on each side of the costa. No. 71. **Polypodium**, § *Selliguea*.

.. Sori parallel to the costa or the margin, 1 on each side.

§. Fertile fronds simple as are the barren ones.

No. 75. **Drymoglossum**.

§.§. Fertile fronds 5-partite, barren ones sagittate.

No. 76. **Christopteris**.

§ 11. ACROSTICHEAE.

1. Fronds dimorphous, the fertile ones dichotomously branched, the barren ones adpressed to the substratum. No. 77. **Platyserium**.

1.1. Fronds uniform or dimorphous, the fertile ones not dichotomous, whether or not contracted, the barren ones not as in **1**.

2. Veins free, or sometimes the apices or bases of the veins united by ± marginal or ± costal lines.

3. Fronds simple, often more or less dimorphous.

No. 78. **Elaphoglossum**.

3.3. Fronds divided, the fertile ones contracted.

4. Rhizome high-scandent, cable-like. Fertile fronds Lomaria-like, with the veins united at the base by costal lines or spurious veins. No. 79. **Stenochlaena**.

4.4. Rhizome erect or creeping, neither high-scandent nor cable-like. Fertile fronds not Lomaria-like, without costal lines or spurious veins. No. 80. **Polybotrya**.

2.2. Veins forming 1—2 rows of areolae against the costae and costulae (as in *Pleocnemia*), otherwise free. No. 81. **Stenosemia**.

2.2.2. Veins anastomosing more copiously, and generally irregularly.

a. Fertile region restricted to the linear or spoon-shaped apex of simple fronds. No. 82. **Hymenolepis**.

a.a. Fertile region restricted to the higher segments of pinnatifid or pinnate fronds.

b. Pinnae free, not articulate to the rachis, ± uniform or the fertile ones rarely linear. No. 83. **Acrostichum**.

b.b. Pinnae free or joining at the base, articulate to the rachis, dimorphous, the fertile ones linear.

No. 84. **Photinopteris.**

a.a.a. Fertile region not restricted as in *a.* or *a.a.* Fronds dimorphous, barren and fertile ones separate.

§. Barren fronds simple or more commonly 2-fid, the lobes divaricating, sometimes 2-fid again. Each segment with 2 or more, stout, longitudinal ribs springing from the base of the frond.

No. 85. **Cheiropleuria.**

§.§. Barren fronds simple or pinnate. Each segment with a single costa.

*. Spores oblong, ovate or 2-lateral.

No. 86a. **Leptochilus.**

.. Spores globoso-tetrahedral. No. 86b. **Platytaenia.**

TRIBE I. WOODSIEAE.

12. DIACALPE, *Blume.*

Sori globose, dorsal on a vein or veinlet, arising from a small, scarcely elevated receptacle. Indusium inferior, globose, sessile, at first entire and enclosing the whole sorus, at length bursting irregularly at the summit. Capsules numerous, nearly sessile.

Rhizome erect. Stipes not articulate to the rhizome. Fronds much divided; veins free. Sori usually 1 to each ultimate segment, on the lowest acropetal veinlet. Hk. Bk., Syn. Fil., tab. I, fig. 7; Diels, in Engl. & Prantl, Nat. Pfl.Fam. I⁴, fig. 87, A—D.

Tropical Asia.

* ***D. aspidioides*, Bl.**, Rac., Flor. Btz., I, 119; Bedd., Ferns S. I., tab. CCLVII; Bau. & Hk., Gen. Fil., tab. IC; *D. pseudo-caenopteris*, Kze, Bot. Zeit., 1846, 457; — var. *membranulosa*, Christ.

Rhizome erect, covered with desiccated stipes and brownish scales. Stipes 15—40 c.M. long, scaly. Fronds deltoid, 25—50 c.M. long, ± as broad, 3—4-pinnate. Pinnæ not quite opposite, deltoid-lanceolate, stalked, to 10 c.M. broad at the base. Ultimate segments obovate, 5—8 m.M. long, 2—5 m.M. broad, sessile, the base narrowed, ± decurrent, the edge crenate or toothed throughout or at the apex only. Texture submembranaceous; surfaces nearly naked, or sparsely chaffy on the veins; colour disposed to blacken in drying. Sori solitary at the base of the ultimate segments; indusium firm.

Var. membranulosa: Fronds smaller; pinnæ close; segments contiguous; indusium weak.

Trop. Asia.

13. DIPLAZIOPSIS, *Christensen.*

Sori dorsal, linear or oblong, running from near the costa to the first areola; indusium sausage-shaped, fastened all round the receptacle, quite enclosing the sorus when young, rupturing irregularly when the spores mature.

Rhizome short. Stipes not articulate to the rhizome. Fronds simply pinnate; veins anastomosing, forming 2—5 rows of \pm hexagonal areolae near the margin. Hk. Bk., Syn. Fil., tab. IV, fig. 39; Diels, in Engl. & Prantl. Nat. Pfl.Fam., I⁴, fig. 121, E. (*Allantodia*).

Paleotropical.

* **D. javanica**, *C. Chr.*, Ind. Fil.; *Allantodia javanica*, *Trev.* (not *Bl.*), *Rac.*, *Flor. Btz.*, I, 252; *A. Brunoniana*, *Wall.*, *Bedd.*, *Ferns S.I.*, tab. CLIX; *Bau. & Hk.*, *Gen. Fil.*, tab. CXX; *Asplenium javanicum*, *Bl.* (oldest name).

Rhizome erect, covered with lanceolate scales. Stipes 15—35 c.M. long, slightly scaly at the base. Fronds oblong-lanceolate, 50—100 c.M. long, 18—35 c.M. broad, pinnate. Pinnae sessile or short-stalked, linear, the apex acuminate, the base truncate, the longest ones 10—15 c.M. long, 2—4 c.M. broad, the margin entire or irregularly crenate. Texture membranaceo-herbaceous; surfaces naked; veins anastomosing, forming several rows of long and short, nearly hexagonal areolae near the margin. Sori confined to the anterior veinlets of the primary forkings.

North. India, Australasia, Polynesia.

14. DENNSTAEDTIA, *Bernhardi*.

Sori submarginal, terminal on the veins, \pm globose; indusium inferior, united with the margin of the frond to form a distinctly cup-shaped, membranaceous involucre to the sorus, the top open, sometimes slightly 2-lipped or 2-valved.

Rhizome mostly creeping, often hairy, rarely erect or arborescent. Stipes not articulate to the rhizome or the caudex. Fronds rather large, at least 2-pinnate, superficially resembling *Dicksonia*; veins free. Hk. Bk., Syn. Fil., tab. II, fig. 13, c (as *Patania*); Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 117, C—E.

Pantropical.

- a.** *Fronde neither scandent nor twining.*
- b.** *Fronde 2-pinnate or 3-pinnatifid.*
- §.** *Pinnulae rhomboidal, entire or lobed.* (1) *D. rhombifolia.*
- §.§.** *Pinnulae lanceolate, pinnatifid.* (2) *D. resinifera.*
- b.b.** *Fronde 3-pinnate.*
- * *Rachis naked; surfaces glaucous.* (5) *D. erythrorachis.*
- ** *Rachis pubescent; surfaces naked.* (4) *D. ampla.*
- *** *Rachis and under surface tomentoso-glandular.* (5) *D. Smithii.*
- b.b.b.** *Fronde at least 4-pinnatifid.*
- c.** *Rachis more or less hairy.*
- d.** *Rhizome creeping. Fronde 4-pinnatifid.*
- e.** *Ultimate lobes entire.* (6) *D. scabra.*
- e.e.** *Ultimate lobes crenato-pinnatifid.* (7) *D. samoensis.*
- d.d.** *Caudex arborescent. Fronde 4-pinnatifid. Lobes toothed.* (8) *D. javanica.*
- c.c.** *Rachis naked.*
- 1.** *Fronde to 90 c.M. broad.*
- 2.** *Texture herbaceous.*
- 3.** *Rachis orange-coloured, slightly scabrous.* (9) *D. remota.*
- 3.3.** *Rachis stramineous, smooth.* (10) *D. cuneata.*
- 2.2.** *Texture coriaceous.* (11) *D. Kingii.*
- 1.1.** *Fronde more than 90 c.M. broad.*
- §.** *Texture herbaceous.* (12) *D. flaccida.*
- §.§.** *Texture coriaceous.*
- * *Stipes ebeneous, muricate.* (15) *D. glabrata.*
- ** *Stipes brownish, smooth.* (14) *D. sumatrana.*
- a.a.** *Fronde scandent or twining.*
- † *Rachis aculeate. Texture rather papyraceous.* (15) *D. scandens.*
- †.† *Rachis aculeate. Texture coriaceous.* (16) *D. moluccana.*
- †.†.† *Rachis smooth.* (17) *D. gomphophylla.*

(1) **D. rhombifolia**, *C. Chr.*, Ind. Fil.; *Dicksonia rhombifolia*, *Bk.*, Journ. of Bot., XXVIII, 105.

Stipe and rachis slender, naked, castaneous. Fronde oblong-lanceolate, 2-pinnate, to 30 c.M. long. Pinnae sessile, lanceolate, cut down to the rachis

into oblong-rhomboidal pinnulae 6—9 m.M. long, 4—5 m.M. broad, which are truncate at the base on the lower side. Both surfaces naked; veins subflabellate. Sori solitary at the upper margin of the pinnulae; indusium cup-shaped, firm.

New Guinea.

(2) ? **D. resinifera**, *Mett.*, Ann. Mus. Bot. L. B., IV, 290. *Cheilanthes resinifera*, *Bl.*, Enum., 158; *Hypolepis resinifera*, *Hk.*, Spec. Fil., II, 65.

Stipes rough. Fronds 2-pinnatifid towards the apex, 3-pinnatifid at the base. Pinnulae sessile, lanceolate, acuminate, deeply pinnatifid, the higher ones confluent. Segments oblong, obtuse, inciso-crenate, subauriculate at the upper base. Texture subcoriaceous; main rachis tomentose above; under surface resinoso-punctate. Sori solitary at the inner edge of the crenulations; indusium dentiform.

Java.

(3) **D. erythrorachis**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 218; *Dicksonia erythrorachis*, *Christ*, Ann. Bot., XV, 86, tab. XIII, fig. 7, a—b.

Stipes erect, polished like the rachises, slightly aculeate or muricate below. Fronds large, deltoid-ovate. Pinnae spreading, straight, the lower ones to \pm 50 c.M. long. Pinnulae linear-lanceolate, acuminate, sessile, to \pm 6 c.M. long and to \pm 2 c.M. broad, rather approximate, the lower part cut down to the rachis into 5—8 close ultimate leaflets, the apex pinnatifid. Leaflets ovate-rhomboidal, erecto-patent, slightly lobed, bearing the sori in the sinuses. Texture rigid; surfaces glaucous, the under one paler coloured than the upper one. Sori small, 1—3 to a leaflet; indusium brownish, glossy, subglobose.

Celebes, New Guinea.

(4) **D. ampla**, *Bedd.*, Journ. of Bot., XXXI, 227; *Dicksonia ampla*, *Bl.* Journ. of Linn. Soc., Bot., XXII, 223.

Rhizome creeping, clothed with dark-coloured scales. Fronds \pm 120 c.M. long or longer, deltoid, 5-pinnate. Pinnae oblong-lanceolate, the lowest ones 60—90 c.M. long, \pm 50 c.M. broad. Pinnulae lanceolate, multijugate, cut down nearly or quite to the costa into oblong, parallel, adnate, close ultimate segments 4—5 m.M. broad. Texture moderately firm; rachis brown, unarmed, pubescent, canaliculate; surfaces naked; costulae pinnate; veins 6—8 on each

side, the lower ones forked, the upper ones simple. Sori small; the outer side of the indusium large, naked, semiorbicular.

Malacca, Borneo.

(5) **D. Smithii**, *Moore*, Ind., 308; *Dicksonia Smithii*, *Hk.*, Hk. Bk., Syn. Fil., 54; Hk., Spec. Fil., I, tab. XXVIII, D.

Lower pinnae 18—50 c.M. long, 7—10 c.M. broad. Pinnulae linear, acuminate, cut down to the rachis throughout into several or more numerous, narrow, obovate or oblong segments. Lower segments (leaflets) $\pm 12\frac{1}{2}$ m.M. long, ± 5 m.M. broad, the base more or less obliquely cuneate, the edges slightly inciso-pinnatifid. Rachis and under surface tomentoso-glandular; texture \pm coriaceous. Sori 2—8 to a leaflet; indusium subglobose, cup-shaped.

Celebes, Philippines.

* (6) **D. scabra**, *Moore*, Christ, Farnkr. d. Erde, 312; *D. deltoidea*, *Moore* Bedd., Ferns S.I., tab. CCLVIII; *Dicksonia scabra*, *Wall.*, Hk. Bk., Syn. Fil., 54; Hk., Spec. Fil., I, tab. XXVIII, B; *Dick. deltoidea*, *Hk.*, l. c., tab. XXVIII, A; *Davallia celebica*, *Christ*, Fil. Saras., IV, 426.

Rhizome creeping. Stipes scattered, reddish, more or less hairy. Fronds 20—80 c.M. long, deltoid or lanceolate. Pinnae lanceolate, the largest 10—20 c.M. or more long. Pinnulae ± 4 c.M. long, 1—2 c.M. broad, cut down nearly or quite to the rachis into several, pinnatifid, oblong or lanceolate-rhomboidal segments. Texture herbaceous; rachis and under surface more or less hairy. Sori small, 2—6 or more to a segment, terminal on the lobes.

Malaya; India, Ceylon, China, Japan, Formosa.

(7) **D. samoensis**, *Moore*, Ind., 307; *Dicksonia samoensis*, *Bk.*, Hk. Bk., Syn. Fil., 462; *Dick. delicata*, *Muell.*, Pap. Pl., IV., 77.

Fronds ample, 4-pinnatifid; rachises thinly pilose. Lower pinnae 30 c.M. or more long, lanceolate-deltoid. Pinnulae and segments close, lanceolate-deltoid, distinctly stalked, cuneate-truncate on lower side at the base, quaternary lobes close, oblong, blunt, $1-2\frac{1}{2}$ m.M. broad, crenato-pinnatifid. Texture membranaceous; ribs beneath hairy. Sori minute, placed at the base of the ultimate sinuses; valves of the indusium equal, hemispherical, glabrous. Rhizome creeping?

New Guinea; Polynesia.

* (8) **D. javanica**, *Christ*, Bull. Herb. Boiss., 2^e Série, IV, 617; *Dicksonia javanica*, *Bl.*, Rac., Flor. Btz., I, 120; C. Chr., Ind. Fil.

Caudex arborescent. Stipe and rachis densely covered with short, adpressed hairs. Fronds lanceolate-oblong, to $1\frac{1}{2}$ M. long, to 60 c.M. broad. Pinnae lanceolate, stalked, erecto-patent, $\pm 10-12\frac{1}{2}$ c.M. broad, the stalks $1-1\frac{1}{2}$ c.M. long. Pinnulae short-stalked, linear-lanceolate, $\pm 2-3$ c.M. broad at the base, the apex acuminate. Tertiary divisions sessile or subsessile, the lowest anterior largest, linear-oblong, $1\frac{1}{2}-2$ c.M. long, to ± 5 m.M. broad, the apex blunt, toothed, the edge cut down nearly or quite to the costa into oblong, blunt, inciso-dentate ultimate segments, the lowest anterior segments the largest. Texture herbaceous; upper surface nearly naked, the under one densely hairy like the secondary and tertiary rachises. Sori small, near the ultimate sinuses; annulus long, the extremities separated by a whorl of radiating, slightly thickened cells with the stalk of the sporangium placed in the centre.

Java.

(9) **D. remota**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 218; *Dicksonia remota*, *Christ*, Ann. Btz., XV, 86, tab. XIV, fig. 8, a—b.

Fronds 3-pinnate to 4-pinnatifid. Pinnae to ± 45 c.M. long and 15 c.M. broad, stalked. Pinnulae lanceolate, acuminate, alternate, ± 5 c.M. apart, short-stalked, ± 2 c.M. broad at the base. Ultimate divisions oblong-ovate, short-stalked, the lower ones rather distant, ± 5 m.M. apart, cuneate at the base, pinnatifid from $\frac{1}{2}$ -way down to quite to the costa, the upper ones more close and less deeply lobed, or even entire. Texture herbaceous but firm; rachis naked, orange-coloured, slightly scabrous. Sori intramarginal, lateral on the lobes, 1—6 to a pinnula, transversally oblong; indusium broadly ovate. Veins conspicuous, pinnate in the lobes.

Celebes.

* (10) **D. cuneata**, *Moore* (not *Christ*), Copel., Polypod. Philipp., 57; *Dicksonia cuneata*, *Hk.*, Hk. Bk., Syn. Fil., 462; *Hk.*, Spec. Fil., I, tab. XXVIII, C.

Fronds ample, subdeltoid, 4-pinnatifid; rachis stramineous, naked. Pinnae lanceolate, 15—55 c.M. long. Pinnulae close, short-stalked, lanceolate, $1-1\frac{1}{2}$ c.M. broad. Tertiary divisions oblong-rhomboidal, $2\frac{1}{2}-5$ m.M. broad, sessile, the base cuneate, more or less cut away on lower side, the edge inciso-pinnatifid. Texture moderately firm; both surfaces naked; veins subflabellate. Sori at the tops of the ultimate sinuses; indusium cup-shaped, naked. — Perhaps this may be the description of a single pinna regarded

by Hooker as a complete frond. A specimen gathered by Mr. Whitford has the fronds 5-pinnatifid, the pinnulae ± 25 (15—35?) c.M. long, etc., the ultimate rachises marginate or rather narrowly winged.

Philippines, Batjan.

(11) **D. Kingi**, *Bedd.*, Journ. of Bot., XXXI, 227; *Dicksonia Kingi*, *Bedd.*, Handb. Ind. Ferns, Suppl., 6.

Stipes stout, erect, ± 45 c.M. long. Fronds 90—120 c.M. long, deltoid-lanceolate, 4-pinnatifid. Pinnae 20—25 c.M. long, the lower ones deltoid-lanceolate, lanceolate to linear-lanceolate upwards. Lower pinnulae ± 6 —7 c.M. long, the others $\pm 2\frac{1}{2}$ c.M., cut down nearly or quite to the rachis into oblong, broadly toothed lobes. Texture coriaceous; surfaces naked. Sori terminal on thickened veins and exserted beyond the margin.

Malacca.

* (12) **D. flaccida**, *Bernh.* (not *Moore*), Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 218; *Dicksonia flaccida*, *Sw.*, Rac., Flor. Btz., I, 121.

Rhizome creeping. Stipes 1—2 M. long. Fronds 3—4-pinnate, ovate-deltoid, to $2\frac{1}{2}$ M. long, to $\pm 1\frac{1}{2}$ M. broad. Pinnae linear, opposite, short-stalked or nearly sessile, lanceolate-acuminate, scarcely narrowed towards the base. Pinnulae sessile, to 8 c.M. long, to 2 c.M. broad, the base \pm truncate. Tertiary divisions close, unequal-sided, to 1 c.M. long, to 4 m.M. broad, pinnatifid $\frac{1}{2}$ — $\frac{3}{4}$ -way down to the costa, the apex toothed, the base decurrent, the lobes crenate or toothed. Texture herbaceous; both surfaces naked. Sori at the tops of the sinuses; indusium subglobose when young, cup-shaped at full maturity.

Java, Sumatra; Polynesia.

(13) **D. glabrata**, *C. Chr.*, Ind. Fil.; *Dicksonia glabrata*, *Ces.*, Fel. Becc. Polin., 2 & 6.

Rhizome tortuose, the scales linear, fibrillose, brown, crisped. Stipe ± 40 c.M. long, rigid, polished, ebeneous like the rachis, the base densely mucronate, the upper part rarely so, the prickles conical, recurved. Fronds ± 150 c.M. long, 3-pinnate. Pinnae spreading, the lower ones ± 60 c.M. long, divided as in *D. flaccida*, *Bernh.* The lower lobes with the margin often undivided. Texture coriaceous; upper surface dark-green, the under one paler.

New Guinea.

* (14) **D. sumatrana**, v. *A. v. R.*

Fronds large, 3-pinnate; stipe and rachis stout, brownish, naked. Pinnæ erecto-patent, 85 c.M. long or longer, \pm 15 c.M. broad at the base. Pinnulae linear-lanceolate, acuminate, \pm 2 c.M. broad at the base, the apex pinnatifid, the rachis very narrowly winged. Tertiary divisions oblong, bluntly rounded, \pm spreading, \pm 5 m.M. broad, the margin cut $\frac{1}{3}$ — $\frac{2}{3}$ -way down, the upper base truncate, slightly auricled, the lower cuneate and decurrent, the lobes close, bluntish or rounded. Texture coriaceous; surfaces naked; veins rather visible. Sori marginal, 1 at the top of each sinus; indusium distinctly cupuliform. — This plant was determined by Baker as a variety of *Davallia Speluncae*, *Bk.*, (Journ. of Bot., XVIII, 210).

Sumatra (Beccari, No. 436).

* (15) **D. scandens**, *Moore*, Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 218; *Dicksonia scandens*, *Bl.*, Rac., Flor. Btz., I, 122; *Dick. Zippeliana*, *Kze*, Bot. Zeit., 1845, 858.

Rhizome creeping. Fronds to 8 M. long, distant, at least 3-pinnate, scandent or twining; main rachis aculeate. Pinnæ opposite, distant, to 1 M. long or longer, scandent, the rachis aculeate. Pinnulae \pm sessile, lanceolate or oblong, acuminate or bluntish, cut down to the rachis into deltoid-oblong or unequally ovate leaflets \pm 2 c.M. long, blunt, the base narrowed, \pm decurrent, the edge more or less deeply pinnatifid into several roundish, blunt, entire or crenate lobes. Texture semicoriaceous or membranaceous, rather flaccid; surfaces naked, or slightly hairy on the veins. Sori small, roundish, near the sinuses; indusium cupuliform.

Malaya; Samoa.

(16) **D. moluccana**, *Moore*, Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 218; *Dicksonia moluccana*, *Bl.*, Hk. Bk., Syn. Fil., 53.

Near *D. scandens*, *Moore*, but firmer. Ultimate leaflets less deeply lobed, the lobes deltoid-oblong. Texture coriaceous. Indusium subglobose.

Malaya; Formosa, Fiji.

(17) **D. gomphophylla**, *C. Chr.*, Ind. Fil; *Dicksonia gomphophylla*, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 223.


Fronds large, naked, deltoid, decomposed; rachises unarmed. Lowest pinnae largest, 45—60 c.M. long, \pm 50 c.M. broad, deltoid, inequilateral. Pinnulae oblong-lanceolate. Tertiary divisions multijugate, oblong, quater-

nary ones multijugate, sessile, to $\frac{3}{4}$ c.M. long, with the upper base parallel to the rachis, the lower cut away, obtuse, deeply pinnatifid. Veins sublabelate. Sori small, placed at the tops of the sinuses of the ultimate lobes: the outer valve of the indusium large, semiorbicular. -- Habit of *D. scandens* & *moluccana*, *Moore*.

Borneo.

TRIBE II. ASPIDIEAE (*)

(*) The genera **Dryopteris** and **Aspidium** have the indusia not rarely rudimentary or caducous and then it is difficult to distinguish the species from **Phegopteris** and **Dictyopteris**.



15. OLEANDRA, *Cavanilles*.

Sori on the back of the veins, rarely \pm medial, generally near the costa. Indusium reniform, firm, fixed by the sinus.

Rhizome creeping, scandent or erect, branched. Fronds simple, articulate to the rhizome; veins free, compact. Hk. Bk., Syn. Fil., tab. V, fig. 46; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 109 (¹).

Pantropical.

-
- 1.** *Rhizome scandent or erect* (²).
- 2.** *Sori medial on the veins.* (1) *O. cuspidata*.
- 2.2.** *Sori near the costa.*
- 3.** *Rhizome scales ciliate, the ciliae cobwebby.* (2) *O. ciliata*.
- 3.3.** *Rhizome scales at best ciliate when young, the ciliae short.*
- 4.** *Stipes articulate at the apex, the joint close to the base of the lamina.* (3) *O. colubrina*.
- 4.4.** *Stipes articulate below the apex or towards the base, the joint not close to the base of the lamina.* (4) *O. neriiformis*.
- 1.1.** *Rhizome creeping.*
- a.** *Costa naked or hairy beneath.*
- b.** *Stipes articulate at the base.*
- c.** *Sori somewhat distant from the costa.* (5) *O. musifolia*.
- c.c.** *Sori close to the costa.* (6) *O. Wallichii*.
- b.b.** *Stipes articulate above the base or towards the apex.*
- §.** *Rhizome scales \pm adpressed. Stipes articulate between the base and the apex.* (7) *O. Cumingii*.
- §.§.** *Rhizome scales spreading. Stipes articulate at the summit.* (8) *O. tricholepis*.
- a.a.** *Costa scaly beneath.* (9) *O. Whitmeei*.
-

(¹) Most of the species are more or less variable and so closely allied that they might be considered as varieties of 1 or 2 species united by several intermediate forms, which are often difficult of determination.

(²) See also *O. musifolia*, *Pr.*

(1) **O. cuspidata**, *Bk.*, Becc., Mal., III, 44.

Rhizome stout, woody, copiously branched, densely clothed with small, adpressed, castaneous, ovate, firm scales, which are pale-coloured at the margin. Stipes short, articulate at the summit. Fronds ligulate, cuspidate, 10—15 c.M. long, narrowed gradually from the middle to the base. Texture rigid; surfaces naked; veins conspicuous. Sori in 1 row on each side of the costa, not below the middle of the veins; indusium small, firm, reniform, persistent.

New Guinea.

(2) **O. ciliata**, *Hl.*, Linn., XXXVI, 126.

Rhizome scandent; the scales ovate and adpressed at the base, the point subulate, spreading, the edge finely ciliate, the ciliae cobwebby. Stipes 8—15 m.M. long, 5—10 m.M. apart. Fronds to \pm 40 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad, acuminate, narrowed gradually at the base. Texture membranaceous or subchartaceous; upper surface naked, \pm glossy, under one covered with scattered hairs. Sori not contiguous, in 1 row on each side near the costa; indusium reniform, both sides unequally rounded, coriaceous, reddish-brown.

Amboyna; New Hebrides, Pinorum Island.

* (3) **O. colubrina**, *Copel.*, Polypod. Philipp., 48; *Blechnum colubrinum*, Blanco, Flor. Filip., 834; — var. *brachypus*; *O. neriiformis*, Cav., var. *brachypus*, Hk., Spec. Fil., IV, 156.

Rhizome woody, stout, erect or suberect; the scales dense, adpressed, the base obovate, peltate, dark-coloured, the margin brown, \pm ciliate when young, the apex acuminate, erect or spreading, deciduous or persistent. Stipes mostly clustered, articulate immediately below the lamina. Fronds \pm 15—20 c.M. long, \pm 2—3 c.M. broad, widening towards the upper part, then narrowed and caudate or acuminate, narrowed gradually towards the base, the edge ciliate with whitish or brownish hairs. Texture papyraceous to coriaceous; costa and surfaces \pm conspicuously clothed with rather long, soft, slightly crisped hairs. Sori in 1 irregular row on each side near the costa.

Var. brachypus: Costa and surfaces pubescent or sparingly villose. *Philippines, Borneo, Java.*

* (4) **O. neriiformis**, *Cav.*, Hk. Bk., Syn. Fil., 302; Bedd., Ferns of Br.I., II, tab. CCLXIV; *O. neriifolia*, Pr. (not Cav.), Rac., Flor. Btz., I, 203, p. p.; — var. *mollis*; *O. mollis*, Pr., Epim., 41; *O. hirtella*, Miq., Kze, Farnkr., II, tab. CXXIX; *O. pilosa*, Hk., Bau. & Hk., Gen. Fil., tab. XLV, B.

Rhizome erect or climbing; the scales dense, adpressed, the base obovate, peltate, dark-coloured, the margin brown, \pm ciliate when young, the apex acuminate, erect or spreading, deciduous or persistent. Stipes scattered or placed in opposite pairs or distant whorls, \pm 1—4 c.M. long, naked, articulate between the apex and the base, often below the middle, the joint in every case remote from the base of the lamina. Fronds 15—40 c.M. long, 1—4 c.M. broad, linear-lanceolate, acuminate or subcaudate, narrowed gradually towards both ends. Texture papyraceous to coriaceous; surfaces naked or partially villous or pubescent on the costa and veins. Sori in 1 mostly irregular row on each side near the costa; indusium oblique, entire or ciliate.

Var. mollis: Fronds more villous with soft, fine hairs.

Malaya; Polynesia, South. America, West. Africa.

* (5) **O. musifolia, Pr.** (not *Kze.*), Hk. Bk., Syn. Fil., 502; *O. neriiiformis, Bedd.* (not *Cav.*), Ferns S.I., tab. XCI; Hk., Fil. Exot., tab. LVIII; *O. neriiifolia, Pr.* (not *Cav.*), Rac., Flor. Btz., I, 203, p. p.

Shoots firm, creeping, scandent or intertwined, clothed with lanceolate-subulate, ciliate, yellow scales, which are dark-coloured and peltate at the base. Stipes scattered, generally single, rarely in whorls or in tufts, \pm 2 $\frac{1}{2}$ —5 c.M. long, naked, jointed close to the base. Fronds elongate-lanceolate, acuminate, 20—50 c.M. long, to \pm 5 c.M. broad. Texture thin-membranaceous to firm-papyraceous; surfaces naked or hairy on the veins and midrib. Sori in 1 irregular row on each side near the midrib.

Malaya; Ceylon.

* (6) **O. Wallichii, Pr.** (not *Hk.*), Hk. Bk., Syn. Fil., 503; *Bedd.*, Ferns Br.I., II, tab. CCLXV; *Aspidium Wallichii, Hk.*, Exot. Flor., I, tab. V.

Rhizome creeping, densely clothed with subulate, ferruginous, often reflexed scales. Stipes distant or sometimes more close, 1—8 c.M. long, sometimes paleaceous, articulate at the base. Fronds 15—40 c.M. long, sub-elliptical-oblong, the sides parallel, the base often obtuse or rounded, the apex shortly and sharply acuminate. Texture soft, membranaceous; surfaces more or less hairy. Sori compact, almost imbricate, in a continuous line on each side and close to the costa; indusium reniform. — Costa naked or scaly.

Malacca; North. India.

* (7) **O. Cumingii, J. Sm.**, Hk. Bk., Syn. Fil., 503; *Bedd.*, Ferns Br.I., I, tab. CXXXV; — var. *Sibaldii, Hk. Bk.*, l. c.; *O. Sibaldii, Grev.*, Ann. Nat. Hist., II¹, 527.

Rhizome horizontally creeping, clothed with \pm adpressed, subulate, ciliate scales. Stipes close or scattered, $2\frac{1}{2}$ —20 c.M. long, articulate within 1—7 c.M. from the base. Fronds 20—45 c.M. long, narrowed gradually towards both ends. Texture herbaceous or membranaceous; midrib and both surfaces slightly hairy. Sori in regular rows very near the midrib; indusium firm, oblique.

Var. Sibaldii: Texture thinner; surfaces more hairy; sori in very irregular, wavy lines, rather distant from the midrib.

Malaya; South. India, Assam, Centr. China, — var. Sibaldii: Sumatra, Java, Timor; Tahiti.

(8) **O. tricholepis**, *Hæe*, Bot. Zeit., 1851, 549.

Rhizome creeping, the scales spreading, subulate, reddish-brown, large. Stipes placed in distant tufts, elongate, slender, articulate with the fronds. Fronds cuneate at the base, slightly hairy. Sori unknown.

Borneo.

* (9) **O. Whitmeei**, *Blk.*, Journ. of Bot., XIV, 11; Engl., Bot. Jahrb., XXIII, tab. V, fig. 2—4; *O. ornata*, *Christ*, Fil. Saras., III, 240.

Rhizome wide-creeping, densely clothed with yellow, spreading, subulate scales. Stipes naked, brown, articulate about the middle. Fronds to \pm 30 c.M. long, 4—5 c.M. broad, narrowed gradually towards the base, the apex finely acuminate. Texture thin, the edge thickened, finely ciliate; upper surface pubescent especially on the midrib; under side of the midrib regularly clothed with 2 opposite rows of spreading, yellow, subulate scales. Sori close, placed in 1 wavy row on each side near the midrib, sometimes in 2 close, irregular rows; indusium pale-yellowish, often subpeltate.

Celebes; Philippines; Samoa.

16. ARTHROPTERIS, *J. Smith.*

Sori roundish, terminal on the anterior veinlets of the once forked veins. Indusium roundish-reniform, fixed by the sinus.

Rhizome creeping or scandent. Stipes articulate to the rhizome. Fronds pinnate; pinnae articulate to the rachis.

A small genus, very near *Nephrolepis*, distinguished from this by the articulated stipes.

Paleotropical.

- *. Surfaces \pm hairy. Indusium fugacious. (1) *A. obliterata*.
 *. *. Surfaces naked. Indusium persistent. (2) *A. glabra*.

* (1) ***A. obliterata***, **J. Sm.**, C. Chr., Ind. Fil.; *A. ramosa*, **Mett.** (not **J. Sm.**), **Copel.**, Polypod. Philipp., 46; *Nephrolepis ramosa*, **Moore**, **Rac.**, Flor. Btz., I, 201; *N. obliterata*, **J. Sm.**, Journ. of Bot., IV, 197; **Bedd.**, Ferns S.I., tab. CCLI; *Nephrodium obliteratum*, **R. Br.**, (oldest name).

Rhizome wide-creeping, slender, branched, scaly. Stipes scattered, short, \pm 2 c.M. long. Fronds 15—45 c.M. long, $2\frac{1}{2}$ —8 c.M. broad, gradually but slightly narrowed towards both ends; rachis slightly flexuose. Pinnae \pm 12—20 on each side, \pm sessile, 1—2 c.M. broad, obliquely rhomboidal, the apex bluntish or subacute, sometimes rounded, the edges entire or more or less distinctly crenate especially in the fertile pinnae, the upper base truncate and auricled, the lower base obliquely narrowed. Texture thin or more or less coriaceous; rachis and both surfaces hairy, sometimes scaly. Sori 1 to each crenation, at a distance of 1—2 m.M. from the edge; indusium roundish or subcordate, fugacious.

Paleotropical.

(2) ***A. glabra***, **Copel.**, Polypod. Philipp., 46.

Rhizome wiry, climbing \pm 5 M. high, sparsely clothed with black, adpressed scales. Stipes \pm 5 m.M. long, articulate 1—2 m.M. beyond the base. Fronds 20—50 c.M. long, 5—7 c.M. broad, pinnate, with a terminal pinna rather larger than the others. Lateral pinnae \pm 8 m.M. broad, obtuse or the higher ones subacute, obscurely crenate, slightly auricled, the upper base abruptly truncate, the lower very oblique. Texture papyraceous; surfaces naked; remaining green when dried. Sori roundish, \pm medial, nearer the margin than the costa; indusium fixed by a very narrow sinus, persistent.

Philippines.

17. NEPHROLEPIS, **Schott.**

Sori *either* round or reniform, submarginal or medial, in 1 row on each side of the costa, terminal on the anterior veinlets of the once forked veins or transversally oblong to linear, marginal, uniting the apices of 2 or more veins.

Indusia of the former sori reniform, attached by the sinus, the lobes sometimes meeting or overlapping and then the indusia becoming round and \pm peltate, opening all round the edge or nearly so, those of the latter ones 2-valved & marginal or transversally oblong to linear & submarginal, attached at the inner side of the sori, opening outwardly, as in *Lindsaya*.

Rhizome erect, creeping or scandent. Stipes adherent to the rhizome. Fronds pinnate; pinnae articulate to the rachis.

Tropical and subtropical.

Arrangement of the sections.

- § 1. **LINDSAYOPSIS.** Sori transversally oblong or linear, marginal or submarginal, uniting the apices of 2 or more veins. Indusium 2-valved & marginal or transversally oblong to linear & submarginal, attached at the inner side of the sorus, opening outwardly, as in *Lindsaya*.
- § 2. **EUNEPHROLEPIS.** Sori round or reniform, intramarginal. Indusium reniform, attached by the sinus, opening all round the edge or nearly so. Hk. Bk., Syn. Fil., tab. V, fig. 45; Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 111.

§ 1. **LINDSAYOPSIS.**

- §. *Fertile pinnae lobed, the lobes small, transversally oblong, forming with the equal-shaped indusia 2-valved involucre to the sori.* (1) *N. dicksonioides*.
- §.§. *Fertile pinnae entire or broadly crenate. Indusia quite as in Lindsaya.*
- *. *Sori and indusia continuous along the margin of the pinnae.*
(2) *N. acutifolia*.
- *.*. *Sori and indusia interrupted especially towards the apex of the pinnae.*
(5) *N. lindsayae*.

(1) **N. dicksonioides**, *Christ*, Fil. Saras., III, 241; Farnkr. d. Erde, fig. 907, n, o, p & q; *Dicksonia nephrolepioides*, *Christ*, l. c.

Fronds pinnate, 1—2 M. long, \pm 20—25 c.M. broad. Pinnae linear, sessile, acuminate, \pm spreading, \pm 1 $\frac{1}{2}$ c.M. broad, the base rounded, truncate or slightly cordate; the higher and lower pinnae commonly sterile, entire, the

central ones fertile, longer and narrower than the barren ones, with the edges shortly and regularly undulate or lobed, the apices of the lobes, which are broader than deep, dilated. Texture \pm coriaceous; veins of the barren pinnae nearly straight, parallel, close, simple or forked, those of the fertile ones oblique, curved, branched in the lobes. Sori marginal, 1 to each lobe, uniting the 2—5 branches of each vein; indusia 2-valved, firm-membranaceous, the outer valves consisting of the lobes of the pinnae.

Celebes.

(2) **N. acutifolia**, *Christ*, Farnkr. d. Erde, 291, fig. 907, r, s & y; *Isoloma lanuginosum*, *J. Sm.*, Bau. & Hk., Gen. Fil., tab. CII; *Lindsaya lanuginosa*, *Hk.*, Hk. Bk., Syn. Fil., 109; *Hk.*, Spec. Fil., I, tab. LXIX, B; *Bedd.*, Ferns Br.I., I, tab. CXL; *L. acutifolia*, *Desv.*, (oldest name).

Rhizome stout, creeping, clothed with fibrillose scales. Stipes erect, stout, 10—15 c.M. long. Fronds 50—60 c.M. long, 7—10 c.M. broad. Pinnae linear-oblong, nearly equilateral, 1—1½ c.M. broad, the base truncate, slightly auricled above, slightly rounded below, the edge entire or slightly toothed towards the apex, which is acute in the fertile, \pm blunt in the barren pinnae. Texture coriaceous; rachis pubescent. Sori and indusia in continuous lines along the edges, as in *Lindsaya*.

Malaya; Burmah, Australia, Trop. Africa.

* **N. lindsayae**, *Christ*, Bull. Herb. Boiss., VI, 837.

Rhizome nodose. Stipes tufted, clothed like the rachis with soft, crisp-ed, filiform scales. Pinnae numerous, close, straight or falcate, unequal-sided, 4 or more c.M. long, \pm 1 c.M. broad, the younger ones slightly tomentose, the sterile ones entire, the fertile ones broadly crenate; the upper base truncate, auricled, the apex \pm gradually narrowed, bluntish or acuminate. Veins numerous, close. Sori 6—8 on each side, often forming a continuous line interrupted towards the apex of the pinna; indusia of the same shape, submarginal, as in *Lindsaya*.

Sumatra, Halmaheira.

§ 2. EUNEPHROLEPIS.

a. Pinnae very small, roundish or flabelliform.

(4) *N. Duffi.*

a.a. Pinnae not very small, varying from oblong to linear, entire or crenate, rarely lobed, pinnatifid or dichotomous.

- b.** *Pinnae (at least the fertile ones) cuneate at the base* (*).
- *. *Fertile pinnae crenate.* (5) *N. floccigera.*
- *.*. *Fertile pinnae more or less deeply and copiously lobed.*
(6) *N. acuminata.*
- b.b.** *Pinnae cordate, truncate, subtruncate or rounded at the base.*
- c.** *Pinnae auriculate at the base on the lower side.* (7) *N. celebica.*
- c.c.** *Pinnae not auriculate at the base, or auriculate on the upper or on both sides.*
- d.** *Surfaces and especially the rachis densely clothed with fimbriate, woolly or plushy scales.* (8) *N. hirsutula.*
- d.d.** *Surfaces and rachis naked or not copiously scaly.*
- e.** *Fully developed fronds mostly under 8 c.M. broad. Pinnae blunt or bluntish.*
- f.** *Fronds to \pm 20 c.M. long; pinnae \pm 3 m.M. broad; indusium subpeltate.* (9) *N. Lauterbachii.*
- f.f.** *Fronds more than 20 c.M. long; pinnae 5—15 m.M. broad; indusium reniform, with a distinct sinus.*
- g.** *Rhizome \pm erect, short, the roots often bearing tubers. Fronds densely tufted; pinnae imbricating with a cordate base.* (10) *N. cordifolia.*
- g.g.** *Rhizome wide-scandent, bearing distant tufts of few fronds; roots without tubers. Pinnae \pm rounded or subtruncate at the base.*
(11) *N. radicans.*
- e.e.** *Fully developed fronds 8 c.M. broad or broader. Pinnae acute or acuminate.*
- §.** *Sori submarginal; pinnae to \pm 7½ c.M. long, distinctly auricled.*
- *. *Indusium orbicular, the lobes overlapping.*
(12) *N. barbata.*
- *.*. *Indusium reniform, with a distinct sinus.*
(13) *N. exaltata.*
- §.§.** *Sori rather distant from the margin; pinnae mostly more than 7½ c.M. long, not or slightly auricled.*
(14) *N. biserrata.*

(*) See also under **N. biserrata**, Schott.

* (4) **N. Duffi**, *Moore*, Nich., Dict. d'Hort., III, 461, fig. 632.

Stipes tufted, erect, chaffy like the rachis, 10—20 c.M. long. Fronds \pm 25—55 c.M. long, linear, 1—1½ c.M. broad, narrowed gradually towards the base, the apex once or more times forked. Pinnae numerous, the lower ones reduced, distant, the central ones approximate, each consisting of 2 oblong or flabellato-orbicular, strongly imbricated leaflets, which are joined at the very base, with the edges entire or irregularly crenate to slightly lobed. Texture herbaceous; under surface clothed with few, scattered, minute, chaffy scales; veins of each leaflet flabellate in the orbicular, pinnate in the oblong forms.

Malaya; North. Australia.

* (5) **N. floccigera**, *Moore*, Hk. Bk., Syn. Fil., 502; Christ, Farnkr. d. Erde, fig. 907, h & i; — var. *monstruosa*, hort.

Stipes furfuraceous. Fronds 60—90 c.M. long, 15—30 c.M. broad. Pinnae 1—2 c.M. broad, the apex acuminate, the edge crenate to a depth of \pm 1½ m.M., the base slightly unequal, \pm cuneate. Texture subcoriaceous; rachis and under surface furfuraceous. Sori submarginal, 1 to each crenation; indusium firm, reniform.

Var. monstruosa: A much cultivated form, with the pinnae very broad, deeply and irregularly divided into numerous, close, imbricate, linear segments with entire or toothed edges.

Java, Celebes, Amboyna.

* (6) **N. acuminata**, *Kuhn*, Ann. Mus. Bot. L. B., IV, 286; *N. davalloides*, *Kze*, Rac., Flor. Btz., I, 202; Hk., Fil. Exot., tab. LX; *Ophioglossum acuminatum*, *Houtt.*, (oldest name).

Caudex short, stoloniferous, densely clothed with black-brown, lanceolate scales. Stipes tufted, \pm 30 c.M. long, \pm scaly at the base. Fronds \pm 60—100 c.M. long, \pm 20—55 c.M. broad, hardly narrowed towards the base, sterile in the lower part. Barren pinnae narrow-lanceolate, 1—3 c.M. broad, the base slightly unequal, the edge crenate, the point acute; fertile pinnae mostly narrower, the highest ones often very narrow, linear, the base \pm cuneate, the edge more or less deeply lobed sometimes nearly to the costa, the sinuses \pm rounded, the lobes blunt. Texture coriaceous; rachis \pm furfuraceous; surfaces \pm naked; veins forked at the base, the barren branch the shortest, often very short. Sori terminal on the lobes, 1 to each lobe.

Malaya.

(7) **N. celebica**, *Zoll.*, Syst. Verz., 44, 49.

Pinnae linear-lanceolate, ensiform, sessile, obtuse, the edge obscurely repand, the base cordate, reniform-auriculate below; the lower pinnae growing shorter. Surfaces naked; colour dark-green; rachis short-hairy. Sori minute.

Celebes.

* (8) **N. hirsutula**, *Pr.*, Rac., Flor. Btz., I, 201; *N. rufescens*, *Wawra* (not *Pr.*), Copel., Polypod. Philipp., 47; *N. acuta*, *Pr.*, var. *rufescens*, *Christ*, Ann. Btz., XV, 140.

Rhizome erect, stout, densely clothed with dark-coloured, small, adpressed scales, stoloniferous. Stipes tufted, \pm 15—60 c.M. long, firm, glabrescent, the base \pm scaly like the rhizome. Fronds 30—120 c.M. long, 8—25 c.M. broad. Pinnae more or less close, not imbricate, 1—1½ c.M. broad, fertile linear, \pm acute, generally serrate or crenato-dentate, often with a minute tooth in the sinuses of the crenations, rounded at the base on lower side, the upper base usually prominently auricled; sterile pinnae shorter than the fertile ones and less acute. Texture herbaceous or subcoriaceous; surfaces (and especially the rachis densely) clothed with \pm deciduous, limbrate, woolly or plushy, pale-brown or reddish scales. Sori submarginal or intramarginal; indusium often becoming orbicular. — *N. rufescens*, *Wawra*, is the broadest form with the sori distant from the margin, united by numerous intermediates with the type, which is the narrowest form with the sori submarginal.

Pantropical.

(9) **N. Lauterbachii**, *Christ*, Bull. Herb. Boiss., 2^e Série, I, 456; *N. cordifolia*, *Pr.*, var. *Lauterbachii*, *Christ*, in *Schum. & Laut.*, Flor. deut. Sch.ggeb., 118.

Near *N. cordifolia*, *Pr.* but the pinnae shorter, narrowly lanceolate, \pm 5 m.M. broad, serrulate or denticulato-crenate. Fronds hardly 20 c.M. long. Sori submarginal; indusium \pm peltate.

New Guinea.

* (10) **N. cordifolia**, *Pr.*, Hk. Bk., Syn. Fil., 300; *N. tuberosa*, *Pr.*, Rac., Flor. Btz., I, 199; *Bedd.*, Ferns S.I., tab. XCII; *Polypodium cordifolium*, *L.*, (oldest name).

Rhizome densely clothed with hyaline, yellow-brown, lanceolate scales; the roots often bearing tubers. Stipes tufted, wiry, 2½—29 c.M. long, glossy, more or less densely scaly. Fronds linear-lanceolate, 25—100 c.M. long,

3—8 c.M. broad. Pinnae numerous, often imbricated, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, the base cordate, or rounded on the lower side, more or less auricled on the upper side, the edge entire or more or less crenated chiefly in the fertile pinnae, the apex \pm narrowed, bluntish. Texture subcoriaceous; rachis yellowish, glossy, slightly scaly; surfaces nearly naked, the upper one often with a row of submarginal, cretaceous dots, the under one sometimes densely villose. Sori submedial, nearer the edge than the midrib; indusium firm, reniform, persistent, often very oblique.

Pantropical, New Zealand, Japan.

* (11) **N. radicans**, *Kuhn*, Ann. Mus. Bot., L. B., IV, 285; *N. volubilis*, *J. Sm.*, Christ, Farnkr. d. Erde, 288; Clarke, Trans. Linn. Soc., Bot., 2^d Series., I, tab. LXXVIII; *Polypodium radicans*, *Burm.*, (oldest name).

Rhizome scandent, indefinite, naked, wiry, bearing distant tufts of \pm 2—3 fronds at the ends of short, scaly branches. Stipes \pm 10 c.M. long, firm, naked or nearly so. Fronds 25—50 c.M. long, \pm 3—6 c.M. broad; rachis firm, scarcely chaffy. Pinnae \pm 5—8 m.M. broad, very obtusely rounded, nearly entire, the upper base slightly auricled, the lower rounded. Texture coriaceous, surfaces naked or the costa minutely scaly above. Sori near the margin; indusium reniform, with a deep sinus.

Trop. Asia.

(12) **N. barbata**, *Copel.*, Polypod. Philipp., 48.

Rhizome short, suberect. Stipes approximate, suberect, strong, 10—15 c.M. long, sparsely scaly at the base, the scales lanceolate, reddish-brown, with a hyaline margin. Fronds linear, 50 c.M. or more long, \pm 8 c.M. broad, pinnate; rachis brown, glossy, clothed in the lower part with minute, scattered, spreading scales. Sterile pinnae \pm $3\frac{1}{2}$ c.M. long, \pm 1 c.M. broad, acute, entire, fertile ones \pm 5 c.M. long, \pm 7 m.M. broad, acuminate, often serrate towards the apex, subfalcate; the lower base rounded, the upper one acutely auricled; lower pinnae reduced, roundish. Texture subcoriaceous; surfaces naked. Sori submarginal; indusium orbicular, black, with a brown margin, the lobes overlapping.

Philippines.

* (15) **N. exaltata**, *Schott*, Rac., Flor. Btz., I, 200; Schott, Gen. Fil., tab. III; Bedd., Ferns S.I, tab. XCHH; — var. *monstruosa*, hort.

Rhizome creeping, bearing distant, short branches with densely tufted

fronds. Stipes 10—15 c.M. long, naked or slightly scaly, the base clothed with narrow, adpressed, linear-lanceolate scales like the rhizome. Fronds 20—80 c.M. long, 8—15 c.M. broad. Pinnae close or the lower ones rather distant, not imbricate, bluntish or subacute, the upper base auricled, the lower rounded or auricled, the edge entire, crenate or serrulate; fertile pinnae linear, acute, often \pm falcate and longer than the barren ones. Texture subcoriaceous; surfaces naked or nearly so. Sori submarginal; indusium reniform, with a deep sinus.

Var. monstrosa is a much cultivated form with 1—3 \times forked pinnae.

Pantropical.

* (14) **N. biserrata**, *Schott*, C. Chr., Ind. Fil.; *N. acuta*, *Pr.*, *Rac.*, Flor. Btz., I, 202; *Bedd.*, Ferns S.I., tab. XCIV; *Aspidium acutum*, *Schk.*, *Muell.*, Pap. Pl., IV, 79; *N. acuminata*, *Pr.*, Tent. Pterid., 79; *N. punctulata*, *Pr.*, l. c.; *Aspidium biserratum*, *Sw.*, (oldest name); — var. *laurifolia*, *Christ*, *Farnkr. d. Erde*, 289, fig. 907, d & e.

Caudex short, erect, stoloniferous, densely clothed with lanceolate, brown, entire or more or less irregularly ciliate scales. Stipes tufted, 10—50 c.M. long, scaly at the base, glabrescent above. Fronds 30—400 c.M. long, 15—40 c.M. broad, pendulous if epiphytical. Pinnae linear-lanceolate, short-stalked or sessile, 1—5 c.M. broad, separated by less than their own breadth, the point acute, the edge entire when barren, crenate, toothed or repand when fertile, the base suddenly narrowed or truncate, not or slightly auricled on the upper on both sides, the lower base sometimes rounded. Texture herbaceous or subcoriaceous; rachis and costae naked or slightly scaly; surfaces naked or nearly so. Sori generally distant from the margin; indusium subreniform, rather becoming orbicular by the meeting of the 2 broad, basal lobes.

Var. laurifolia: Pinnae broad, ovate-oblong or broadly linear-lanceolate, the edges subentire or sometimes irregularly inciso-lobate towards the base, the base truncate, rounded or broadly cuneate. Sori medial between the midrib and the margin.

Pantropical; — var. *laurifolia*: *in a few malayan islands.*

18. CYCLOPELTIS, *J. Smith.*

Sori round, placed in 1—4 rows on each side of the midrib. Indusium peltate.

Rhizome oblique, decumbent. Stipes not articulate to the rhizome. Fronds pinnate; pinnae articulate to the rachis; veins free, erecto-patent, forked or branched, bearing the sori dorsal or sometimes terminal. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 96, A—B.

Tropical Asia and America.

*. *Pinnae with the lower base not hamato-recurvate.*

(1) *C. semicordata.*

.. *Pinnae with the lower base hamato-recurvate.*

(2) *C. Presliana.*

* (1) **C. semicordata, J. Sm.**, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 183, fig. 96, A; *Aspidium semicordatum, Sw.*, Hk. Bk., Syn. Fil., 249; *Polystichum semicordatum, Moore*, Bedd., Ferns Br.I., I, tab. XXXV; — var. *crenata*, Hk., Spec. Fil., IV, 17; — var. *subdimorpha*, Christ, in Warb., Mons., I, 76; — var. *Kingii*, Christ in Ann. Bot., XV, 126; *A. Kingii, Hance.*

Rhizome strong; the scales subulate, dense. Stipes approximate or scattered, 10—30 c.M. long, fibrillose at the base. Fronds 60—100 c.M. long, 20—50 c.M. broad, oblong, acuminate, simply pinnate. Pinnae close, articulate, not rarely contiguous, sessile or subsessile, 1—2 c.M. broad, \pm falcate, acute or acuminate, the edge nearly entire, the base truncate or cordate. Texture papyraceo-herbaceous or subcoriaceous; surfaces naked; rachis more or less scaly; veins pinnate, the lower ones ending short from the margin. Sori in 1—4 rows on each side, the inner row the most constant and regular, often close to the midrib or nearly so.

Var. crenata: Superior base of the pinnae broadly cuneate, lower one variable.

Var. subdimorpha: Pinnae of fertile fronds distant, very narrow.

Var. Kingii: Small form. Pinnae blunt, the base bluntly rounded. Sori in 1 row.

Malacca; Burmah, Trop. America; — var. *crenata*: *Borneo*; — var. *subdimorpha*: *Celebes*; — var. *Kingii*: *Celebes; Caroline Islands.*

* (2) **C. Presliana**, *Berk.*, C. Chr., Ind. Fil.; *Aspidium Preslianum*, *Mett.*, *Farnng.*, IV, *Phegop. & Aspid.*, 56; *Polystichum Preslianum*, *Moore*, *Copel.*, *Polypod. Philipp.*, 16; — var. *biauriculata*.

Like *C. semicordata*, *J. Sm.*, but: Stipes clothed with long, spreading scales, which are brown and subulate. Pinnae numerous, distant or approximate, spreading, the edge entire, crenato-sinuate or subrepand, highest pinnae often confluent with the terminal one, the upper base subcordate or rounded to slightly auricled, the lower base conspicuously auricled with an oblong, rounded, hamato-recurvate lobe which is adpressed on the rachis.

Var. biauriculata: Upper base with a conspicuous, deltoid, \pm deflexed auricle, which has the apex obliquely truncate.

Malaya; Burmah, Tonkin; — var. *biauriculata*: *New Guinea* (Peekel).

19. POLYSTICHUM, *Roth.*

Sori roundish, dorsal or terminal on the veins. Indusium normally round and peltate, exceptionally reniform and attached by the sinus.

Rhizome creeping or erect. Stipes not articulate to the rhizome. Fronds at least pinnate, usually tough or rigid, the leaflets mostly unequal-sided, i. e. the lower side less developed, or the upper base broadly cuneate to truncate and often auricled, the lower one obliquely cut away; the edges commonly mucronate or spinulose; veins free. *Hk. Bk.*, *Syn. Fil.*, tab. V, fig. 43, a—b; *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 99.

Tropical and extra-tropical.

a. *Fronds simply pinnate, at best 2-pinnatifid at the base.* (1) *P. auriculatum*.

a.a. *Fronds at least 2-pinnate at the base.*

b. *Main rachis densely scaly or fibrillose.*

c. *Texture membranaceous or herbaceous.* (2) *P. diaphanum*.

c.c. *Texture rather coriaceous.*

d. *Pinnulae distinct, short-stalked, not joined at the base.*

e. *Scales of stipe and rachis ferrugineous.*

f. *Teeth aristate or spinulose.* (3) *P. aculeatum*.

f.f. *Teeth blunt or adpresso-mucronate.* (4) *P. obtusum*.

e.e. *Principal scales of stipe and rachis \pm black.*

(5) *P. biaristatum*.

- d.d.** Pinnulae decurrent, joined at the base. (6) *P. lobatum*.
- b.b.** Main rachis naked, or slightly scaly or fibrillose.
- 1.** Sori submarginal. (7) *P. amabile*.
- 1.1.** Sori distant from the margin.
- 2.** Sori costal. (8) *P. platypus*.
- 2.2.** Sori \pm medial.
- 3.** Lowest pinnae deeply 2-pinnatifid at the base on the lower side. (9) *P. varium*.
- 3.3.** Lowest pinnae at least 2-pinnate at the base on the lower side.
- 4.** Rhizome creeping; the scales hair-like or \pm setiform. (10) *P. aristatum*.
- 4.4.** Rhizome erect; the scales long, linear-lanceolate. (11) *P. coniiifolium*.

(1) **P. auriculatum**, *Pr.*, Tent. Pterid., 85; *Aspidium auriculatum*, Sw., Hk. Bk., Syn. Fil., 251; — var. *nervosum*, Christ, in Bull. Herb. Boiss., VI, 192; *Phegopteris nervosa*, Fée, Mém., VI, tab. II, fig. 4.

Stipes tufted, 10—15 c.M. long, scaly below or throughout. Fronds 30—45 c.M. long, 5—10 c.M. broad. Pinnae numerous, subsessile, usually close, 2½—5 c.M. long, 1—1½ c.M. broad, ovate-rhomboidal, falcate, the point acute, the edge spinoso-serrate, not lobed; the upper base with a prominent auricle, the lower one truncate. Texture subcoriaceous; rachis stramineous, slightly scaly. Sori in 2 rows.

Var. nervosum: Stipes decumbent; pinnae small, crenate, not aristate; sori scattered, small; indusium hardly present.

Malacca; Ceylon, Br. India; — var. nervosum: Philippines; Formosa.

* (2) **P. diaphanum**, *Moore*, C. Chr., Ind. Fil.; *Aspidium diaphanum*, Zoll. & Mor., Rac., Flor. Btz., I, 175.

Rhizome short, erect, densely clothed with ferruginous, lanceolate, acuminate, ciliate scales. Stipes 5—6 c.M. long, densely scaly. Fronds \pm 10—15 c.M. long, \pm 2½—5½ c.M. broad. Pinnae sessile or the lower ones stalked, oblong-rhomboidal, the base obliquely cuneate, the edge toothed to lobed and toothed. Lowest pinnae pinnate with few, small, obovate pinnulae at the base. Texture thin; rachis densely scaly; the edges and both surfaces hairy; midribs wavy. Sori 1—6 on each side of the costae.

Java.

* (5) **P. aculeatum**, *Schott* (not *Roth*), *Copel.*, *Polypod. Philipp.*, 17; *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 100, A; *Aspidium aculeatum*, *Sw.*, *Hk. Bk.*, *Syn. Fil.*, 252; — var. *vulcanicum*; *A. vulcanicum*, *Bl.*, *Enum.*, 165; — var. *subamoenum*, *Christ*, in *Warb., Mons.*, I, 77; — var. *mucronifolium*; *A. mucronifolium*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 172; — var. *moluccense*; *A. moluccense*, *Bl.*, l. c., 168; — var. *batjanense*, *Christ*, l. c.; *A. batjanense*, *Christ*, *Bull. Herb. Boiss.*, 2e Série, VI, 926; *P. horizontale*, *Copel.*, l. c., 18.

Rhizome erect or suberect, strong, scaly. Stipes tufted, scaly, 10—100 c.M. long; the scales of 2 forms, the smaller ones slender, resembling hairs or fibrils, the larger ones ovate-lanceolate, thin, both ferruginous. Fronds 50—80 c.M. long, 18—40 c.M. broad, 2-pinnate, sometimes 5-pinnate, ovate-lanceolate to deltoid, acuminate. Pinnae close, numerous, ± linear-lanceolate, the lowest commonly the longest, 2—4 c.M. broad near the base. Pinnulae subrhomboidal, close, regular, the upper base truncate, ± auricled, the lower base cuneate, the margin toothed, the apex and the teeth mucronate or aristate. Texture ± coriaceous; rachis densely, ± deciduously scaly, the scales linear-lanceolate, ferruginous, acuminate, often mixed with white or reddish, ± hair-like smaller ones; under surface naked or slightly fibrillose. Sori in 1 row on each side, ± medial or subcostal. — A variable species. —

a. *Pinnulae at best pinnatifid, the auricle not free.*

b. *Pinnulae not pinnatifid.*

Forma typica: Pinnulae aristate. Sori ± medial or subcostal, not solitary. — *Cosmopolitan.*

Var. vulcanicum: Pinnulae rather small, subimbricate, spinuloso-dentate, obscurely auricled. Sori solitary. — *Java, Sumatra.*

b.b. *Pinnulae incised.*

Var. subamoenum: *Philippines.*

a.a. *Pinnulae more or less divided, at least the auricle ± free.*

Var. mucronifolium: Under surface slightly paleaceo-pilose; pinnulae small, at best once pinnate. — *Malaya; Br. India.*

Var. moluccense: Under surface sparingly furfuraceous; pinnulae once pinnate. — *Moluccas, Philippines.*

Var. batjanense: The whole plant densely clothed with cinnamon-coloured scales; pinnulae large, ± 5 c.M. long, the lowest pinnate or 2-pinnatifid. — *Batjan, Philippines.*

* (4) **P. obtusum**, *J. Sm.* (not *Pr.*), *Copel.*, *Polypod. Philipp.*, 17; *Aspidium obtusum*, *Kze* (not *Mett.*), *Hk. Bk.*, *Syn. Fil.*, 252, *Mett.*, *Farg.*, IV,

Phegop. & Aspid., 51; Hk., Spec. Fil., IV, tab. CCXXI; *A. aculeatum*, Sw., var. *obtusum*; Christ, Bull. Herb. Boiss., VI, 192; — var. *densum*; *A. densum*, Zipp., MS. in Herb. Bog.

Stipes 10—15 c.M. long, scaly; the scales copious, ferrugineous, ovate, long-pointed, soft, mixed with narrow, subulate ones which continue upon the rachis, becoming there more or less fibril-like, both forms ciliate. Fronds \pm 20—50 c.M. long, ovate-lanceolate, acuminate. Lower pinnae distant, subspreading, 5—15 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad. Pinnulae rhomboid-ovate, obtusely auriculate, obtuse, the teeth blunt or adpresso-mucronate. Texture coriaceo-membranaceous; rachis densely & deciduously scaly or fibrillose; under surface more or less scaly, the scales small, partially filiform, partially stellato-fimbriate. Sori \pm medial; indusium peltate, thin, fringed at the margin. — Very near *P. aculeatum*, Schott.

Var. densum: Fronds longer and broader, copiously 2-pinnate.

Luzon; — var. *densum*:? (Zippelius).

* (5) **P. biaristatum**, Moore, Ind., 86; *P. aculeatum*, Schott, var. *biaristatum*, Bedd., Handb. Ind. Ferns, 209; C. Chr., Ind. Fil.; *Aspidium biaristatum*, Bl., Enum., 164; Hk., Spec. Fil., IV, 29; *A. amabile*, Rac. (not Bl.), Flor. Btz., I, 172.

Stipes tufted, 20—40 c.M. long, very paleaceous below with long, subulate, ciliate, castaneous scales (sometimes having a black, central line) mixed with large, broad-lanceolate, intensely black ones, which are brown at the margin, the rest of the stipe and the rachis clothed with more or less deciduous and copious, black-brown, crinite scales, at length often glabrous. Fronds 30—60 c.M. long, from a broad base ovate-oblong, suddenly acuminate at the apex. Pinnae sessile or subsessile, 10—20 c.M. long, acuminate, serrate towards the apex, $2\frac{1}{2}$ —4 c.M. broad at the base. Pinnulae subrhomboidal, acute, serrate, obsoletely auriculate, the serratures mucronato-spinose. Texture coriaceous; under surface paleaceo-pilose. Sori medial or \pm submarginal; indusium rather small, peltate. — ? Different from *P. vestitum*, Pr. (*A. vestitum*, Sw., Mett. Farng. IV, Phegop. & Aspid, 45).

Malaya; *Br. India*.

(6) **P. lobatum**, Pr., Tent. Pterid., 83; Bau. & Hk., Gen. Fil., tab. XLVIII, C; *Aspidium lobatum*, Sw., Christ, Farnkr. d. Erde, 258; — var. *discretum*, Christ, in Ann. Btz., XV, 125; *A. discretum*, Don, Prod. Flor. Nepal., 4; *P. discretum*, Diels in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 191.

Like *P. aculeatum*, *Schott*, but the pinnulae decurrent, confluent at the base. Fronds lanceolate, narrowed towards both ends.

Var discretum: Fronds deltoid or triangular. Pinnae cut down to $\frac{1}{3}$ -way to the costa in the outer half.

Europe and South. Africa to Japan; Hawaii; — var. discretum: Malaya; North. India, Japan.

* (7) ***P. amabile*, J. Sm.**, *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, 195; *Aspidium amabile*, *Bl.*, *Hk. Bk., Syn. Fil.*, 254; *Hk., Spec. Fil.*, IV, tab. CCXXV; *Lastraea amabilis*, *Moore*, *Bedd., Ferns S.I.*, tab. CIX.

Rhizome creeping, the scales linear-lanceolate, ferruginous. Stipes scattered, 15—50 c.M. long, slightly scaly below, slender, glossy. Fronds \pm 50 c.M. long, 15—25 c.M. broad, with a long terminal pinna and 2—6 similar lateral ones on each side. Pinnae erecto-patent, the lower ones 10—20 c.M. long, 2—4 c.M. broad, sometimes slightly compound at the base. Pinnulae subrhomboidal, 1—2 $\frac{1}{2}$ c.M. long, $\frac{1}{2}$ —1 c.M. broad, the upper and outer edge cut $\frac{1}{3}$ — $\frac{1}{2}$ -way down to the costa into rounded lobes which terminate into a long bristle or a narrow, sharp tooth; the upper base subtruncate, the lower one obliquely cut away. Texture subcoriaceous; rachis and surfaces naked. Sori submarginal; indusium sometimes reniform.

Malaya; North. India, South. China, Japan.

(8) ***P. platypus*, Diels**, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, 195; *Nephrodium platypus*, *Hk.*, *Hk. Bk., Syn. Fil.*, 286; *P. aristatum*, *Pr., var., C. Chr., Ind. Fil.*

Stipes 30—90 c.M. or more long, smooth, glossy, with a dense tuft of bright-coloured scales at the base. Fronds 45—60 c.M. long, 50 c.M. and more broad, deltoid. Lower pinnae 15—25 c.M. long, 7—10 c.M. broad. Pinnulae deltoid, obliquely truncate at the lower base. Lower segments often free, $\frac{1}{2}$ —1 $\frac{1}{2}$ c.M. broad, ovate-oblong, the teeth mucronate. Texture sub-coriaceous; rachis and both surfaces naked and glossy. Sori large, copious, in rows close to the midribs; indusium reniform.

Java; Moulmein, Khasia.

(9) ***P. varium*, Pr.**, *Copel., Polypod, Philipp.*, 19; *Aspidium varium*, *Sw.*, *Hk. Bk., Syn. Fil.*, 254; *Hk., Spec. Fil.*, IV, tab. CCXXVI.

Rhizome short, subrepent. Stipes 50—60 c.M. long, densely fibrillose below. Fronds 30—50 c.M. long, 20—30 c.M. broad, lanceolate-deltoid. Lowest pin-

nae much the largest, subdeltoid, unequal-sided, 7—10 c.M. broad. Pinnulae lanceolate, imbricate, those at the lower base larger than the others, the lowest deeply pinnatifid. Segments oblong, blunt, faintly toothed, the teeth not awned. Texture subcoriaceous; both surfaces naked; rachis and midribs slightly fibrillose. Sori principally medial; indusium peltate or reniform.

Philippines; Assam, China, Korea, Japan.

* (10) **P. aristatum, Pr.**, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 195; *Aspidium aristatum, Sw.*, Christ, Farnkr. d. Erde, 240; *Lastraea aristata, Moore*, Bedd., Ferns S.I., tab. CI; — Var. *Hamiltonii, Spr.*, Syst., IV, 108.

Rhizome creeping, densely clothed with short, setiform or hair-like scales. Stipes scattered, 50—70 c.M. long, fibrillose or scaly at least near the base. Fronds 50—70 c.M. long, deltoid, 20—50 c.M. broad, with a long, pinnate apex and below this \pm 5—6 pinnate pinnae on each side. Lowest pinnae 2-pinnate to 5-pinnatifid at the base, the lower basal pinnulae the longest. Leaflets subrhomboidal, the upper base \pm truncate and auricled, the lower base cuneate, the edges more or less lobed with the lobes copiously toothed, the teeth aristate. Texture \pm coriaceous; surfaces naked; rachis nearly so. Sori nearer the midrib than the edge; indusium peltate or reniform.

Var. Hamiltonii: Largest form; ultimate segments large, ovate-deltoid, slightly lobed. —

The rhizome of the type is figured in Bedd., Handb. Ind. Ferns, fig. 118, A.

Malaya; Natal, India, China, Japan, Polynesia; — Var. *Hamiltonii: ? Malacca; Nepal, Khasia, Ceylon.*

(11) **P. conifolium, Pr.**, Copel., Polypod. Philipp., 18; *P. carvifolium, C. Chr.*, Ind. Fil.; *Aspidium curvifolium, Kze.*, Bot. Zeit., 1848, 285; *A. carvifolium, Kze, Linn.*, XXIV, 292; *A. conifolium, Wall.*, Christ, Farnkr. d. Erde, 240; *A. aristatum, Sw.*, var. *conifolium, Hk. Bk.*, Syn Fil., 255; *Lastraea conifolia, Moore*, Bedd., Ferns Br.I., II, tab. CCLXI; — var. *Sarasinorum; A. aristatum, Sw.*, var. *Sarasinorum, Christ, Ann. Btz.*, XV, 124.

Near *P. aristatum, Pr.*, but the rhizome erect, the scales long, linear-lanceolate, Stipes tufted, the base clothed with narrow, blackish scales. Fronds large, 60—90 c.M. long, (50—50 c.M. long, conf. Copel.), 4—5-pinnate. Ultimate pinnules and segments small, less coriaceous, ovate or rhomboidal, aristate. Sori large or small; indusium peltate, sometimes reniform. — The creeping rhizome figured in the head of the mentioned table is that of *P. aristatum, Pr.*

Var. Sarasinorum: Small form. Fronds \pm 20 c.M. long: segments short, deeply incised, bluntish or short-acuminate, not aristate.

Malaya; Polynesia, Australia, Japan, China, North. India, Natal; — var. *Sarasinorum: Celebes.*

20. DIDYMOCHLAENA, *Desvoux.*

Sori oblong, terminal on the veins, but not marginal. Indusium the same shape of the sorus, fixed by its longitudinal axis, opening on all sides.

Rhizome erect. Stipes not articulate to the rhizome. Fronds 2-pinnate. Leaflets unequal-sided, subdimidiate, the basiscope half being obliquely cut away; veins free. Hk. Bk., Syn Fil., tab, IV, fig. 42; Diels, in Engl & Prantl., Nat. Pfl.Fam., I⁴, fig. 95.

Pantropical.

* ***D. lunulata, Desv.,*** Rac., Flor. Btz., I, 197; Bedd., Ferns Br.I., I, tab. XV; Hk., Gard. Ferns, tab. XVII; *D. sinuosa, Desv.,* Bau. & Hk., Gen. Fil., tab. VIII; *D. truncatula, J. Sm.,* Journ. of Bot., IV, 196; C. Chr., Ind. Fil; *Adiantum lunulatum, Houtt.,* (oldest name).

Caudex erect, subarborescent. Stipes densely tufted, several d.M. long, paleaceous. Fronds 60—180 c.M. long, 50—60 c.M. broad, 2-pinnate, oblong to lanceolate. Pinnae numerous, sessile or short-stalked, erecto-patent, linear, acuminate, 2—4 c.M. or more broad. Pinnulae numerous, close, articulate, subrhomboidal, about half as broad as long, the upper base broadly and the lower narrowly cuneate, the edge entire or slightly sinuate. Texture subcoriaceous; main rachis scaly, the scales narrow, fimbriate; secondary rachises slightly scaly, sometimes hairy; upper surface slightly covered with subulate bristles. Sori 2—6 to a pinnula.

Pantropical.

21. PLEOCNEMIA, *Presl.*

Sori roundish, medial or terminal on the free veins or veinlets, rarely on the netted veins. Indusium reniform, attached by the sinus, rarely round and peltate.

Rhizome generally erect. Stipes not articulate to the rhizome. Fronds often ample and compound; lower veins of contiguous groups united and those

of the same group slightly so, forming 1—2 rows of irregular areolae near the costae and costulae without included free veinlets, the rest free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 97, D. (*Aspidium*, p.p., C. Chr., Ind. Fil).
Tropical Asia, Polynesia.

-
- a.** *Fronds simple, lobed or pinnatifid.* (1) *P. heterophylla*.
a.a. *Fronds at least 2-pinnatifid.*
b. *Texture rather coriaceous.*
1. *Lower pinnae pinnatifid.*
2. *Surfaces slightly hairy on the veins.* (2) *P. excellens*.
2.2. *Surfaces naked, or the upper one slightly scaly.* (5) *P. megalocarpa*.
1.1. *Lower pinnae pinnate, at least on the lower side.* (4) *P. Leuzeana*.
b.b. *Texture rather herbaceous or membranaceous.*
c. *Lower pinnae pinnatifid.* (5) *P. gigantea*.
c.c. *Lower pinnae pinnatifid with the lowest basiscopical segment pinnatifid again.* (6) *P. membranifolia*.
c.c.c. *Lowest pinnae pinnate at the base, the lowest basiscopical segment pinnatifid or pinnate at the base.*
d. *Indusium cordato-remiform.* (7) *P. Bakeri*.
d.d. *Indusium peltate.* (8) *P. devexa*.
-

(1) **P. heterophylla**, v. **A. v. R.**; *Nephrodium heterophyllum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 295; *Aspidium heterophyllum*, *Mett.*, *Copel.*, *Polypod. Philipp.*, 39., *Hk.*, *Icon. Pl.*, X, tab. CMXX; *Dryopteris canescens*, *C. Chr.*, var., *C. Chr.*, *Ind. Fil.*

Rhizome creeping. Stipes 5—10 c.M. long, densely villose, scaly towards the base. Fronds 12—15 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, simple, acuminate, the edge lobed $\frac{1}{4}$ — $\frac{1}{3}$ -way down into numerous, oblong, blunt, slightly falcate lobes. Texture coriaceous; surfaces villose especially on the edges and the midribs beneath; main veins pinnate in the lobes, those of the barren fronds forming 1—2 rows of costal and constular areolae, those in the fertile fronds anastomosing \pm as in *Nephrodium*, 5—6 on each side. Sori medial on the veins; indusium small, fugacious.

Philippines.

(2) **P. excellens**, v. **A. v. R.**; *Nephrodium excellens*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 295; *Aspidium excellens*, *Bl.*, *Enum.*, 160; *C. Chr.*, *Ind. Fil.*

Stipes naked, greyish. Fronds large. Lower pinnae to 50 c.M. long, 7—10 c.M. broad, deeply pinnatifid into oblong-lanceolate, acuminate, entire or distantly crenate lobes. Texture subcoriaceous; rachis, veins and margins slightly hairy on both sides; veins of the lobes slightly pinnate, and mostly forming costal arches. Sori small, copious, principally in irregular rows near the costulae.

Java.

(3) **P. megalocarpa**, *Bedd.*, Handb. Ind. Ferns, Suppl, 48; *Polypodium megalocarpum*, *Hk.*, Hk. Bk, Syn. Fil., 518; *Nephrodium oligodictyon*, *Bk.*, Ann. of Bot., V, 328, *Aspidium megalocarpum*, *C. Chr.*, Ind. Fil.

Rhizome erect, scaly at the crown. Stipes stout, 40—60 c.M. long, scaly towards the base. Fronds oblong-deltoid, 60—90 c.M. long, the upper part deeply pinnatifid with lanceolate, pinnatifid lobes. Pinnae in ± 6 pairs; the lowest largest, equilateral, 15—25 c.M. long, 5—9 c.M. broad, cut more than $\frac{1}{2}$ -way down to the costa into narrow, oblong, crenated segments to $\pm 3\frac{1}{2}$ c.M. long, ± 5 —8 m.M. broad. Texture subcoriaceous, opaque; surfaces naked or the upper one furnished with scattered, small, jointed, transparent scales; venation pinnate in the lobes, forming areolae near the costae only. Sori medial or nearly so, in 2 rows on each side of the costa along the wing and in a single row on each side of the costulae; indusium fugacious.

Java, Malacca.

* (4) **P. Leuzeana**, *Pr.*, Rac., Flor. Btz., I, 194; *Bedd.*, Ferns Br.I., I, tab. CXXXIV; *Nephrodium Leuzeanum*, *Hk.*, Hk. Bk., Syn. Fil., 295; *Aspidium Leuzeanum*, *Kze.*, Copel., Polypod. Philipp., 39; *C. Chr.*, Ind. Fil. — var. *hemiteliiformis*, *Rac.*, l. c.; — var. *digitata*.

Caudex erect, short, to ± 12 c.M. high, or subarborescent and to ± 1 M. high, the apex clothed with long, linear-lanceolate, ciliate, woolly, brown scales. Stipes 50—100 c.M. long, often striated, scaly at the base. Fronds subdeltoid, 2—3-pinnatifid, ± 1 —2 M. long, 30—90 c.M. broad. Pinnae numerous, close, pinnatifid, or at least the lowest pinnate on the lower side, short-stalked. Pinnulae sessile or short-stalked, to ± 25 c.M. long, to ± 8 c.M. broad, the lowest 1—2 \times pinnatifid. Lobes numerous, separated by narrow sinuses with roundish tops, oblong, bluntish, subfalcate, entire, toothed or sinuated. Texture coriaceous; surfaces naked. Sori usually in 1 row on each side, nearer the edge than the costulae; indusium early fugacious, (often wanting?).

Var. hemiteliiformis: Pinnulae 6—9 c.M. long, 8—12 m.M. broad; edges ciliate; sori close.

Var. digitata: Fronds digitate.

Malaya; Polynesia, South. China, North. India; — var. hemiteliiformis: *Java;* — var. digitata: *Celebes.*

* (5) **P. gigantea, Pr.,** *Rac., Flor, Btz., I, 195; P. Trimeni, Bedd., Handb. Ind. Ferns, 225; Sagenia gigantea, Moore, Bedd., Ferns S.I., tab. LXXX; Aspidium giganteum, Bl., Enum., 159; C. Chr., Ind. Fil., p.p.*

Caudex erect, to \pm 8 c.M. high, clothed with ovate-lanceolate, rigid, entire scales. Stipes to 50—60 c.M. long, scaly at the base. Fronds 50—90 c.M. long, to 45 c.M. broad, 2-pinnatifid. Pinnae sessile or short-stalked, lanceolate to linear, the edge cut to $\frac{2}{3}$ — $\frac{5}{6}$ -way down into erecto-patent, subfalcate, entire, crenate or sinuate lobes. Lowest pinnae to 12 c.M. broad, the apex acuminate, the base obliquely truncate or cuneate, medial pinnae to 5 c.M. broad, the higher ones decurrent. Texture herbaceous to submembranaceous; veins forming about 3—4 rows of costal areolae and about 1 row of costular ones. Sori numerous, in 1 or rarely 2 rows on each side of the costulae.

Malaya; Ceylon, South. India.

(6) **P. membranifolia, Bedd.,** *Handb. Ind. Ferns, 255, fig. 115; Nephrodium membranifolium, Pr., Hk., Spec. Fil., IV, tab. CCLXI; Aspidium membranifolium, Kze. (not Pr.), Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 188; C. Chr., Ind. Fil.*

Stipes tufted, clothed with black, linear, subulate scales, but more copiously towards the base, \pm 50 c.M. long. Fronds to 50 c.M. or more long, deltoid to ovate, pinnate, the apex pinnatifid. Lateral pinnae in few pairs, \pm opposite, 5—10 c.M. long, 3—4 c.M. broad, more or less deeply pinnatifid, the segments rounded, nearly entire; lowest pair of pinnae largest, deltoid, pinnatifid or subpinnate, with the lower basal segments elongate and pinnatifid. Rachis and surfaces \pm softly hairy; texture softly herbaceous. Fertile fronds similar to the barren ones or \pm contracted. Sori generally terminal on free veinlet, in 1 row on each side of the costulae; indusium reniform.

Malacca, (? Malaya).

* (7) **P. Bakeri, v. A. v. B.,** *Nephrodium giganteum, Bl., Hk. Bk., Syn Fil., 503; Aspidium giganteum, Copel. (not Bl.), Polypod. Philipp., 58; (? Bl., C. Chr., Ind. Fil., p.p.).*

Stipes 50—60 c.M. long, brown, glossy. Fronds 50—60 c.M. long, broadly deltoid, with a pinnatifid apex and \pm 4—5 pairs of pinnae on each side, the higher ones decurrent, the lower stalked. Lowest pinnae largest, deltoid, the pinnulae on the lower side prolonged, deeply pinnatifid; highest pinnae bluntly lobed. Texture moderately firm; surfaces nearly naked; veins anastomosing at least near the costae. Indusium subcordate. — A plant, gathered in Borneo and cultivated in the Buitenzorg gardens, which may be a form of this species has the fronds under 50 c.M. long, the ultimate lobes linear, gradually narrowed towards the apex, entire or broadly toothed and the sori submarginal.

Philippines, ? Borneo.

* (8) **P. devexa**, v. **A. v. B.**; *P. membranacea*, *Bedd.*, *Handb. Ind. Ferns*, 225; *Aspidium membranaceum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 259; *A. devexum*, *Kze.*, (oldest name), *Bot. Zeit.*, 1848, 259; *C. Chr.*, *Ind. Fil.*; — var. *minor*; *A. giganteum*, *Bl.*, var. *minor*, *Hk.*, *Spec. Fil.*, IV, 50; *Sagenia gigantea*, *Bedd.*, var. *minor.*, *Bedd.*, *Ferns S.I.*, tab. CCXLIII.

Stipes tufted, to about 50 c.M. long, nearly naked, with a few spreading, linear scales below. Fronds to 45—60 c.M. long, \pm 50 c.M. or more broad. Lower pinnae much the largest, subdeltoid, 15—20 c.M. long, 7—10 c.M. broad. Pinnulae on lowest side much larger than the others, pinnatifid (pinnate at the base) with lanceolate, pinnatifid segments, with small, oblong, sinuated lobes. Texture thinly herbaceous; colour pale-green; veins anastomosing principally in costal arches, but sometimes also as in *Aspidium*. Sori copious, terminal on the veins or veinlets; indusium peltate.

Var. minor: A less divided form; pinnae pinnatifid, the lowest ones pinnate at the base; pinnulae pinnatifid, the edges of the lobes entire or \pm slightly lobed.

Malaya; Formosa, South. China, Ceylon.

22. DRYOPTERIS, *Adanson.*

Sori roundish, commonly \pm medial on the veins. Indusium reniform, attached by the sinus, rarely peltate and round or cordate.

Rhizome creeping or erect. Stipes not articulate to the rhizome. Fronds commonly pinnate or sometimes more compoundly divided; veins free or the lower ones of adjacent groups anastomosing and then forming on each side of the midribs 1 row of regular, deltoid, costal areolae with an excurrent

veinlet springing from the top, free or joined in the angle of the next superior anastomosing veins, which often form 1 or more rows of regular, rhomboidal areolae between the costal ones and the margin. Segments mostly equal-sided or nearly so; surfaces often \pm hairy; texture commonly herbaceous. — The species are difficult of determination unless the whole stipe and the rhizome are present; the frond only is of little value.

Tropical and extra-tropical.

Arrangement of the sections.

§ 1. **LASTRAEA.** Veins free. Indusium reniform. Hk. Bk., Syn. Fil., tab. V, fig. 44, c—d.

§ 2. **NEPHRODIUM.** Veins anastomosing. Indusium reniform. Hk. Bk., l. c., fig. 44, a—b.

§ 3. **CYCLOSORUS.** Like § 2, but the indusium peltate, and round or cordate.

§ 1. **LASTRAEA.**

a. Fronds pinnate. Pinnae entire or lobed less than $\frac{1}{3}$ -way down to the costa.

*. Fronds sessile. Pinnae entire or slightly crenate. (1) *D. Warburgii*.

.. Fronds moderately long-stalked. Pinnae lobed $\frac{1}{4}$ — $\frac{1}{3}$ -way down.

(2) *D. hirtipes*.

a.a. Fronds pinnate. Pinnae cut more than $\frac{1}{3}$ -way down to quite to the costa. Secondary segments entire or crenate to repand (1).

1. Central ribs of the lobes pinnate; veins simple.

2. Lower pinnae slightly, if at all, reduced, not dwindling down into mere auricles.

3. Fronds small, under 10 c.M. long (2). (3) *D. grammitoides*.

3.3. Fronds more than 10 c.M. long.

4. Texture \pm chartaceous or coriaceous.

5. Fronds under 75 c.M. long.

6. Surfaces naked, at best pubescent on the costae.

7. Rachis naked (3).

(1) See also *D. phegopteroidea*, C. Chr.

(2) See also *D. diversifolia*, v. A. v. R.

(3) See also *D. pectinata*, v. A. v. R.

S. *Fronds* ± 15 c.M. long; *veins* ± 8 on each side in the secondary segments.

(4) *D. Koordersii*.

S.S. *Fronds* more than 50 c.M. long; *veins* ± 4—5 on a side.

(5) *D. phaeostigma*.

7.7. *Rachis* pubescent or villose.

✦. *Pinnae* under 6 c.M. long, cut down to the costa. (6) *D. gymnopoda*.

✦.✦. Longest *pinnae* more than 6 c.M. long, cut to $\frac{2}{3}$ -way down to the costa. (7) *D. crassifolia*.

6.6. *Under surface* pubescent.

! . *Stipes* scaly, at length scabrous. *Fronds* more than 20 c.M. broad. (8) *D. echinata*.

! .! . *Stipes* naked or pubescent. *Fronds* under 20 c.M. broad.

* . *Stipes* pubescent, subtufted. *Pinnae* close, not dotted with white above.

(9) *D. Teuscheri*.

* .* . *Stipes* naked, scattered. *Pinnae* sub-remote, dotted with white above.

(10) *D. pectinata*.

5.5. *Fully developed fronds* more than 75 c.M. long.

S. *Rachis* pubescent or scaly, *under surface* pubescent on the veins.

* . *Stipes* naked, grey. (11) *D. ligulata*.

* .* . *Stipes* hairy and scaly, brown.

(12) *D. Hallieri*.

S.S. *Rachis* naked.

✦. *Under surface* glandular. (13) *D. attenuata*.

✦.✦. *Under surface* naked. (14) *D. Beccariana*.

4.4. *Texture* ± membranaceous or herbaceous.

α. *Sori* medial, or lower on the veins.

β. *Fronds* mostly under 60 c.M. long.

γ. *Stipes* naked.

δ. Lower secondary segments not very small. (15) *D. Creaghii*.

- $\delta.\delta.$ Lower secondary segments very small.
(16) *D. vulcanica*.
- $\gamma.\gamma.$ Stipes more or less finely villose or pubescent, at least on the upper side (1).
! . Under surface villous.
(17) *D. calcarata*.
- !:. Under surface naked or glandular.
‡. Densely glandular.
(18) *D. viscosa*.
- ‡.‡. Naked or slightly glandular.
(19) *D. gracilescens*.
- $\beta.\beta.$ Fully developed fronds more than 60 c.M. long.
!. Stipes densely hairy. Secondary segments with a single tooth in the sinus on the anterior side.
(20) *D. monodonta*.
- !:. Stipes scaly. Secondary segments without tooth. (21) *D. polytricha*.
- $\alpha.\alpha.$ Sori nearer the margin, or terminal on the veins (2).
 β' . Sori immersed. (22) *D. immersa*.
 $\beta'\beta'$. Sori superficial.
- γ' . Lowest pinnae with the lowest anterior segment toothed or pinnatifid, the other segments entire. (23) *D. Peekeli*.
- $\gamma'\gamma'$. All the secondary segments similar, entire or crenate.
 δ' . Surfaces naked. (24a) *D. Dayi*.
 $\delta'\delta'$. Surfaces more or less villous.
 ϵ' . Fronds \pm small.
(24b) *D. diversifolia*.
 $\epsilon'\epsilon'$. Fronds \pm large. (25) *D. patens*.
- 2.2.** Lower pinnae conspicuously reduced or dwindling down into mere auricles.
- 3.** Sori medial, or lower on the veins.

(1) See also under *D. hispidula*, O. Ktze.

(2) See also *D. gracilescens*, O. Ktze.

- 4.** *Pinnae cut \pm $\frac{1}{2}$ -way down to the costa (*)*.
(26) *D. exigua*.
- 4.4.** *Pinnae cut down \pm to the costa.* (27) *D. conterminoides*.
- 3.3.** *Sori near the margin, or terminal on the veins.*
- ✧. *Rhizome short. Veins 8—12 on a side. Fronds rather large.* (28) *D. Luerssenii*.
- ✧✧. *Rhizome wide-creeping. Veins 4—5 on a side. Fronds rather small.* (29) *D. Beddomei*.
- 1.1.** *Central ribs pinnate, lateral veins forked.*
- !. *Rachis scaly.* (30) *D. adnata*.
- !!. *Rachis naked, hairy or fibrillose.*
- ✧. *Fronds under 75 c.M. long.*
- ?. *Stipes dark-coloured. Surfaces naked. Rachis naked.*
(31) *D. sagenioides*.
- !?. *Stipes grey. Surfaces finely pubescent. Rachis pubescent*
(32) *D. singalanensis*.
- ✧✧. *Fronds more than 75 c.M. long when fully developed. Rachis fibrillose.* (35) *D. Wigmanii*.
- 1.1.1.** *Central ribs pinnate, lower lateral veins subpinnate.* (34) *D. filix mas*.
- a.a.a.** *Fronds pinnate. Pinnae cut down nearly or quite to the costa. Secondary segments toothed or lobed.*
- b.** *Secondary segments toothed.*
- !. *Sinuses with a tooth in the top, just between each pair of lobes.*
(55) *D. syrmatia*.
- !!. *Sinuses without tooth in the top. Indusia red when young.*
(36) *D. erythrosora*.
- b.b.** *Secondary segments lobed, not pinnatifid.*
- c.** *Fronds under 75 c.M. long.*
- d.** *Under surface naked. Texture \pm chartaceous or coriaceous.*
- e.** *Ultimate lobes entire, with \pm 6 pairs of veins. Sori subcostular.* (37) *D. Preslii*.
- e.e.** *Ultimate lobes crenate or toothed, with \pm 3—4 pairs of veins. Sori medial on the lower veins.*
(38) *D. chaerophyllifolia*.
- d.d.** *Under surface fibrillose on the main veins. Texture herbaceous. Sori costal.* (59) *D. edulenta*.

(*) See also *D. hispidula*, O. Ktze.

d.d.d. Under surface villous. Texture herbaceous. Sori medial.

(40) *D. flaccida*.

c.c. Fronds more than 75 c.M. long.

(41a) *D. splendens*.

b.b.b. Secondary segments pinnatifid.

(41b) *D. phegopteroidea*.

a.a.a.a. Fronds at least deeply 3-pinnatifid at the base.

b. Tertiary segments entire, crenulate or slightly toothed.

c' Rachis naked or nearly so.

!. Tertiary segments entire or crenulate.

!. Texture herbaceous. Lobes crenulate. (42) *D. dissecta*.

!.!. Texture subcoriaceous. Lobes entire. (45) *D. papuana*.

!.!. Tertiary segments slightly & \pm sparingly toothed. (44) *D. sparsa*.

c'.c' Rachis more or less pubescent or pilose.

d' Sori nearer the edge than the costulae.

*. Ultimate segments separated by an intermediate vein, which unites the costa with the top of the sinus.

(45) *D. Raciborskii*.

.. Intermediate veins wanting.

(46) *D. recedens*.

d'.d' Sori medial, or lower on the veins.

e' Rhizome wide-creeping.

(47) *D. borneensis*.

e'.e' Rhizome short-creeping or erect.

f' Sori close to the costulae.

(48) *D. Millettii*.

f'.f' Sori medial.

(49) *D. padangensis*.

c'.c'.c' Rachis scaly.

§. Texture coriaceous. Sori occupying nearly the whole under surface of the small ultimate segments. (50) *D. vilis*.

§.§. Texture herbaceous or moderately firm Sori medial, or lower on the veins.

!. Fronds at utmost \pm 15 c.M. broad.

!. Rhizome short-creeping.

(51) *D. intermedia*.

!.!. Caudex erect.

(52) *D. sarawakensis*.

!.!. Fully developed fronds more than \pm 20 c.M. broad.

!.!. Ultimate segments to 1 c.M. long. (55) *D. pulvinulifera*.

!.!.!. Ultimate segments 2—2 $\frac{1}{2}$ c.M. long.

(54) *D. aciculata*.

b'.b' Tertiary segments at least conspicuously toothed (*).

1. Rachis naked, or slightly scaly.

(*) See also *D. sparsa*, O. Ktze & *Raciborskii*, v. A. v. R.

- §. *Tertiary segments toothed. Sori subcostular.* (55) *D. Boryana.*
- §§. *Tertiary segments at least pinnatifid. Sori ± medial.*
(56) *D. purpurascens.*
- 1.1. *Rachis hairy or fibrillose, at least on the upper side.*
2. *Surfaces densely hairy like the rachis.* (57) *D. crenata.*
- 2.2. *Under surface slightly hairy or scaly on the veins.*
3. *Rachis slightly fibrillose, the fibrils deciduous.*
(58) *D. rhodolepis.*
- 3.3. *Rachis pubescent on the upper side.* (59) *D. setigera.*
- 1.1.1. *Rachis scaly or bristly.*
- ‡. *Scales of the stipe ovate; under surface ± glandular.*
(60) *D. zeylanica.*
- ‡‡. *Scales of the stipe bristle-like; under surface naked or hairy.*
(61) *D. multisetata.*
- ‡‡‡. *Scales of the stipe subulate; under surface scaly.*
(62) *D. subarborea.*

(1) **D. Warburgii**, C. Chr., Ind. Fil.,; *Aspidium Warburgii*, Kuhn & Christ, Warb., Mons., I, 81; — var. *reticulata*, Christ, l. c.

Stipes tufted, very short, nearly wanting. Fronds ± 10 c.M. long, ± 2 c.M. broad; rachis hairy. Pinnae ligulate, bluntish, the upper base truncate, the lower one cuneate, the margin entire or slightly crenate. Veins simple or forked. Sori minute, medial, in 1 row on each side of the midribs, 8—10 to a row; indusium small, fugacious.

Var. reticulata: Pinnae ovate; veins ± anastomosing.

New Guinea.

* (2) **D. hirtipes**, C. Chr., Ind. Fil.; *Aspidium hirtipes*, Bl., Rac., Flor. Btz., I, 168; *A. atratum*, Wall., Mett., Farng., IV, Phegop. & Aspid., No. 126; *Lastrea hirtipes*, Moore, Bedd., Ferns S.I., tab. XCVI; *Nephrodium hirtipes*, Hk., Hk. Bk., Syn. Fil., 261.

Rhizome short-creeping; the scales dense, long, lanceolate-subulate, dark-coloured. Stipes subtufted, 15—35 c.M. long, scaly; the scales black-brown, the lower ones large, lanceolate-subulate, the higher ones linear-subulate or filiform. Fronds 15—60 c.M. long, 8—35 c.M. broad, pinnate, the apex pinnatifid. Pinnae 1½—2½ c.M. broad, short-stalked, the highest sessile, the lowest longest, linear-lanceolate, the base truncate, the edge lobed from

$\frac{1}{4}$ — $\frac{1}{3}$ on the way down to the costa, the point long-acuminate, toothed. Texture herbaceous to semicoriaceous; rachis fibrillose with black, filiform scales; surfaces naked, or the midribs slightly scaly; main veins pinnate in the lobes; lower veins not reaching the edge. Sori medial on the veins. — This resembles *D. filix mas*, *Schott* in general habit (Bk.).

Malaya; Polynesia, South. China, North. India.

(3) **D. grammitoides**, *C. Chr.*, Ind. Fil.; *Aspidium grammitoides*, *Christ*, Bull. Herb. Boiss., VI, 193; *Nephrodium grammitoides*, *Copel.*, Polypod. Philipp., 21.

Rhizome short-creeping, scaly. Stipes weak, 3—4 c.M. long, stramineous, clothed like the whole plant, and especially the costae and veins, with short, rigid, whitish hairs. Fronds 5—7 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, broadly oblong, narrowing from the base to the caudate, pinnatifid apex. Pinnae \pm 6 on each side, lowest largest, distant, sessile, deltoid, acuminate, cut down to the costa at the base, the upper portion pinnatifid. Lobes \pm 4 m.M. long, deltoid, crenate. Main veins pinnate in the lobes. Sori minute, 3—4 on each side of the main veins; indusium small and villose, or suppressed.

Luzon.

(4) **D. Koordersii**, *C. Chr.*, Ind. Fil.; *Aspidium Koordersii*, *Christ*, Ann. Bot., XV, 128, tab. XV, fig. 17, a—b.

Rhizome short, erect, woody, clothed with blackish, acuminate scales. Stipes tufted, polished, rufo-stramineous, partially spotted, firm, \pm 15 c.M. long. Fronds \pm 15 c.M. long, \pm 10 c.M. broad, ovate-deltoid, 2-pinnate. Pinnae \pm 8 on each side, erecto-patent, the lower ones not shorter, ovate towards the base, subsessile, the apex caudate to acuminate, the rachis narrowly winged. Secondary segments \pm $1\frac{1}{2}$ c.M. long, distant, linear-lanceolate, acute, slightly crenate, narrowed at the base. Texture thinly coriaceous; colour dark-green; rachis and both surfaces naked; veins black, erecto-patent, simple, \pm 8 on each side. Sori small, medial on the veins; capsules few; indusium laterally fixed, crustaceous, convex, brown, glossy, the margin decurved.

Celebes.

(5) **D. phaeostigma**, *C. Chr.*, Ind. Fil.; *Aspidium phaeostigma*, *Ces.*, Fel. Becc. Polin., 4 & 7.

Stipes erect or suberect, 60—75 c.M. long, the base densely clothed with long, linear-subulate, brown scales, the rest naked. Fronds 45—60 c.M. long, \pm 20—25 c.M. broad, lanceolate, pinnate. Pinnae spreading, pinnatifid

quite to the costa. Segments linear, falcate, confluent at the base, hardly repand, subacute. Surfaces naked, opaque; veins 4—5 on each side. Sori solitary on the veins; indusium coriaceous, moderately large, at length deciduous.

New Guinea.

(6) **D. gymnopoda**, *C. Chr.*, Ind. Fil.; *Nephrodium gymnopodum*, *Bk.*, Trans. Linn. Soc., Bot., 2^d Series, IV, 252.

Stipes slender, naked, glossy, reddish-brown, 10—15 c.M. long. Fronds oblong-lanceolate, 2-pinnate, \pm 15 c.M. long. Pinnae \pm 20-jugate, the lowest largest, deflexed, lanceolate, $5\frac{1}{2}$ — $4\frac{1}{2}$ c.M. long, \pm 1 c.M. broad, cut down to the costa. Segments linear-oblong, entire, \pm 2 m.M. broad. Texture subcoriaceous; surfaces naked; rachis pubescent; veins distant, simple, erectopatent. Sori solitary at the base of the segments, medial; indusium small.

Borneo.

* (7) **D. crassifolia**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium crassifolium*, *Bl.*, Rac., Flor. Btz., I, 169; *Lastreaa nephrodioides*, *Bedd.*, Ferns Br.I., II, tab. C1C; *L. crassifolia*, *Moore*, *Bedd.*, Handb. Ind. Ferns, 238; *Nephrodium crassifolium*, *Hk.*, Hk. Bk., Syn. Fil., 266; — var. *Motleyanum*; *A. Motleyanum*, *Christ*, Ann. Btz., XX, 105; *N. Motleyanum*, *Hk.*, Hk. Bk., l. c.; *D. Motleyana*, *C. Chr.*, Ind. Fil.

Rhizome creeping, the scales lanceolate. Stipes approximate, scaly below, hairy above, 30—50 c.M. long. Fronds 50—60 c.M. long, 15—50 c.M. broad, deltoid-lanceolate. Pinnae 2— $4\frac{1}{2}$ c.M. broad, the fertile commonly narrower than the barren ones and sometimes much contracted, sessile or short-stalked, the lowest stalked, cut $\frac{1}{2}$ — $\frac{2}{3}$ -way down to the costa into blunt or acute, falcate, entire lobes 5—8 m.M. broad, the lowest of which are \pm reduced in size. Texture subcoriaceous; rachises and midribs finely or obscurely hairy; veins 10—12 on each side of the main ones, the lowest sometimes joining or terminating in the tops of the sinuses. Sori commonly 1 to each vein, the higher ones close to the main veins, the lowest often divaricating.

Var. Motleyana: Veins anastomosing in the barren fronds; sori medial. Pinnae less numerous, distant.

Malaya; Burmah; — var. Motleyana: Borneo.

* (8) **D. echinata**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium echinatum*, *Mett.*, Ann. Mus. Bot. L. B., I, 230; *Nephrodium echinatum*, *Bk.*, Hk. Bk., Syn. Fil., 266.

Stipes \pm 30 c.M. long, clothed with long, spreading, brown, lanceolate-subulate scales, scabrous when they fall. Fronds \pm 30 c.M. long, 20—25 c.M. broad, lanceolate-deltoid. Lowest pinnae stalked, \pm 12 $\frac{1}{2}$ c.M. long, 15—20 m.M. broad, cut down to a narrowly winged rachis into close, linear-oblong, entire lobes. Texture coriaceous; rachis like the stipe; under surface thinly pubescent and glandular; veins 7—8 on a side, the lowest sometimes joining. Sori nearer the midrib than the edge.

Sumatra, Borneo, Celebes, New Guinea.

* (9) **D. Teuscheri**, v. A. v. B.

Rhizome scaly; the scales brown, linear-lanceolate, acuminate, ciliate. Stipes subtufed, \pm 5—10 c.M. long and like the rachis greyish or brownish, densely short-hairy. Fronds linear-lanceolate, acuminate, pinnate, \pm 15—20 c.M. long, \pm 5—4 c.M. broad, with a pinnatifid apex and an elongate, crenate point. Higher pinnae oblong, entire, sessile; following pinnae subsessile; central pinnae close, short-stalked, erecto-patent, falcate, \pm 1 c.M. broad, the base truncate, auricled on the upper side, the edge pinnatifid \pm $\frac{1}{3}$ on the way to the costa, but more deeply and nearly to the costa towards the base, the apex oblong, blunt, entire or crenate; lower pinnae shorter, deflexed. Lobes blunt, subentire. Texture semicoriaceous; under surface densely glandular and short-hairy; veins forked, or pinnate in the lobes. Sori medial on the veins, in 1 row on each side of the costulae; indusium densely hairy. — Fronds with the general aspect of *D. pilosiuscula*, *C. Chr.*, but the pinnae more deeply lobed and the upper surface naked.

Borneo (Teuscher).

(10) **D. pectinata**, v. A. v. B.; *D. orientalis*, *C. Chr.*, Ind. Fil.; *Aspidium pectinatum*, *Christ*, Farnkr. d. Erde, 250; *Nephrodium albo-punctatum*, *Desv.*, Hk. Bk., Syn. Fil., 264; Hk., Fil. Exot., tab. LXXXIX; *N. pectinatum*, *Hier.*, Engl. Jahrb., XXVIII, 341; *N. Webbianum*, *Hk.*, Spec. Fil., IV, 85; *Polypodium pectinatum*, *Forsk.*, (oldest name).

Rhizome wide-creeping, scandent. Stipes naked, greyish-blackish, distant, \pm 10 c.M. long. Fronds 20—30 c.M. long, 10—15 c.M. broad, ovate. Pinnae spreading, subdistant, sessile, lanceolate, blunt, 1—1 $\frac{1}{2}$ c.M. broad, cut $\frac{1}{2}$ -way down or more into oblong, entire, subspreading lobes. Texture firm; surfaces and rachis more or less pubescent; upper surface with white cretaceous dots; main veins pinnate, with 3—4 veins on each side. Sori small, near the edge; indusium hairy. *N. Webbianum*, *Hk.* has the fronds

quite naked, the pinnae less deeply lobed, the upper base auriculate and the sori between the main veins and the margin.

Amboyna, New Guinea; Fiji, Mascaren Islands, Trop. Africa.

(11) **D. ligulata**, **O. Ktze**, C. Chr., Ind. Fil.; *Aspidium ligulatum*, Kze, Mett., Farng., IV, Phegop. & Aspid., No. 213; *Nephrodium ligulatum*, Hk., Hk. Bk., Syn. Fil., 264.

Stipes grey, glossy, naked. Fronds 90—120 c.M. long, 30—45 c.M. broad. Pinnae 15—25 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, cut down very nearly to the costa into linear, blunt, entire, erecto-patent lobes $\pm 2\frac{1}{2}$ m.M. broad and with more than their own space between them, lower pinnae slightly reduced. Texture subcoriaceous; rachis and under surface finely pubescent; veins ± 10 —12 on a side. Sori submarginal.

Philippines.

* (12) **D. Hallieri**, **C. Chr.**, Ind. Fil.; *Aspidium Hallieri*, Christ, Ann. Btz., XX, 106.

Stipes thick, brown, opaque, clothed like the rachis with short hairs and like the base of the rachis with lanceolate-subulate, spreading, brown scales. Fronds ovate-lanceolate, ± 100 c.M. long or longer, ± 35 c.M. broad, acuminate, hardly narrowed at the base. Pinnae of the barren fronds close, sessile, spreading, caudate-acuminate, the margin incised $\frac{3}{4}$ -way down to the costa, the base ± 3 c.M. broad. Segments pectinate, close, ± 35 below the serrate apex, ± 1 c.M. long, ± 4 —5 m.M. broad, falcate, lanceolate, bluntish, the margin subentire or subcrenulate, short-ciliate, slightly inflexed. Costae, costulae and veins pubescent; veins 12—15 on each side, very prominent. Fertile pinnae 2—2 $\frac{1}{2}$ c.M. broad, the lower ones distant; segments 2—3 $\frac{1}{2}$ m.M. broad. Sori minute, medial or subcostular, ± 15 on each side; indusium small or wanting (suppressed?).

Borneo.

* (13) **D. attenuata**, **O. Ktze**, Rev. Gen. Pl., II, 812; *D. stenobasis*, C. Chr., Ind. Fil.; *Aspidium attenuatum*, Kze, Mett., Farng., IV, Phegop. & Aspid., 96; *Nephrodium attenuatum*, Fée (not Bk.), Hk. Bk., Syn. Fil., 263.

Stipes 30 c.M. or more long, firm, glossy, naked. Fronds 90—120 c.M. long, ± 50 —60 c.M. broad. Pinnae ± 2 c.M. broad, acuminate to caudate at the apex, the lower ones short-stalked, the edge cut $\frac{1}{2}$ — $\frac{3}{4}$ of the way down into entire, spreading, blunt lobes 2 $\frac{1}{2}$ —4 m.M. broad, the basal lobes

reduced. Texture subcoriaceous; rachis naked; under surface \pm glandular; veins numerous. Sori in close rows near the midribs of the lobes. — General habit of *Mesochlaena polycarpa*, *Bedd.* (Bk.).

Philippines, Celebes, Amboyna, New Guinea.

(14) **D. Beccariana**, *C. Chr.*, *Ind. Fil.*; *Nephrodium Beccarianum*, *Ces.*, *Fel. di Bor.*, 23.

Near *D. attenuata*, *O. Ktze.* Especially the lower pinnae moderately long-stalked, long-acuminate, divaricating. — Stipes lucid, the base clothed with linear-subulate, brown, entire, lucid scales. Surfaces naked. Sori less numerous; indusium coriaceous.

Borneo.

(15) **D. Creaghii**, *C. Chr.*, *Ind. Fil.*; *Nephrodium Creaghii*, *Bk.*, *Kew Bull.*, 1898, 230.

Stipes stramineous, naked, 35—40 c.M. long. Fronds oblong-lanceolate, 2-pinnatifid, \pm 45 c.M. long, \pm 15 c.M. broad at the middle. Pinnae multijugate, sessile, lanceolate, deeply pinnatifid, 2—2½ c.M. broad, the lower ones not reduced. Lobes linear-oblong, entire, obtuse, \pm 2½ m.M. broad. Texture moderately firm; upper surface naked, under one obscurely pubescent; veins simple, erecto-patent, 7—8-jugate. Sori small, medial; indusium naked, persistent.

Borneo.

(16) **D. vulcanica**, *C. Chr.*, *Ind. Fil.*; *Nephrodium vulcanicum*, *Bk.* *Ann. of Bot.*, VIII, 127.

Stipes naked, slender. Fronds oblong-lanceolate, \pm 45 c.M. long, deeply 2-pinnatifid. Pinnae sessile, multijugate, lanceolate, acuminate, cut down nearly to the costa, the lower ones 6½—7½ c.M. long, 1½—2 c.M. broad, not reduced. Lobes linear-oblong, \pm 2½ m.M. broad, entire, the lower ones very small. Rachis slender; surfaces naked except on the costae; main veins pinnate; veins 6—8-jugate, simple, immersed, obscure. Sori small, medial; indusium naked, persistent, reniform, with a deep sinus.

Java.

* (17) **D. calcarata**, *O. Ktze.*, *C. Chr.*, *Ind. Fil.*; *Aspidium calcaratum*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 170; *Lastrea calcarata*, *Moore*, *Bedd.*, *Ferns S.I.*, tab. CCXLVI; *L. falciloba*, *Hk.*, *Bedd.*, l. c., tab. CV; *Nephrodium falcilobum*,

Hk., Spec., Fil., IV, 108; *N. calcaratum* *Hk.*, *Hk. Bk.*, Syn. Fil., 264; — var. *serecia*; *L. calcarata*, *Moore*, var. *sericea*, *Bedd.*, Handb. Ind. Ferns. 257; *L. sericea*, *Schott*, *Bedd.*, Ferns Br.I., II, tab. CCCVIII.

Rhizome erect, clothed with few, broad, lanceolate scales. Stipes tufted, 4—10 c.M. long, hairy on the upper side, deciduously scaly near the base. Fronds 10—60 c.M. long, 5—20 c.M. broad, ovate, the apex acuminate, pinatifid with linear-oblong, suberect lobes. Pinnae spreading or erecto-patent, sessile or stalked, 1—2 c.M. broad, the lower ones hardly shorter, the apex acute or acuminate, the base cuneate to truncate, the edge cut down $\frac{2}{3}$ or more to the costa into oblong, acute or bluntish, falcate or suberect lobes 1—5 m.M. broad, the lowest lobe on the anterior side not rarely elongate, \pm parallel to the rachis. Texture herbaceous but firm; rachis and surfaces villous; veins simple, 3—8 on a side. Sori medial; indusium hairy or naked, persistent. — A well-marked plant by the often nearly erected lobes. *N. falcilobum*, *Hk.* is a form with broader pinnae and deeper and more spreading lobes than usual.

Var. sericea: Pinnae short, oblong, $2\frac{1}{2}$ —5 c.M. long, 1—2 c.M. broad, obtuse at the apex or ending in a sudden point, never cordate. Indusium naked.

Malaya; Br. India, South. China, Samoa, — var. *sericea*: *Malacca, North. & South. India.*

* (18) **D. viscosa.**, **O. Ktze**, C. Chr., Ind. Fil., *Nephrodium viscosum*, *Bk.*, *Hk. Bk.*, Syn. Fil., 264; *Lastrea viscosa*, *J. Sm.*, *Bedd.*, Ferns Br.I., II, tab. CCCXXXIV.

Stipes 15 c.M. or more long, firm, erect, reddish-brown, slightly scaly, finely villose. Fronds 20—50 c.M. long, 10—15 c.M. broad, oblong-lanceolate. Pinnae close, 5—8 c.M. long, 1— $1\frac{1}{2}$ c.M. broad, cut nearly to the costa into close, spreading, linear-oblong lobes under $2\frac{1}{2}$ m.M. broad. Texture herbaceous; rachises villose; at least the under surface densely glandular; veins 5—6 on a side. Sori medial. — Lower pinnae a little shorter and deflexed.

Malacca, Sumatra, Borneo, Philippines.

* (19) **D. gracilescens**, **O. Ktze**, C. Chr., Ind. Fil.; *Aspidium gracilescens*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 170., *Lastrea gracilescens*, *Moore*, *Bedd.*, Ferns Br.I., II, tab. CCLIII; *Nephrodium gracilescens*, *Hk.*, *Hk. Bk.*, Syn. Fil., 262; — var. *decipiens*, *Bedd.*, Handb. Ind. Ferns, Suppl., 51; var. *glanduligerá*, *Bedd.*, l. c.; *A. glanduligerum*, *Kze.*, *Mett.*, *Farnng.*, IV, *Phegop. & Aspid.*, No. 207.

Rhizome short-creeping; the scales few, linear-lanceolate, entire. Stipes approximate, 15—40 c.M. long, slender, slightly scaly and short-hairy. Fronds lanceolate-oblong, 15—40 c.M. long, 8—24 c.M. broad, acuminate. Pinnae sessile, spreading, straight or falcate, acuminate, $\pm 7\frac{1}{2}$ —15 m.M. broad, deeply pinnatifid nearly to the costa, the lowest rather distant, not or not much shorter, \pm deflexed. Lobes oblong, blunt, straight or subfalcate, approximate, to ± 3 m.M. broad. Texture thin; rachis naked, or hairy above; surfaces naked, or glandular beneath, sometimes setulose on the veins; costulae pinnate in the lobes; veins 4—6 on a side. Sori in 1 row on each side of the costulae, medial or submarginal; indusium naked or minutely pilose.

Var. decipiens: Fronds shorter, deltoid; pinnae rather broader.

Var. glanduligera: Rhizome wide-creeping; stipes distant.

Malaya; Br. India; — var. *decipiens:* *Malacca; Br. India;* — var. *glanduligera:* *Malacca; Br. India, China, Japan.*

(20) **D. monodonta, C. Chr.,** Ind. Fil.; *Lastreae unidentata, Bedd.,* Handb. Ind. Ferns, Suppl., 53

Stipes nearly 60 c.M. long, densely villose, as is the rachis, with fine hair-like, jointed scales. Fronds 60—90 c.M. long, pinnate. Pinnae $17\frac{1}{2}$ c.M. long, $\pm 2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, cut down nearly to the costa into narrow-oblong segments, which are entire or nearly so except a large tooth-like lobe on one side at the base near the sinus (most prominent in the lower pinnae), lower pinnae not smaller, deflexed. Texture herbaceous to subcoriaceous; costae densely strigose above, furnished beneath with long, white, weak hairs; costulae slightly hairy on both sides; under surface covered with glandular dots; veins in 10—12 pairs, simple or the lower forked. Sori medial on the veins; indusium persistent.

Malacca.

* (21) **D. polytricha, v. A. v. R.;** *D. trichopoda, C. Chr.,* Ind. Fil.; *Nephrodium polytrichum, Blk,* Journ. of Bot., XXIX, 107; — var. *Hallieri.*

Stipes stout, 45—60 c.M. long, densely clothed with large, spreading, lanceolate-acuminate, brown scales. Fronds lanceolate-oblong, 2-pinnatifid, 90—120 c.M. long, ± 60 c.M. broad. Pinnae lanceolate, sessile, spreading, the lowest the largest, $\pm 3\frac{1}{2}$ —4 c.M. broad, cut down to a narrow wing into entire, linear-oblong segments ± 4 m.M. broad. Texture moderately firm; under surface pubescent; rachis destitute of scales except a few in the lower part; veins simple, 15—20-jugate, distinct, arcuate. Sori nearer the midrib than the edge; indusium small, fugacious.

Var. Hallieri: Fronds smaller; pinnae narrower, the lower ones narrowed towards the base; rachis deciduously scaly like the stipe; pinnae pubescent on the costa and slightly so at the margin; sori superficial, the lower ones divaricating. (Hallier No. 248, determined by Burck as *Aspidium immersum*, Bl.).

Borneo.

* (22) **D. immersa**, O. Ktze, C. Chr., Ind. Fil.; *Aspidium immersum*, Bl., Rac., Flor. Btz., I, 169; *Lastraea immersa*, Moore, Bedd., Handb. Ind. Ferns, 234; *Nephrodium immersum*, Hk., Hk. Bk., Syn. Fil., 265.

Rhizome short-creeping. Stipes approximate, 70—120 c.M. long, glossy, slightly scaly at the base. Fronds 80—200 c.M. long, 30—70 c.M. broad, ovate-oblong. Pinnae sessile, 2—4 c.M. broad, almost horizontal, acuminate or nearly caudate, cut down nearly to the costa into \pm close, spreading or nearly spreading, straight or subfalcate, subacute, linear lobes 2—5 m.M. broad. Texture herbaceous; upper surface naked or provided with distant, inconspicuous hairs; under surface and rachis provided with minute, yellow, glandular dots or stiff, white hairs. Sori near the margin, 1 to each vein, much immersed as to project on the upper surface; indusium persistent, reniform, the basal lobes overlapping.

Malaya.

* (23) **D. Peekeli**, v. A. v. R.

Stipes \pm 10 c.M. long, naked or slightly glanduloso-pilose. Fronds lanceolate or obovate, \pm 30—40 c.M. long, 15—20 c.M. broad. Pinnae 18—22 on each side below the deeply pinnatifid apex, linear-lanceolate, the central ones the largest, rather spreading, acuminate, 1—1½ c.M. broad, sessile or short-stalked, cut down nearly to the costa into linear-oblong, erecto-patent or subspreading, entire, bluntish segments 1½—2 m.M. broad, the lowest lobes not or hardly shorter. Lowest pinnae shorter, deflexed, with the lowest segments free and only those on the upper base broader and deeply toothed or pinnatifid. Texture thin; rachis and surfaces finely short-hairy; veins 8—10-jugate, simple. Sori submarginal, solitary at the base of the lobes or numerous along the edges; indusium villose.

New Guinea (Peekel, No. 44).

(24a) **D. Dayi**, C. Chr., Ind. Fil.; *Aspidium Dayi*, Bedd., Journ. of Bot., XXVI, 4; *Lastraea Dayi*, Bedd., Handb. Ind. Ferns, Suppl., 54; *Nephrodium Dayi*, Bedd., Journ. of Bot., XXV, 323.

Stipes 30—60 c.M. long, stramineous, naked, or with a few hair-like scales towards the base. Fronds 45—60 c.M. long. Pinnae 12—15 c.M. long, 2—2½ c.M. broad, the lower ones scarcely shorter, cut down nearly to the costa into linear-oblong lobes 3½—4 m.M. broad. Texture papyraceo-herbaceous, surfaces naked except the costa, which is finely puberulous above; veins 6—8 on each side, simple, not reaching the margin. Sori terminal on the veins; indusium persistent.

Malacca.

* (24b) **D. diversifolia, v. A. v. R.**

Rhizome erect or ascending. Stipes tufted or approximate, 3—6 c.M. long, pale-brown, copiously pubescent with more or less, short, whitish, spreading hairs, scaly at the base, the scales brown, subulate, ciliate. Fronds acuminate, ± 6—12 c.M. long, ± 4—9 c.M. broad, triangular with the lowest pinnae longest or obovate-lanceolate with the central pinnae longest and then the lower pinnae deflexed and gradually growing shorter (dwarfed but not reduced into mere auricles). Pinnae in ± 6—10 pairs below the pinnatifid apex, close or imbricating, the longest spreading, linear-oblong, 7—12 m.M. broad, the apex entire, blunt, the edge pinnatifid ± ¾-way down or more, or even pinnate at the very base. Secondary segments oblong, close or slightly imbricating, blunt, entire, the lowest ones not rarely free and subdistant. Texture herbaceous; rachis pubescent like the stipe; surfaces hairy on the veins and costae and minutely so between the veins beneath; costulae pinnate in the lobes; veins simple. Sori small, confined to the lobes, in 1 row on each side nearer the margin than the costulae; indusium minutely pubescent, the margin beset with globose, yellow glands, or the surface and the margin covered with yellow glands and powder.

Batu Islands (Raap, No 27).

(25) **D. patens, O. Ktze**, C. Chr., Ind. Fil.; *Aspidium patens*, Sw., Christ, Farnkr. d. Erde, 251; *A. deversum*, Kze, Mett., Farngr., IV, Phegop. & Aspid., No. 209; *Nephrodium polytrichum*, Schrad., Gött. Anz., 869; *N. patens*, Desv., Hk. Bk., Syn. Fil., 262.

Rhizome oblique, slightly creeping. Stipes 30 c.M. or more long, naked or slightly hairy, pale-coloured. Fronds 60—100 c.M. long, 20—50 c.M. broad, linear-oblong. Pinnae close, spreading, sessile with a broad base, 1—2 c.M. broad, finely acuminate, cut down ± ¾ of the space to the costa. Lobes linear-oblong, subfalcate, bluntish, entire or slightly crenate. Lowest pinnae not

reduced. Texture herbaceous; colour pale-green; rachis and surfaces more or less villose; veins simple, 6—12 on a side. Sori nearer the edge than the midrib; indusium reniform, persistent, hairy. — General aspect of *D. parasitica*, *O. Ktze*, but much larger.

New-Guinea; Polynesia, Japan, China, Trop. Africa, Trop. & Subtrop. America.

(26) ***D. exigua***, *O. Ktze*, Rev. Gen. Pl., II, 812; *D. philippina*, *C. Chr.*, Ind. Fil.; *Aspidium exiguum*, *Kze*, Mett., Farng., IV, Phegop. & Aspid., No. 180; *Nephrodium exiguum*, *Ilk*, Hk. Bk., Syn. Fil., 267.

Stipes tufted or approximate, 12—25 c.M. long, slender, grey, villosotomentose with short hairs like the rachis. Fronds 20—25 c.M. long, 3½—5 c.M. broad. Pinnae numerous, approximate, ovate-falcate, ½—1 c.M. broad, blunt, the base cordate, sometimes subauriculate on the upper side, the edge cut ± ½-way down or more to the costa into close, blunt lobes; lowest pinnae reduced. Texture papyraceo-herbaceous; veins obscure, 2—5 on each side of the main veins (constulae). Sori near the constulae.

Philippines.

(27) ***D. conterminoides***, *C. Chr.*, Ind. Fil.; *Nephrodium simulans*, *Bk.*, Journ. of Bot., XXVIII, 106 (not Journ. of Bot., XXVI, 525, nor Hk. Bk., Syn. Fil., 500).

Stipes short, naked. Fronds lanceolate-oblong, 2-pinnate, to 50 c.M. long, ± 7½ c.M. broad at the middle, narrowed gradually to the base. Pinnae lanceolate, 1—1½ c.M. broad, the central ones 2½—4 c.M. long, cut down ± to the costa into linear-oblong segments ± 2 m.M. broad. Texture moderately firm; rachis densely crinite; veins simple, erecto-patent, 5—6-jugate. Sori subcostal; indusium persistent.

New Guinea.

(28) ***D. Luerssenii***, *C. Chr.*, Ind. Fil.; *Aspidium Luerssenii*, *Christ*, Bull. Herb. Boiss., VI, 195; *Nephrodium Luerssenii*, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 29.

Stipes clustered, firm, shining, light-purplish. Fronds 45—75 c.M. long, 15—25 c.M. broad, oblong. Pinnae 10—20 c.M. long, 2¼—4 c.M. broad, lanceolate, acuminate, jointed at the base, cut down nearly or sometimes quite to the costa into numerous linear or oblanceolate lobes 2½—4 c.M. long, 2½—4 m.M. broad. Lower pinnae very small. Texture membranaceous; surfaces naked; veins 8—12 on a side, simple, free, distant. Sori terminal on the veins.

Philippines.

(29) **D. Beddomei**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium Beddomei*, Prantl, Verh. Zool. Bot. Ges., XXXI, 119; *Nephrodium Beddomei*, Bk., Hk. Bk., Syn. Fil., 267; *Lastreaa gracilescens*, Bedd., Ferns S.I., tab. CX.

Rhizome slender, wide-creeping. Stipes 15—25 c.M. long, slender, glossy. Fronds 15—50 c.M. long, 7—10 c.M. broad, oblong-lanceolate. Central pinnae the longest, 3½—5 c.M. long, ± 1 c.M. broad, ± spreading, cut down to the costa into close, rather acute, entire lobes under 2½ m.M. broad. Lower pinnae distant, dwindling down gradually. Texture firm; main veins villous beneath; veins 4—5 on a side. Sori close to the incurved margin.

Java, Philippines; South. China, South. India, Ceylon.

* (50) **D. adnata**, *v. A. v. R.*; *D. filix mas*, Schott., var., C. Chr., Ind. Fil.; *Aspidium paleaceum*, Rac. (not Bl.), Flor. Btz., I, 171; *A. adnatum*, Bl., Enum., 162.

Rhizome suberect, densely clothed with very long, yellow-brown, thin, lanceolate, entire scales. Stipes 25—55 c.M. long, the lower half scaly like the rhizome, the upper half like the rachis. Fronds linear-lanceolate, 55—100 c.M. long, 15—50 c.M. broad. Pinnae sessile, linear, 1½—2½ c.M. broad, the apex acuminate, toothed, the edge cut down nearly to the costa; lower pinnae shorter, but not reduced. Lobes linear-oblong, close, blunt, toothed at the apex, 3½—5 m.M. broad, the edge entire or slightly crenate. Texture coriaceous; rachis scaly; veins forked, bearing the sori at the base of the anterior branches, in 1 row on each side of the costulae.

Java.

(31) **D. sagenioides**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium sagenioides*, Mett., Christ, Farnkr. d. Erde, 255; *Lastreaa fuscipes*, Moore, Bedd., Handb. Ind. Ferns, 243; *L. melanopus*, Bedd., Ferns Br.I., I, tab. XXXVIII; *Nephrodium melanopus*, Hk., Spec. Fil., IV, 110; *N. sagenioides*, Bk., Hk. Bk., Syn. Fil., 271.

Rhizome short, erect. Stipes tufted, slender, 15—50 c.M. long, dark-coloured, polished, scaly at the base. Fronds 50—60 c.M. long, 15—30 c.M. broad, oblong-lanceolate. Pinnae ± 8—10 on each side below the pinnatifid apex, ± 5—4 c.M. broad, the base subtruncate, the apex short-acuminate, the edge cut down nearly to the midrib into linear-oblong, crenate or entire lobes; the lowest ones deflexed, rather shorter than the others with some of the lobes on the lower side sometimes prolonged. Texture papyraceo-herbaceous; surfaces naked; rachis polished; veins 6—10 on a side, forked, or

the higher ones simple. Sori \pm terminal on the acropetal veinlets, \pm medial between the costulae and the edge.

Malaya; Burmah.

(52) **D. singalanensis**, C. Chr., Ind. Fil.; *Lastraea singalanensis*, Bedd., Handb. Ind. Ferns, Suppl., 54; *Nephrodium singalanense*, Bk., Journ. of Bot., XVIII, 212.

Stipes densely tufted, grey, slender, to \pm 30 c.M. long, with a few small, linear, brown scales near the base. Fronds lanceolate-oblong, 2-pinnate, 45—65 c.M. long, 13—50 c.M. broad. Pinnae lanceolate, sessile, the central ones $6\frac{1}{2}$ —17 c.M. long, 2— $2\frac{1}{2}$ c.M. broad, the lower a little reduced, cut down nearly to the costa into lanceolate, entire or crenate lobes 2—4 m.M. broad. Texture rather thin; rachis substramineous and minutely pubescent; surfaces finely pubescent; veins 8—10-jugate, distinct, chiefly forked. Sori small, medial.

Sumatra, Malacca.

* (55) **D. Wigmanii**, C. Chr., Ind. Fil.; *Aspidium Wigmanii*, Rac., Bull. Ac. Crac., 1902, 61; *Nephrodium Wigmanii*, Rac., MS. in Herb. Bog.

Caudex erect, clothed like the lower part of the stipes with castaneous, lanceolate-subulate, deflexed scales. Stipes \pm 50 c.M. long, canaliculate, densely clothed like the whole frond (costae, veins and parenchyma) with erected or spreading, pale-coloured or whitish, glutinose fibrils. Fronds 100—150 c.M. long, 30—40 c.M. broad. Pinnae numerous, rather approximate, opposite, linear-lanceolate, acuminate, sessile or stalked, erecto-patent, to 25 c.M. long, the lower deflexed, $2\frac{1}{2}$ — $4\frac{1}{2}$ c.M. broad in the barren, $5\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad in the fertile fronds, deeply pinnatifid. Lobes close, 4—5 m.M. broad, crenate or serrate, linear-oblong, the base often slightly narrowed. Texture herbaceous; main veins pinnate with 4—8 pairs of mostly forked veins. The lowest posterior lobe mostly shorter than the lowest anterior one. — Fertile fronds with the lobes deeper, narrower, more distant, repando-crenate. Sori numerous, 1—2 to a crenation; indusium reniform.

Aru Islands.

* **D. filix mas**, Schott, C. Chr., Ind. Fil.; *Aspidium Filix-mas*, Sw., Christ, Farnkr. d. Erde, 256; *Lastraea Filix-mas*, Pr., Bedd., Handb. Ind. Ferns, 248; *L. odontoloma*, Moore, Bedd., Ferns S.I., tab. CXIV; *Nephrodium Filix-mas*, Rich., Hk. Bk., Syn. Fil., 272; — var. *parallelogramma*; var. *fibrillosa*, Christ, in Ann. Bot., XV, 127; *A parallelogrammum*, Kze, Linn., XIII, 146; —

var. *cochleata*, Bedd., Handb. Ind. Ferns, 250; *L. cochleata*, Moore, Bedd., Ferns S.I., tab. CXV; *D. cochleata*, C. Chr., Ind. Fil.

Rhizome oblique, stout, densely clothed with pale-brown, broadly lanceolate scales. Stipes tufted, \pm 15—20 c.M. long, scaly like the rachis, the scales lanceolate. Fronds 30—100 c.M. or more long, 20—50 c.M. broad, ovate-lanceolate. Pinnae lanceolate, acuminate, 2—4 c.M. broad, cut down nearly or quite to the rachis into numerous, blunt, close, entire, crenate or serrulate segments 5—5 m.M. broad; lower pinnae rather shorter than the others and more distant. Texture herbaceous; surfaces naked; lower veins subpinnate. Sori large, numerous, in 1 row on each side, nearer the costulae than the margin; indusium reniform, \pm convex, naked, often persistent.

Var. parallelogramma: Segments of the pinnae slightly crenate; rachis scaly, the scales linear, black-brown.

Var. cochleata: Fronds dimorphous, barren ones subdeltoid, 2-pinnatifid, fertile ones linear, pinnate, with the pinnae oblong, simple, toothed, blunt. Sori very large, completely covering the under surface of the fertile fronds.

Inhabiting almost all the temperate climates of the world except Australia; — var. *parallelogramma*: *Celebes; South. India, Columbia, Mexico; —* var. *cochleata*: *Malacca; Br. India.*

* (35) **D. syrmatica**, O. Ktze, C. Chr., Ind. Fil.; *Aspidium syrmaticum*, Willd., Rac., Flor. Btz., I, 171; *Lastrea syrmatica*, Moore, Bedd., Handb. Ind. Ferns, 243; *L. spectabilis*, J. Sm., Bedd., Ferns S.I., tab. CVIII; *Nephrodium spectabile*, Hk., Spec. Fil., IV, 115; *N. syrmaticum*, Desv., Hk. Bk., Syn. Fil., 272.

Rhizome erect, densely clothed with entire, lanceolate scales. Stipes tufted, 30—60 c.M. long, scaly at the base. Fronds 50—150 c.M. long, ovate-oblong. Pinnae \pm 10—30 c.M. long, 2—4 c.M. broad, linear-lanceolate, cut $\frac{3}{4}$ — $\frac{4}{5}$ -way down to a broadly winged rachis into ovate, bluntish, slightly falcate, toothed lobes 4—8 m.M. broad, the sinuses rounded, with a small tooth in the top. Lower pinnae not much smaller, short-stalked. Texture subcoriaceous, slightly pellucid; surfaces naked; veins 7—15 on a side, nearly all forked. Sori small, medial on the anterior veinlets, nearer the edge than the costulae.

Malaya; North. India.

(36) **D. erythrosora**, O. Ktze, C. Chr., Ind. Fil.; *Nephrodium erythrosorum*, Hk., Hk. Bk., Syn. Fil., 273; *N. prolificum*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 173.

Stipes tufted, 15—25 c.M. long, more or less densely clothed with long, lanceolate and linear scales. Fronds 50—45 c.M. long, 20—50 c.M. broad, ovate-lanceolate. Pinnae lanceolate, the lowest the largest, 15—15 c.M. long, $\pm 5\frac{1}{2}$ —4 c.M. broad, cut down quite to the rachis below into oblong, bluntish pinnulae 5—8 m.M. broad, the edge slightly (sometimes spinosely) toothed. Texture herbaceous, but firm; surfaces naked; rachis slightly scaly. Sori in rows of 6—9 to a pinnula, near the midribs; indusium red when young.

Philippines; China, Japan.

(57) **D. Preslii**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium Preslii*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 272.

Stipes 15—25 c.M. long, slender, deciduously fibrillose. Fronds 15—25 c.M. long, 10—15 c.M. broad, lanceolate-deltoid. Upper pinnae lanceolate, close, lowest pair deltoid. Upper pinnulae ± 5 m.M. broad, blunt, entire, lower ones pinnatifid with similar lobes to a broad, uncut centre. Texture subcoriaceous; rachis fibrillose: both surfaces naked. Sori about 6 to the lower lobes, dorsal on the veins, nearer the midribs than the edge.

Philippines.

(58) **D. chaerophyllifolia**, *C. Chr.*, Ind. Fil.; *Aspidium chaerophyllaeifolium*, *Zipp.*, *Ann. Mus. Bot. L. B.*, I, 228.

Caudex oblique, clothed like the base of the stipes with large, ferruginous scales. Stipes ± 50 c.M. long. Fronds to ± 45 c.M. long, ovate, acuminate. Pinnae subopposite, short-stalked, ± 15 c.M. long, ovate-oblong, acuminate, the lower ones spreading. Pinnulae spreading, $\pm 5\frac{1}{4}$ —4 c.M. long, ± 2 c.M. broad, approximate, the higher ones confluent, the lower ones short-stalked, ovate-lanceolate, the apex obtuse, serrate, the base truncate on the upper, subcordate on the lower side. Lobes not deep, with a distinct, sharp sinus, oblong, subtruncate, crenate or toothed. Texture thinly chartaceous; rachis yellowish or reddish, furnished with minute, reddish-brown, lanceolate or subulate, scattered scales; surfaces naked, the upper one opaque, the under one yellowish; main veins pinnate in the lobes with 5—4 pairs of simple veins. Sori medial on the lower veins.

Java; Aneitum.

(59) **D. edulenta**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium edulentum*, *Kze*, *Bot. Zeit.*, 1846, 474; *Nephrodium edulentum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 279.

Stipes stramineous. Fronds oblong, acute, 50 c.M. or more long and

nearly as broad. Lower pinnae 10—15 c.M. long, more than 5 c.M. broad, lanceolate-oblong, acuminate, erecto-patent. Pinnulae rather distant, spreading, connected at the base, oblong, obtuse, \pm 5 c.M. long, with oblong, obtuse, entire lobes. Texture thinly herbaceous; colour deep-green; under surface fibrillose on the main veins, the rest naked; veins 1 to each lobe. Sori close to the costae, 1—2 to a lobe; indusium thin, glabrous, persistent.

Java.

* (40) **D. flaccida**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium flaccidum*, *Bl.*, Rac., Flor. Btz., I, 176; *Nephrodium flaccidum*, *Hk.*, Hk. Bk., Syn. Fil., 274.

Rhizome erect, densely clothed with linear-lanceolate scales. Stipes tufted, 15—50 c.M. or more long, slender, stramineous, slightly scaly below, naked or villose above. Fronds linear-lanceolate, slightly narrowed towards the base, \pm 50—45 c.M. long, 15—20 c.M. broad. Pinnae sessile, linear-lanceolate, $2\frac{1}{2}$ —4 c.M. broad, cut down nearly to the rachis with a narrow, distinct, straight wing into linear-oblong lobes \pm 5 m.M. broad. Lobes lobed again \pm $\frac{1}{2}$ -way down or more to the costa. Ultimate segments entire, bluntish or acute, the edge \pm hairy. Lowest pinnae distant and often deflexed. Texture herbaceous; rachis and both surfaces villose, the under surface commonly copiously; veins forked, or subpinnate in the lower lobes. Sori about midway between the edge and the midribs.

Malacca, Java, Philippines; South. China, North. India, Ceylon.

(41a) **D. splendens**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium splendens*, *Christ*, Farnkr. d. Erde, 259; *Lastrea splendens*, *Bedd.*, Ferns Br.I., I, tab. XLII; *Nephrodium splendens*, *Hk.*, Hk. Bk., Syn. Fil., 282,

Very large.* Rhizome creeping (?). Stipes 60—100 c.M. long, stout, slightly scaly, polished, dark-brown or nearly black like the rachises. Fronds 90—120 c.M. long, 45—60 c.M. broad. Lowest pinnae stalked, to \pm 50 c.M. long, \pm 5—6 c.M. broad, numerous, elongate-lanceolate. Pinnulae numerous, unequal-sided, oblong, blunt, the lowest short-stalked, the base dilated and slightly auricled, the edge lobed, the lobes blunt, often toothed. Texture subcoriaceous; rachis naked like both surfaces; veins forked. Sori large, near the midribs; indusium coriaceous, brown.

Malacca; North. India, South. China.

(41b) **D. phegopteroidea**, *C. Chr.*, Ind. Fil.; *Hypodematium phegopteroideum*, *Kuhn*, Forsch. Gaz., IV, Farne, 8.

Stipes to 11 c.M. long, stramineous like the rachis, the base clothed with scattered, cystopteroid, lanceolate, ferruginous, acuminate scales to 5 m.M. long. Fronds 7—10 c.M. long, 6—7 c.M. broad, triangular-ovate, acuminate, 2-pinnatisect. Pinnae erecto-patent, petiolate, the lowest subopposite, triangular-ovate. Secondary segments oblong with a broad base, obtuse, entire or pinnatifid, the apex obscurely crenate. Texture membranaceous, thin; colour bright-green; both surfaces densely pilose; veins free, forked, reaching the sinuses of the crenations. Sori roundish, in 2 rows on the ultimate segments indusium membranaceous, pale-coloured, setose and ciliate; sporangia naked; spores globose or ovate, smooth.

Timor.

* (42) **D. dissecta**, **O. Ktze**, C. Chr., Ind. Fil.; *Aspidium dissectum*, Christ, Ann. Bot., XV, 156; *Lastrea dissecta*, Carr., Bedd., Handb. Ind. Ferns, 260; *L. membranifolia*, Bedd., Ferns S.I., tab. CII; *Nephrodium dissectum*, Desv., Hk. Bk., Syn. Fil., 282, p.p.; *N. conjugatum*, T. & B., MS. in Hcb. Bog.

Rhizome erect. Stipes to 50 c.M. long or longer, \pm scaly at the base. Fronds deltoid to ovate, 50—150 c.M. long, 15—90 c.M. broad, pinnate. Higher pinnae 15—25 c.M. long by 5½ c.M. broad, pinnatifid $\frac{2}{3}$ towards the costa; lower pinnae pinnate, some of the lower basal pinnulae stalked, much elongated, deeply pinnatifid, and similar to some of the upper primary pinnae. Ultimate segments crenulate. Texture herbaceous; upper surface hairy with scale-like hairs, under one naked or also hairy; veins forked. Sori terminal on the anterior veinlets, forming a single row on each side near the margin; indusium reniform, naked. — Fertile portions of the fronds not rarely more or less contracted.

Malaya; Polynesia, Trop. Australia, North. India, Madagascar.

(43) **D. papuana**, **C. Chr.**, Ind. Fil.; *Nephrodium dissitifolium*, Bk., Kew Bull., 1899, 120.

Stipes naked, glossy, pale-brown. Fronds lanceolate-oblong, 5-pinnate 45—60 c.M. long. Pinnae lanceolate, the lowest largest, short-stalked, 10—13 c.M. long. Pinnulae ovate-oblong, sessile, obtuse, the base cuneate, pinnatifid or pinnate. Segments obtuse, erecto-patent. Texture subcoriaceous; rachis naked, slender, pale-brown; veins pinnate. Sori in 1 row on each side of the costae, not contiguous; indusium naked, fugacious.

New Guinea.

* (44) **D. sparsa**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium sparsum*, Spr., Rac., Flor. Btz., I, 175; *Lastrea sparsa*, Moore, Bedd., Ferns S.I., tab. CIII; *Nephrodium sparsum*, Don, Hk. Bk., Syn. Fil., 276, p.p.; — var. *Raapii*.

Rhizome suberect; the scales dense, light-brown, entire, thin, oblong. Stipes 15—40 c.M. long, \pm scaly. Fronds 30—60 c.M. long, 10—50 c.M. broad, 3—4-pinnatifid at the base. Pinnae distant, short-stalked, $3\frac{1}{2}$ —10 c.M. broad, acuminate. Highest pinnae simple, oblong, blunt, obliquely cuneate at the base. Lowest pinnulae largest, short-stalked, blunt or acute, $1-2 \times$ pinnatifid at the base, those on the lower side prolonged. Ultimate free segments obliquely cuneate at the base, the apex mostly blunt; ultimate lobes oblong, blunt, the margin toothed with few, distant, sharp, small teeth, the teeth not rarely solitary on the anterior side of the larger lobes. Texture firm, membranaceous to papyraceous; rachis stramineous or brownish like the stipe, naked like both surfaces. Sori \pm near the costae, often 1 to each ultimate lobe.

Var. Raapii: Lower ultimate segments truncate or subcordate at the base. Each lobe with a single, small tooth on the upper side, often nearly in the top of the sinus. (*)

Malaya; South. China, North. India, Ceylon, Mauritius; — var. *Raapii*: *Tiobloug* (Raap, No. 3).

* (45) **D. Raciborskii**, *v. A. v. R.*; *Aspidium sagenioides*, Rac. (not *Mett.*), Flor. Btz., I, 178.

Stipes long, clothed with lanceolate scales and gland-like hairs. Fronds deltoid-lanceolate, to 100 c.M. long, to 60 c.M. broad. Pinnae large, distinctly stalked, deltoid, the stalks to ± 4 c.M. long. Pinnulae lanceolate or lanceolate-oblong, short-stalked, the lowest posterior ones the longest, the higher ones oblong, sessile, decurrent at the base, the highest ones united at the base by short, intermedial lobes, the longest to 18 c.M. long, to 8 c.M. broad, deeply pinnatifid or rarely subpinnate at the base. Tertiary segments to 4 c.M. long, $1-1\frac{1}{2}$ c.M. broad, bluntish or acute, the smaller ones entire, the larger ones toothed or not deeply lobed. Texture herbaceous; rachises finely short-hairy; both surfaces sparingly fibrilloso-pilose; main veins (and veins) forked or pinnate in the ultimate lobes; between each 2 lowest veins of contiguous groups is a free, simple, forked or pinnate vein which unites the midrib with the top of the sinus. Sori nearer the edge than the main veins.

Java.

(*) See also **D. purpurascens**, *v. A. v. R.*

* (46) **D. recedens**, *O. Ktze*, C. Chr. Ind. Fil.; *Lastraea recedens*, *J. Sm.*, Bedd., Ferns S.I., tab. XCVIII; *Nephrodium recedens*, *Hk.*, Hk. Bk., Syn. Fil., 281.

Rhizome wide-creeping. Stipes about 30 c.M. long, erect, villose, clothed at the base with squarrose, linear scales. Fronds 45—60 c.M. long, 50 c.M. or more broad, deltoid. Lower pinnae much the largest, 7—15 c.M. broad. Pinnulae of the lowest side the largest, often $12\frac{1}{2}$ c.M. long, to 5 c.M. broad, with distinct, lanceolate segments. Segments decurrent, with close, entire or slightly toothed, acute, linear-oblong lobes. Texture firm; rachises villose; under surface naked or villose. Sori small, 6—8 round the edge of the larger lobes.

Java, Philippines; Ceylon, South. India.

(47) ? **D. borneensis**, *O. Ktze*, C. Chr., Ind. Fil.; *Lastraea borneensis*, *Hk.*, Icon. Pl., X, tab. XMIII; *Nephrodium borneense*, *Hk.*, Hk. Bk., Syn. Fil., 275; *Davallia borneensis*, *J. Sm.*, Mett., Ann. Mus. Bot. L. B., IV, 286.

Rhizome wide-creeping, stout, densely fibrillose. Stipes \pm 15 c.M. long, firm, erect, brown, naked. Fronds 30—45 c.M. long, 10—15 c.M. broad, lanceolate. Pinnae 5—8 c.M. long, 2— $2\frac{1}{2}$ c.M. broad, cut down to a broadly winged rachis into linear-oblong, deeply pinnatifid lobes. Texture subcoriaceous; rachis and veins beneath finely villose; veins about 6 on a side. Sori \pm midway on the veins; indusium coriaceous, persistent. — Habit of *D. flaccida*, *O. Ktze*, but different in texture and mode of growth, in which it approximates *Davallia* (Bk.).

Borneo.

(48) **D. Millettii**, *C. Chr.*, Ind. Fil.; *D. setosa*, *O. Ktze*, Rev. Gen. Pl., II, 815; *Nephrodium setosum*, *Bk.*, Hk. Bk., Syn. Fil., 274.

Stipes tufted, 10—15 c.M. long, densely clothed with fibrillose scales. Fronds 50—45 c.M. long, 15—25 c.M. broad. Lowest pinnae largest, deltoid, 5—8 c.M. broad. Pinnulae lanceolate, $5\frac{1}{2}$ —5 c.M. long, cut down to the rachis below into erecto-patent, entire, linear segments. Texture herbaceous; rachises densely fibrillose; both surfaces naked; veins of the segments 5—6 on a side, the lower ones forked. Sori in a line close to the costulae.

Java.

(49) **D. padangensis**, *C. Chr.*, Ind. Fil.; *Lastraea padangensis*, *Bedd.*, Handb. Ind. Ferns, Suppl., 60.

Rhizome erect. Stipes 20—25 c.M. long, furnished with long, hair-like scales as are the rachises, costae and costulae. Fronds 30—60 c.M. long, narrowly deltoid-lanceolate. Lower pinnae largest, deltoid, with the lower pinnulae considerably elongated; upper pinnae $7\frac{1}{2}$ —10 c.M. long, pinnulae $1\frac{1}{4}$ —2 c.M. long, the lower ones cut down nearly to the costa into lanceolate, entire or slightly crenated lobes. Texture firm-herbaceous; surfaces sparingly covered with minute hairs; costulae of ultimate segments wavy; veins distant, simple or forked, not reaching the margin. Sori medial; indusium?

Malacca.

* (50) **D. vilis**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium vile*, Kze, Rac., Flor. Btz., I, 173; *Nephrodium vile*, Bk., Hk. Bk., Syn. Fil., 499.

Rhizome suberect, densely clothed with rigid, pale, linear-lanceolate, ciliate scales. Stipes 10—45 c.M. long, clothed like the rachis with subulate, blackish, brownish or pale, spreading scales, which become shorter towards the apex of the frond. Fronds ovate-deltoid, 5-pinnatifid (rarely 2-pinnate), 25—60 c.M. long, 15—40 c.M. broad. Pinnae lanceolate-deltoid, short-stalked, lowest largest. Pinnulae oblong to lanceolate, lowest sessile or subsessile, to $5\frac{1}{2}$ c.M. long, to 1 c.M. broad, deeply pinnatifid. Segments close, oblong-deltoid, the margin recurved and crenated. Texture coriaceous; under surface villose. Sori numerous, occupying nearly the whole under surface of the small lobes; indusium minute, hairy.

Java.

* (51) **D. intermedia**, *O. Ktze*, Rev. Gen. Pl., II, 813; (? *D. rhodolepis*, C. Chr., Ind. Fil.); *Aspidium intermedium*, Bl., Rac., Flor. Btz., I, 176.

Rhizome short-creeping, clothed with linear-lanceolate, pale-brown, entire scales. Stipes 3—15 c.M. long, furnished with broadly obovate, suddenly acuminate, blackish-brown, spreading scales. Fronds 2—5-pinnatifid, obovate-lanceolate, 5—55 c.M. long, 5—15 c.M. broad. Lower pinnae short-stalked, lanceolate or deltoid, unequal-sided (the lower side produced), to 3 c.M. broad; higher pinnae sessile. Lobes 2—5 m.M. broad, the lowest ones entire or pinnatifid. Texture herbaceous; rachis scaly like the stipe; main veins naked or slightly scaly beneath; veins simple. Sori medial on the veins, 1—3 on each side of the main veins (or costae). — See the information in Rac., Flor. Btz., I, 177.

Malaya.

(52) **D. sarawakensis**, *v. A. v. R.*; *D. rhodolepis*, *C. Chr.*, *var.*, *C. Chr.*, *Ind. Fil.*; *Nephrodium sarawakense*, *Bk.*, *Journ. of Linn. Soc., Bot.*, XXII, 225.

Caudex erect. Stipes tufted, \pm 15 c.M. long, clothed throughout with small, squarrose, rigid, dark-brown scales like the rachis. Fronds ovate-lanceolate, nearly 30 c.M. long, 8—10 c.M. broad at the base, 5-pinnate. Lowest pinnae largest, deltoid, opposite, short-stalked, the lower side produced, the other ones alternate, sessile, oblong-lanceolate. Pinnulae multijugate, oblong-lanceolate, the lower ones pinnate. Tertiary divisions 6—7 m.M. long, 5—5½ m.M. broad, adnate, contiguous, entire, with the central ribs pinnate. Texture thin, obscurely pubescent on the rachises and main veins; veins erecto-patent, \pm 4 on a side, the lower ones forked. Sori small, fugacious.

Borneo.

* (53) **D. pulvinulifera**, *O. Ktze*, *C. Chr.*, *Ind. Fil.*; *Aspidium pulvinuliferum*, *Rac.* (not *Bedd.*), *Flor. Btz.*, I, 177; *Lastraea pulvinulifera*, *Bedd.*, *Ferns Br.I.*, II, tab. CCCXXXIII; *Nephrodium pulvinuliferum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 500.

Caudex erect, densely clothed with linear-lanceolate, entire, yellowish or reddish-brown scales. Stipes 15—75 c.M. long, stramineous to brownish, the base scaly like the rhizome, the rest scaly like the rachises. Fronds 30—90 c.M. long, 20—40 c.M. broad, 3—4-pinnatifid, deltoid. Lowest pinnae largest, subdeltoid, stalked, \pm produced on the lower side. Pinnulae lanceolate, lower ones stalked, higher ones decurrent at the base. Ultimate lobes ligulate-oblong, the largest ones to 8 m.M. long, 3½—5 m.M. broad. Texture herbaceous or moderately firm; rachises scaly, the scales becoming shorter towards the apices of the fronds and segments, broad at the base and suddenly narrowed into a long, bristle-like, firm, spreading, dark-brown or blackish point; surfaces naked; veins pinnate in the lobes. Sori medial or subcostal, 3—5 on each side of the costulae, terminal on short veinlets. — Resembling *D. sparsa*, *O. Ktze*, but the rachises scaly.

Java; Sikkim.

(54) **D. aciculata**, *C. Chr.*, *Ind. Fil.*; *Nephrodium aciculatum*, *Bk.*, *Journ. of Linn. Soc., Bot.*, XXII, 226.

Stipes 30—45 c.M. long, densely clothed throughout like the rachises with squarrose or spreading, subulate, brown scales. Fronds oblong-deltoid,

\pm 60 c.M. long and \pm 50 c.M. broad, 3-pinnate. Lowest pinnae largest, deltoid, with the pinnulae on the lower side produced. Pinnulae multijugate, lanceolate, deeply pinnatifid, or pinnate at the base. Ultimate segments oblong, obtuse, entire, contiguous, 2—2½ c.M. long, 4—4½ m.M. broad. Texture moderately firm; surfaces naked; central veins of the segments pinnate, with 6—8 erecto-patent, obscure, hidden veins on each side. Sori small, many to a segment, subcostular; indusium small, membranaceous, fugacious.

Borneo.

* (55) **D. Boryana**, *C. Chr.*, Ind. Fil.; *Aspidium Boryanum*, Willd., Rac., Flor. Btz., I, 178; *Lastraea Boryana*, Moore, Bedd., Handb. Ind. Ferns, 266; *L. divisa*, Moore, Bedd., Ferns S.I., tab. XCVII; *Nephrodium divisum*, Hk., Copel., Polypod. Philipp., 26; *N. Boryanum*, Bk., Hk. Bk., Syn. Fil., 284.

Caudex \pm erect, clothed with thin, ovate-lanceolate scales. Stipes 60—100 c.M. long, stramineous or brownish, sometimes blackish, the base winged or dilated, deciduously scaly. Fronds lanceolate to deltoid, 120—300 c.M. long, 60—100 c.M. broad. Lower pinnae longest, \pm 10—15 c.M. broad, on stalks to 3 c.M. long. Pinnulae lanceolate, sessile, cut down to a narrowly winged rachis into blunt, spreading, sharply toothed lobes 5—15 m.M. long, 2—4 m.M. broad, which are sometimes lobed and toothed again. Texture herbaceous; rachis and surfaces naked; veins simple or forked. Sori in rows of 3—4 near the midribs of the ultimate segments; indusium fugacious.

Malaya; Japan, South. China, North. India, Reunion.

* (56) **D. purpurascens**, *v. A. v. R.*; *D. sparsa*, O. Ktze., var., *C. Chr.*, Ind. Fil.; *Aspidium purpurascens*, Bl., Rac., Flor. Btz., I, 174; *Nephrodium purpurascens*, Hk., Spéc. Fil., IV, 152, p. p.

Rhizome erect, densely clothed with thin, entire, long, linear-lanceolate scales. Stipes tufted, to 100 c.M. long, the base densely scaly, the rest slightly so. Fronds deltoid-ovate, acuminate, 60—120 c.M. long, to 80 c.M. broad, 4-pinnatifid, or rarely 5-pinnatifid at the base. Pinnae stalked, linear-oblong, the apex acuminate, inciso-serrate. Pinnulae spreading, oblong, acuminate, short-stalked, \pm 5—6 c.M. long, \pm 2—3 c.M. broad or broader at the base. Tertiary divisions trapezio-oblong, \pm 3—5 m.M. broad or broader, the higher decurrent and confluent, the lower sessile, with the base obliquely subtruncate, pinnatifid especially in the lower half with oblong, blunt, entire or slightly emarginate lobes, the superior basal one mostly larger than the others. Texture herbaceous to semicoriaceous; main rachis nearly naked,

reddish- or blackish-brown; surfaces naked; secondary rachises slightly scaly beneath; veins 4—6 on each side. Sori \pm medial, mostly 1 to each lobe. — Resembling *D. sparsa*, *O. Ktze*, but larger and more divided.

Malaya.

* (57) **D. crenata**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium crenatum*, *Kuhn*, Christ, Farnkr. d. Erde, 262; *Lastraea crenata*, *Bedd.*, Handb. Ind. Ferns, 258; *L. criocarpa*, *Pr.*, *Bedd.*, Ferns S.I., tab. XCX; *Nephrodium crenatum*, *Bk.*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 175; *N. odoratum*, *Bk.*, *Hk. Bk.*, Syn. Fil., 280; *Polypodium crenatum*, *Forsk.*, (oldest name).

Rhizome erect, stout, short; the scales dense, bright-brown, lanceolate-subulate, \pm 5 c.M. long. Stipes tufted, 20—50 c.M. long, pale-coloured, glossy, the base scaly like the rhizome. Fronds 20—40 c.M. long, deltoid. Lowest pinnae much the longest, \pm 10—25 c.M. long, 5—10 c.M. broad. Pinnulae lanceolate, often imbricate. Tertiary divisions oblong, decurrent, pinnatifid, the lobes rounded. Texture herbaceous; colour pale-green; rachis and surfaces densely furnished with greyish hairs. Sori copious; indusium large, pale-grey, hairy.

Malacca, Philippines; Cape Verde Islands to China.

* (58) **D. rhodolepis**, *C. Chr.*, Ind. Fil.; *Aspidium intermedium*, *Christ* (not *Bl.*), Farnkr. d. Erde, 263; *Lastraea Blumei*, *Moore*, *Bedd.*, Ferns S.I., tab. CCIL; *Nephrodium Blumei*, *Hk.*, Spec. Fil., IV, 155, p.p.; *N. intermedium*, *Bk.*, *Hk. Bk.*, Syn. Fil., 285; *N. rhodolepis*, *Clarke*, Trans. Linn. Soc., Bot., 2^d Series, I, tab. LXXII.

Stipes 50—60 c.M. long, stout, erect, densely clothed at the base with long, bright-brown, silky fibrils. Fronds 60—100 c.M. long, 50—50 c.M. broad, subdeltoid. Lower pinnae lanceolate, often 50 c.M. long, 10—15 c.M. broad. Pinnulae close, lanceolate, with distinct, oblong-lanceolate segments, with small, ligulate, subentire lobes to about 2½ m.M. broad. Texture herbaceous, firm; rachises slightly fibrillose like the stipe; under surface slightly hairy or scaly on the midribs. Sori small, copious, medial or nearer the midribs than the edge; indusium thin, fugacious.

Trop. Asia; China, Japan.

* (59) **D. setigera**, *O. Ktze*, C. Chr., Ind. Fil.; *Aspidium setigerum*, *Kuhn*, Rac., Flor. Btz., I, 178; *Lastraea tenericaulis*, *Moore*, *Bedd.*, Handb. Ind. Ferns, 266; *Nephrodium tenericaule*, *Hk.*, Spec. Fil., IV, tab. CCLXIX; *N. setigerum*, *Bk.*, *Hk. Bk.*, Syn. Fil., 284; — var. *pallida*; *Polypodium pallidum*, *Brack.*, *Hk. Bk.*, l. c.; *D. ornata*, *C. Chr.*, var., C. Chr., Ind. Fil.

Rhizome erect or suberect. Stipes tufted or approximate, 25—60 c.M. long, often glaucous, scaly throughout or at the base only. Fronds deltoid-lanceolate, 50—100 c.M. long. Lower pinnae largest, often 20—30 c.M. long, 10—15 c.M. broad, sessile or short-stalked, linear-lanceolate, with the rachis often narrowly winged above. Pinnulae close, ovate to linear-lanceolate, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, cut down nearly or quite to the rachis into close, pinnatifid lobes $1\frac{1}{2}$ — $2\frac{1}{2}$ m.M. broad. Texture herbaceous; main rachis hairy on the upper side; rachises and under surface finely villose, the hairs whitish, scattered. Sori minute, copious, 4—6 on each side of the costulae; indusium fugacious.

Var. pallida: Indusium suppressed and undeveloped, visible only by 50—100 \times magnifying power, soon hidden between the much larger capsules.

Br. India across China and Japan, Polynesia and Trop. Australia.

* (60) **D. zeylanica, v. A. v. R.**; *D. obtusiloba*, O. Ktze (not C. Chr.), Rev. Gen. Pl., II, 813; *D. peranemiformis*, C. Chr., var.; C. Chr., Ind. Fil.; *Aspidium obtusilobum*, Prantl (not Bk.), Christ, Ann. Bot., XV, 150; *Lastrea obtusiloba*, Bedd., Ferns Br.I., II, tab. CCXCVI; *Nephrodium obtusilobum*, Bk., Hk. Bk., Syn Fil., 284.

Stipes 30—60 c.M. long, firm, erect, densely clothed with large, ovate scales. Fronds 90—120 c.M. long, 45—60 c.M. broad. Lower pinnae longest, to 15 c.M. broad. Pinnulae lanceolate; most of the segments of the lower pinnae distinct, ovate, oblong, cut down to the rachis into oblong-rhomboidal, \pm entire lobes, the largest to 5—6 m.M. long, to $2\frac{1}{2}$ —3 m.M. broad. Texture herbaceous; rachises more or less scaly, the upper ones, and especially the under surface more or less glandular; lower veins of the lobes forked. Sori medial.

Celebes; Ceylon.

* (61) **D. multiseta, C. Chr.**, Ind. Fil.; *Nephrodium multisetum*, Bk., Journ. of Linn. Soc., Bot., XXII, 226.

Stipes tufted, long, pale, densely clothed like the rachises with long, spreading or squarrose, rigid, subulate, castaneous bristles. Fronds deltoid, decomposed, \pm 120 c.M. long. Lower pinnae oblong-lanceolate, 50—45 c.M. long, 10—15 c.M. broad. Pinnulae multijugate, lanceolate, 1—2 c.M. broad, the largest with 12—15 adnate, distinct, pinnatifid divisions on each side below the pinnatifid apex. Tertiary divisions linear-oblong, \pm 2 m.M. broad; the lobes small, deltoid. Texture herbaceous; veins forked in the largest

lobes. Sori small, medial, 1—2 to a final segment; indusium membranaceous, fugacious.

Borneo.

(62) **D. subarborea**, C. Chr., Ind. Fil.; *Aspidium megaphyllum*, Christ, Bull. Herb. Boiss., VI, 194; *A. Filix-Mas*, Sw., var. *megaphyllum*, Christ, Ann. Btz., XV, 127; *Polypodium subarboreum*, Bk., Journ. of Linn. Soc., Bot., XXIV, 259; *Nephrodium megaphyllum*, Bk., l. c., XXII, 227; — var. *Warburgii*, Christ, in Warb, Mons., I, 81; *Alsophila Warburgii*, Christ, Ann. Btz., XV, 80, tab. XIII, fig. 5, a—e.

An enormous fern. Stipes rough, clothed with adpressed, subulate, brown scales like the rachis. Fronds large, deltoid, 4-pinnate. Pinnae 45—75 c.M. long, \pm 15 c.M. broad, stalked, acuminate, oblong-lanceolate. Pinnulae ovate-oblong, remote, sessile, alternate. Tertiary divisions oblong, \pm sessile, 2—5 c.M. long, the rachis narrowly winged. Ultimate segments \pm 4 m.M. long, toothed. Texture herbaceous; surfaces furfuraceous. Sori single at the base of the ultimate lobes; indusium membranaceous, fugacious.

Var. Warburgii: Scales of the stipe dark-reddish-brown, long-acuminate from a broad base, the higher ones obliterate, mixed with rough, short prickles. Stipes brown, rachis furfuraceous, polished; secondary rachises densely clothed with reddish-brown, ovate scales like the veins beneath.

Malaya; — var. *Warburgii*: *Batjan, Celebes.*

§ 2. NEPHRODIUM.

1. Fronds simple or pinnate, the terminal pinna much exceeding the lateral ones. (65) **D. Bakeri**.
- 1.1. Fronds pinnate or 2-pinnatifid; the terminal pinna not or hardly exceeding the largest lateral ones.
2. Rhizome wide-creeping ⁽¹⁾.
3. Lower pinnae slightly, if at all, reduced; the fronds occasionally with a single pair of dwarfed or reduced pinnae at the base ⁽²⁾.
4. Texture \pm herbaceous.
5. Pinnae entire, or cut less than $\frac{1}{2}$ -way down tho the costa.

⁽¹⁾ See also **D. parasitica**, O. Ktze.

⁽²⁾ See also **D. invisa**, O. Ktze.

6. *Pinnae entire or crenate.* (64) *D. lineata.*
- 6.6. *Pinnae lobed.*
7. *Sori ± marginal.* (65) *D. pteroides.*
- 7.7. *Sori not marginal.* (66) *D. otaria.*
- 5.5. *Pinnae cut more than 1/2-way down to the costa.*
- α. *Rachis ± scaly and hairy.* (67) *D. alatella.*
- α.α. *Rachis naked or ± hairy, but not scaly.*
- β. *Sori ± terminal on the veins.* (68) *D. extensa.*
- β.β. *Sori ± medial on the veins.*
- γ. *Pinnae ± 1 1/2 c.M. broad, cut down nearly to the costa.* (69) *D. procurrens.*
- γ.γ. *Pinnae 2—2 1/2 c.M. broad, cut 2/3-way down to the costa.* (70) *D. longipes.*
- 4.4. *Texture ± coriaceous.* (71) *D. gongylodes.*
- 3.3. *Lower pinnae conspicuously reduced or dwindling down into mere auricles.*
- p. *Pinnae cut to ± 1/3-way down to the costa.*
- q. *Lower pinnae conspicuously dwarfed only. (*)* (72) *D. invisā.*
- q.q. *Lower pinnae reduced into mere auricles.*
- r. *Sori ± medial on the veins; veins resinous beneath; lower pinnae ± gradually reduced.* (73) *D. arida.*
- r.r. *Sori subterminal on the veins; under surface glandular; lower pinnae suddenly reduced.* (74) *D. cucullata.*
- p.p. *Pinnae cut 1/3—1/2-way down to the costa.*
- *. *Fronde generally under 50 c.M. long. Under surface glandular.* (75) *D. appendiculata.*
- *.*. *Fully developed fronds more than 50 c.M. long.*
- †. *Sori ± terminal on the veins. Under surface pubescent.* (76) *D. Haenkeana.*
- †.†. *Sori ± medial on the veins.*
- !. *Stipes naked. Under surface pilose on the ribs.* (77) *D. eminens.*
- !.!. *Stipes villose. Under surface glandular.* (78) *D. hirsuta.*
- p.p.p. *Pinnae cut more than 1/2-way down to the costa. Surfaces naked.* (79) *D. ochtodes.*
- 2.2 *Rhizome short-creeping or erect.*
- α. *Pinnae entire, crenate, toothed or slightly lobed.*

(*) See also under 3.

- b.** Lower pinnae slightly, if at all, reduced, not dwindling down into mere auricles ⁽¹⁾; the fronds occasionally with a single pair of dwarfed or reduced pinnae at the base.
- c.** Fronds under 30 c.M. long. Under surface glandular and ± hairy. (80) *D. pilosiuscula*.
- c.c.** Fully developed fronds more than 30 c.M. long.
- d.** Fronds rather dimorphous; fertile pinnae distinctly narrower than the barren ones. Under surface short-hairy and glandular. (81) *D. glandulosa*.
- d.d.** Fronds not dimorphous.
- e.** Sori contiguous, forming transverse bands of brown lines close to the main veins. (82) *D. cyatheoides*.
- e.e.** Sori not contiguous, rather distant from the main veins.
- f.** Veins 8—14 on each side of the main veins. (85) *D. urophylla*.
- f.f.** Veins 15—20 on each side of the main veins. (84) *D. moulemeinensis*.
- b.b.** Lower pinnae conspicuously reduced or dwindling down into mere auricles.
- c'.** Surfaces naked.
- d'.** Veins 3—5 on each side of the main veins. (85) *D. latipinna*.
- d'.d'.** Veins 6—8 on each side of the main veins. (86) *D. abortiva*.
- c'.c'.** Under surface ± hairy, at least on the veins.
- s.** Lower pinnae suddenly reduced. (87) *D. megaphylla*.
- s.s.** Lower pinnae gradually reduced.
- ‡. The 2—3 lower veins anastomosing.
- *. Fronds 10—20 c.M. broad. (88) *D. arbuscula*.
- *.*. Fronds 20—30 c.M. broad. (89) *D. amboinensis*.
- ‡.‡. The 5—6 lower veins anastomosing. (90) *D. multilineata*.
- a.a.** Pinnae cut from $\frac{1}{4}$ — $\frac{2}{3}$ -way down to the costa.
- b'.** Texture ± coriaceous.
- c'.** Lower pinnae slightly, if at all, reduced; the fronds occasionally with a single pair of dwarfed or reduced pinnae at the base ⁽²⁾.
- d'.** Stipe and rachis purveyed with large, distant glands, those of the rachis placed on the under side at the base of the pinnae. (91) *D. callosa*.

(¹) See also *D. lineata*, C. Chr.

(²) See also *D. crassifolia* & *echinata*, O. Ktze.

- d.d.** *Stipe and rachis without glands.*
- e.** *Stipes naked.*
- f.** *Veins ± 7 on each side of the main veins. (92) D. laevis.*
- f.f.** *Veins 10—15 on each side of the main veins.*
(93) D. brachyodus.
- e.e.** *Stipes hairy or bristly. (94) D. ferox.*
- c.c.** *Lower pinnae conspicuously reduced or dwindling down into mere auricles.*
- s.** *Sori submarginal. (95) D. Loheriana.*
- s.s.** *Sori \pm medial.*
- ***. *Costae with a large gland at the base beneath.*
(96) D. latebrosa.
- *.*.** *Costae without gland at the base beneath.*
- †.** *Reduced pinnae few, distant. Surfaces rather naked.*
(97) D. piloso-squamata.
- †.†.** *Reduced pinnae numerous, close. Surfaces conspicuously hairy.*
(98) D. sagittifolia.
- b.b.** *Texture herbaceous to firm-papyraceous.*
- a.** *Lower pinnae hardly, if at all, reduced; the fronds occasionally with a single pair of dwarfed or reduced pinnae at the base.*
- β .** *Pinnae cut less than $\pm \frac{1}{2}$ -way down to the costa.*
- γ .** *Sori medial on the veins.*
- δ .** *Surfaces naked. (99) D. caudiculata.*
- $\delta.\delta$.** *Surfaces hairy, at least on the ribs beneath.*
- ϵ .** *Lowest pinnae sharply deflexed. (100) D. simillima.*
- $\epsilon.\epsilon$.** *Lowest pinnae not sharply deflexed.*
- †.** *Lowest 1—2 pairs of veins anastomosing.*
- ***. *Pinnae cut $\frac{1}{4}$ — $\frac{1}{3}$ -way down to the costa. The 2 lower veins anastomosing. (101) D. pseudoreptans.*
- *.*.** *Pinnae cut $\pm \frac{1}{2}$ -way down to the costa. Only the lowest veins anastomosing. (102) D. parasitica.*
- †.†.** *Lower 5—8 pairs of veins anastomosing.*
(103) D. indica.
- $\gamma.\gamma$.** *Sori near the main veins. Upper surface \pm hairy, the under one \pm glandular. (104) D. sakayensis.*
- $\beta.\beta$.** *Pinnae cut more than $\pm \frac{1}{2}$ -way down to the costa. (*)*

(*) See also **D. parasitica**, O. Ktze.

- ‡. *Sori submarginal, or terminal on the veins.*
 *. *Pinnae ± 6 c.M. long, cut $\frac{1}{2}$ — $\frac{2}{3}$ -way down.* (105) *D. Benoitiana.*
 *. *. *Pinnae ± $7\frac{1}{2}$ —9 c.M. long, cut $\frac{2}{3}$ -way down or more.* (106). *D. didymosora.*
 ‡. ‡. *Sori medial, or lower on the veins.* (107) *D. microchlamys.*
- α.α. Lower pinnae conspicuously reduced or dwindling down into mere auricles.*
β' Lower pinnae gradually reduced.
 *. *Stipe and rachis glaucous.* (108) *D. glaucostipes.*
 *. *. *Stipe and rachis not glaucous.*
 ‡. *Fronde under 40 c.M. long.* (109) *D. perakensis.*
 ‡. ‡. *Fully developed fronds more than 100 c.M. long.* (110) *D. sumatrana.*
- β'.β'. Lower pinnae suddenly reduced.*
γ'. Pinnae cut less than $\frac{1}{2}$ -way down to the costa ()*
δ'. The lowest veins anastomosing. Pinnae provided with a short, thick gland at the base beneath. (111) *D. stipellata.*
δ'δ' The 2—3 lower veins anastomosing. Pinnae without gland. (112) *D. truncata.*
γ'.γ'. Pinnae cut more than $\frac{1}{2}$ -way down to the costa.
 *. *Veins 3—4 on each side of the costulae.* (113) *D. hispidula.*
 *. *. *Veins 6—10 on each side of the costulae.*
 ×. *Stipe and rachis hairy.* (114) *D. heterocarpa.*
 ×. ×. *Stipe and rachis scaly.* (115) *D. crinipes.*

(65). **D. Bakeri**, *Copel.*, Philipp. Journ., II, 405; *D. canescens*, *C. Chr.*, var., *C. Chr.*, Ind. Fil.; *Aspidium canescens*, *Christ*, Ann. Bot., XV, 150, p.p.; *Nephrodium Bakeri*, *Harr.*, *Copel.*, Polypod. Philipp., 27; *Hk.*, Icon. Pl., XVII, tab. MDCLXIV.

Stipes clustered, light-brown, villose or scaly, 5—15 c.M. long. Fronds 7—15 c.M. long, 1— $2\frac{1}{2}$ c.M. broad, oblong-oblongeolate, obtuse, coarsely crenate, the segments extending about $\frac{1}{3}$ the distance to the midrib, with usually 2—4 distant, oblong-ovate, sessile pinnae at the base, rather smaller than the

(*) See also **D. abortiva** *O. Ktze.*, & **megaphylla**, *C. Chr.*

lobes. Texture papyraceous; surfaces villose throughout. Veins usually $\pm 6-8$ on each side, of which $\pm 4-5$ unite with the opposite veins of adjacent lobes. Sori small, medial on the veins; indusium hairy, often absent.

Philippines.

* (64) **D. lineata**, C. Chr., Ind. Fil.; *Nephrodium lineatum*, Pr., Rac., Flor. Btz., I, 186; *N. glandulosum*, J. Sm., Hk. Bk., Syn. Fil., 291, p. p.; *Aspidium lineatum*, Bl., Enum., 144.

Rhizome creeping, bearing on regular distances groups of tufted or approximate fronds, the scales brown, narrow-lanceolate. Stipes 15—50 c.M. long, those of the fertile fronds much elongated. Fronds $\pm 25-45$ c.M. long, lanceolate. Pinnae in $\pm 12-14$ pairs, the barren ones $5\frac{1}{2}-7$ c.M. long, 1—2 c.M. broad, sessile, opposite or alternate, sometimes very close and overlapping, lanceolate, acute, erecto-patent, the base truncate, auricled on the upper side and sometimes on the lower side at the base, the margin entire or more or less crenate particularly towards the apex, the lower pinnae not rarely deflexed; fertile pinnae shorter and generally contracted, distant. Texture firm-herbaceous; rachis strigose; surfaces nearly naked but verruculose; veins 5—5 on each side, all anastomosing. Sori terminal on the veins, elongated, soon confluent and becoming meniscioid; indusium villose or glandular, at length deciduous.

Malaya.

* (65) **D. pteroides**, O. Ktze, C. Chr., Ind. Fil.; *Nephrodium pteroides*, J. Sm., Rac., Flor. Btz., I, 185; *N. terminans*, J. Sm., Bedd., Ferns S.I., tab. XC; *Aspidium pteroides*, Sw., Christ, Farnkr. d. Erde, 247.

Rhizome creeping, the scales narrow, lanceolate. Stipes distant, stramineous or greyish, 50—60 c.M. long, scaly at the base. Fronds 50—120 c.M. long, 20—40 c.M. broad, deltoid-oblong. Pinnae 6 to many on each side, spreading or erecto-patent, the lowest largest, short-stalked, $1-2\frac{1}{2}$ c.M. broad, acuminate, the edges cut $\frac{1}{3}-\frac{1}{2}$ -way down. Lobes oblong or subdeltoid, straight or subfalcate, to ± 7 m.M. long, 4—5 m.M. broad. Texture \pm herbaceous; rachis and both surfaces nearly naked except on the \pm villose costae and veins; main veins pinnate in the lobes with 5—12 veins on each side, the 2—3 lower veins anastomosing. Sori submarginal, nearly confined to the lobes.

Malaya; China, Burmah, Queensland, Polynesia.

* (66) **D. otaria**, O. Ktze, C. Chr., Ind. Fil.; *Nephrodium otaria*, Bk., Hk. Bk., Syn. Fil., 288; *N. aristatum*, Hk., Spec. Fil., IV, tab. CCXXXVIII;

Pleocnemia aristata, *Bedd.*, Ferns S.I., tab. LXXXIII; *Aspidium otaria*, *Kze*, *Christ*, *Farnkr. d. Erde*, 249.

Rhizome creeping. Stipes to 15—50 c.M. long, substramineous. Fronds 15—50 c.M. or more long, with a linear-oblong, terminal pinna 5—15 c.M. long, 2—4 c.M. broad, the apex acuminate, the margin with finely serrated, lanceolate lobes 4—8 m.M. long. Below this 5—6 distant, spreading, similar pinnae on each side, the lower ones stalked. Texture thinly herbaceous; rachis and both surfaces naked; main veins pinnate in the lobes; veins 4—8 on each side, with sometimes a sorus on each.

Philippines; Assam, South. India, Ceylon.

(67) **D. alatella**, **C. Chr.**, *Ind. Fil.*; *Nephrodium alatellum*, *Christ*, *Schum. & Laut., Flor. deut. Sch. geb.*, 112; *Aspidium alatellum*, *Christ*, *Bull. Herb. Boiss.*, 2^e Série, I, 454.

Rhizome not very wide-creeping. Stipes \pm 60 c.M. long, dark-yellow, smooth, clothed near the base with ovate, acuminate, broad, brown scales, the rest sparingly scaly and hairy like the rachis. Fronds 35—40 c.M. long, ovate-oblong, \pm 15 c.M. broad, not narrowed towards the base. Pinnae erecto-patent, 20—25 on each side, sessile, not adnate, 10—25 c.M. long, \pm 1 $\frac{1}{2}$ c.M. broad, linear-lanceolate, caudate-acuminate, the point nearly entire, the margin cut $\frac{2}{3}$ -way down to the costa into numerous (\pm 50 on each side), truncate, coarsely crenate lobes \pm 5 m.M. broad. Texture flaccido-herbaceous; costae and main veins naked or slightly hairy; main veins pinnate in the lobes; veins simple, the lower ones anastomosing. Sori medial on the veins; indusium rudimentary, thin, fugacious, not rarely wanting.

New Guinea.

* (68) **D. extensa**, **O. Ktze**, *C. Chr.*, *Ind. Fil.*; *Nephrodium extensum*, *Moore* (not *Hk.*), *Hk. Bk., Syn. Fil.*, 289; *N. punctatum*, *Bedd.*, *Ferns Br.I.*, I, tab. CXXXI; *Aspidium extensum*, *Bl.*, *Christ*, *Farnkr. d. Erde*, 247; *A. conioneuron*, *Fée*, *Mett., Farng., IV, Phegop. & Aspid.*, No. 245.

Rhizome creeping. Stipes 50—60 c.M. long, naked or slightly pubescent. Fronds 45—120 c.M. long, 20—50 c.M. broad. Pinnae 1—2 c.M. broad, cut down \pm $\frac{2}{3}$ on the way to the costa into linear-oblong, falcate or straight, slightly crenulate lobes; lower pinnae scarcely shorter than the others. Texture papyraceo-herbaceous; rachis nearly naked; upper surface naked or with a few distant hairs; under surface hairy, and glandular with minute glands between or on the veins; main veins pinnate in the lobes; veins 6—12 on

each side, often only the lowest united. Sori nearly terminal on the veins, present on all the veins, or on all except the 1—2 lower ones, not rarely immersed and visible as punctiform dots on the upper surface.

Malaya; Ceylon, South. India, Burmah.

(69) **D. procurrens**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium procurrens*, *Bk.*, Hk. Bk., Syn. Fil., 290; *Aspidium procurrens*, *Mett.*, Ann. Mus. Bot. L. B., I, 251.

Rhizome wide-creeping, clothed sparingly with deciduous, linear scales. Stipes approximate, up to 45—60 c.M. long, slightly and deciduously scaly towards the base. Fronds deltoid to deltoid-lanceolate, 45 c.M. or more long, nearly 50 c.M. broad; rachis more or less hairy or nearly glabrous. Pinnæ 12—30 on each side, patent, the lowest sometimes a little smaller and deflexed, narrow-linear, acuminate, $\pm 1\frac{1}{2}$ c.M. broad, cut down nearly to the costa into narrow, oblong, entire lobes, the lowest of which often suddenly reduced in size. Texture papyraceo-membranaceous, more or less hairy or nearly glabrous, often furnished beneath with golden pubescence in the way of close, pressed hairs or oblong dots more on the paranchyma than on the veins; veins in 6—12 pairs, all simple, the lowest pair only anastomosing. Sori medial on the veins; indusium naked or hairy. — Very like *D. parasitica*, *O. Ktze*, (Bk.).

Java, Borneo, Celebes; North. India.

* (70) **D. longipes**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium longipes*, *Moore*, Rac., Flor. Btz., I, 182; *Aspidium longipes*, *Bl.*, Christ, Ann. Btz., XV, 133.

Rhizome creeping, the extremity clothed with few, small scales. Stipes 1—8 c.M. apart, 80—120 c.M. long, livid, naked, glossy. Fronds linear-lanceolate, to 100 c.M. long, 20—45 c.M. broad. Pinnæ numerous (20—35 on each side), 2—2 $\frac{1}{2}$ c.M. broad, sessile, slightly narrowed towards the base, linear-lanceolate, acuminate, the edge cut down $\frac{2}{3}$ of the way to the costa into oblique, oblong, bluntish or acute lobes. Lobes straight or subfalcate, 10—14 m.M. long, ± 4 m.M. broad, with a distinct sinus between them. Texture herbaceous; the edges and both surfaces more or less hairy on the costae and veins; veins 8—16 on a side, often only the lowest ones anastomosing. Sori ± 6 —9 on each side of the main veins, medial, not occupying the higher veins.

Java, Celebes.

* (71) **D. gongylodes**, **O. Ktze**, C. Chr., Ind. Fil.; *Nephrodium unitum*, R. Br., Rac., Flor. Btz., I, 182; *N. gongylodes*, Schott, Gen. Fil., tab. X; *Aspidium unitum*, Mett. (not Sw.), Christ, Farnkr. d. Erde, 246; *A. gongylodes* (*goggilodes*), Schk., (oldest name); — var. *propinqua*; *N. propinquum*, R. Br., Hk., Spec. Fil., IV, 79; Bedd., Ferns S.I., tab. LXXXIX.

Rhizome wide-creeping, the extremity clothed with ovate-deltoid scales. Stipes 20—50 c.M. long, brownish, naked. Fronds 50—60 c.M. or more long \pm 10—20 c.M. broad. Pinnae numerous, erecto-patent, 5—12 $\frac{1}{2}$ c.M. long, \pm $\frac{1}{2}$ —1 $\frac{1}{2}$ c.M. broad, the edges cut $\frac{1}{4}$ — $\frac{1}{2}$ -way down into \pm erecto-patent, triangular, acute, entire or toothed lobes; lower pinnae not or a little smaller than the others. Texture coriaceous; veins 5—8 on each side of the main veins, the 1—2 lowest ones anastomosing; rachis and surfaces naked. Sori nearly terminal on the veins, at length confluent and forming a zigzag line along the margin; indusium small.

Var. propinqua: Rachis and under surface hairy.

Pantropical.

* (72) **D. invis**a, **O. Ktze**, C. Chr., Ind. Fil.; *Nephrodium invisum*, Carr., Hk. Bk., Syn. Fil., 290; *Aspidium dissectum*, Mett., Ann. Mus. Bot. L. B., I, 252; *A. invisum*, Christ, Engl. Jahrb., XXIII, 350.

Rhizome stout, creeping. Stipes 25—30 c.M. long or longer, stout, brownish, villose. Fronds 40—60 c.M. long, \pm 20—50 c.M. broad. Pinnae numerous, \pm 1—1 $\frac{1}{2}$ c.M. broad, cut \pm $\frac{1}{3}$ -way down to the costa into sharp, triangular, \pm falcate lobes; lower pinnae dwarfed, distant. Texture rather coriaceous; rachis and under surface villose, upper surface \pm short-pubescent; main veins pinnate in the lobes; veins 6—8 on each side. Sori in 1 row on each side, \pm close to the midribs or more numerous and then in pairs close to the intermedial veins; sporangia setose. — Fronds with the aspect of *D. cucullata*, Christ, but the lower pinnae conspicuously dwarfed only, not reduced into mere auricles.

Philippines, New Guinea; Polynesia.

* (75) **D. arida**, **O. Ktze**, C. Chr., Ind. Fil.; *Nephrodium obscurum*, Moore, Rac., Flor. Btz., I, 185; *N. aridum*, J. Sm. (not Bk.), Hk., Bk., Syn. Fil., 291; Bedd., Ferns Br.I., II, tab. CCXCVII; *Aspidium aridum*, Don, (oldest name), Christ, Farnkr. d. Erde, 247.

Rhizome creeping. Stipes 15—30 c.M. long or longer, erect, naked or pubescent. Fronds 40—120 c.M. long, 20—45 c.M. broad, deltoid-lanceolate.

Pinnae \pm 1—2 $\frac{1}{2}$ c.M. broad, linear, the base \pm truncate, the edges cut \pm $\frac{1}{4}$ — $\frac{1}{3}$ -way down to the costa, the apex acuminate. Lobes numerous, subtriangular, \pm acute, slightly recurved at the edge. Lower pinnae more or less gradually reduced into mere auricles. Texture firm; rachis naked or pubescent; surfaces pubescent or pilose on the costae and veins; veins \pm 8—11 on each side of the main veins, resinous beneath, the secretions minute, oblong, brown or sometimes yellow, pellucid, the 3—6 lower veins anastomosing. Sori small, \pm medial on the veins; indusium slightly hairy, \pm fugacious; sporangia naked.

Malaya; North. India.

* (74) **D. cucullata**, *Christ*, Philipp. Journ., II, 194; *D. unita*, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium cucullatum*, *Bk.*, Rac., Flor. Btz., I, 184; *N. unitum*, *Bory*, Bél., Voy. Bot., II, 61; *Bedd.*, Ferns S.I., tab. LXXXVIII; *Aspidium cucullatum*, *Bl.*, Christ, Farnkr. d. Erde, 247; *A. unitum*, *Sieb.*, *Mett.*, Farng., IV, Phegop. & Aspid., No. 257.

Rhizome creeping. Stipes distant, 10—55 c.M. long, strong, erect, brownish, more or less villose like the rachis, scaly at the base. Fronds ovate-lanceolate, 55—80 c.M. long, 18—50 c.M. broad. Pinnae rather close, \pm 1—2 c.M. broad, linear, erecto-patent, the base \pm truncate, the edges cut \pm $\frac{1}{3}$ -way down to the costa, the apex acuminate. Lobes triangular or oblong, bluntish or caute, subfalcate, the margin slightly recurved. Lower pinnae dwindling down suddenly into mere auricles. Texture coriaceous; costae hairy; under surface hairy on the veins, glandular between or on the veins, the glands minute, globose, brown or more commonly yellow or pale-yellow, impellucid; veins 7—10 on each side of the main veins, the 3—5 lower ones anastomosing. Sori subterminal on the veins; indusium naked.

Malaya; Polynesia, Ceylon, South. India, Mascarenes, Seychelles.

* (75) **D. appendiculata**, *C. Chr.*, Ind. Fil.; *Nephrodium appendiculatum*, *Rac.*, Flor. Btz., I, 186; *Gymnogramme appendiculata*, *Bl.*, Flor. Jav., II, tab. XXXIX.

Rhizome creeping, clothed at the extremity with narrow, lanceolate, ciliated scales. Stipes 15—18 c.M. long, densely villose. Fronds oblong-lanceolate, 20—40 c.M. long, 10—16 c.M. broad. Pinnae linear-lanceolate, sessile, to 15 m.M. broad, commonly cut \pm $\frac{1}{2}$ -way down into entire, oblong, subfalcate, bluntish or acute lobes to \pm 5 m.M. long and 5 $\frac{1}{2}$ m.M. broad. The lower, not-reduced pinnae deflexed, the reduced ones \pm 2 c.M. apart,

occupying nearly the whole stipe. Texture subcoriaceous; under surface slightly villose on the veins and covered with globose glands, upper one clothed with adpressed hairs, the edges hairy; veins 5—9 on a side, the lowest ones only anastomosing. Sori 5—6 on a side, medial on the veins; indusium hairy, fugacious; capsules hairy.

Java.

(76) **D. Haenkeana**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium Haenkeanum*, *Pr.*, Hk. Bk., Syn. Fil., 290; *N. multilineatum*, *Pr.*, Tent. Pterid., 81.

Rhizome creeping. Stipes scattered, 30—60 c.M. long, strong, erect, slightly pubescent. Fronds 90—120 c.M. long, 30—40 c.M. broad. Pinnae \pm 2 c.M. broad, cut $\frac{1}{3}$ — $\frac{1}{2}$ -way down to the costa into acute, subfalcate lobes. Texture subcoriaceous; rachis and under surface finely pubescent; veins 10—12 on a side. Sori subterminal on the veins; indusium very fugacious.

Malaya; Polynesia.

(77) **D. eminens**, *C. Chr.*, Ind. Fil.; *Nephrodium eminens*, *Bk.* Journ. of Bot., XVIII, 213.

Rhizome creeping. Stipes stout, naked, \pm 120 c.M. long or longer, the upper half occupied by the much reduced pinnae. Fronds 2-pinnatifid, 75—90 c.M. long in the normal portion, half as broad. Normal pinnae distinctly stalked, the central ones lanceolate, 15—50 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad, cut \pm $\frac{1}{2}$ -way down to the costa into oblong, obtuse, contiguous, entire lobes \pm 4—5 m.M. broad. Lower pinnae dwarfed, deltoid, sessile, the lowest changed into glands. Texture subcoriaceous; surfaces finely pilose on the ribs; veins distinct, 10—12-jugate, several lower ones anastomosing. Sori small, medial; indusium minute, fugacious.

Sumatra, Malacca.

(78) **D. hirsuta**, *O. Ktze*, Rev. Gen. Pl., II, 815; *D. adenophora*, *C. Chr.*, Ind. Fil.; *Nephrodium hirsutum*, *J. Sm.*, Hk. Bk., Syn. Fil., 291; Hk., Spec. Fil., IV, tab. CCXL, B; *Aspidium hirsutum*, *Kze* (not *Mett.*), Christ, Ann. Btz., XV, 154.

Rhizome creeping. Stipes 30 c.M. long or longer, grey, villose. Fronds 90—120 c.M. long, \pm 45 c.M. broad. Pinnae very close and numerous, spreading or more or less erecto-patent, the largest \pm 25 c.M. long, \pm 2 c.M. broad, cut

down $\frac{1}{2}$ -way to the costa into very numerous, entire, linear-oblong, subfalcate lobes. Lower pinnae with a distinct gland beneath at the base. Texture subcoriaceous; rachis densely clothed with soft, spreading hairs; under surface glandular; veins 6—8 on a side, generally the 5 lower anastomosing. Sori medial on the veins.

Celebes, Philippines.

(79) **D. ochtodes**, *C. Chr.*, Ind. Fil.; *Nephrodium multijugum*, *Bk.*, *Hk. Bk.*, Syn. Fil., 291; *N. ochtodes*, *Hk.*, Spec. Fil., IV, 109; *Aspidium multijugum*, *Christ* (not *Wall.*), Ann. Btz., XV, 155; *A. ochtodes*, *Kze.* (oldest name).

Rhizome creeping. Stipes 50—60 c.M. long, subglabrous, strong, erect. Fronds 90—120 c.M. long, 50—45 c.M. broad. Pinnae \pm 2—2 $\frac{1}{2}$ c.M. broad, cut down \pm $\frac{2}{3}$ of the way to the costa into close, acute, subfalcate lobes 2 $\frac{1}{2}$ —3 $\frac{1}{2}$ m.M. broad. Texture subcoriaceous; rachis and both surfaces \pm naked; veins 10—12 on a side. Sori nearer the main veins than the edge.

Malacca, New Guinea, Celebes; Japan, South. China, Br. India.

* (80) **D. pilosiuscula**, *C. Chr.*, Ind. Fil.; *Nephrodium pilosiusculum*, *Zipp.*, Rac., Flor. Btz., I, 189.

Rhizome short-creeping or suberect, the extremity clothed with lanceolate, ciliate scales. Stipes approximate, 12—28 c.M. long, densely pubescent above, polished below. Fronds ovate-lanceolate, long-acuminate, 15—20 c.M. long, 4—7 c.M. broad. Pinnae close, lanceolate, erecto-patent or falcate, sessile, the base truncate, the edge slightly crenate, the apex blunt, the lower base rounded, the upper slightly auricled. Texture semicoriaceous; rachis pubescent; upper surface densely clothed with adpressed hairs, under one covered with numerous, globose glands, hairy on the veins; veins 4—5 on a side, the 2 lower ones commonly anastomosing. Sori 2—5 on a side; indusium fugacious, densely villose.

Java.

* (81) **D. glandulosa**, *O. Ktze.*, *C. Chr.*, Ind. Fil.; *Nephrodium glandulosum*, *Moore* (not *Sm.*), Rac., Flor. Btz., I, 185; *Hk. Bk.*, Syn. Fil., 291, p.p.; *Aspidium glandulosum*, *Bl.*, *Christ*, *Farnkr. d. Erde*, 245.

Rhizome short-creeping or erect, slightly clothed with lanceolate, short-ciliated hairs. Stipes \pm tufted, 10—60 c.M. long, naked or pubescent, scaly at the base. Fronds 50 c.M. or more long, 10—15 c.M. broad or rarely broader, with \pm 2—10 pairs of spreading pinnae on each side below the

terminal one. Pinnae sessile or short-stalked, more or less close or the largest ones distant, the barren ones the largest, $1\frac{1}{2}$ —5 c.M. broad, the apex acute or short-acuminate, the margin entire or slightly lobed, the base \pm truncate, not rarely \pm auricled above; fertile pinnae generally considerably contracted, but sometimes the larger uncontracted pinnae soriferous. Texture papyraceo-herbaceous; surfaces \pm short-hairy, the under one glandular with numerous, dot-like, yellowish glands; veins to 12 pairs in the larger pinnae, 4—8 in the narrower ones, nearly all anastomosing except the highest ones. Sori medial on the veins, eventually elongating, meeting, and covering the whole of the veins so as to become meniscioid; indusium fugacious, ciliate.

Malaya.

* (82) **D. cyatheoides**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium cyatheoides*, Pr. (not Klf.), Hk. Bk., Syn. Fil., 295; Hk., Spec. Fil., IV, tab. CCXLI, A; *Aspidium cyatheoides*, Klf., Enum. Fil., 254.

Rhizome erect or suberect. Stipes 30—60 c.M. long, strong, naked, glossy, greyish. Fronds 60—90 c.M. long, 50—60 c.M. broad. Pinnae \pm close, spreading or \pm oblique, 2—4 c.M. broad, the apex acuminate, the edge irregular, furnished with acute, forward pointing teeth or lobes to a depth 2—5 m.M., the lowest pinnae not shorter than the next. Texture papyraceous, herbaceous or subcoriaceous; rachis and both surfaces naked; main veins pinnate with 10—12 veins on a side. Sori contiguous, close to the main veins, the lower ones close to the costae.

Sumatra, New Guinea; Hawaii.

* **D. urophylla**, *C. Chr.*, Ind. Fil.; *Nephrodium urophyllum*, *Keys*. (not *Bedd.*), *Rac.*, *Flor. Btz.*, I, 184; *N. lineatum*, *Bedd.*, *Ferns Br.I.*, I, tab. CXXXIII; *Aspidium urophyllum*, *Christ* (not *Bedd.*), *Farnkr. d. Erde*, 248; *Polypodium urophyllum*, *Wall.*, *Hk. Bk.*, *Syn. Fil.*, 314; — var. *Pinwillei*, *Bedd.*, *Handb. Ind. Ferns, Suppl.*, 72.

Rhizome short-creeping or erect, clothed with small, ovate-lanceolate scales at the extremity. Stipes approximate, 20—60 c.M. long, naked or short-hairy, stout, erect. Fronds \pm 45—120 c.M. long, 25—45 c.M. broad, with a terminal pinna and below this 2—7 lateral ones on each side. Pinnae 2—5 c.M. broad or broader, sessile or short-stalked, the base truncate or narrowed, the edge entire, crenate, serrate or sharply toothed, the apex acuminate. Texture thinly coriaceous; rachis and both surfaces naked or nearly so, surfaces verruculose; veins conspicuous, \pm 8—14 on each side of the main

ones, nearly all anastomosing. Sori medial on the veins; indusium fugacious or undeveloped.

Var. Pinwillei: Pinnae very broad in the middle, considerably attenuated at both ends, often long-caudate at the apex.

Malaya; North. India, South. China, Trop. Australia, Polynesia; — var. *Pinwillei: Malacca.*

* (84) **D. moulmeinensis, C. Chr.,** Ind. Fil.; *Nephrodium moulmeinense, Bedd.,* Handb. Ind. Ferns, 275; *Polypodium multilineatum, Wall., Hk. Bk., Syn. Fil.,* 516; *Aspidium multilineatum, Christ, Farnkr. d. Erde,* 249; *A. moulmeinense, Christ, Ann. Btz.,* XIX, 59; *Goniopteris lineata, Bedd., Ferns Br.I.,* I, tab. III; *G. multilineata, Bedd., l. c.,* II, tab. CCXXXI.

Rhizome short-creeping. Stipes erect, 60—120 c.M. long, naked. Fronds 60—120 c.M. long, 40—60 c.M. broad. Pinnae broadly linear, usually more than 6 on each side, 20—40 c.M. long, \pm 3—4½ c.M. broad, the apex acuminate to subcaudate, the edges subparallel, entire, crenate or with sharp, falcate teeth reaching about 2—5 m.M. deep, the base broadly cuneate or subtruncate. Texture coriaceous or subcoriaceous; rachis and surfaces naked or nearly so, \pm verruculose; veins prominent, \pm 15—20 on each side of the main veins, nearly all anastomosing. Sori medial on the veins; indusium fugacious.

Celebes, Philippines; Fiji, China, Br. India.

(85) **D. latipinna, O. Ktze,** Rev. Gen. Pl., II, 815; *D. parasitica, O. Ktze, var., C. Chr.,* Ind. Fil.; *Nephrodium latipinna, Hk., Hk. Bk., Syn. Fil.,* 292; *Aspidium latipinna, Hance, Journ. of Linn. Soc., Bot.,* XIII, 141.

Rhizome \pm erect. Stipes tufted, 10—15 c.M. long, slender, naked. Fronds 20—30 c.M. long, 7—13 c.M. broad, the upper ½ or ⅓ lanceolate, pinnatifid with broad, oblong lobes. Below this several pinnae on each side, the largest 5—7 c.M. long, 1—2 c.M. broad, cut to \pm ¼ of the way down into broad, oblong, subfalcate lobes. Lower pinnae much reduced and distant. Texture papyraceo-herbaceous; rachis and both surfaces naked; main veins pinnate in the lobes with 3—5 distant veins on each side. Sori medial on the veins. — Hardly different from *D. parasitica, O. Ktze.*

Java, Philippines; Hongkong.

(86). **D. abortiva, O. Ktze,** C. Chr., Ind. Fil.; *Nephrodium abortivum, J. Sm., Hk. Bk., Syn. Fil.,* 292; *N. pennigurum, Moore, Bedd.,* Handb.

Ind. Ferns, 277, p. p.; *Aspidium abortivum*, Bl., Enum., 154; — var. *puberula* & *tomentosa*; var. *B & C*, Bl., l. c.

Rhizome \pm erect. Stipes tufted, 15—30 c.M. long, firm, erect, naked. Fronds 50—60 c.M. long or longer, 15—50 c.M. broad. Central pinnae 7—15 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, acuminate, cut to $\pm \frac{1}{3}$ of the way down into close, truncate lobes; lobes oblong, entire, or crenulate at the apex, the lower ones sub-abbreviate, $\pm \frac{1}{4}$ — $\frac{1}{3}$ c.M. broad. Lower pinnae distant, reduced suddenly into mere auricles. Texture firm-herbaceous; rachis naked, or puberous on the upper side; costae puberous above; veins \pm 6—8 on each side of the main veins. Sori medial.

Var. puberula: Pinnae long-acuminate. Rachis puberulous throughout.

Var. tomentosa: Pinnae short and narrow. Rachis tomentose and sparingly paleaceous above.

Malaya.

* (87) **D. megaphylla**, C. Chr., Ind. Fil.; *Nephrodium pennigerum*, Moore (not Hk.), Rac., Flor. Btz., I, 190; *N. abruptum*, Bedd., Ferns S.I., tab. LXXXVI; *Aspidium pennigerum*, Bl., Christ, Farnkr. d. Erde, 248; *A. megaphyllum*, Mett., (oldest name), Ann. Mus. Bot. L. B., I, 253.

Rhizome short-creeping or suberect, clothed with firm, blackish-brown, lanceolate, entire scales. Stipes approximate, stout, 15—100 c.M. long, finely or minutely villose, scaly at the base. Fronds 50—120 c.M. long, 20—60 c.M. broad, linear-lanceolate or oblong. Pinnae numerous, oblique, 1—4 c.M. broad, the apex acuminate, the edge cut $\frac{1}{6}$ — $\frac{1}{4}$ or rarely to $\frac{1}{3}$ of the way down to the midrib into oblong, falcate lobes, the base broadly cuneate; lower pinnae suddenly dwarfed, distant, occupying \pm the upper $\frac{1}{2}$ — $\frac{2}{3}$ of the stipe. Texture herbaceous to papyraceous; rachis and under surface villose, or the under surface slightly hairy on the costae and veins only; veins 6—10 on each side, the 5—4 lower anastomosing. Sori medial on the veins, 5—7 to a row; sporangia often setose.

Malaya; North. India, Ceylon, Comores, South. Africa.

* (88) **D. arbuscula**, O. Ktze, C. Chr., Ind. Fil.; *Nephrodium Arbuscula*, Desv., Rac., Flor. Btz., I, 189; Bedd., Ferns S.I., tab. LXXXVII; *Aspidium Hookeri*, Wall., Hk., Icon. Pl., X, tab. CMXXII; *A. Arbuscula*, Willd., Christ, Farnkr. d. Erde, 249.

Rhizome short-creeping or erect, clothed with narrow, lanceolate scales.

Stipes approximate or tufted, 10—25 c.M. long, naked or slightly pubescent. Fronds lanceolate, narrowing towards both extremities, 50—65 c.M. long, 10—20 c.M. broad. Pinnæ often close and numerous, linear-lanceolate, acuminate, the base truncate, slightly cordate or broadly cuneate, the edge cut down $\frac{1}{4}$ or less into blunt lobes; 2—10 lower pairs distant, gradually reduced. Texture herbaceous; rachis pubescent or naked; under surface short-hairy; veins pubescent above and beneath, 5—6 on a side, the 2—3 lower ones commonly anastomosing. Sori medial or terminal, on the veins, forming close rows; indusium often naked, persistent.

Malaya — *Polynesia*; *Br. India*, *Mascarenes*.

(89) **D. amboinensis**, *O. Ktze*, *Rev. Gen. Pl.*, II, 812; *D. parasitica*, *O. Ktze*, *var.*, *C. Chr. Ind. Fil.*; *Nephrodium amboinense*, *Pr.*, *Hk. Bk.*, *Syn. Fil.*, 292; *N. molle*, *R. Br.*, *var. amboinense*, *Bedd.*, *Handb. Ind. Ferns*, 278; *N. extensum*, *Moore*, *var. minor*, *Bedd.*, *Ferns Br. I.*, II, tab. CCI.

Rhizome short-creeping or erect. Stipes tufted, to 20 c.M. long, greyish, nearly naked. Fronds 60 c.M. or more long, 20—30 c.M. broad. Pinnæ spreading, 7—25, alternate or opposite, the longest 1—2 c.M. broad, sometimes caudate, the edge entire, slightly crenate or shallowly lobed, the lobes blunt; lower pinnæ gradually reduced into mere auricles, often sagittate or butterfly-shaped, rather distant, the normal ones rather close. Texture thinly herbaceous to papyraceous; rachis and under surface slightly pubescent; veins 4—6 rarely 7—8 on a side, the 2—3 lower pairs anastomosing. Sori medial on the 1—3 lower veins, the lowest often close to the midribs.

Malaya; *Br. India*.

(90) **D. multilineata**, *O. Ktze*, *Rev. Gen. Pl.*, II, 811; (? *D. megaphylla*, *C. Chr.*, *Ind. Fil.*); *Nephrodium multilineatum*, *Bedd.*, *Handb. Ind. Ferns*, *Suppl.*, 80.

Rhizome erect. Stipes furnished with a few lanceolate scales at the base. Fronds 90 c.M. or more long, $\pm 4\frac{1}{2}$ c.M. broad, pinnate, gradually dwindling down nearly to the base into small, rounded auricles. Pinnæ sessile or short-stalked, $\pm 2\frac{1}{2}$ c.M. broad, the apex very acuminate, the edge cut down only about $\frac{1}{6}$ — $\frac{1}{5}$ towards the costa into rounded lobes, the base truncate. Texture herbaceous to papyraceous; under surface finely villose on the costae and veins; veins ± 7 on each side, the 5—6 lower anastomosing. Sori medial; indusium persistent.

Malacca.

* (91) **D. callosa**, *C. Chr.*, Ind. Fil.; *Nephrodium callosum*, *Keys.*, *Rac.*, *Flor. Btz.*, I, 192; — var. *sumatrana*.

Rhizome erect, to 10 c.M. long, mucous at the apex. Stipes naked, glossy, to 100 c.M. long, provided with large, distant glands sometimes 1 c.M. long and subulate. Fronds lanceolate, acuminate, 100—200 c.M. long, 40—80 c.M. broad. Pinnae linear, sessile, to 2 c.M. broad, the base \pm truncate, the edge cut \pm $\frac{1}{4}$ — $\frac{1}{3}$ -way down to the costa. Lobes 3—5 m.M. long, to 5 m.M. broad, falcate, entire, crenate or toothed. Texture coriaceous; surfaces naked, not rarely minutely verruculose; rachis hairy above, provided on the under side (at the bases of the pinnae) with glands similar to those of the stipe; veins pale-coloured, 8—14 on each side of the main veins, the 5—6 lower anastomosing. Sori close, 8—15 on a side, divaricating from the apices of the lobes towards the costa.

Var. sumatrana: Fronds small, \pm 20—25 c.M. long, \pm 5—7 c.M. broad, on stipes to 15 c.M. long. Pinnae cut \pm $\frac{1}{4}$ -way down to the costa, the lower ones gradually dwarfed; veins not pale-coloured, 6—8 on each side, the 3 lower ones anastomosing. Sori confluent.

Java, Sumatra; — var. *sumatrana*: *Sumatra* (Burck).

(92) **D. laevis**, *C. Chr.*, Ind. Fil.; *Aspidium laeve*, *Kze*, *Mett.*, *Farngr.* IV, *Phegop.* & *Aspid.*, 104.

Stipes stramineous, naked. Fronds \pm 90 c.M. long, ovate, the base abruptly narrowed, pinnate. Pinnae \pm 22 c.M. long, 15—18 m.M. broad, sessile, linear, pinnatifid caudato-acuminate, the base truncate, the apex serrate; lower pinnae shorter, the lowest \pm 10 c.M. long, oblong, remote. Lobes ovate, acute, repando-denticulate, those on the upper base the largest. Texture subcoriaceous; surfaces glossy, scabrous; costae thinly pubescent; veins \pm 7 on each side, the 1—2 lowest anastomosing. Sori medial; indusium naked.

Luzon.

(95) **D. brachyodus**, *O. Ktze*, *C. Chr.*, Ind. Fil.; *Nephrodium brachyodon*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 295; *Aspidium brachyodus*, *Gris.*, *Christ*, *Farnkr. d. Erde*, 248.

Rhizome \pm erect. Stipes tufted, 50—60 c.M. long, greyish, naked. Fronds 60—90 c.M. long, 30—45 c.M. broad. Pinnae $3\frac{1}{2}$ —4 c.M. broad, cut from $\frac{1}{3}$ — $\frac{1}{2}$ the way down into blunt, oblong, slightly falcate lobes; lower pinnae not much smaller than the others. Texture coriaceous; rachis and both surfaces \pm naked; costae and costulae slightly hairy; main veins

pinnate in the lobes with 10—15 veins on a side, with a sorus on each about midway between the main veins and the edge; indusium small, fugacious.

Malaya; Galapagos Islands, Trop. America.

* (94) **D. ferox**, **O. Ktze**, C. Chr. Ind. Fil.; *Nephrodium ferox*, Moore, Rac., Flor. Btz., I, 192; Bedd., Ferns Br.I., I, tab. CXXIX; *Aspidium ferox*, Bl., Christ, Farnkr. d. Erde, 247; *Polypodium barbatum*, Hk., Sepe. Fil., V, 11; *Phegopteris barbata*, Mett., Farng., IV, Phegop. & Aspid., No. 48.

Caudex to 10—20 or sometimes to 100 c.M. high Stipes tufted, 50—150 c.M. long, clothed like the rachis with long, subulate, reddish or blackish, spreading bristles leaving the stipes rough when they fall. Fronds 90—300 c.M. long, to 60—100 c.M. broad, lanceolate-oblong. Pinnae rather close, linear-lanceolate, 17—30 m.M. broad, the apex acuminate, the edge cut $\pm \frac{1}{3}$ -way down into numerous, entire, deltoid-oblong, straight or falcate lobes, the base truncate, on stalks to ± 1 m.M. long or longer. Texture subcoriaceous; surfaces glossy; midribs villose beneath; main veins pinnate in the lobes; veins ± 10 —15 on each side, the 5—7 lower anastomosing. Sori often numerous, close to the main veins, the lower close to the midribs; indusium fugacious.

Malaya.

(95) **D. Loheriana**, **C. Chr.**, Ind. Fil.; *Nephrodium Loherianum*, Copel., Polypod. Philipp., 50; *Aspidium Loherianum*, Christ, Bull. Herb. Boiss., VI, 191.

Rhizome short-creeping. Stipes approximate, the base clothed with few, ovate, brown scales, the rest short-haired like the rachis, costae and veins, dark-stramineous, firm, ± 50 c.M. long, the upper part bearing distant, reduced, auriculiform pinnae. Fronds short-haired, deltoid-ovate, ± 50 c.M. long, 18—20 c.M. broad, acuminate. Lowest not-reduced pinnae the longest, ± 2 c.M. broad, lanceolate, patent, acuminate, sessile, cut down to a distance of ± 3 m.M. from the costa into falcate, oblong, pectinate, obtuse, entire lobes ± 3 m.M. broad. Texture subcoriaceous; veins ± 12 -jugate. Sori submarginal, close, small; indusium villose.

Philippines.

(96) **D. latebrosa**, **C. Chr.**, Ind. Fil.; *Aspidium latebrosus*, Kze, Mett., Farng., IV, Phegop. & Aspid., 104.

Stipes nodulose below. Fronds ± 90 c.M. long, linear-lanceolate, pinnate.

Pinnae \pm 9 c.M. long, \pm 2 c.M. broad, sessile, elongate-oblong. the base broadly truncate, narrowed gradually towards the acuminate apex, the margin deeply pinnatifid: lower pinnae gradually reduced, obtuse, pinnatifid, the lowest abortive, reaching nearly the base of the stipe. Lobes ovate-oblong, subfalcate, obtuse, the lowest ones the largest. Texture rigid; upper surface clothed with adpressed bristles, under surface covered with minute, sessile, scattered glands: costae with a large tubercular gland beneath at the base; veins \pm 8 on each side, the 1—2 lowest anastomosing. Sori \pm 8 on a side, medial or nearer the margin than the midrib; indusium subcoriaceous, naked.

Java, Sumatra.

* (97) **D. piloso-squamata**, *v. A. v. R.*, Hort. Bog.

Rhizome short, clothed with linear-lanceolate, brown, acuminate, hair-covered scales. Stipes approximate, \pm 55 c.M. long, stout, erect, firm, green, inconspicuously short-hairy, scaly towards the base. Fronds ovate-lanceolate, \pm 50 c.M. long, \pm 50 c.M. broad. Pinnae sessile, linear-lanceolate, 2—2 $\frac{1}{2}$ c.M. broad, acuminate, the edge cut \pm $\frac{1}{3}$ — $\frac{1}{2}$ -way down to the costa into oblong, rounded or subtruncate, subentire lobes \pm 2 $\frac{1}{2}$ m.M. broad, the base truncate or subtruncate, the lowest not reduced pinnae the longest, deflexed; below this \pm 4—5 distant, much reduced, deflexed auricles, occupying the upper half of the stipe. Texture \pm coriaceous, rachis inconspicuously hairy; surfaces minutely hairy on the costae and costulae, rather naked; veins 8—10 on each side of the main veins, the 5—5 lower anastomosing. Sori numerous, medial on the veins, afterwards becoming subconfluent and densely covering the under surface of the pinnae; indusium reniform, with the surface covered with very short hairs and yellow, globose glands. — The description is taken from a specimen cultivated in the Buitenzorg gardens.

Skroe (New Guinea).

* (98) **D. sagittifolia**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium sagittaeifolium*, *Moore*, Rac., Flor. Btz., I, 191; *Aspidium sagittaeifolium*, *Bl.*, Enum., 155

Rhizome erect or suberect, the extremity sparingly clothed with delioid scales. Stipes tufted, 50—60 c.M. long, stout, densely villose, scaly at the base. Fronds 90—120 c.M. long, 50 c.M. or more broad. Pinnae numerous, spreading, straight or falcate, \pm 2 c.M. broad, the apex acuminate, the base truncate and often auricled on the upper side, the edge cut $\frac{1}{3}$ — $\frac{1}{2}$ -way down into oblong-falcate lobes; lower pinnae suddenly dwarfed and deflexed, close,

occupying nearly the whole stipe. Texture subcoriaceous; rachis and under surface densely villose; main veins pinnate in the lobes with 7—10 veins on a side, the 2—3 lower anastomosing. Sori midway on the veins, 6—7 on a side; capsules setose or naked; indusium densely villose.

Malacca, Java, Borneo.

(99) **D. caudiculata**, v. **A. v. B.**; *D. basilaris*, C. Chr., Ind. Fil.; *Nephrodium philippinense*, Bk., Copel., Polypod. Philipp., 51; *N. basilare*, Pr., Epim., 258, *N. caudiculatum*, J. Sm., (oldest name).

Rhizome erect. Fronds oblong-lanceolate, 2-pinnatifid, 60—90 c.M. long, 50—45 c.M. broad. Pinnae lanceolate, acuminate, 20—25 c.M. long, 1—1½ c.M. broad, cut less than ½-way down to the costa into oblong, erecto-patent lobes 5—3½ m.M. broad. Lower pinnae not reduced. Texture moderately firm; rachis and surfaces naked; veins simple, 8—9-jugate. Sori medial; indusium firm, naked, persistent.

Philippines.

(100) **D. simillima**, C. Chr., Ind. Fil.; *Nephrodium simulans*, Bk., Journ. of Bot., XXVI, 525 (not Hk. Bk., Syn. Fil., 500, nor Journ. of Bot. XXVIII, 106).

Rhizome erect, Stipes tufted, naked, nearly 50 c.M. long. Fronds oblong-lanceolate, 2-pinnatifid, 50—45 c.M. long, 5—8 c.M. broad. Pinnae sessile, linear-oblong, obtuse, shallowly pinnatifid, ± 1—1½ c.M. broad, the lower ones sharply deflexed. Texture moderately firm; both surfaces finely pubescent; main veins pinnate; veins 4—5-jugate. Sori medial on all the veins; indusium obscure, minute, fugacious.

Borneo.

* (101) **D. pseudoreptans**, C. Chr., Ind. Fil.; *Nephrodium debile*, Bk., Journ. of Bot., XVIII, 212.

Rhizome short-creeping. Stipes approximate, slender, grey, naked, 20—50 c.M. long. Fronds lanceolate, 20—50 c.M. long, 5—7 c.M. broad. Pinnae sessile, oblong-lanceolate, 2½—4 c.M. long, 1—1½ c.M. broad, cut ¼—⅓ of the space down to the costa into oblong, obtuse lobes. Lower pinnae slightly reduced or dwarfed. Texture moderately firm; upper surface strigillose, under one pubescent on the ribs; veins 4—6-jugate, the 2 lower joining. Sori small, medial; indusium persistent.

Sumatra.

* (102) **D. parasitica**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium molle*, R. Br., (not *Desv.*), Rac., Flor. Btz., I, 188; Bedd., Ferns S.I., tab. LXXXIV; *N. parasiticum*, *Desv.* (not *Bk.*), Copel., Polypod. Philipp., 52; *Aspidium parasiticum*, *Sw.* (not *Mett.*), Christ, Farnkr. d. Erde, 243; *Polypodium parasiticum*, *L.*, (oldest name).

Rhizome erect or suberect, clothed with dark-brown, entire, lanceolate scales. Stipes tufted or subremote, 15—50 c.M. or more long, slender, densely and deciduously hairy; hairs short. Fronds ovate-lanceolate, to 30—60 c.M. long, $12\frac{1}{2}$ —50 c.M. broad. Pinnae spreading, approximate, $\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, cut $\pm \frac{1}{2}$ -way down to the midrib into scarcely falcate, blunt or acute lobes, the base not narrowed, the apex acuminate. Lower pinnae distant, not rarely shorter than the next in order, the lowest sometimes slightly dwarfed. Texture herbaceous, rachis and surfaces finely hairy; main veins pinnate in the lobes with 6—9 veins on each side, only the lowest anastomosing. Sori medial on the veins; capsules naked; indusium villose. — A form found in Borneo (*forma caudata*) has the fronds caudate and subglabrous, and a form gathered in Sumatra (*forma repens*) has the rhizome wide-creeping.

Tropical and subtropical regions all round the world.

* (103) **D. indica**, *v. A. v. R.*; *N. pennigerum*, *Bedd.*, Handb. Ind. Ferns, Suppl., 73; — var. *malayensis*, *Bedd.*, l. c., 74.

Rhizome erect or short-creeping. Stipes hairy or nearly naked, to 120 c.M. long. Pinnae numerous, spreading, 20—25 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, the fertile sometimes much contracted, the apex acuminate, the edge cut down only $\pm \frac{1}{4}$ towards the costa into oblong, falcate lobes; lowest pinnae not reduced. Texture herbaceous; rachis hairy; upper surface strigose on the costae, otherwise naked or more or less strigose particularly near the edges, under surface hairy on the veins; veins 8—12 pair, 5—8 pair anastomosing except in contracted, fertile pinnae. Sori medial on the veins; indusium generally persistent.

Var. malayensis: Pinnae rather deeply cut; surfaces rather densely clothed with short hairs; only 2—3 lower pairs of veins anastomosing. *Burmah, Br. India, Ceylon*; — var. *malayensis*: *Malacca, Tenasserim*.

(104) **D. sakayensis**, *C. Chr.*, Ind. Fil.; *Nephrodium sakayense*, *Zeill.*, Bull. Soc. Bot. Fr., XXXII, 75.

Rhizome erect. Fronds densely tufted, bipinnatifid, 100—120 c.M. long,

25—30 c.M. broad, ovate-lanceolate, with a long, terminal, pinnatifid pinna. Pinnae erecto-patent, alternate, $\pm 2\frac{1}{2}$ —3 c.M. apart, linear-lanceolate, 15—20 c.M. long, 15—20 m.M. broad, the apex narrowed to an acute point, the margin cut $\frac{1}{3}$ — $\frac{2}{5}$ -way down into subacute (rather bluntish), slightly falcate lobes, obscurely crenate at the apex. Texture papyraceous; rachis dark-reddish-brown, slightly canaliculate and hairy and scaly on the upper side, the scales large, pale-brown; upper surface covered with pale or whitish hairs on the costae and rarely on the costulae and veins; under surface finely glandular; veins ± 8 —10 on each side, only the 5 lowest ones anastomosing. Sori costular, confined to the 4—5 lowest veins; indusium very fugacious.

Malacca.

(105) **D. Benoitiana**, *v. A. v. R.*; *Nephrodium Benoitianum*, *Pr.*, Tent. Pterid., 81; *C. Chr.*, Ind. Fil.; *Polystichum Benoitianum*, *Gaud.*, Freyc., Voy. Ur. Phys. Bot., 351; *Aspidium Benoitianum*, *Gaud.*, Freyc., l. c., sub tab. XI.

Rhizome \pm erect. Stipes tufted, to ± 20 c.M. long, short-hairy, scaly at the base. Fronds ovate-lanceolate, acuminate, ± 55 c.M. long, ± 12 c.M. broad, with ± 12 pairs of lateral pinnae below the deeply pinnatifid apex. Highest pinnae erecto-patent, central ones largest, lanceolate, slightly acuminate, the point bluntish, the edge cut $\frac{1}{2}$ — $\frac{2}{3}$ -way down into oblong, rounded, \pm erecto-patent lobes, the base truncate or broadly cuneate, not much narrower than the centre, which is $\pm 1\frac{1}{2}$ c.M. broad, lowest pinnae shorter, deflexed. Rachis and veins densely short-hairy beneath; veins ± 7 on each side, only the lowest ones anastomosing. Sori 5—6 on a side, submarginal; indusium short-hairy.

Moluccas.

(106) **D. didymosora**, *C. Chr.*, Ind. Fil.; *Nephrodium didymosorum*, *Par.*, *Bedd.*, Ferns Br.I., II, tab. CC; *N. tectum*, *Bedd.*, Handb. Ind. Ferns, Suppl., 79.

Rhizome strong, erect. Stipes densely tufted, a span or more long. Fronds deltoid or somewhat lanceolate, 50—40 c.M. long, 15—18 c.M. broad. Pinnae cut down $\frac{2}{3}$ or more towards the costa into entire, slightly falcate lobes, lowest pinnae not or slightly reduced, deflexed. Texture firm-herbaceous; rachis, midribs and lamina on both sides rather densely covered with long, white, needle-like hairs; veins 7—8 pair, only the lowest anastomosing. Sori terminal, on nearly all the veins or not rarely confined to the lowest

pair; indusium often very hairy. — Some specimens have the lowest pinnae suddenly reduced and much deflexed, but not distant from the others.

Malacca.

(107) **D. microchlamys**, *v. A. v. R.*; *Nephrodium microchlamys*, *Bk.*, Journ. of Linn. Soc., Bot., XV, 107; *Mesochlaena polycarpa*, *Bedd.*, var., *C. Chr.*, Ind. Fil.

Rhizome \pm erect. Stipes \pm 15 c.M. long, pale-brown, scaleless, finely pilose. Fronds oblong-lanceolate, 60—90 c.M. long, 30—40 c.M. broad, 2-pinnatifid. Pinnae in \pm 40 pairs below the small, entire, terminal one, sessile, linear-ligulate, the largest, 20—22 $\frac{1}{2}$ m.M. broad, the lower not reduced, cut down to a broad wing into close, blunt, entire, rather ascending lobes \pm 3—3 $\frac{1}{2}$ m.M. broad, the lowest anterior lobe slightly elongated. Texture membranaceous; surfaces finely villose and densely covered with minute, pale-brown glands; veins 12—16-jugate, the lowest anastomosing, the next meeting at the sinus. Sori round, minute, in regular rows nearer the main veins than the edge.

Kei Islands.

(108) **D. glaucostipes**, *C. Chr.*, Ind. Fil.; *Nephrodium glaucostipes*, *Bedd.*, Handb. Ind. Ferns, Suppl., 80.

Rhizome erect or suberect. Stipes scaly at the base, glaucous as is the main rachis. Fronds 90—120 c.M. long, pinnate. Central pinnae 12—15 c.M. long, 2—2 $\frac{1}{2}$ c.M. broad, cut down $\frac{1}{2}$ -way into oblong, rounded lobes, lower pinnae very gradually reduced in size, becoming at first small and hastate or semihastate, and at length mere auricles down to almost the base of the stipe. Texture rather papyraceous; surfaces naked but the costae minutely villose above; veins in \pm 10 pairs, the 2 lower ones anastomosing, all soriferous. Sori medial.

Malacca.

(109) **D. perakensis**, *C. Chr.*, Ind. Fil.; *Nephrodium perakense*, *Bk.*, Ann. of Bot., V, 319; *Aspidium perakense*, *Bedd.*, Journ. of Bot., XXVI, 4.

Rhizome small, erect. Stipes slender, villous. Fronds pinnate, 30—35 c.M. long, 5—15 c.M. broad, oblong-lanceolate, the lower pinnae gradually reduced. Central pinnae largest, 1—1 $\frac{1}{2}$ c.M. broad, cut \pm $\frac{1}{2}$ -way down to the costa into close, rather acute lobes. Texture softly herbaceous; rachis and surfaces copiously furnished with long, white, soft hairs; veins 4—5 on

each side of the main veins, the lowest pair anastomosing. Sori subterminal on the veins.

Malacca.

(110) **D. sumatrana**, *v. A. v. R.*; *N. molle*, *R. Br.*, var. *major*, *Bedd.*, *Handb. Ind. Ferns*, *Suppl.*, 76.

Rhizome erect or somewhat creeping. Fronds large, often 120—150 c.M. long, 20—50 c.M. broad. Pinnae very numerous, cut down $\frac{1}{3}$ — $\frac{1}{2}$ -way towards the costa, the lower ones gradually reduced to butterfly-like auricles down the base of the stipe. Texture herbaceous; rachis and surfaces not very hairy, often quite glabrous except the costae above; veins 7—8 pair, the 2—3 lower pair anastomosing. Sori medial, generally on all the veins except the 1—2 highest ones.

Sumatra, Malacca; Ceylon, Br. India.

* (111) **D. stipellata**, *O. Ktze*, *C. Chr.*, *Ind. Fil.*; *Nephrodium stipellatum*, *Moore* (not *Hk.*), *Rac.*, *Flor. Btz.*, I, 188; *Aspidium stipellatum*, *Bl.*, *Enum.*, 152.

Rhizome short-creeping or erect. Stipes tufted, 30—100 c.M. long, hairy above, naked below. Fronds linear-lanceolate, 60—100 c.M. long, 15—25 c.M. broad. Pinnae sessile, linear-lanceolate, the apex acuminate, the base truncate or broadly cuneate, 1—2 c.M. broad, provided with a short, thick gland beneath, the edge cut about $\frac{1}{3}$ — $\frac{1}{2}$ -way down into oblong, subfalcate lobes; reduced pinnae numerous, often occupying nearly the whole stipe. Texture herbaceous to semicoriaceous; rachis, midribs and apices of the veins villose beneath; main veins pinnate in the lobes; veins 6—9 on a side, only the lowest ones anastomosing. Sori in rows, medial or nearer the main veins than the edge.

Sumatra, Java, Celebes.

* (112) **D. truncata**, *O. Ktze*, *C. Chr.*, *Ind. Fil.*; *Nephrodium truncatum*, *Pr.*, *Rac.*, *Flor. Btz.*, I, 190; *N. eusorum*, *Bedd.*, *Ferns Br.I.*, I, tab. CXXX; *N. abruptum*, *J. Sm.*, *Hk.*, *Spec. Fil.*, IV, tab. CCXLI, B; *Aspidium truncatum*, *Gaud.* (not *Pr.*), *Christ*, *Farnkr. d. Erde*, 248.

Rhizome erect or suberect. Stipes tufted, stout, erect, 60—80 c.M. long, greyish, naked, or slightly villose above. Fronds 60—120 c.M. long, 30—45 c.M. broad or broader, lanceolate-oblong. Pinnae sessile, 2—5 c.M. broad, linear-lanceolate, the apex acuminate, the base more or less broadly cuneate

to subtruncate, the edge cut $\frac{1}{3}$ — $\frac{1}{2}$ -way down into blunt or subtruncate, spreading or suberecto-patent lobes \pm 5 m.M. broad, subentire, or crenate or toothed at the apex, lower pinnae \pm narrowed at the base; reduced pinnae few, distant. Texture papyraceo-herbaceous; under surface naked or nearly so, verruculose; rachis slightly villose; main veins pinnate in the lobes; veins 6—11 on each side, the 2—3 lower ones anastomosing. Sori medial or lower on the veins, to \pm 6 on a side.

Malaya; Polynesia, Trop. Australia, Br. India, Madagascar, Mascarenes.

* (115) **D. hispidula**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium hispidulum*, *Bk.*, *Hk. Bk.*, Syn. Fil., 293; *N. angustifolium*, *Pr.*, *Hk.*, Spec. Fil., IV, 69; *Aspidium hispidulum*, *Dec.*, Nouv. Ann. Mus., III, 346.

Rhizome \pm erect. Stipes tufted, 15—45 c.M. long, villose at least on the inner side. Fronds 20—45 c.M. long, 8—25 c.M. broad. Pinnae distant, spreading, 5—15 c.M. long, 5—15 m.M. broad, the apex acuminate, the edge cut \pm $\frac{1}{2}$ -way down into oblong, more or less bluntish, slightly falcate lobes, the base obtuse, lower pinnae reduced to mere auricles. Rachis villose; texture herbaceous to semicoriaceous; under surface finely villose; main veins pinnate in the lobes; veins 5—4 on a side, the lowest pair anastomosing. Sori nearly terminal or medial on the veins. — A plant gathered in Borneo by Dr. Nieuwenhuis and determined by Dr. Christ (*Ann. Btz.*, XX, 107) has the pinnae erecto-patent, with the base cuneate, the surfaces decidedly hispid especially on the costae and edges, the lowest veins joining at or a little below the tops of the sinuses, the lobes strongly erect as in *D. calcarata*, *O. Ktze*, but less deep and the rhizome short-creeping.

Malaya; Samoa.

* (114) **D. heterocarpa**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium heterocarpon*, *Moore*, *Rac.*, *Flor. Btz.*, I, 187; *Aspidium heterocarpon*, *Bl.*, *Christ*, *Ann. Btz.*, XV, 133.

Rhizome short-creeping or suberect, clothed with ovate-lanceolate, brown scales. Stipes \pm tufted, 30—60 c.M. long, densely (deciduously?) villose, the base glossy and scaly. Fronds ovate-lanceolate, 30—90 c.M. long, 20—40 c.M. broad. Pinnae linear-lanceolate, sessile, acuminate, more or less narrowed towards the base, 1—2 c.M. broad, the edge cut $\frac{1}{2}$ — $\frac{3}{4}$ down into linear-oblong, subfalcate lobes to 4 m.M. broad, lower not-reduced pinnae deflexed; below this several distant, abruptly dwarfed pinnae occupying the upper half of the stipe, sometimes changed into glands. Texture herbaceous, firm, rachis

and under surface hairy especially on the veins; under surface densely covered with globose, yellow glands; main veins pinnate in the lobes; veins 8—10 on a side, only the lowest anastomosing. Sori few or to 6—9 on a side, the lowest close to the midribs; indusium hairy or naked; capsules naked.

Malaya.

* (115) **D. crinipes**, *O. Ktze*, C. Chr., Ind. Fil.; *Nephrodium crinipes*, *Hk.*, Hk. Bk., Syn. Fil., 294; *Bedd.*, Ferns Br.I., II, tab. CCLXIII.

Rhizome \pm erect. Stipes 30—45 c.M. long, stout, erect, densely clothed with lanceolate-linear, brown scales. Fronds \pm 60—90 c.M. long, \pm 20—35 c.M. broad. Pinnae horizontal, \pm 10—17½ c.M. long, 1—2 c.M. broad, cut down ½—¾-way to the costa into oblong, scarcely acute lobes, the upper basal one often somewhat enlarged; lower pinnae dwarfed into auricles. Texture herbaceous; rachis scaly like the stipe, the scales \pm deciduous; costae, costulae and veins sparingly hairy above; veins 6—8 on each side of the main veins, minutely hairy beneath, the lowest pair anastomosing. Sori medial on the veins, but a little nearer the apex than the base; indusium persistent.

Celebes, Malacca; North India.

Species of an indefinite systematical place ().*

(116) **D. penangiana**, *C. Chr.*, Ind. Fil.; *Nephrodium costatum*, *Bedd.*, Handb. Ind. Ferns, 275; Ferns Br.I., II, tab. CCXX; *Polypodium lineatum*, *Colebr.*, Hk. Bk., Syn., Fil., 316; *P. penangianum*, *Hk.*, Spec. Fil., V, 15., *Goniopteris penangiana*, *Bedd.*, l.c., II, tab. CCXXXII.

Rhizome indefinite. Stipes \pm 45 c.M. long or longer, rather stout, scaleless, glossy and as well as the rachis strongly tinged with red, or rarely stramineous. Fronds broad-oblong or -lanceolate, 30—90 c.M. long, pinnate. Pinnae numerous, rather distant, spreading, sessile, 12—20 c.M. long, 1—2½ c.M. broad, the lower ones wether or not subpetiolate, lanceolate or elongate-oblong, finely acuminate, the edge coarsely and sharply submucronate-serrate, the base obliquely cuneate. Texture coriaceo-submembranaceous; surfaces naked, or slightly hairy on the veins; costae prominent beneath, generally reddish; veins \pm 6—8-jugate, the 3—6 lower anastomosing. Sori in 2

(*) Rhizome unknown or indefinite.

rows between the main veins, generally near the base of the veins; indusium very fugacious or undeveloped, only to be detected on very young fronds.

Penang; North India, China.

(117) **D. oosora**, **C. Chr.**, Ind. Fil.; *Nephrodium oosorum*, **Bk.**, Kew Bull., 1896, 41.

Rhizome unknown. Stipes \pm 30 c.M. long, grey, slightly pubescent. Fronds oblong-lanceolate, 45—60 c.M. long, 15—20 c.M. broad, 2-pinnatifid. Pinnae multijugate, $1\frac{1}{2}$ —2 c.M. broad, pinnatifid $\frac{1}{2}$ -way down to the costa, the lower ones not reduced. Lobes ovate-lanceolate, crenulate, slightly falcate. Texture subcoriaceous; rachis pubescent; both surfaces naked; veins simple, erecto-patent, conspicuous, 10—12-jugate. Sori oblong, medial; indusium membranaceous, pubescent, persistent.

Borneo.

* (118) **D. Riedleana**, **v. A. v. R.**; *Nephrodium Riedlianum*, **Moore**, Ind., 102; *Polystichum Riedleanum*, **Gaud.**, Freyc., Voy. Ur. Phys. Bot., 327; **C. Chr.**, Ind. Fil.; *Aspidium Riedleanum*, **Gaud.**, Freyc., l. c., 326.

Rhizome unknown. Stipe and rachis dark-red, nearly black, (brown when dry?), pubescent. Fronds pinnate. Pinnae lanceolate, pinnatifid, subacuminate, the base truncato-cuneate, lower pinnae gradually reduced, distant. Lobes ovate, obtuse, subfalcate, the lowest subauriculiform. Surfaces hairy. — Different from *D. parasitica*, **O. Ktze** by the rather gradually reduced lower pinnae.

Timor.

(119) **D. producta**, **C. Chr.**, Ind. Fil.; *Nephrodium productum*, **Klf.**, **Mett.**, Farng., IV, Phegop. & Aspid., 109.

Rhizome unknown. Stipes hirsute. Fronds 30—60 c.M. long, oblong, acuminate, the base suddenly narrowed. Pinnae sessile, 7—15 c.M. long, 1—2 c.M. broad, linear-lanceolate, acuminate with a long, entire point, the edge pinnatifid, the base truncate, lowest pinnae suddenly reduced, distant, \pm $1\frac{1}{2}$ c.M. long, semiorbicular, obtuse, 3-fid. Texture rigid; rachis canaliculate and pubescent on the upper side; surfaces covered with inconspicuous hairs on the costae and veins; veins 6—7 on each side, the 2—3 lowest anastomosing, all soriferous. Sori medial on the veins; indusium persistent.

Luzon.

§ 3. CYCLOSORUS.

*. Fronds 10—30 c.M. long. Rachis hairy. (120) *D. Blumei*.

.. Fronds 60—90 c.M. long. Rachis naked. (121) *D. nephrodioides*.

(120) ***D. Blumei***, v. ***A. v. R.***; *D. canescens*, *C. Chr.*, Ind. Fil., p. p.; *Nephrodium Blumei*, *J. Sm.*, Journ. of Bot., III, 411; *N. subdimorphum*, *Christ*, Fil. Saras., I, 16; *N. canescens*, *Christ*, Copel., Polypod. Philipp., 27, p. p.; *Aspidium canescens*, *Christ*, forma *nephrodiiformis*, *Christ*, Ann. Btz., XV, 150—151.

Rhizome short-creeping or erect, nearly naked. Stipes 5—20 c.M. long, scaly at the base, \pm hairy above. Fronds 10—30 c.M. long, with a long, acuminate, terminal pinna 5—10 c.M. long, cut to \pm $\frac{1}{2}$ -way down into blunt lobes. Lateral pinnae few or to 14 on each side, alternate, 2—6 c.M. long, 1—3 c.M. broad, the apex blunt, the edge slightly lobed, the base \pm adnate to the rachis; lower pinnae deflexed, sometimes much reduced. Texture papyraceous to herbaceous; surfaces naked, or hairy on the veins; rachis hairy; only the lowest veins anastomosing. Fertile fronds often on the longest stipes, with the pinnae sometimes shorter and narrower than in the barren ones. Sori near the main veins or \pm medial on the veins.

Celebes, Philippines; Fiji.

(121) ***D. nephrodioides***, ***O. Ktze***, Rev. Gen. Pl., II, 811; *D. orbicularis*, *C. Chr.*, Ind. Fil.; *Aspidium Hookeri*, *Bk.*, Hk. Bk., Syn. Fil., 257; *A. nephrodioides*, *Hk.*, Spec. Fil., IV, tab. CCXXXV.

Stipes 30 c.M. or more long, naked, erect. Fronds 60—90 c.M. long, \pm 30 c.M. broad. Pinnae 15—20 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, cut down to a broadly winged rachis into nearly close, spreading, entire, linear-oblong lobes \pm $3\frac{1}{2}$ m.M. broad. Texture herbaceous, rachis naked, substramineous; veins slightly hairy, about 10 on a side, the lowest pair uniting. Sori nearer the edge than the midribs; indusium membranaceous, reddish in the centre, scariose in the outer half.

Malaya.

23. MESOCHLAENA, *R. Brown.*

Sori oblong, medial on the veins. Indusium oblong, fastened by its longitudinal axis, opening all round the edge.

Rhizome short, \pm erect. Stipes not articulate to the rhizome. Fronds large, 2-pinnatifid; veins anastomosing as in *Nephrodium*. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 84; Christ, Farnkr. d. Erde, fig. 793.

Malaya, Polynesia.

*. Pinnae $1\frac{1}{2}$ —2 c.M. broad, cut to $\frac{1}{2}$ -way down to the costa. (1) *M. polycarpa*.
 *. *. Pinnae 3—5 c.M. broad, cut to $\frac{1}{4}$ -way down to the costa. (2) *M. larutensis*.

* (1) ***M. polycarpa***, *Bedd.*, Handb. Ind. Ferns, 199; *M. javanica*, *R. Br.*, *Bedd.*, Ferns Br.I., II, tab. CCCXLIV; *Didymochlaena polycarpa*, *Bk.*, *Rac.*, *Flor. Btz.*, I, 197; *Nephrodium javanicum*, *Hk.*, *Fil. Exot.*, tab. LXII; *Aspidium polycarpum*, *Bl.*, *Christ*, *Farnkr. d. Erde*, 249; *Sphaerostephanos asplenoides*, *J. Sm.*, *Bau. & Hk.*, *Gen. Fil.*, tab. XXIV.

Rhizome erect, densely clothed with linear-lanceolate, ciliate scales. Stipes tufted. Fronds 60—200 c.M. long, pinnate, acuminate. Pinnae spreading, close, linear-lanceolate, gradually narrowing from the \pm truncate base towards the acuminate apex, the largest 15—25 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, the edge cut down to $\frac{1}{2}$ -way or more to the costa into close, bluntish, oblong lobes, lower pinnae reduced into several or numerous mere auricles occupying the largest part of the stipe. Texture herbaceous; rachis and surfaces hairy; main veins pinnate in the lobes; veins numerous, close, the 2—3 lower anastomosing. Sori numerous, medial on the veins, ultimately confluent; indusium villose and glandular.

Malaya; Polynesia.

* (2) ***M. larutensis***, *v. A. v. R.*; *Nephrodium larutense*, *Bedd.*, *Handb. Ind. Ferns*, *Suppl.*, 75; *Dryopteris larutensis*, *C. Chr.*, *Ind. Fil.*; — var. borneensis.

Stipes 20—50 c.M. long, furnished with very regular auricles (reduced pinnae) below the pinnae, densely villous with short hairs as in the rachis. Fronds 60—90 c.M. long. Pinnae 25—30 c.M. long. 3—5 c.M. broad, finely acuminate at the apex, cut down $\pm \frac{1}{4}$ towards the costa into short, sub-falcate lobes, the base truncate or broadly cuneate. Texture herbaceous or subcoriaceous; upper surface hairy on the costae, veins and parenchyma, under surface hairy on the costae and veins, the lamina furnished with minute resinous dots; veins \pm 10 pairs, ascending, the 5—6 lower pairs anastomosing. Sori medial; indusium hairy.

Var. borneensis: Upper surface naked except on the costae; under surface copiously and minutely short-hairy; indusium covered with yellow, minute, globose glands.

Malacca; — var. *borneensis*: *Borneo* (Hallier No. 2704).

24. LUERSSENIA, *Kuhn*.

Sori oblong, terminal on included, excurrent, free veinlets, alternate, in several rows on each side of the costa. Indusium oblong, attached by its longitudinal axis, opening all round the edge.

Rhizome erect. Stipes not articulate to the rhizome. Fronds simple, slightly dimorphous, the fertile ones a little narrower and longer stalked than the barren ones; veins anastomosing, forming several rows or areolae; areolae with free veinlets, which are more copious in the barren than in the fertile fronds. Sori solitary in the areolae of the fertile fronds. Diels, in Engl. & Prantl, *Nat. Pfl.Fam.*, I⁴, fig. 95, A—B.

Sumatra.

L. Kehdingiana, *Kuhn*, *Bot. Centralbl.*, XI, 77; Diels, in Engl. & Prantl, *Nat. Pfl.Fam.*, I⁴, fig. 95, A & B; *Aspidium Kehdingianum*, *Luerss.*, *Christ, Farnkr. d. Erde*, 255.

Barren fronds \pm 50 c.M. long, \pm 5 c.M. broad, lanceolate, the apex acute, acuminate or bluntish, the base gradually narrowed, the stipe 4—5 c.M. long. Texture papyraceous or chartaceous; main veins flexuose, distinct to the edge; areolae with free veinlets. Fertile fronds linear-lanceolate, \pm 20—25 c.M. long, \pm 2 c.M. broad, narrowed gradually towards both ends, the base decurrent, the stipe \pm 20 c.M. long; no distinct main veins; areolae numerous, hexagonal. Sori terminal on the excurrent, free veinlets; indusium elliptical or oblong:

Sumatra.

25. ASPIDIUM, *Swartz*.

Sori roundish, rarely oblong, compital (*) or on free veinlets. Indusium round or reniform, attached by a central point or at the apex of a \pm deep

(*) On connected veinlets.

sinus, very rarely oblong and attached by a short, longitudinal axis, opening all round the edge or nearly so.

Rhizome creeping or erect. Stipes not articulate to the rhizome. Fronds uniform or nearly so, often inclined to a triangular shape, simple or divided, if divided the lowest pinnae often inclined to a stronger development of the basiscopic side; texture often herbaceous; veins anastomosing \pm copiously and irregularly; areolae with or without included, free veinlets.

Pantropical.

Arrangement of the sections.

§ 1. **TECTARIA** (*Eu-aspidium*). Indusium round, peltate. Hk. Bk., Syn. Fil., tab. V, fig. 43, d; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 96, C—D.

§ 2. **SAGENIA**. Indusium reniform and attached at the top of the sinus, very rarely oblong and attached by a short, longitudinal axis. Diels, l. c., fig. 97, A—C.

§ 1. **TECTARIA** (1).

- a.** *Fronds simple* (2). (1) *A. singaporianum*.
- a.a.** *Fronds palmately 3—5-fid.*
- * *Segments lanceolate.* (2) *A. tricuspe.*
- *.* *Segments triangular.* (5) *A. platanifolium.*
- a.a.a.** *Fronds ternate, or pinnate to 3-pinnatifid.*
- b.** *Sori arranged along the margin of the ultimate segments.* (4) *A. calcareum.*
- b.b.** *Sori scattered irregularly* (3).
- S.** *Stipe and rachis glossy, ebeneous.* (5) *A. angulatum.*
- S.S.** *Stipe and rachis minutely pubescent.* (6a) *A. Kunstleri.*
- b.b.b.** *Sori in rows between the main veins or costulae* (3).
- c.** *Fronds under 55 c.M. long.*
- d.** *Lowest pinnae simple or forked.*
- e.** *Sori dorsal on the veinlets.*

(1) See also under **A. coadunatum**, Wall.

(2) See also **A. angulatum**, J. Sm.

(3) See also **A. multicaudatum** & **polymorphum**, Wall., & **A. ologophyllum**, Rst.

f. *Lateral pinnae in 2 pairs.* (6b) *A. oligophyllum.*

f.f. *Lateral pinnae in 4 or more pairs.*

(7) *A. Haenkei.*

e.e. *Sori terminal on free veinlets.*

(8) *A. ternatense.*

d.d. *Lowest pinnae 2-pinnatifid.*

(9) *A. Beccarianum.*

c.c. *Fully developed fronds more than 55 c.M. long.*

(10) *A. repandum.*

* (1) **A. singaporianum**, *Wall.*, *Christ, Farnkr. d. Erde*, 227; *Bedd., Ferns Br.I., II, tab. CLXVIII*; *Nephrodium singaporianum*, *Bk., Hk. Bk., Syn. Fil., 296*; *Hk. & Grev., Ic. Fil., tab. XXVI.*

Rhizome short-creeping or erect. Stipes subtufted, those of the fertile fronds 15—30 c.M. long in the unwinged part, scaly below. Fronds to 30 c.M. long or longer, 5—10 c.M. broad, the fertile ones not rarely more or less smaller than the barren ones, oblong, entire, acuminate, narrowed rather suddenly in the lower part and then decurrent at the base so as to form a not very long wing to the highest part of the stipe. Texture papyraceo-herbaceous to subcoriaceous; main veins distinct nearly to the edge; areolae copious, subquadrangular, divided into irregular-shaped smaller ones, with free veinlets. Sori numerous, often ranged in lines which are subparallel to the costa, 4—6 in a line between the main veins.

Malaya; China.

(2) **A. tricuspe**, *Bedd.*, *Handb. Ind. Ferns, Suppl.*, 44.

Rhizome stout, erect. Stipes 20—45 c.M. long, naked or nearly so, winged upwards. Fronds 25—30 c.M. long, 15—20 c.M. broad, deeply 3-lobate. Lobes lanceolate, long-acuminate, the lateral ones 12—15 c.M. long, 4—5 c.M. broad, the central one \pm 25 c.M. long, $6\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad; margins entire. Texture papyraceo-herbaceous; surfaces slightly pilose on the costae and veins beneath; main veins prominent nearly to the margin, connected by prominent transverse veins, between which are copious small areolae with free veinlets. Sori small, copious; indusium rather persistent.

Malacca.

* (3) **A. platanifolium**, *Mett.*, *Hk. Bk., Syn. Fil.*, 258; *A. angulatum*, *Hk., Spec. Fil.*, IV, 44.

Rhizome erect, the scales dense, dark-brown, linear-subulate. Stipes tufted,

15—40 c.M. long, ebeneous, naked or covered with numerous, minute, adpressed, glandlike, brown hairs, scaly at the base. Fronds 10—25 c.M. each way, palmately and more or less deeply 3—5-fid, cordate at the base. Terminal lobe \pm triangular, acuminate, lateral ones similar but smaller, the basal ones sometimes blunt. Texture papyraceo-herbaceous to subcoriaceous; each lobe with a prominent midrib; main veins distinct to the edge, connected by copious areolae, with free veinlets. Sori small, scattered, abundant; indusium peltate or reniform, often absent.

Malaya.

(4) **A. calcareum**, *Pr.*, Hk. Bk., Syn. Fil., 259; Mett., Farng., IV, Phegop. & Aspid., tab. XVIII, fig. 1—5; *Tectaria calcarea*, *Copel.*, Philipp. Journ., II, 415.

Rhizome subrepent. Stipes approximate, naked, 10—15 c.M. long. Fronds 30 c.M. or more long, 15—25 c.M. broad, with 4—6 stalked, distant, acuminate pinnae on each side below the pinnatifid apex, the lowest triangular, to 15 c.M. long, 7—10 c.M. broad, cut down to the rachis below in stalked, lanceolate, deeply and bluntly lobed pinnulae, the upper part lobed or more or less pinnatifid. Texture semicoriaceous; areolae very irregular, with free veinlets. Sori submarginal; indusium small, peltate or reniform.

Philippines.

* (5) **A. angulatum**, *J. Sm.*, Hk. Bk., Syn. Fil., 258.

Stipes 30—60 c.M. long, glossy, ebeneous, scaly below. Fronds 30—60 c.M. long, sometimes simple, usually with a large, cordate, entire or trifold terminal pinna and 1—2 lateral ones on each side, the lowest 15—30 c.M. long, 5—10 c.M. broad, often forked at the base with 2—3 branches; which are often unequal. Texture papyraceo-herbaceous; midribs glossy, ebeneous; main veins distinct to the edge; areolae fine, copious, with free, included veinlets. Sori small, scattered, abundant; indusium minute, fugaceous.

Malaya.

(6a) **A. Kunstleri**, *Bedd.*, Handb. Ind. Ferns, Suppl., 44.

Stipe and rachis minutely pubescent. Fronds 30—45 c.M. long, deltoid-ovate, pinnate. Upper pinnae narrow-lanceolate, \pm $12\frac{1}{2}$ c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, entire or with scalloped margins, lowest pair broad-deltoid, the lower side produced towards the base into 3—4 long lobes, the largest lobes

10—13 c.M. long, the upper side deeply scalloped. Texture rather coriaceous; surfaces naked except the main veins which are zigzag and tolerably prominent nearly to the margin; areolae rarely with free veinlets. Sori very small, numerous, scattered, generally on connected veinlets.

Malacca.

(6b) **A. oligophyllum**, *Rst.*, Fed., Rep., V, 15.

Stipes tufted, 1 M. long or longer, straight, erect, clothed towards the base with lanceolate, long-acuminate, ferruginous scales 1 c.M. long. Fronds to 50 c.M. long, 20 c.M. broad, pinnate (quinate). Pinnae simple, elongate-ovate, the lateral ones erecto-patent, more or less incurved, the base cuneate, the margin entire or obscurely repand, the apex long-cuspidate, lowest pinnae subopposite, \pm 30 c.M. long, \pm 7 c.M. broad, short-stalked, the stalks slightly marginate, highest pinnae sessile, smaller than the lowest ones, the very base decurrent, terminal pinna largest, the base long-decurrent, sometimes joining with the next following ones. Texture membranaceo-chartaceous; rachis naked, livid-brown, glossy, canaliculate above; surfaces naked; costae and veins stramineous, \pm prominent beneath; main veins erecto-patent, nearly straight, upcurved towards the apex; primary areolae subquadrangular, in \pm 9 rows on each side, divided into subhexagonal or irregular smaller ones, with copious included free veinlets. Fertile pinnae narrower than the barren ones. Sori large, numerous, compital, in \pm 18 longitudinal rows, 4—5 to a row (in 4—5 rows between the main veins?); indusium persistent, afterwards corrugate.

Sumatra.

* (7) **A. Haenkei**, *Pr.*, Mett., Ann. Mus. Bot. L. B., I, 258.

Rhizome? Stipes \pm 15 c.M. long or longer, stramineous like the rachis. Fronds 30 c.M. long or longer, \pm 20 c.M. broad, broadly oblong, pinnate with 4 or more pinnae on each side below the terminal one. Terminal pinna oblong, sinuate or subpinnatifid, the base cuneato-rotundate, lateral pinnae erecto-patent, to 15 c.M. long, \pm 3 c.M. broad, lanceolate-oblong, acuminate, falcate, narrowed at the base, or the upper base cuneate or truncato-rotundate and the lower one rounded, higher pinnae adnate to the rachis, the lower ones short-stalked, unequal-sided, sub-2-fid. Texture rigidly membranaceous; surfaces naked, or the upper one short-hairy on the costae only; main veins prominent, erecto-patent, upcurved; areolae in 4—5 rows on each side of the

costae, translucent, divided in smaller secondary ones. Sori in 2 rows, 2 or rarely 3 to each primary areola, dorsal on the veinlets; indusium persistent.

Timor, (? Malaya); Murianne Islands.

* (8) **A. ternatense, v. A. v. B.**

Stipes 30—45 c.M. long, naked, firm. Fronds \pm 45 c.M. long. Terminal pinna lanceolate, \pm 10—15 c.M. long, \pm 2 c.M. broad, gradually narrowed towards both ends, the edge entire or irregularly sinuate, the apex acute or acuminate, lateral pinnae 6—8 on each side, suberect, similar to the terminal one, the highest sessile, decurrent, free or joining with the terminal one, the lower stalked, the lowest forked at the base. Texture herbaceous-membranaceous; surfaces naked; main veins fine, erecto-patent; areolae subquadrangular, in 2 rows between the main veins, each with a single, included, excurrent, free veinlet; costae prominent beneath. Sori copious, slightly immersed, 1 to each areola, terminal on the free veinlets, forming 2 regular rows between the main a veins, with \pm 5—7 sori to a row; indusium persistent. Barren fronds not seen.

Ternate (Teysmann).

(9) **A. Beccarianum, Bk.,** Becc., Mal., III, 43; *A. Cesatianum* C. Chr., Ind. Fil.

Rhizome erect, densely clothed with firm, linear, acuminate, brown scales. Stipes tufted, brown, firm, \pm 50 c.M. long, sparingly paleaceous above and below the centre. Fronds oblong-deltoid, \pm 50 c.M. long, 2—3-pinnatifid. Pinnae in \pm 3 opposite pairs, the lower ones short-stalked, inequilateral, deltoid, produced on the lower side. Lower pinnulae deeply pinnatifid, only adnate at the base. Segments oblong. Texture firm; surfaces naked like the castaneous rachis; main veins straight from the costa to the margin; veins immersed, obscure; areolae copious, hexagonal. Sori in 2 rows, parallel to the main veins; indusium large, glabrous, persistent.

New Guinea.

(10) **A. repandum, Willd.,** Hk. Bk., Syn. Fil., 258; *A. platyphyllum, Pr.,* Epim., 65; *Sagenia repanda, Moore, Ind.,* 102; *Tectaria crenata, Cav.,* Copel., Philipp. Journ., II, 414, p. p.

Stipes 50—75 c.M. long, furnished with a few hair-like scales near the base, naked upwards. Fronds 60—90 c.M. long, pinnatifid at the apex. Pinnae 4—8 on each side, 15—30 c.M. long, 3—5 c.M. broad, linear-oblong,

acuminate, slightly and bluntly sinuate, the base narrowed, the lowest stalked, 2-partite. Texture subcoriaceous; main veins distinct to the edge; areolae copious, with free veinlets. Sori large, in 2 rows between the main veins. — Resembling *A. pachyphyllum*, *Kze* & *polymorphum*, *Wall.* (Bk.)

Malacca, Borneo, Philippines, New Guinea.

§ 2. SAGENIA (*).

1. *Sori scattered irregularly.*
2. *Fronde pinnatifid.*
 - *. *Stipes not winged.* (11) *A. Everetti.*
 - *.*. *Stipes long-winged.* (12) *A. vastum.*
- 2.2. *Fronde simple or more generally pinnate, 3—∞-foliate.*
3. *Simple frond or terminal pinna entire or sinuate.*
 - 4. *Fronde simple or 3-foliate.*
 - §. *Stipes winged nearly to the base.* (13) *A. pteropodum.*
 - §.§. *Stipes not winged, or winged in the upper half only.*
 - ‡. *Rhizome wide-creeping.* (14) *A. ternatum.*
 - ‡.‡. *Rhizome ± erect.* (15) *A. subcaudatum.*
 - 4.4. *Fronde 5—∞-foliate.*
 - 5. *Lowest pinnae simple, or forked at the base.*
 - 6. *Edges entire, crenate or repand.*
 - 7. *Rhizome erect.* (16) *A. psilopodum.*
 - 7.7. *Rhizome creeping.* (17) *A. polymorphum.*
 - 6.6. *Edges sinuate.* (18) *A. menyanthidis.*
 - 5.5. *Lowest pinnae 3-partite or 3-nate.*
 - *. *Lateral pinnae ± 2 on each side. Texture ± coriaceous.* (19) *A. Lobbii.*
 - *.*. *Lateral pinnae ± 5 on each side. Texture ± herbaceous.* (20) *A. semibipinnatum.*
- 3.3. *Simple frond or terminal pinna deeply lobed at the base.*
 - a. *Rhizome erect or ascending.*
 - b. *Stipes black.*
 - c. *Fronde mostly under 25 c.M. long.* (21) *A. Weinlandii.*

(*) See also *A. platanifolium*, *Mett.* & *calcareum*, *Pr.*

c.c. Fully developed fronds more than 25 c.M. long.
(22) *A. melanocaulon*.

b.b. Stipes not black.

×. Stipes greyish, finely villose. (23) *A. irriguum*.

×.×. Stipes castaneous, slightly scaly at the base.

(24) *A. nebulosum*.

a.a. Rhizome creeping.

(25) *A. subtriphylum*.

1.1. Sori in 2 rows between the main veins (*), in 1 row on each side of the costae in No. 29.

α. Fronds palmatifid or pinnatifid.

β. Stipes winged nearly or quite to the base.

γ. Fronds ± 20—30 c.M. long, the edges serrate or toothed.

(26) *A. heterodon*.

γ.γ. Fronds ± 60—120 c.M. long, the edges entire, repand or sinuate.

(27) *A. decurrens*.

β.β. Stipes not winged.

(28) *A. palmatum*.

α.α. Fronds 3-nate, each division 3-partite.

(29) *A. stenophyllum*.

α.α.α. Fronds simple or generally 3—∞ foliate, the divisions simple, pinnatifid, or even pinnate at the base.

β'. Texture herbaceous to papyraceous.

γ'. Stipes more or less densely hairy.

(30) *A. Keckii*.

γ'.γ'. Stipes naked, or scaly at the base.

δ'. Lateral pinnae in ± 1—6 pairs below the terminal one.

ε'. Lowest pinnae pinnatifid or pinnate at the base.

×. Stipes brown or rufous.

*. Central pinnae sinuate or slightly lobed.

(31) *A. variolosum*.

*. *. Central pinnae pinnatifid.

(32) *A. coadunatum*.

×.×. Stipes ebeneous.

(33) *A. latifolium*.

ε'.ε'. Lowest pinnae forked at the base. Fronds 60—90 c.M. long.

(34) *A. grande*.

δ'.δ'. Lateral pinnae in ± 9—11 pairs, the 2—4 lowest pairs 2—3-furcate.

(35) *A. nudum*.

γ'.γ'.γ'. Stipes more or less scaly throughout.

(36) *A. multicaudatum*.

(*) See also *A. polymorphum*, Wall.

$\beta'.\beta'$. *Texture generally chartaceous to coriaceous.*

§. *Surfaces pubescent. Texture moderately firm.*

(37) *A. melanorachis.*

§.§. *Surfaces naked. Texture \pm coriaceous except in No. 40.*

*. *Lowest pinnae simple. Sori dense, superficial.*

(38) *A. persoriferum.*

.. *Lowest pinnae forked, lobed or pinnatifid. Sori superficial.*

×. *Fronds under 50 c.M. long. Sori in close rows, often confluent, on connected veins. (39) A. siifolium.*

×.×. *Fully developed fronds often more than 50 c.M. long. Sori not in close rows, on free veinlets. Texture herbaceous to chartaceous.*

(40) *A. pachyphyllum.*

..*. *Lowest pinnae pinnate. Sori copious, immersed.*

(41) *A. brachiatum.*

(11) **A. Everetti**, **C. Chr.**, Ind. Fil.; *Nephrodium Everetti*, **Bk.**, Kew Bull., 1896, 41.

Stipes slender, 15—20 c.M. long, naked, castaneous. Fronds deltoid, 20—25 c.M. each way, cut down nearly to the costa. Lobes 2 on each side lanceolate or oblong-lanceolate, acuminate, entire, the lowest largest, $2\frac{1}{2}$ — 3 c.M. broad. Texture membranaceous; both surfaces naked; primary veins arcuate, reaching the margin; areolae small, copious, with free veinlets. Sori copious, scattered, small, superficial; indusium membranaceous, persistent.

Natuna Islands.

* (12) **A. vastum**, **Bl.**, Christ, Farnkr. d. Erde, 250; *A. alatum*, Wall., Hk. & Grev., Ic. Fil., tab. CLXXXIV; *Sagenia vasta*, Moore, Rac., Flor. Btz., I, 196; *S. alata*, Bedd., Ferns. Br.I., II, tab. CLXIX; *Tectaria vasta*, Copel., Philipp. Journ., II, 411.

Rhizome creeping or erect, clothed with rigid, brown, lanceolate, subentire scales. Stipes scattered or approximate, narrowly winged often nearly or quite to the base, 30—60 c.M. long. Fronds ovate, 50—120 c.M. long, 15—60 c.M. broad, cut down to a broadly winged rachis into entire, distant, linear-oblong segments 7—50 c.M. long, 2—5 c.M. broad, acute or acuminate, about 4—6 on a side. Texture papyraceo-herbaceous or subcoriaceous; main veins distinct, reaching to $\pm \frac{3}{4}$ of the way or nearly to the edge,

connected with copious areolae with included free veinlets. Sori numerous, densely crowded, often confluent; indusium small, persistent.

Malaya; Burmah, North. India.

(15) **A. pteropodum**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 183; *Nephrodium pteropodum*, *Bk.*, Journ. of Bot., XXVI, 325; *Sagenia pteropoda*, *Christ*, Ann. Bot., XX, 108; (? *Polypodium Scortechinii*, *Bk.*, C. Chr., Ind. Fil.).

Rhizome wide-creeping. Stipes with a narrow wing beginning \pm 5 c.M. from the base, \pm $\frac{1}{2}$ — $\frac{3}{4}$ c.M. broad, 30—45 c.M. above it. Fronds oblong-lanceolate, acuminate, entire, \pm 30 c.M. long, 7—10 c.M. broad, narrowed at the base gradually. Main veins distinct nearly to the edge; areolae with copious free veinlets. Sori minute, scattered irregularly, sometimes confluent; indusium minute, obscure.

Borneo.

* (14) **A. ternatum**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam. I⁴, 184; *Nephrodium ternatum*, *Bk.*, Hk. Bk., Syn. Fil., 296.

Rhizome wide-creeping, scaly. Stipes 15—25 c.M. long, slender, stramineous. Fronds sometimes simple, 20—25 c.M. long, 5—8 c.M. broad, oblong, entire, narrowed gradually to the apex and base, but more usually trifoliate, the lateral pinnae like the terminal one but smaller. Texture papyraceo-herbaceous; main veins distinct nearly to the edge; areolae copious, but rather large, with few free veinlets. Sori distant, in 2 distinct rows and others scattered between; indusium distinct.

Borneo.

* (15) **A. subcaudatum**, *v. A. v. R.*

Rhizome erect or suberect, the scales linear-lanceolate, \pm acuminate, dark-brown. Stipes \pm tufted, erect, naked, or scaly at the base, 6—12 c.M. long, the upper $\frac{1}{2}$ — $\frac{1}{5}$ narrowly winged by the decurrent base of the frond except in the truly trifoliate forms. Fronds simple, deeply 5-fid or 3-foliate, with the central leaflet similar to the simple fronds, lanceolate, 15—25 c.M. long, 4—5 c.M. broad below the middle, the base cuneate, \pm gradually narrowed, the margin entire or repand, the apex long-acuminate, subcaudate; lateral leaflets similar but smaller and narrower, erecto-patent, slightly upcurved, free or joined at the base with the terminal one. Texture papyraceous; surfaces naked; costae and main veins prominent; main veins

distinct nearly to the edge, slightly upcurved; areolae distinct, fine, with numerous free veinlets. Sori not copious, scattered irregularly, some of them sometimes confluent; indusium fugacious.

Borneo (Teuscher).

(16) **A. psilopodum**, *C. Chr.*, *Ind. Fil.*; *Nephrodium subdigitatum*, *Bk.*, *Journ. of Linn. Soc., Bot.*, XXIV, 259; *Sagenia subdigitata*, *Christ*, *Ann. Btz.*, XX, 100.

Rhizome erect. Stipes densely tufted, 7—10 c.M. long, with a few, spreading, black-brown scales near the base. Fronds small, deltoid, subdigitate. Pinnae 5—7, lanceolate, acuminate, the edge entire or repand, the lower ones usually forked at the base, the terminal one the largest, distinctly stalked, \pm 7—8 c.M. long, 1—1½ c.M. broad at the middle, distinctly repand, narrowed gradually to both ends. Surfaces naked; no distinct main veins; areolae copious. Sori not very small, irregularly scattered; indusium firm, persistent.

Borneo.

* (17) **A. polymorphum**, *Wall.*, *Christ*, *Farnkr. d. Erde*, 229; *Bedd.*, *Ferns S.I.*, tab. CXVI; *A. grandifolium*, *Mett.*, *Farngr.*, IV, *Phegop. & Aspid.*, 124; *A. angulatum*, *Christ* (not *J. Sm.*), *Bull. Herb. Boiss.*, 2^e Série, VI, 1003; *Nephrodium polymorphum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 297; *Sagenia polymorpha*, *Christ*, *Ann. Btz.*, XX, 108; *Tectaria polymorpha*, *Copel.*, *Philipp. Journ.*, II, 413; — var. *Wightii*: *N. Wightii*, *Clarke*, *Trans. Linn. Soc., Bot.*, 2^d Series, I, 538, tab. LXXVI; — var. *subvariegatum*, *Hort. Bog.*

Rhizome creeping. Stipes seriate, to 30 c.M. long or longer, brown, naked or \pm scaly. Fronds 30—120 c.M. long, 25—45 c.M. broad. Terminal pinna lanceolate-oblong, 10—50 c.M. long, 2—8 c.M. broad, entire, acuminate, \pm rounded at the base; below this 2—6 similar, lateral pinnae on each side, the higher ones \pm sessile or adnate to the rachis, mostly distinct, rarely joined with the terminal one, the lower ones \pm stalked, often forked at the base on the lower side. Texture papyraceous to coriaceous; rachis brownish, naked or nearly so; surfaces naked; areolae fine, with copious free veinlets. Fertile pinnae not contracted. Sori small, abundant, scattered, placed on the connected veins; indusium mostly present, reniform or peltato-cordate to orbicular, the edge not rarely irregularly fimbriate.

Var. Wightii: Fertile pinnae contracted; sori in 2 rows between the main veins.

Var. subvariegatum: Like var. *Wightii* but the fronds inconspicuously variegated with green and pale-green.

Malaya; North. India, Burmah, Ceylon.

(18) **A. menyanthidis**, *Pr.*, Copel., Polypod. Philipp., 38; *Nephrodium Menyanthidis*, *Bk.*, Hk. Bk., Syn. Fil., 297; *Tectaria Menyanthidis*, Copel., Philipp. Journ., II, 414.

Rhizome creeping. Stipes 15—30 c.M. long, naked, brownish. Fronds 30—60 c.M. long, 30 c.M. or more broad, with an acuminate, oblong-lanceolate, sinuate, terminal pinna 15—30 c.M. long, 2½—4 c.M. broad; below this 2—4 similar pinnae on each side, the lowest stalked, sometimes forked. Texture subcoriaceous; main veins close, conspicuous, reaching the edge; areolae fine, with copious free veinlets. Sori small, on the connected veins.

Philippines, New Guinea; Salomon Islands.

(19) **A. Lobbii**, *Hk.*, Spec. Fil., IV, 49, tab. CCXXXII; *Nephrodium Lobbii*, *Bk.*, Hk. Bk., Syn. Fil., 297; *Sagenia Lobbii*, *Christ*, Ann. Btz., XX, 100.

Stipes 15 c.M. or more long, naked, substramineous. Fronds ± 30 c.M. each way, subdeltoid, with an entire terminal pinna 10—15 c.M. long, 1—2 c.M. broad, tapering gradually from the middle to each end, and 2 similar lateral ones on each side, the lowest long-stalked and tripartite or ternate. Texture subcoriaceous; veins obscure, the main ones close and very oblique; areolae without free veinlets. Sori copious.

Borneo.

* (20) **A. semibipinnatum**, *Wall.*, Hk., Spec. Fil., IV, 59, tab. CCXXXI; *Nephrodium semibipinnatum*, *Bk.*, Hk. Bk., Syn. Fil., 297.

Stipes 50 c.M. or more long, pale-brown, naked. Fronds 50—45 c.M. long, 15—25 c.M. broad, with a long terminal pinna 15—25 c.M. long, ± 2½ c.M. broad, entire and narrowed towards both ends; below this ± 3—6 lateral pinnae on each side, the lowest 1—2 pairs tripartite or ternate. Texture papyraceo-herbaceous; veins inconspicuous, the main ones not reaching the edge; areolae fine, close, with free veinlets. Sori minute, confined to the connected veins.

Malacca, Banca, Borneo.

(21) **A. Weinlandii**, *Christ*, Bull. Herb. Boiss., 2^o Série, I, 453.

Rhizome suberect. Stipes subtufted, 10—20 c.M. long, slender. Rachis, main veins (and partially also the veins) blackish. Fronds \pm 10—20 c.M. long, 10—15 c.M. broad, ternate. Terminal division long-stalked, cuneate or subcordate, long-acuminate, 3-lobed, the lobes produced, with the margin broadly lobato-dentate, lateral divisions falcate, short-stalked, lobed, the lowest lobes on the lower side much produced. Texture herbaceous; surfaces naked; veins fine but conspicuous, reaching the margin; areolae with free veinlets. Sori central in the lobes, or scattered, deciduous, minute; indusium fugacious.

New Guinea.

* (22) **A. melanocaulon**, *Bl.*, Copel., Polypod. Philipp., 57; *Sagenia melanocaulon*, Moore, Rac., Flor. Btz., I, 193; *Nephrodium melanocaulon*, *Bk.*, Hk. Bk., Syn. Fil., 296; *Tectaria melanocaulon*, Copel., Philipp. Journ., II, 416.

Rhizome erect or ascending, clothed with black-brown, lanceolate scales to 10 m.M. long. Stipes 50—50 c.M. long, ebeneous, glossy, scaly at the base. Fronds 30—90 c.M. long, 15—50 c.M. broad, ovate- or lanceolate-deltoid, with a large, deeply pinnatifid terminal pinna and 1—4 lateral ones on each side, the highest ones entire or lobed, the central ones mostly deeply pinnatifid with oblong, acuminate lobes, the lowest stalked, 8—30 c.M. long, to 7—15 c.M. broad, often again pinnate at the base. Texture herbaceous-papyraceous; rachises ebeneous, glossy, naked or the upper side pubescent or glandular; both surfaces naked; main veins reaching the edge; areolae numerous, fine, with free veinlets. Sori numerous, minute; indusium small, fimbriate, fugacious.

Malaya; East. China, Tonkin.

* (23) **A. irriguum**, *J. Sm.*, Copel., Polypod. Philipp., 38; *Nephrodium irriguum*, *Bk.*, Hk. Bk., Syn. Fil., 297; *Tectaria irrigua*, Copel., Philipp. Journ., II, 415, p. p.

Stipes tufted, 10—15 c.M. long, greyish, finely villose. Fronds 15—50 c.M. long, 10—20 c.M. broad, subdeltoid, with a large, oblong, terminal pinna with a pair of deep, lanceolate lobes at the base; below this 1—2 pairs of slightly sinuate lateral pinnae, the lowest forked at the base. Texture papyraceo-herbaceous; main veins flexuose; areolae fine, copious, with free veinlets. Sori small, often confluent; indusium small, fugacious.

Philippines, Amboyna.

* (24) **A. nebulosum**, **C. Chr.**, Ind. Fil.; *Nephrodium nebulosum*, **Bk.**, Journ. of Bot., XVIII, 215.

Stipes tufted, castaneous, \pm 45 c.M. long, with only a few, small, brown, lanceolate scales at the base. Fronds deltoid, 30—40 c.M. long, rather less broad, with a deeply pinnatifid, large terminal pinna with short, obtuse upper lobes and large, lanceolate lower ones; below this 2—4 lanceolate, free pinnae, the lowest ones unequal-sided and forked at the base on the lower side. Texture moderately firm; both surfaces naked; main veins distinct to the edge, slender, erecto-patent, with very copious areolae between them, with free veinlets. Sori very abundant, minute; indusium minute, very fugacious.

Sumatra.

* (25) **A. subtriphyllum**, **Hk.**, Christ, Farnkr. d. Erde, 227; *Nephrodium subtriphyllum*, **Bk.**, Hk. Bk., Syn. Fil., 296; *Sagenia subtriphylla*, **Bedd.**, Ferns Br.I., I, tab. XLVIII; Ferns S.I., tab. CCXLII; *Tectaria subtriphylla*, **Copel.**, Philipp. Journ., II, 410.

Rhizome creeping. Stipes 30 c.M. long or longer, substramineous or brownish. Fronds 30—90 c.M. long and 20—75 c.M. broad, rarely smaller, subdeltoid, with a large, often deeply pinnatifid apex with entire or broadly toothed, lanceolate lobes; below this 1—2 pinnae on each side, the lowest stalked, distant, forked at the base or pinnatifid, the apex lanceolate. Texture herbaceous to semicoriaceous; surfaces somewhat hairy beneath and on the main veins above; main veins distinct to the edge; areolae fine, with copious free veinlets. Sori often moderately large, confined to the united veins.

Malaya; Polynesia, China, Burmah, Ceylon.

(26) **A. heterodon**, **Copel.**, Polypod. Philipp., 36; *A. Copelandii*, **C. Chr.**, Ind. Fil.; *Tectaria decurrens*, **Copel.**, Philipp. Journ., II, 412, p. p.

Rhizome erect, woody. Stipes 15—20 c.M. long, naked or nearly so, winged, the wing \pm 5 m.M. broad on each side, narrowed gradually towards the base. Fronds 20—30 c.M. long, 3—5-partite. Terminal (or central) segment oblanceolate, 4—5 c.M. broad, acuminate, irregularly serrate or toothed, the teeth 0—1 c.M. long, lateral segments smaller, erecto-patent, obscurely serrate. Texture herbaceous; surfaces naked; main veins reaching the margin; areolae irregular, with free veinlets. Sori large, in rows near the main veins; indusium persistent

Philippines.

* (27) **A. decurrens**, *Pr.*, Christ, Farnkr. d. Erde, 229; *A. pteropus*, Kze, Bedd., Ferns S.I., tab. LXXXII; *Sagenia decurrens*, *Houlst.* (not *Moore*), Rac, Flor. Btz., I, 196; *Nephrodium decurrens*, *Bk.*, Hk. Bk., Syn. Fil., 299; *Tectaria decurrens*, *Copel.*, Philipp. Journ., II, 412, p. p.; — var. *mamillosum*; *A. mamillosum*, *C. Chr.*, Ind. Fil.; *S. mamillosa*, *Moore*, in Gard. Chron., N. S., XXVI, 330.

Rhizome creeping or erect, scaly. Stipes 20—30 c.M. long, winged often nearly or quite to the base. Fronds 60—120 c.M. long, 30—60 c.M. broad, cut down to a winged rachis into 3—8 pairs of entire, repand or sinuate linear-oblong, acuminate laciniae 15—30 c.M. long, 2—6 c.M. broad, terminal segment similar, the lowest ones often bifurcate. Texture herbaceous or subcoriaceous; surfaces naked; main veins distinct to the edge, connected by copious areolae with free veinlets. Sori rather large, in 2 rows between the main veins; indusium persistent. — Resembling *A. vastum*, *Bl.*

Var. mamillosum: Sori deeply immersed so as to form prominent papilliae on the upper surface of the frond.

Malaya; *North. India*, *South. China*, *Polynesia*; — var. *mamillosum*: *Malaya*.

(28) **A. palmatum**, *Mett.*, Ann. Mus. Bot. L. B., I, 258; *Sagenia palmata*, *Christ*, Ann. Btz., XX, 108.

Rhizome short, erect; the scales lanceolate, ciliate or dentate, ferruginous, nearly glossy. Stipes densely tufted, 30—40 c.M. long, glossy, purpureous. Fronds 15—25 c.M. long, 20—25 c.M. broad, palmatipartite, cordate at the base. Central segment ovate-oblong, acuminate, the lateral ones 2-fid with a large, falcate, erecto-patent, anterior lobe and a smaller, spreading posterior one. Texture membranaceous or thinly chartaceous; surfaces naked; costae and veins prominent; primary areolae in 6—8 rows on each side, divided into nearly regular smaller ones. Sori in 2 regular rows between the main veins; indusium rigid, roundish-reniform, subpeltate, with a short-toothed margin.

Sumatra, *Borneo*.

(29) **A. stenophyllum**, *Diels*, Engl. & Prantl., Nat. Pfl.Fam., I⁴, 185; *A. Hosei*, *C. Chr.*, Ind. Fil.; *Nephrodium Hosei*, *Bk.*, Ann. of Bot., V, 350; *N. stenophyllum*, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 227, tab. XI.

Stipes slender, tufted, naked, castaneous, 10—13 c.M. long. Fronds ternate, 10—13 c.M. each way, the lateral segments nearly as long as the central one, each made up of 5 linear, entire, acuminate free or joined, secondary

segments 5—8 c.M. long and 6—7 m.M. broad at the middle, tapering gradually to the base. Surfaces naked; veins faint and indistinct; areolae copious, hexagonal. Sori in a single, irregular row on each side of the central vein of the segments; indusium large, naked, persistent.

Borneo.

(30) **A. Keckii**, *Luerss.*, Bot. Centralbl., XI, 76; *Nephrodium Keckii*, *Bk.*, Ann. of Bot., V, 550.

Stipes \pm 40 c.M. long, red-brown, \pm densely clothed with small, ferruginous hairs, \pm rough below, slightly canaliculate. Fronds \pm 48 c.M. each way, deltoid-ovate, 2-pinnatifid. Terminal pinna deeply pinnatifid, the lobes \pm falcate, entire, acuminate, the lowest lobe bluntly lobed; below this 2 pairs of lateral pinnae, which are \pm falcate, acuminate, with the lower side more or less deeply lobed, the lobes blunt or acuminate, falcate, lowest pinnae on stalks \pm 5 c.M. long. Texture papyraceous; rachis, costae and main veins hairy like the stipe; main veins distinct nearly or quite to the edge; areolae moderately large, with free veinlets. Sori few, in short, regular rows, large; indusium large, membranaceous, naked, persistent.

Sumatra.

* (31) **A. variolosum**, *Wall.*, Christ, Farnkr. d. Erde, 228; *A. Zollingerianum*, *Bedd.*, Ferns Br.I., II, tab. CCLI; *Nephrodium variolosum*, *Bk.*, Hk. Bk., Syn. Fil., 298.

Stipes to 50 c.M. or more long, naked, or scaly near the base, brownish. Fronds 15—45 c.M. long, 10—50 c.M. broad, subdeltoid, the apex deeply pinnatifid with lanceolate, acuminate, entire or sinuate lobes; below this \pm 2—3 pairs of lateral, lobed or sinuate pinnae, the lowest much the largest, often cut down nearly or quite to the rachis at the base with more or less deeply pinnatifid pinnulae, the higher ones less divided. Texture papyraceo-herbaceous; surfaces naked or somewhat puberulous; main veins distinct; areolae rather large, with free veinlets. Sori large, in rows near the main veins, terminating free veinlets. — Resembling *A. subtriphylum*, *Hk.*

Malaya; North. India, Burmah.

(32) **A. coadunatum**, *Wall.*, Christ, in Philipp. Journ., II, 187; *Mett.*, Farng., IV, Phegop. & Aspid., 118, p. p.; *Hk. & Grex.*, Ic. Fil., tab. CCII; *A. cicutarium*, *Christ*, Farnkr. d. Erde, 250, p. p.; (*Sw.*, C. Chr., Ind.

Fil., p p.); *Tectaria Christii*, Copel., Philip. Journ., II, 416; *Sagenia coadunata*, J. Sm., Bedd., Ferns S.I., tab. LXXXI.

Stipes rufous, \pm naked, sparingly scaly at the base, the scales acute, flaccid, brown. Fronds 50—90 c.M. long, pinnate, the rachis naked, smooth, glossy, flavido-rufous. Pinnae pinnatifid, \pm 1—4 on each side below the deeply pinnatifid terminal one, the lowest largest, ovate-lanceolate, acute, deeply pinnatifid, the lobes broadly ovate, obtuse or subacute, crenate or lobed. Texture thin, diaphanous; upper surface clothed with short, white pubescence, the margins ciliate; costulae slightly pubescent; veins copiously anastomosing, with copious, included, clavate, free veinlets. Sori in 1 row on each side of the costulae; indusium orbicular-reniform, with a pale margin. — This species was often confounded with the american *A. cicutarium*, Sw., which is quite another plant. — *Trop. Asia*.

Christ describes another form in Philipp. Journ., II, 187: *Aspidium malayense*, Christ (*Tectaria malayensis*, Copel., l. c., 416), distinguished from this by the scaly stipes, the naked fronds, the less copiously anastomosing veins, the rather wanting included veinlets and the peltate indusia. — *Malacca, Singapore, Luzon*.

(33) **A. latifolium**, J. Sm., Journ. of Bot., III, 410; *Nephrodium latifolium*, Bk., Hk. Bk., Syn. Fil., 298; (? *Tectaria melanocaulon* or *suifolia* Copel., Philip. Journ., II, 418); *T. latifolia*, Copel., l. c., 416.

Stipes 50 c.M. or more long, glossy, ebeneous. Fronds 60—90 c.M. long, 50—45 c.M. broad, the apex deeply pinnatifid with lanceolate-acuminate lobes, the lower lobes sinuated; below this 5—5 pinnae on each side, the lowest much the largest, 20—50 c.M. long, 15—20 c.M. broad, deeply pinnatifid above, pinnate at the base, lobes acute. Texture papyraceo-herbaceous; rachis glossy like the stipe; main veins distinct to the edge; areolae fine, with copious free veinlets. Sori in irregular rows near the main veins, mainly or entirely on the connected veins.

Celebes, Amboyna, New Guinea; Polynesia.

(34) **A. grande**, J. Sm., Copel., Polypod. Philipp., 36; *A. grandifolium*, Pr., C. Chr., Ind. Fil.; *Nephrodium grande*, Bk., Ann. of Bot., V, 350; *Tectaria grandifolia*, Copel., Philip. Journ., II, 415.

Stipes rufescent, shining. Fronds 60—90 c.M. long, ovate-oblong, pinnate. Lateral pinnae in 5—6 pairs below the deeply 3-fid apex, \pm 25 c.M. long, 6—6½ c.M. broad, oblong-lanceolate, caudato-acuminate, slightly sinuate, the

lowest ones stalked, cuneately rounded at the base, decurrent upon the petiole, unequally 2-partite. Texture membranaceous; surfaces naked; venation as in *Drynaria*; primary areolae in several rows on each side of the costae. Sori in 2 rows between the main veins and approximate to them, 2 in each primary areola; indusium with a short sinus.

Philippines.

(35) **A. nudum**, *Diels*, in Schum. & Laut., Flor. deut. Sch. geb., 116; *Nephrodium nudum*, *Bk.*, Journ. of Bot., XVII, 41.

Stipes \pm 60 c.M. long, pale-brown, naked. Fronds oblong-deltoid, 45—60 c.M. long, 15—50 c.M. broad. Terminal pinna lanceolate, 15—30 c.M. long, 2—2½ c.M. broad, much acuminate, the edge entire, the base gradually narrowed; below this 9—11 distant, sessile, similar but smaller pinnae, the lowest stalked, the 2—4 lower bi- or trifurcate from the base. Texture membranaceous; rachis and surfaces naked; main veins erecto-patent, distinct to the edge; areolae copious, small, irregular, with free veinlets. Sori in regular rows near the main veins; indusium naked, persistent.

Borneo; Bismarck Islands.

(36) **A. multicaudatum**, *Wall.*, Bedd., Handb. Ind. Ferns, 222; *Nephrodium multicaudatum*, *Clarke*, Trans. Linn. Soc., Bot., 2^d Series, I, tab. LXXVII; *Sagenia Griffithii*, *Bedd.*, Ferns Br.I., II, tab. CCCXXXVII.

Stipes 30—90 c.M. long, brownish, often densely clothed (nearly its whole length) with linear-subulate, brown, persistent scales. Fronds 90—120 c.M. long, pinnate. Terminal pinna large, deeply pinnatifid, the lobes lanceolate-acuminate; below this several lateral pinnae on each side, the lowest deltoid, \pm 30 c.M. or more long and nearly as broad, deeply pinnatifid above and pinnate below. Texture firm, herbaceous; rachises rusty beneath; main veins distinct to the edge; areolae with copious free veinlets. Sori rather large, in 2 rows between the main veins or more numerous and scattered, generally terminal on free veinlets; indusium reniform or peltate.

Malacca; North. India, Burmah.

(37) **A. melanorachis**, *C. Chr.*, Ind. Fil.; *Nephrodium melanorachis*, *Bk.*, Journ. of Bot., XXVI, 325.

Stipes black, \pm 45 c.M. long, scaly on the lower half, but more densely at the base, the scales linear, spreading, brown. Fronds oblong-deltoid, 2-pinnatifid, 45—60 c.M. long, \pm 30 c.M. broad. Pinnae \pm 3 on each side, the terminal one pinnatifid, the lowest much the largest, deltoid, produced on the

lower side, deeply pinnatifid, the lobes lanceolate. Texture moderately firm; both surface finely pubescent; main veins distinct to the margin; areolae with copious free veinlets. Sori 2-serial, approximate to the main veins; indusium small, membranaceous, \pm persistent.

Borneo.

* (38) **A. persoriferum**, *Copel.*, Polypod. Philipp., 36; ? *Tectaria crenata*, *Cav.*, *Copel.*, Philipp. Journ., II, 414, p.p.

Rhizome short, ascending. Stipes 40—60 c.M. long, stramineous or brown, the base clothed with lanceolate, arcuate scales \pm 12 m.M. long, the rest naked like the rachis. Fronds 30—50 c.M. long, ovate, pinnate, the barren ones the largest. Terminal pinna \pm 25 c.M. long, \pm 7 c.M. broad, acuminate, entire or subrepand; below this 3—5 lateral pinnae on each side, which are sessile, erecto-patent, straight or subarcuate, lanceolate, 15—20 c.M. long; fertile pinnae smaller, \pm 10 c.M. long, \pm 2 c.M. broad, obscurely crenate, the lowest stalked. Texture subcoriaceous; surfaces naked; main veins almost reaching the margin; areolae irregular, with included, free veinlets. Sori large, in 2 rows between the main veins, almost covering the entire lamina; indusium persistent. — A form from Borneo, cultivated in the Buitenzorg gardens has the pinnae of both barren and fertile fronds \pm 2 \times as broad as in the philippine plant (forma latipinna).

Mindanao, Borneo.

* (39) **A. siifolium**, *Mett.*, Ann. Mus. Bot. L.B., I, 257; *A. meniscocarpum*, *Mett.*, Farng., IV, Phegop. & Aspid., No. 287; *A. biserialum*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1002; *Nephrodium siifolium*, *Bk.*, *Hk.*, *Bk.*, Syn. Fil., 299; *Tectaria siifolia*, *Copel.*, Philipp. Journ., II, 414; — var. *Teysmannianum*, *Christ*, Ann. Bot., XV, 158; *A. Teysmannianum*, *Hk.*, Spec. Fil., IV, tab. CCXXXVI; — var. *subtrifoliatum*, *Christ*, l. c.

Rhizome oblique, the scales black-brown. Stipes few, tufted, 15—45 c.M. long, brownish, naked, or scaly at the base. Fronds 30—45 c.M. long, 15—20 c.M. broad, with a broad, oblong, acuminate, terminal pinna and 2—4 lateral ones on each side. Lateral pinnae short-stalked or sessile, the lower ones stalked, 7—15 c.M. long, 4—6 c.M. broad, bipartite, the margin rather deeply lobed. Texture subcoriaceous; main veins close, distinct to the edge; areolae fine, copious, with free veinlets. Sori large, copious, in 2 close rows, the contiguous pairs often confluent, on connected veins; indusium ciliate. — Fertile pinnae 7—10 c.M. long, lanceolate, acuminate.

Var. Teysmannianum: Pinnae wavy, broadly crenate or sinuate.

Var. subtrifoliatum: Barren fronds with a broad, ovate, terminal pinna to \pm 16 c.M. long, to \pm 8 c.M. broad and below this commonly 1 pair of similar lateral pinnae. Fertile pinnae small, subentire, in 2—3 pairs.

Malaya; Polynesia; — var. *Teysmannianum:* *Sumatra, Celebes;* — var. *subtrifoliatum:* *Celebes.*

* (40) **A. pachyphyllum**, *Kze*, Christ, Farnkr. d. Erde, 229; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 97, A—C; *A. siifolium*, *Mett.*, var. *pachyphyllum*, Christ, Ann. Bot., XV, 158; *Sagenia pachyphylla*, *Moore*, Rac., Flor. Bot., I, 193; *Nephrodium pachyphyllum*, *Bk.*, Hk. Bk., Syn. Fil., 299; *Tectaria crenata*, *Cav.*, Copel., Philipp. Journ., II, 414, p. p.

Caudex erect, to 10 c.M. high, the scales ovate-lanceolate, dark-brown. Stipes 50—50 c.M. long, naked, brownish, scaly at the base. Fronds 60—100 c.M. long, \pm half as broad, with an oblong-lanceolate, entire or sinuate terminal pinna 10—50 c.M. long, 2—5 c.M. broad, often deeply pinnatifid at the base; below this 2—8 erecto-patent, entire or repand lateral pinnae on each side, the lowest lobed, forked or pinnatifid on the lower side. Fertile pinnae often narrower, or even \pm contracted. Texture herbaceous to chartaceous; main veins straight or slightly curved, distinct nearly to the edge; areolae fine, with copious free veinlets. Sori large, in regular rows, distant, on free veinlets; indusium entire.

Malaya; Polynesia.

(41) **A. brachiatum**, *Zoll. & Mor.*, Nat. Geneesk. Arch., I, 599; *A. Zollingerianum*, *Kze*, Bot. Zeit., 1846, 462; *A. impressum*, *Hk.*, Spec. Fil., IV, 58; *Nephrodium Zollingerianum*, *Bk.*, Hk. Bk., Syn. Fil., 298.

Stipes 20—30 c.M. long, naked. Fronds 20—30 c.M. long, 15—20 c.M. broad, subdeltoid. Terminal pinna deeply pinnatifid; below this 5—4 lateral pinnae on each side, the lowest largest, long-stalked, deltoid, cut down below into distinctly separated, deeply pinnatifid pinnulae with blunt, linear-oblong lobes. Texture subcoriaceous; veins beneath prominent, anastomosing principally in large costular arches; areolae large, with included free veinlets, depressed so as to form cavities in which on the ends of the free veinlets the large, copious sori are placed.

Java.

26. CYRTOMIUM, *Presl.*

Sori round, terminal on the included free veinlets. Indusium round, peltate.

Rhizome erect. Stipes not articulate to the rhizome. Fronds pinnate with the pinnae whether or not \pm auricled at the base, the base \pm rounded; veins pinnate from a central costa, angularly anastomosing, forming \pm hexagonal areolae within which are 1—3 included, excurrent free veinlets. Bedd., Handb. Ind. Ferns, fig. 206. (*Polystichum* p. p., C. Chr., Ind. Fil.).

Asia and Africa.

* **C. falcatum, Pr.**, Bedd., Handb. Ind. Ferns, 211; *Aspidium falcatum*, Sw., Hk. Bk., Syn. Fil., 257; Hk., Fil. Exot., tab. XCH; *Polystichum falcatum*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 194; C. Chr., Ind. Fil.; — var. *caryotideum*, Bedd., Handb. Ind. Ferns, 211; *C. caryotideum*, Pr., Tent. Pterid., 86; Bau. & Hk., Gen. Fil., tab. XLIX, C; Bedd. Ferns S.I., tab. CXIX; *A. caryotideum*, Wall., Hk., Gard. Ferns, tab. XIII; Hk. & Grev., Ic. Fil., tab. LXIX.

Rhizome short, erect, clothed like the base of the stipes with large, dark-coloured scales. Stipes tufted, 15—50 c.M. long. Fronds 50—60 c.M. long, 15—25 c.M. broad, ovate. Pinnac \pm 10—12 on each side, ovate-acuminate, subfalcate, \pm erecto-patent, 10—15 c.M. long, 2½—5 c.M. broad, the point acuminate, the edge entire or slightly undulate, the upper base suddenly narrowed or broadly rounded, sometimes auricled, the lower base narrower. Texture coriaceous; surfaces naked. Sori copious, scattered.

Var. caryotideum: Pinnae sharply toothed, slightly lobed, sometimes auricled on 1 or on both sides at the base.

Malaya; Ceylon, Br. India, China, Japan, Sandwich Islands, South. Africa, Madagascar.

TRIBE III. DAVALLIEAE.

27. CYSTOPTERIS, *Bernhardi*.

Sori roundish, dorsal on a vein or veinlet. Indusium membranaceous, round and cucullate or lanceolate, convex, covering the sorus, but at length more or less reflexed or fugacious, attached at the inner side of the receptacle, opening towards the apex or the margin of the segment.

Rhizome creeping or erect. Stipes not articulate to the rhizome. Fronds bipinnate or more compoundly divided; veins free, forked or pinnate. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 89, B—C.

Ferns inhabiting the temperate zones and the alpine regions of the tropics.

*. *Stipes* dark-purplebrown. *Indusium* ligulate, fugacious. (1) *C. tristis*.

.. *Stipes* stramineous. *Indusium* semiorbicular or transversally oblong.

(2) *C. tenuisecta*.

* (1) ***C. tristis***, *Mett.*, Ann. Mus. Bot. L. B., I, 241; *Davallia tristis*, *Rac.*, Flor. Btz., I, 151; *Alsophila tristis*, *Bl.*, Hk. Bk., Syn. Fil., 44; *C. Chr.*, Ind. Fil.; *Aspidium tristis*, *Bl.*, Enum., 169.

Rhizome erect, short, densely clothed with entire, lanceolate, brown scales. Stipes 50—90 c.M. long, \pm 7 m.M. thick at the scaly, tuberculato-asperous base, purple-brown, villose towards the rachis. Fronds ovate-lanceolate to deltoid, \pm 45—90 c.M. long, \pm 40—60 c.M. broad, 3-pinnate, Pinnae ovate to lanceolate, acuminate, short-stalked, the largest to 10 c.M. broad. Pinnae lanceolate, short-stalked. Ultimate segments sessile, narrowed at the base, 5—12 $\frac{1}{2}$ m.M. long, to \pm 4 m.M. broad, oblong, obtuse, incised to \pm $\frac{1}{2}$ -way down to the costa, more or less decurrent. Texture subcoriaceous; rachises and costae pilose or fibrillose especially beneath; veins simple or forked. Sori 1 to each lobule, covered by a small, at length reflexed, fugacious, ligulate indusium; receptacle elevated.

Java.

(2) ***C. tenuisecta***, *Mett.*, Ann. Mus. Bot. L. B. I, 241; *Alsophila tenuisecta*, *Bl.*, Moore, Ind., 58; *Aspidium tenuisectum*, *Bl.*, Enum., 170.

FILICES.

Rhizome short, oblique, clothed like the young fronds with large, membranaceous, ferrugineous, ovate-lanceolate, acuminate scales. Stipes stramineous, to 45 c.M. long, scaly at the base, purveyed like the rachis with articulated hairs and minute tubercles. Fronds to 45 c.M. long, ovate-lanceolate. Pinnæ sessile, subopposite, spreading, subdivaricating, 15—30 c.M. long, oblong-lanceolate, the lower ones distant. Pinnulae numerous, approximate, spreading, 5—8 c.M. long, 15—25 m.M. broad, oblong-lanceolate. Tertiary divisions oblong, obtuse, the lower ones sessile, pinnatifid, the higher ones adnate, shortly toothed or crenate. Texture membranaceous, flaccid or firm; costae and veins sparingly hairy; under surface glandular; veins fine. Indusium transversely oblong or semiorbicular, with the margin fimbriato-glandular, at length reflexed.

Java.

28. ODONTOSORIA, *Presl.*

Sori roundish or transversally oblong, terminal or subterminal on the ultimate segments of compoundly divided fronds, solitary on the vein tips or uniting the tips of \pm 2—3 veins. Indusium of the same shape, attached at the inner side of the sorus, opening outwardly.

Rhizome short. Stipes not articulate to the rhizome. Fronds compound; ultimate segments more or less narrowly cuneate, the apex \pm truncate. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 116, D.

Pantropical.

Arrangement of the sections.

- § 1. **EU-ODONTOSORIA.** Fronds small or middle-sized, erect, not climbing.
 § 2. **STENOLOMA.** Fronds long, climbing. (Not in Malaya).

§ 1. **EU-ODONTOSORIA.**

- §. *Sori solitary on the vein tips, 1—2 to a lobe.*
 *. *Lower pinnæ not caudate at the apex.* (1) *O. chinensis.*
 .. *Lower pinnæ caudate at the apex.* (2) *O. biflora.*
 §.§. *Sori uniting the apices of 2 or more veins or veinlets, occupying the whole breadth of the lobes.* (3) *O. retusa.*
-

* (1) **O. chinensis**, *J. Sm.*, Copel., Polypod. Philipp., 57; *Davallia tenuifolia*, *Sw.*, Rac., Flor. Btz., I, 156; *Bedd.*, Ferns S.I., tab. XVI; *Lindsaya tenuifolia*, *Mett.*, Christ, Farnkr. d. Erde, 296; *L. chinensis*, *Burck.*, Ann. Btz., IV, 90; *Stenoloma chinensis*, *Bedd.*, Handb. Ind. Ferns, 70; *Trichomanes Chinense*, *L.*, (oldest name).

Rhizome creeping, the scales dense, brown, hair-like. Stipes 2—20 m.M. apart, polished, naked, erect, \pm 10—50 c.M. long. Fronds 3-pinnatifid to 4-pinnate, \pm 10—50 c.M. long, 4—25 c.M. broad, lanceolate or ovate. Lower pinnae ovate to deltoid-lanceolate, erecto-patent, 5—15 c.M. long, $1\frac{1}{2}$ —8 c.M. broad. Pinnulae \pm oblong or lanceolate. Ultimate segments cuneate, the apex \pm truncate, entire or toothed, 1—4 m.M. broad. Texture subcoriaceous; surfaces naked; veins dichotomous. Sori terminal, roundish, 1—2 to a lobe.

Trop. Asia, Polynesia, Japan, China, Comore & Mascarene Islands, Madagascar.

(2) **O. biflora**, *C. Chr.*, Ind. Fil.; *Davallia biflora*, *Klf.*, Enum. Fil., 221.

Like *O. chinensis*, *J. Sm.*, but the fronds 2—3-pinnate, the apices of the lower pinnae caudate, the ultimate segments cuneate, broader; indusium suborbicular.

Luzon.

* (3) **O. retusa**, *J. Sm.*, Copel., Polypod. Philipp., 57; *Davallia retusa*, *Cav.*, Hk., Spec. Fil., I, tab. LII, A; *D. decipiens*, *Ces.*, Fel. Becc. Polin., 3 & 7; *Lindsaya retusa*, *Mett.*, Hk. Bk., Syn. Fil., 110; *L. cuneifolia*, *Pr.*, Hk., l. c., 219.

Rhizome short-creeping, strong. Stipes strong, erect, reddish like the rachises, to \pm 100 c.M. long. Fronds ovate-deltoid, 3-pinnatifid, to \pm 90 c.M. long. Pinnae stalked, the lower ones 20—60 c.M. long, 10—20 c.M. broad, deltoid. Pinnulae stalked, lanceolate-deltoid, pinnate or more commonly pinnatifid below, the apex pinnatifid. Segments cuneate, the lower ones to \pm $1\frac{1}{2}$ c.M. long, to \pm 1 c.M. broad, the base cuneate, the outer margin irregularly flabelliform, cut down towards the base into 5—6 cuneate lobes often different in length, 1—6 m.M. deep, 5—6 m.M. broad at the apex. Texture herbaceous, slightly fleshy; veins flabellate, several times forked. Sori terminal on the lobes, occupying the whole breadth.

Malaya; Melanesia.

29. LINDSAYA, *Dryander.*

Sori submarginal, solitary and terminal on the veins or uniting the apices of 2 or more veins or veinlets, roundish, oblong or linear. Indusium \pm of the same shape as the sorus, attached at its inner side, opening outwardly.

Rhizome short or creeping. Stipes not articulate to the rhizome. Fronds at least simply pinnate. Leaflets dimidiate, i. e. the half on the lower side almost or quite undeveloped, the »midrib" therefore running along the lower margin, which is always destitute of sori; veins free or anastomosing. Hk. Bk., Syn. Fil., tab. II, fig. 20, a; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 119, D, E, G, H; Christ, Farnkr. d. Erde, fig. 926—928.

Tropical and subtropical.

Arrangement of the sections.

- § 1. **ODONTOLOMA.** Sori solitary on the vein tips, but sometimes contiguous and with a tendency to become confluent. Christ, l. c.
- § 2. **EULINDSAYA.** Sori uniting the apices of 2 or more veins or veinlets. Veins free. Hk. Bk. & Diels, l. c.
- § 3. **SYNAPHEBIUM.** Sori as in § 2. Veins anastomosing.

§ 1. ODONTOLOMA.

- a.** Fronds not branched, i.e. pinnate or 2-pinnatifid.
- b.** Leaflets (*pinnae*) entire or crenate. (1) *L. repens.*
- b.b.** Leaflets (*pinnae*) coarsely toothed or lobed.
- c.** Fronds to $1\frac{1}{2}$ c.M. broad.
- ***. Leaflets toothed or lobed with 2—4 teeth or lobes. (2) *L. gracilis.*
- *.*.** Leaflets 2-fid. (3) *L. blanda.*
- c.c.** Fronds more than $1\frac{1}{2}$ c.M. broad.
- d.** Fronds sessile. (4) *L. delicatula.*
- d.d.** Fronds stalked, the stipes \pm 2—20 c.M. long.
- e.** Fronds rather short, under 35 c.M. long.
- f.** Leaflets shallowly lobed. (5a) *L. adiantoides.*
(5b) *L. Havicei.*

f.f. Leaflets rather deeply incised (6a) *L. apoensis*.

(6b) *L. cyathicola*.

e.e. Fronds rather long, more than 35 c.M. long.

(7) *L. Merilli*.

b.b.b. Leaflets deeply pinnatifid or even pinnate, the segments linear-cuneate.

*. Lobes truncate at the apex, which is slightly rounded and entire or somewhat erose. (8) *L. hymenophylloides*.

.. Lobes as in *, but the apex purveyed with 2—3 distinct teeth. (9) *L. capillacea*.

a.a. Fronds branched, i.e. 2-pinnate or 3-pinnatifid.

§. Leaflets (pinnulae) deeply lobed to a broad, leafy rachis, the segments broad. (10) *L. tenuifolia*.

§.§. Leaflets (pinnulae) cut down to a filiform rachis, the segments filiform. (11) *L. Blumeana*.

* (1) ***L. repens***, *Bedd.*, Handb. Ind. Ferns, 74; Ferns S.I., tab. CCIX; Christ, Farnkr. d. Erde, fig. 926; *Davallia repens*, *Desv.*, Hk. Bk., Syn. Fil., 93; *Odontoloma Boryanum*, *J. Sm.*, Bau. & Hk., Gen. Fil., tab. CXIV, B; — var. *hemiptera*; *D. hemiptera*, *Bory*, Bel., Voy. Bot., II, 73, tab. VII, fig. 2.

Rhizome wide-creeping, climbing, scaly. Fronds simply pinnate, 20—45 c.M. long, 2—4 c.M. broad. Leaflets 1—2 c.M. long, \pm half as broad, the lower line often slightly upcurved, the point acute or rounded, the upper edge straight or \pm rounded, subentire or crenato-lobate. Texture pellucido-herbaceous. Sori submarginal, large, \pm as broad as long, 1 to each crenation, often confluent in the subentire forms when the apices of the veins are joining. — Scarcely different from *L. pectinata*, *Bl.*

Var. hemiptera: Barren fronds dimorphous (as in *Teratophyllum*), the abnormal ones deeply pinnatifid on the upper and outer side, the lobes very fine, narrowly linear, simple or 1—2 \times forked.

Trop. Asia, Polynesia, Mauritius.

* (2) ***L. gracilis***, *Bl.*, Rac., Flor. Btz., I, 140.

Rhizome creeping, slender, very long, brown, glossy, sparingly provided with lanceolate, spreading scales. Stipes 1—5 c.M. apart, naked, polished, \pm 1—8 c.M. long. Fronds 15—30 c.M. long, 8—15 m.M. broad, linear-lanceo-

late, simply pinnate. Leaflets \pm triangular, erecto-patent, \pm 3 m.M. broad, close, sessile or short-stalked, the base \pm cuneate, the lower edge often curved upwards, the apex rounded or bluntish, the upper margin toothed or lobed with 2—4 entire, rounded or truncatate teeth or lobes. Texture membranaceous; veins simple or forked, often joining. Sori 1—2 to each lobe, the pairs often confluent when the veins are joining. (*Forma concinna* = *L. concinna*, *J. Sm.*).

Trop. Asia, New Caledonia.

(3) **L. blanda**, *Mett.*, *Linn.*, XXXVI, 80.

Rhizome long, slender, branched, clothed with fugacious scales. Stipes \pm 5 c.M. long, slender, stramineous like the rachis. Fronds \pm 15 c.M. long, \pm 15 m.M. broad, linear, acuminate, simply pinnate. Leaflets numerous, imbricate, erecto-patent, sessile, 7—10 m.M. long, cuneate or obovate-cuneate, bifid, the outer lobe commonly the broadest. Texture membranaceous; surfaces naked; veins simple or forked, free or joining at the apex. Indusium reniform or transversally oblong.

Java.

(4) **L. delicatula**, *Christ*, *C. Chr.*, *Ind. Fil.*; *Davallia delicatula*, *Christ*, *Ann. Bot.*, XV, 79, tab. XIV, fig. 11, a—b.

Rhizome slender, filiform, wide-creeping, densely clothed with yellow-brown scales. Fronds sessile, close, but not tufted, to 12 c.M. long and to 2½ c.M. broad, lanceolate, simply pinnate, narrowing gradually towards both ends. Leaflets \pm 20 on each side of the rachis, to \pm 1½ c.M. long and to \pm 1 c.M. broad, the lower edge straight or slightly curved upwards, the outer edge cut into \pm 5 rounded lobes \pm 4 m.M. deep, the sinuses very sharp. Texture thin, hyaline, flaccid; both surfaces naked; colour bright-light-green. Sori placed in the lobes on a short distance below the apices, blackish; indusium broad, subreniform, reaching to the edge.

Celebes.

* (5a) **L. adiantoides**, *Kuhn*, *Ann. Mus. Bot. L. B.*, IV, 278; *L. pulchella*, *Mett.*, *Linn.*, XXXVI, 81; *Davallia pulchella*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 93; *Hk.*, *Spec. Fil.*, I, tab. LIII, B; *Aspidium adiantoides*, *Bl.*, (oldest name).

Rhizome slender, wide-creeping. Stipes slender, wiry, erect, naked, 5—7½ c.M. long. Fronds 10—30 c.M. long, 1½—2½ c.M. broad, simply pinnate. Leaflets slightly stalked, 7—12 m.M. long, to \pm 5 m.M. broad, the

lower margin more or less curved upwards, the upper cut into 2—5 blunt, short lobes. Texture herbaceous; rachis and both surfaces naked. Sori small, placed in the lobes at a short distance from the edge; veins sometimes joining at the apex and then their sori becoming confluent.

Malaya; Fiji, Samoa.

(5b) **L. Havicei**, *Copel.*, Philipp. Journ., I, Suppl., 149.

Rhizome scandent, slender, fuscous, clothed with caducous, minute scales of the same colour. Stipes distant, 2—4 c.M. long, fuscous, naked. Fronds 12—15 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, acuminate, simply pinnate. Leaflets short-stalked, $\pm 1\frac{1}{4}$ — $1\frac{1}{2}$ c.M. long, ± 4 m.M. broad at the truncate base, falcate, narrowed gradually towards the acuminate apex, the upper edge finely lobed. Texture membranaceous; rachis fine, stramineous; veins free. Sori minute, terminal, more broad than deep.

Mindanao.

(6a) **L. apoensis**, *Copel.*, Polypod. Philipp., 64; Perk., Fragm. Philipp., tab. IV, A.

Rhizome scandent, ferrugineous, nearly naked. Stipes straight or flexuose, firm, 12—18 c.M. long, reddish-brown near the base, the rest green, nearly naked as is the rachis. Fronds ± 20 c.M. long, ± 6 c.M. broad, simply pinnate. Leaflets short-stalked, the largest ± 4 c.M. long, ± 7 m.M. broad, the lower edge curved, the upper one more or less deeply incised with the higher lobes mostly linear. Texture herbaceous; surfaces naked; veins free or confluent at the apex. Sori small, terminal on the lobes; indusium suborbicular.

Mindanao.

(6b) **L. cyathicola**, *Copel.*, Philipp. Journ., I. Suppl., 149, tab. V.

Rhizome scandent, purpureous, glossy, sparingly fulvo-paleaceous. Stipes distant, 4—5 c.M. long, slender, naked, glossy, purpureous below, greenish above like the rachis. Fronds simply pinnate, 15—20 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, narrowed towards both ends. Leaflets sometimes cuneate, the largest ± 9 m.M. long, ± 5 m.M. broad, obtuse, the acroscopical edge cut hardly $\frac{1}{2}$ -way down into 2—3 lobes. Texture membranaceous; surfaces naked; veins free except when united by the sori. Sori small, distant from the margin, more broad than deep, simple or confluent.

Luzon.

* (7) **L. Merrilli**, *Copel.*, Polypod. Philipp., 61.

Rhizome strong, scandent, reddish-brown, glossy, nearly naked or clothed with spreading, ferrugineous, lanceolate, entire scales. Stipes \pm 2—5 c.M. long, strong, slightly scaly, brownish or stramineous. Fronds 40—80 c.M. long, 5—4 c.M. broad, simply pinnate, narrowed towards both ends. Leaflets sessile, subsessile or short-stalked, the largest to 22 m.M. long, 5—9 m.M. broad near the rachis, the base cuneate, the lower edge straight or curved especially near the base, the upper edge inciso-crenate, the apex acute or bluntish; crenations \pm acute, the higher ones sometimes suddenly elongated and forked so as to form abnormal, linear, simple or bifid lobes. Texture membranaceous; rachis stramineous and like the surfaces naked; veins free, 1 to each crenation, the apices sometimes uniting. Sori submarginal, the tips of the lobes projecting beyond them, solitary in the crenations or occasionally confluent in pairs.

Philippines.

* (8) **L. hymenophylloides**, *Bl.*, Christ, Farnkr. d. Erde, 294; *L. Lapeyrousii*, *Bk.*, Hk. Bk., Syn. Fil., 106; *Davallia hymenophylloides*, *Bk.*, Hk. Bk., l. c., 93; Hk., Icon. Pl., XVII, tab. MDCXXIII; *D. Lapeyrousii*, *Hk.*, 2^d Cent. of Ferns, tab. LVI.

Rhizome slender, wide-creeping, scaly. Stipes short, to 5 c.M. long. Fronds simply pinnate, lanceolate, 10—40 c.M. long, 2—4 c.M. broad. Leaflets close, oblong, \pm 5—5 m.M. broad, the lower edge entire, straight or slightly curved, the upper margin deeply pinnatifid nearly to the lower edge into 5—8 narrow, linear-cuneate, truncate lobes. Texture thinly herbaceous; veins 1 to each lobe, simple or forked; surfaces naked. Sori transversally oblong, subterminal on the lobes, solitary, or rarely double and confluent.

Malaya; Polynesia.

* (9) **L. capillacea**, *Christ*, Bull. Herb. Boiss., VI, 144, tab. IV, fig. 7, a—c; (? *L. pectinata*, *Bl.*, Christ, l. c., 2^e Série, VI, 1006).

Rhizome creeping, filiform, intertwined. Stipes filiform, brownish, naked. Fronds \pm 10 c.M. long, \pm 10 m.M. broad, linear-lanceolate, acuminate, simply pinnate. Leaflets alternate, \pm 5 m.M. long, triangular-oblong, the lower edge entire, the upper edge cut nearly or quite down into 3—4 segments, which are \pm spreading, linear-cuneate, the lower ones forked, the apices of all truncate, 2—5-dentate. Texture thin, weak, flaccid; rachis green, naked; surfaces naked; 1 central vein to each lobe. Sori

small, subterminal and solitary on the segments, flanked on each side by the acute teeth.

Luzon.

* (10) **L. tenuifolia**, *Bl.*, Enum., 219; *L. triquetra*, *Mett.*, Christ, Farnkr. d. Erde, 294; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 119, H; *Davallia triquetra*, *Bk.*, Hk. Bk., Syn. Fil., 93; *D. Blumeana*, *Ces.*, Fel. Becc. Polin., 3.

Rhizome creeping, fibrillose. Stipes strong, erect, 10—15 c.M. long. Fronds 2-pinnate, with a terminal branch and 2—7 pairs of spreading or erecto-patent lateral ones, which are 5—10 c.M. long, to 5½ c.M. broad, linear or lanceolate. Leaflets oblong, ± ½ × as broad as long, cut down on the upper side into linear or oblong, simple or 2-fid lobes to a nearly straight, leafy rachis. Texture pellucido-herbaceous; surfaces naked. Sori small, terminal on the segments; indusium suborbicular.

Malaya; Polynesia.

* (11) **L. Blumeana**, *Kuhn*, Ann. Mus. Bot. L. B., IV, 277; *Davallia Blumeana*, *Hk.*, Hk. Bk., Syn. Fil., 93; *Hk.*, Spec. Fil., I, tab. LIV, A.

Rhizome creeping. Stipes strong, erect, 10—50 c.M. long. Fronds 20—50 c.M. long, branched. Branches rigid, erecto-patent, 7½—15 c.M. long, to ± 1½ c.M. broad, pinnate, acuminate. Leaflets cut down on the upper side to a filiform, decurved rachis into very slender, filiform, simple or forked segments. Texture thinly herbaceous. Sori small, subterminal on the dilated, entire or crenated apices of the segments.

Sumatra, Celebes, Philippines, New Guinea.

§ 2. EULINDSAYA.

1. Fronds unbranched, i.e. pinnate or 2-pinnatifid. (*)

2. Fronds to 1½ c.M. broad.

a. Leaflets (pinnae) entire, crenate or slightly lobed.

b. Leaflets 4—5 m.M. long, ± 3 m.M. broad.

(12) *L. gracillima*.

b.b. Leaflets 6—7 m.M. long, ± 5 m.M. broad. (13) *L. concinna*.

a.a. Leaflets (pinnae) deeply lobed.

(14) *L. Loheriana*.

(*) See also *L. orbiculata*, *Mett.* & *lancea*, *Bedd.*

- 2.2.** Fronds $1\frac{1}{2}$ c.M. broad or broader when fully developed ⁽¹⁾.
- 3.** Fronds to 15 c.M. long.
- §. Indusium continuous or interrupted, not undulate.
- *. Fronds small, 3—6 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad. (15) *L. montana*.
- *.*. Fronds larger, 10—15 c.M. long, ± 2 —5 c.M. broad.
- ×. Leaflets blunt. (16a) *L. humilis*.
- ×.×. Leaflets acute. (16b) *L. falcata*.
- §.§. Indusium continuous, undulate. (17) *L. crispa*.
- 3.3.** Fully developed fronds more than 15 c.M. long.
- 4.** Rhizome short-creeping.
- *. Texture membranaceous. (18a) *L. cultrata*.
- *.*. Texture coriaceous. (18b) *L. crassipes*.
- 4.4.** Rhizome firm, scandent.
- 5.** Leaflets crenate; sori interrupted. (19) *L. pectinata*.
- 5.5.** Leaflets entire; sori continuous. (20) *L. scandens*.
- 1.1.** Fronds branched, i.e. 2-pinnate or 3-pinnatifid ⁽²⁾.
- 2'.** Veins flabellate.
- *. Leaflets (pinnulae) 5—8 m.M. broad. (21) *L. orbiculata*.
- *.*. Leaflets (pinnulae) 10—15 m.M. broad. (22) *L. gomphophylla*.
- 2.2'.** Veins not flabellate.
- 3.** Branches of the fronds mostly under 2 c.M. broad.
- §. Leaflets (pinnulae) entire, crenate or toothed.
- ‡. Sori roundish or oblong, terminal on the leaflets. (23) *L. tricrenata*.
- ‡.‡. Sori in a continuous submarginal line. (24) *L. borneensis*.
- §.§. Leaflets (pinnulae) bluntly, not deeply lobed. (25) *L. rigida*.
- 3.3'.** Branches mostly 2—4 c.M. broad; leaflets (pinnulae) not entire.
- 4.** Leaflets dimorphous. (26) *L. impressa*.
- 4.4'.** Leaflets uniform.
- 5.** Fronds rather small, to ± 30 c.M. long, the branches ± 3 —4-jugate, to 15 c.M. long. (27) *L. natunae*.
- 5.5'.** Fronds rather large, to ± 80 c.M. long or longer, the branches ± 5 —6-jugate, to 40 c.M. long. (28) *L. longissima*.
- 3.3.3'.** Branches mostly 4 c.M. broad or broader; leaflets (pinnulae) entire. (29) *L. lancea*.

⁽¹⁾ See also *L. conciuna*, J. Sm.

⁽²⁾ See also *L. pectinata*, Bl. & *scandens*, Hk.

* (12) **L. gracillima**, *Copel.*, Polypod. Philipp., 61.

Rhizome short. Stipes densely tufted, erect, brown, nearly naked, \pm 5 c.M. long. Fronds simply pinnate, linear, \pm 52 c.M. long, \pm 1 c.M. broad. Leaflets alternate, petiolate, deltoid or rhomboidal, \pm 4 m.M. long, \pm 5 m.M. broad, the lower ones \pm reduced, the lower edge straight, upper edge rounded, entire, or when sterile crenate. Texture herbaceous; surfaces naked; veins flabellate. Sori continuous or subinterrupted.

Luzon.

* (15) **L. concinna**, *J. Sm.*, Hk. Bk., Syn. Fil., 105; Hk., Spec. Fil., I, tab. LXI, B; *L. Lobbiana*, *Hk.*, l. c., tab. LXII, C; (? *L. gracilis*, *Bl.*, C. Chr., Ind. Fil.).

Rhizome short-creeping. Stipes wiry, flexuose or straight, erect, 5—10 c.M. long. Fronds 15—50 c.M. long, \pm 12—15 m.M. or rarely 2 c.M. broad, simply pinnate. Leaflets \pm 5 m.M. broad, oblong; the lower edge straight or curved commonly upwards, the upper and outer margin rounded, entire or slightly lobed with shallow, broad, truncate lobes. Texture pellucido-herbaceous. Sori in a continuous line except when interrupted by the lobes.

Malaya.

(14) **L. Loheriana**, *Christ*, Bull. Herb. Boiss., VI, 144, tab. IV, fig. 6, a—b.

Rhizome short-creeping, firm. Stipes fasciculate or approximate, terete, slender, firm, erect, \pm 2 c.M. long, purple at the base. Fronds \pm 12 c.M. long, \pm 1 c.M. broad, linear-lanceolate, acuminate, simply pinnate. Leaflets alternate, triangular-cuneate, on stalks \pm 2 m.M. long, the lower edge entire, the upper one deeply lobed with 2—5 lobes. Lobes cuneate, truncate at the apex. Texture firm; rachis stramineous; veins few, conspicuous, forked in the lobes. Sori subterminal on the lobes, single or double; indusium inflated.

Luzon.

(15) **L. montana**, *Copel.*, Polypod. Philipp., 62; *L. Copelandi*, *C. Chr.*, Ind. Fil.; (? *L. orbiculata*, *Mett.*, p. p., *Copel.*, Philipp. Journ., I, Suppl., 148).

Rhizome short, creeping, the scales ferruginous. Stipes tufted, 1—3 c.M. long, slender, flexuose, reddish-brown below, stramineous upwards, nearly naked. Fronds 5—6 c.M. long, \pm 1½ c.M. broad, simply pinnate. Leaflets stalked, \pm 9 m.M. long, \pm 6 m.M. broad, semiorbicular, the lower edge deflexo-

arcuate, the upper edge rounded, entire or incised, the higher leaflets erecto-patent, the lower much deflexed. Texture membranaceous to coriaceous; rachis nearly naked; veins flabellate. Sori often continuous.

Philippines.

* (16a) **L. humilis**, *Kuhn*, Ann. Mus. Bot. L. B., IV, 278; *L. adiantoides*, *J. Sm.*, Hk. Bk., Syn. Fil., 104; Hk., Spec. Fil., I, tab. LXI, C.

Stipes nearly tufted, often blackish, polished, wiry, $2\frac{1}{2}$ —5 c.M. long. Fronds 10—15 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, simply pinnate. Leaflets to $1\frac{1}{2}$ c.M. long, to 7 m.M. broad, the upper ones imbricated, the lower edge straight or slightly curved, the upper rounded and broadly lobed to $\pm \frac{1}{3}$ of the way down. Texture pellucido-herbaceous. Sori subterminal on the lobes.

Java, Celebes, Philippines.

(16b) **L. falcata**, *Dry.*, Trans. Linn. Soc., Bot., III, 41, tab. VII, fig. 2; *L. trapeziformis*, *Dry.*, Hk. Bk., Syn. Fil., 107, p. p.; *L. Leprieurii*, *Hk.*, Spec. Fil., I, 298, tab. LXII, D.

Rhizome filiform, creeping. Stipes dark-brown, tetragonal, almost black and glossy. Fronds oblong, at utmost $22\frac{1}{2}$ c.M. long inclus. the stipe, simply pinnate. Leaflets half-ovate, $\pm 2\frac{1}{2}$ c.M. or more long, acuminate, deflexo-falcate, truncate at the base, which is parallel to the obscurely winged rachis, the edge entire, the apex rather acuminate, the terminal pinna formed of 3 combined pinnae with a main nerve running through the centre. Texture membranaceous; rachis like the stipe. Sori continuous, distant from the margin, indusium narrow.

Negros; Trop. America.

(17) **L. crispa**, *Bk.*, Journ. of Bot., XVII, 59; Hk., Icon. Pl., XVII, tab. MDCXXVII.

Rhizome short-creeping. Stipes wiry, naked, castaneous, $2\frac{1}{2}$ —25 c.M. long. Fronds lanceolate, simply pinnate, ± 7 —15 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad. Leaflets not crowded, sessile, deltoid, truncate on the inner and spreading or erecto-patent on the lower side, the upper edge irregularly incisocrenate. Texture membranaceous; surfaces naked; colour bright-green; veins distinct, \pm flabellate. Indusium undulate, with 2 distinct, nearly equal, cartilaginous valves.

Borneo.

* (18a) **L. cultrata**, Sw., Rac., Flor. Btz., I, 141; Hk., Fil. Exot., tab. LXVII; Bedd., Ferns S.I., tab. XXIII; *L. apiculata*, Kze, Farnkr., I, tab. LXXXV, fig. 2; — var. *varia*, Copel., Polypod. Philipp., 62.

Rhizome creeping, more or less densely clothed with linear, brown, setiform scales. Stipes approximate, 7–15 c.M. long, naked, polished, often reddish, the base clothed with setiform scales. Fronds linear-lanceolate, 15–50 c.M. long, \pm 2–4 c.M. broad, simply pinnate. Leaflets to 1 c.M. broad, not imbricate, triangular, the lower margin straight or slightly curved (usually upwards), the upper edge crenate, so that the continuity of the line of fructification is broken, sometimes nearly entire, lower leaflets short-stalked. Texture firm-membranaceous. Sori 1–5 m.M. broad, in an intramarginal line, only broken by the crenations, slightly concave.

Var. varia: Fronds shorter and narrower, membranaceous.

Trop. Asia, North. Australia, East. Africa; — var. *varia*: *Philippines*.

(18b) **L. crassipes**, Rst., Fed., Rep., V, 56.

Rhizome short-creeping, branched, the scales minute, linear-lanceolate. Stipes dense, flexuose, black-brown, polished, slender, naked, to nearly 10 c.M. long. Fronds linear-lanceolate, to 18 c.M. long, to 2 c.M. broad at the centre, simply pinnate. Leaflets alternate, subremote, trapezio-ovate, on stalks 1–1½ m.M. long, the midial ones nearly 1 c.M. long, ½ c.M. broad, narrowed at the apex, with the upper edge 2-crenate, the lower ones smaller, rounded at the apex, 5–5-crenate, the higher ones gradually growing smaller, entire, narrowed at the apex. Texture coriaceous; surfaces naked, the upper one dark-green, the under one somewhat paler; rachis black-brown, flavido-marginate above; veins slightly visible, dichotomous. Sori interrupted by the crenations.

New Guinea.

* (19) **L. pectinata**, Bl., Hk. Bk., Syn. Fil.; *L. oblongifolia*, Reinw., Hk., Spec. Fil., I, tab. LXI, D; *L. repens*, Kze, Rac., Flor. Btz., I, 142, p. p.

Rhizome long, creeping or scandent, densely clothed with short, lanceolate scales. Fronds sessile or short-stalked, ½–4 c.M. apart, lanceolate-ligulate, narrowed towards both ends, 20–60 c.M. long, 2–5 c.M. broad, simply pinnate, rarely 2-pinnate. Leaflets 7–10 c.M. broad, the lower ones reduced, the central ones linear-oblong, closely placed, the lower edge nearly straight, the upper margin rounded, \pm crenate, the apex bluntish to bluntly

rounded. Texture pellucido-herbaceous. Sori in an interrupted submarginal line along the upper edge and the apex.

Tropical Asia.

* (20) **L. scandens**, *Hk.*, Hk. Bk., Syn. Fil., 106; Hk., Spec. Fil., I, tab. LXIII, B; *L. repens*, *Kze*, Rac., Flor. Btz., I, 142, p. p.

Like *L. pectinata*, *Bl.*, but the lower edge of the leaflets slightly decurved the upper and outer edges entire, broadly rounded. Sori in a continuous submarginal line.

Malaya.

* (21) **L. orbiculata**, *Mett.*, Bedd., Handb. Ind. Ferns, 75; *L. flabellulata*, *Dry.*, Hk. Bk., Syn. Fil., 107; *Dry.*, Trans. Linn. Soc., Bot., III, tab. VIII, fig. 2; Bedd., Ferns S.I., tab. CCXVI; *Adiantum orbiculatum*, *Lam.*, (oldest name); — var. *polymorpha*; *L. polymorpha*, *Wall.*, Hk. & Grev., Ic. Fil., tab. LXXV; — var. *gigantea*, Hk., Spec. Fil., I, tab. LXIII; — var. *tenera*, Bedd., Ferns S.I., tab. XXIV; *L. tenera*, *Dry.*, l. c., tab. X; *L. striata*, *Bl.*, Enum., 220.

Rhizome short-creeping, the scales brown, dense, small, lanceolate-subulate. Stipes ± approximate, 5—30 c.M. long, erect, naked, glossy, often black or brown, the fertile fronds on the longest stipes. Fronds 10—25 c.M. long, the fertile ones longer than the barren, simply pinnate, or 2-pinnate with a ± long, terminal branch and 1 to several, spreading or suberecto-patent branches on each side various in length. Leaflets various in shape, roundish, flabellate, oblong, sub-quadrangular or trapeziform, 7—15 m.M. long, 5—8 m.M. broad, the lower line nearly straight or decurved, the upper and outer edges bluntish or rounded, entire, crenate or lobed, the lobes entire, blunt or 2—5-dentate in the barren, ± truncate in the fertile leaflets, which are often the smallest. Texture ± pellucid, rather rigid; rachis and surfaces naked; veins flabellate, dichotomous. Sori in a continuous line except when interrupted by the lobes. — Variable.

Var. polymorpha: Some of the branches reduced to large, irregularly incised or subdivided leaflets.

Var. gigantea: Some of the leaflets of the central rachis $2\frac{1}{2}$ —5 c.M. long, lengthened to an acute point.

Var. tenera: Texture thin.

Trop. Asia & Australia.

* (22) **L. gomphophylla**, *Bk.*, Ann. of Bot., V, 204.

Rhizome creeping, the scales lanceolate, acute or acuminate. Stipes naked, castaneous. Fronds 2-pinnate, \pm 15 c.M. long, branched in the lower half, the lowest branches much the longest. Leaflets orbicular-cuneate, \pm 1—1½ c.M. broad, the base cuneate, the upper and outer edge deeply crenate or rather deeply lobed and crenate, the lobes cuneate-flabellate or cuneate-oblong. Texture thin, membranaceous or more rigid; surfaces naked; veins flabellate. Sori interrupted by the lobes and crenations; indusium narrow.

Borneo.

(23) **L. tricrenata**, *Bk.*, Journ. of Bot., XXVIII, 106.

Stipes 10—13 c.M. long, like the rachises slender, naked, castaneous. Fronds deltoid, 2-pinnate. Branches few, 15—20 c.M. long, 8—9 m.M. broad. Leaflets sessile, cuneate, 4—4½ m.M. broad, crenate on the outer edge. Texture moderately firm; surfaces naked; veins \pm 3 to a pinnula, forked. Sori roundish or oblong, terminal on the pinnulae; indusium narrow, firm.

New Guinea.

* (25) **L. borneensis**, *Hk.*, Hk. Bk., Syn. Fil., 107.

Rhizome short-creeping, paleaceous. Stipes strong, erect, polished, 20—35 c.M. long. Fronds 2-pinnate, 30—45 c.M. long, 20—30 c.M. broad, with a long, terminal branch and 6—10 pairs of lateral ones, which are 15—25 c.M. long. Leaflets 7—10 m.M. long, 3—5 m.M. broad, closely placed, often slightly imbricate, the upper edge entire, the apex broadly rounded, entire or finely toothed. Texture pellucido-herbaceous to subcoriaceous; veins prominent. Sori in a continuous, submarginal line.

Borneo, Sumatra, Malacca.

(25) **L. rigida**, *J. Sm.*, Hk. Bk., Syn. Fil., 108; Hk., Spec. Fil., 1, tab. LXIII, A.; Bedd., Ferns Br.I., II, tab. CLXVI.

Rhizome wide-creeping. Stipes 10—15 c.M. long, rigid, erect, prickly towards the base. Fronds 2-pinnate, with a long terminal branch and 1—4 pairs of flexuose, lateral ones 10—20 c.M. long. Leaflets 6—10 m.M. long, 4—5 m.M. broad, the lower edge often falcate, the upper 3—4 times bluntly, not deeply lobed, placed close together, not imbricate. Texture very thick and coriaceous; veins prominent. Sori subterminal on the lobes. — The colour of the nature frond is sepia-brown and the pale veins stand out from the ground-work in relief (Bk).

Malacca, Batjan, Celebes, New Guinea, Philippines,

* (26) **L. impressa**, *Christ*, Ann. Btz., XX, 132.

Rhizome creeping, brown, the scales subulate, brown. Stipes approximate, 5—20 c.M. long castaneous, firm, naked. Fronds 1—2 × pinnate, 20—40 c.M. long. Branches 5—5, sessile, erecto-patent, ± 15 c.M. long, ± 5 c.M. broad, acuminate. Leaflets approximate, spreading, dimorphous, the barren ones ± 15 m.M. long, ± 6 m.M. broad, sessile, ovate-oblong, with the apex obtuse, narrowed, the upper and outer edge toothed or nearly lobed, the lobules ± serrate, the fertile ones smaller, acute, triangular, entire. Texture subcoriaceous, fragile; rachis and surfaces naked; veins blackish, conspicuous, deeply impressed, forked, some of them sometimes anastomosing. Sori narrow, linear, continuous, occupying the upper edge of the fertile leaflets.

Borneo.

* (27) **L. natunae**, *Bk.*, Kew Bull., 1896, 40.

Rhizome short-creeping. Stipes approximate, elongate, naked, pale-brown, canaliculate. Fronds deltoid, ± 50 c.M. long, 2-pinnate. Branches ± 5—4-jugate, 8—12½ c.M. long, 2—2½ c.M. broad, lanceolate, subsessile. Leaflets multijugate, close, the central ones the largest, ± 5 m.M. broad, sessile, the lower edge straight, the upper one lobed. Texture membranaceous; rachis naked, pale-brown; surfaces naked; veins simple, distant, conspicuous, the lower ones forked. Sori oblong, terminal on the lobes. — A plant found in Lingga by Mr. Teysmann and which may be a form of this species has the leaflets nearly close, the veins forked, the sori continuous except when interrupted by some shallow, broad, lobe-like crenations. Branches of the fronds 1—2-jugate.

Natuna Islands, ? Lingga Islands.

* (28) **L. longissima**, *Christ*, Ann. Btz., XX, 151.

Stipes castaneous, slender, nearly glossy, naked, ± 40 c.M. long. Fronds to 80 c.M. long or longer, 2-pinnate. Branches distant, alternate, sessile, acuminate, arcuato-nutant, ± 5—6 below the terminal one, ± 60 c.M. long, 2—2½ c.M. broad, pectinato-pinnate. Leaflets very numerous, close, sessile, rhomboidal, obtuse, ± 5 m.M. broad, the base cuneate, the lower edge ± decurved, the upper and outer edges broadly lobed, the lobes shallow, obtuse, crenate. Texture herbaceous, opaque; surfaces naked; veins not conspicuous, forked, some of them sometimes anastomosing. Sori few, submarginal, terminal on the outer lobes, 2—5 to a pinnula; indusium greyish.

Borneo.

* (29) **L. lancea**, *Bedd.*, Handb. Ind. Ferns, 75; *L. trapeziformis*, *Dry.*, Hk. Bk., Syn. Fil., 107, p. p.; Trans. Linn. Soc., Bot., III, 42, tab. 9; Bau. & Hk., Gen. Fil., tab. LXIII, A; *Adiantum lanceum*, *L.*, (oldest name); — var. *caudata*; *L. caudata*, *Hk.*, Spec. Fil., I, 215; *Bedd.*, Ferns S.I., tab. CCXVII; *Hk.*; Icon. Pl., X, tab. CMLVIII.

Rhizome short-creeping, more or less provided with brown, lanceolate scales. Stipes strong, erect, 20—40 c.M. long, dark-coloured. Fronds 20—40 c.M. long, simply pinnate or more commonly 2-pinnate with a long terminal branch and 1—4 similar, erecto-patent, lateral ones on each side. Leaflets 2—3½ c.M. long, 7—15 m.M. broad, rarely smaller, oblong, the lower line ± de-curved or nearly straight, the upper and outer edges entire, rounded. Texture pellucido-herbaceous. Sori in a continuous line round the upper margin.

Var. caudata: A large form with more branches than usually.

Malaya; Ceylon, Trop. America.

§ 3. SYNAPHLEBIUM.

a. *Sori continuous.*

(30) *L. azurea.*

a.a. *Sori interrupted.*

b. *Leaflets entire, toothed or shallowly incised enough to interrupt the line of the sori.*

×. *Rhizome short-creeping, sparingly scaly.* (31a) *L. decomposita.*

×.×. *Rhizome wide-creeping, naked.* (31b) *L. Werneri.*

b.b. *Leaflets lobed 1/3—1/2-way down.*

c. *Leaflets dimidiate throughout, under 2 c.M. long, lobed ± 1/3-way down.*

d. *Lobes rounded.* (32) *L. davallioides.*

d.d. *Lobes truncate.* (33) *L. ambigens.*

c.c. *Leaflets dimidiate in the lower half only, more than 2 c.M. long, lobed ± 1/2-way down.* (34) *L. Sarasinorum.*

(30) **L. azurea**, *Christ*, Ann. Btz., XV, 101, tab. XIV, fig. 12, a—b.

Stipes rigid, dark-reddish, ± 30 c.M. long. Fronds 2-pinnate, with 2—3 lateral branches on each side below the terminal one. Leaflets ± 6 m.M.

long, 4 \pm m.M. broad, the lower edge straight or curved, the upper and outer ones rounded, entire. Texture subcoriaceous; surfaces naked, the upper one with a distinct, bluish reflexion of light; veins inconspicuous. Sori continuous, occupying the upper and outer margin.

Celebes.

* (31a) **L. decomposita**, *Willd.*, Spec., V, 425; *L. lobata*, *Poir.*, Rac., Flor. Btz., I, 142; *L. recurvata*, *Wall.*, Hk., Spec. Fil., I, 222; *L. propinqua*, *Hk.*, l.c., 223, tab. LXVI, B.; *L. nitens*, *Bl.*, Enum., 217; *Schizoloma lobata*, *Bedd.*, Handb. Ind. Ferns, 77; *S. recurvata*, *Bedd.*, Ferns S.I., tab. XXVII; *Synaphlebium recurvatum*, *J. Sm.* (not *Bl.*), Bau & Hk., Gen. Fil., tab. CI.

Rhizome short-creeping, the scales scattered, linear-lanceolate, brown. Stipes approximate, firm, erect, naked, 5—25 c.M. long. Fronds simply pinnate or 2-pinnate with a long unbranched apex and 1—6 pairs of lateral branches 7—25 c.M. long. Leaflets 1—2 c.M. long, \pm half as broad, the lower often strongly decurved principally at the base, the upper and outer margin rounded, entire or shallowly incised enough to interrupt the sori, toothed or double-toothed. Texture thinly herbaceous, pellucid. Sori submarginal.

Trop. Asia & Australia, Polynesia.

(31b). **L. Wernerii**, *Rst.*, Fed., Rep., V, 37.

Rhizome wide-creeping, ferruginous, glossy, rather naked. Stipes distant, subquadrangular, firm, 4—10 c.M. long, stramineous. Fronds linear-lanceolate, to 40 c.M. long, \pm 6 $\frac{1}{2}$ c.M. broad, simply pinnate. Leaflets 15—25-jugate, to 3 $\frac{1}{2}$ c.M. long, \pm 1 $\frac{1}{4}$ c.M. broad, short-stalked, the lower ones slightly decurved, the rest rather spreading, the inner edge parallel to the rachis, the upper and outer edges irregularly crenate, the apex bluntly rounded or obliquely truncate. Texture membranaceous; surfaces naked. Sori linear, interrupted, occupying the crenations, rather distant from the margin; indusium very narrow, chartaceous, naked.

New Guinea.

* (32) **L. davallioides**, *Bl.*, Rac., Flor. Btz., I, 143; *Hk.*, Spec. Fil., I, tab. LXVIII, A; *Davallia Kunzeana*, *Hk.*, l.c., 177; *Schizoloma davallioides*, *Moore*, *Bedd.*, Ferns Br.I., I, tab. CXLI.

Rhizome short-creeping, the extremity clothed with short, lanceolate scales. Stipes rather approximate, 6—20 c.M. long, firm, erect. Fronds 2-pinnate, with a long terminal branch and 1—5 pairs of spreading, curved, lateral branches

4—20 c.M. long. Leaflets 1—1½ c.M. long, 5—7½ m.M. broad, the lower margin straight or slightly curved, the upper with 5—6 regular, oblong, rounded lobes reaching $\pm \frac{1}{3}$ -way down, placed close together, but not imbricate. Texture pellucido-herbaceous. Sori terminal on the lobes. — High alpine forms not rarely unbranched.

Malaya.

(33) **L. ambigenus**, *Ces.*, Fel. Becc. Polin., 3 & 7; (? *L. decomposita*, *Willd.*, C. Chr., Ind. Fil.).

Like *L. davallioides*, *Bl.*, but the lobes of the leaflets truncate.

New Guinea.

(34) **L. Sarasinorum**, *Christ*, Ann. Bot., XV, 101, tab. XIV, fig. 13, a—b.

Rhizome short-creeping. Stipes clustered, reddish-brown, ± 12 c.M. long, provided like the rachis with scattered, brown scales. Fronds ± 8 —12 c.M. long, simply pinnate, or 2-pinnate with a pair of lateral branches below the central one. Leaflets dimidiate near the base, becoming \pm equilateral towards the apex, ± 3 c.M. long, narrowly oval or rhomboidal, strongly bent outwards, the upper and outer edges rounded with 4—5 lobes, which are broad, obtuse, often lobed again; primary lobes $\pm \frac{1}{2}$ as deep as the breadth of the leaflets. Texture herbaceous; surfaces naked. Sori thick, brown, terminal on the lobes.

Celebes, Philippines.

30. SCHIZOLOMA, *Gaudichaud.*

Sori linear or linear-oblong, uniting the apices of many veins. Indusium of the same shape, continuous, attached at the inner side of the sorus, opening outwardly.

Rhizome mostly short. Fronds simple or generally pinnate; leaflets equal-sided or unequal-sided, but *not* dimidiate; veins free or anastomosing. Diels, in Engl. & Prantl, Nat. Pfl Fam., I⁴, fig. 118.

Tropical and extra-tropical.

Arrangement of the sections.

§ 1. **ISOLOMA.** Veins free. Fronds always pinnate.

§ 2. **EUSCHIZOLOMA.** Veins anastomising. Fronds sometimes simple.

§ 1. **ISOLOMA.**

a. Pinnae roundish or oblong, 5—12 $\frac{1}{2}$ m.M. long.

b. Fronds to 20 c.M. long. Pinnae 4—6 m.M. broad.

c. Pinnae \pm equal-sided, cordate or truncate at the base.

(1a) *S. jamesonioides*.

(1b) *S. angustum*.

c.c. Pinnae unequal-sided, the upper base truncate, the lower one cuneate or rounded.

(2) *S. ovatum*

b.b. Fully developed fronds to 30 c.M. long. Pinnae 7—10 m.M. broad.

(3a) *S. induratum*.

(3b) *S. fuligineum*.

a.a. Pinnae linear-oblong or linear, 1 $\frac{1}{2}$ —10 c.M. long.

S. Pinnae spreading, 1 $\frac{1}{2}$ —2 $\frac{1}{2}$ c.M. long.

‡. Sori not reaching the apex of the pinnae. (4) *S. pellaeiforme*.

‡‡. Sori continuous along the apex of the pinnae. (5) *S. divergens*.

S.S. Pinnae erecto-patent, 3 $\frac{1}{2}$ —10 c.M. long. (6) *S. Walkerac.*

* (1a) ***S. jamesonioides***, *Copel.*, Philipp. Journ., I, Suppl., 252; *Lindsaya Jamesonioides*, *Bk.*, Journ. of Bot., XVII, 39; *Hk.*, Icon. Pl., XVII, tab. MDCXXVI; *C. Chr.*, Ind. Fil.

Rhizome short-creeping, the scales minute subulate, rigid, nearly black. Stipes wiry, nearly black, 2 $\frac{1}{2}$ —15 c.M. long. Fronds linear, pinnate, 7—20 c.M. long, 1—2 c.M. broad, composed of sessile, contiguous, roundish or oblong, alternate, spreading, deciduous, entire leaflets 4—6 m.M. broad, with the apex rounded, the base rounded, broadly cuneate or truncate, those of the upper $\frac{1}{3}$ of the frond growing gradually smaller. Rachis wiry, dark-brown; texture coriaceous; surfaces naked, the upper one wrinkled; veins hidden. Sori continuous all round the edge or occupying the outer half of the pinnae; indusium broad, chartaceous, persistent.

Borneo, Celebes.

(1b) **S. angustum**, *Copel.*, Philipp. Journ., I, Suppl., 252, tab. I, B.

Rhizome short-creeping, clothed with minute, narrow, castaneous scales. Stipes approximate, 5—8 c.M. long, black-purpureous, sometimes flexuose, sparingly clothed towards the base with scattered hairs of the same colour, the rest naked, glossy, triangular like the rachis. Fronds 10—18 c.M. long, ± 1 c.M. broad, simply pinnate. Pinnae sessile, spreading, oblong, slightly convex, ± 5 m.M. long, ± 4 m.M. broad, at length deciduous, the apex rounded, the margin entire, the base truncate or subcordate, often overlapping the rachis, lower pinnae somewhat distant, deflexed. Texture coriaceous; veins hidden, flabellate. Sori broad, continuous. — Near *S. ovatum*, *Copel.*

Philippines.

* (2) **S. ovatum**, *Copel.*, Philipp. Journ., I, Suppl., 252; *S. Guerinianum*, *Gaud.*, var., *C. Chr.*, Ind. Fil.; *Lindsaya ovata*, *J. Sm.*, Hk. Bk., Syn. Fil., 105; Hk., Spec. Ftl., I, tab. LXIV, A.

Rhizome short-creeping. Stipes 5—10 c.M. long, wiry, flexuose, black. Fronds 10—15 c.M. long, ± 2 c.M. broad, simply pinnate. Pinnae ± 5 m.M. broad, not imbricate, sessile, the lower ones with their own breadth between them, horizontally oblong, the apex broadly rounded, the lower base rounded or cuneate, the upper one truncate, slightly auricled. Texture \pm coriaceous. Sori in a continuous marginal line.

Philippines, Borneo, Celebes.

(3a) **S. induratum**, *C. Chr.*, Ind. Fil.; *Lindsaya indurata*, *Bk.*, Journ. of Bot., XXVI, 324.

Rhizome creeping, the scales, minute, spreading, linear, brownish-black. Stipes wiry, naked, castaneous, 20—25 c.M. long. Fronds lanceolate, pinnate, ± 30 c.M. long, 2—2½ c.M. broad. Pinnae oblong, obtuse, sessile, sub-equilateral, ± 8 —9 m.M. broad, the edges entire, rather reflexed. Texture firm; surfaces naked; costae distinct; veins immersed, obscure. Sori continuous all round the sides and the apex of the pinnae; indusium narrow.

Borneo.

(3b) **S. fuliginum**, *Copel.*, Philipp. Journ., I, Suppl., 252, tab. I, A.

Rhizome short, creeping, clothed with minute, filiform, castaneous scales. Stipes caespitose, 7—10 c.M. long, scaly at the base, the rest glossy, naked, castaneous like the rachis. Fronds 20—30 c.M. long, 2½—5 c.M. broad, narrowed gradually towards the apex. Pinnae subsessile, articulate to the

rachis, the lowest deflexed, the others spreading, obtuse, entire, $\pm 1\frac{1}{4}$ — $1\frac{1}{2}$ c.M. long, ± 7 — 10 m.M. broad, the upper base truncate, auriculate, the auricle sometimes acute. Texture coriaceous; surfaces naked, the upper one black-castaneous, the under one fuliginous; veins free. Sori continuous. — About one pinna in ten shows an anastomosis of veinlets.

Mindanao.

(4) **S. pellaeiforme**, *C. Chr.*, Ind. Fil.; *Lindsaya pellaeiformis*, *Christ*, Ann. Btz., XV, 102, tab. XIV, fig. 14, a—b.

Rhizome short, strong, slightly creeping, clothed with black-brown, spreading, subulate scales. Stipes approximate, subtufted, blackish-red, polished, naked or punctate, 10—20 c.M. long. Fronds ligulate-linear, ± 30 c.M. long, 3—4 c.M. broad, simply pinnate. Terminal pinna oblong, slightly lobed. Lateral pinnae inequilateral, numerous (± 40 on each side), close, spreading, sessile, the lower ones short-stalked, ovate-rhomboidal, the base truncate & \pm auricled above, cuneate & commonly slightly rounded below, the apex blunt, the margin entire or finely crenate. Texture papyraceous, pellucid, dark-green; veins erecto-patent, simple or forked, the tips thickened. Sori occupying the upper and lower margin, forming a continuous line on each side of the costa.

Celebes.

* (5) **S. divergens**, *Kuhn* (not *Diels*), *Copel.*, Polypod. Philipp., 58; *Vittaria divergens*, *Rb.*, Wall., List, No. 2191; *Lindsaya divergens*, *Hk. & Grev.* (not *Wall.*), *Hk. Bk.*, Syn. Fil., 109; *Bedd.*, Ferns Br.I., II, tab. CCL; *Hk. & Grev.*, Ic. Fil., tab. CCXXVI. — var. *auriculata*, *Christ*, MS. in Herb. Bog.

Stipes approximate, wiry, erect, blackish, polished, 5—15 c.M. long. Fronds 10—30 c.M. long, 3—5 c.M. broad, lanceolate, pinnate. Pinnae close, spreading, $3\frac{1}{2}$ — $7\frac{1}{2}$ m.M. broad, the apex blunt, the edge entire or nearly so, the upper base truncate, slightly auricled, the lower base rounded or cuneate, lower pinnae \pm reduced, \pm deflexed. Texture subcoriaceous; rachis naked, polished; surfaces naked; veins forked, inconspicuous; costa present. Sori continuous along both sides and the apex.

Var. auriculata: Pinnae distinctly auricled at the base on the upper side.

Malaya.

* (6) **S. Walkerae**, *Kuhn*, *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam. I⁴, 218; *Lindsaya Walkerae*, *Hk.*, *Hk. Bk.*, Syn. Fil., 109; *Hk.*, Spec. Fil., I, tab. LXIX, A; *Bedd.*, Ferns S.I., tab. CCXV.

Rhizome creeping, ferrugineous, with scale-like hairs. Stipes wiry, erect, 15—30 c.M. long and as well as the rachis dark-coloured, glossy. Fronds 15—30 c.M. long, with a terminal pinna and 6—9 pairs of similar lateral ones. Pinnae $3\frac{1}{2}$ —10 c.M. long, $\frac{1}{2}$ —1 c.M. broad, distant, erecto-patent, gradually narrowing from a cuneate base towards the bluntish or acute apex, the highest ones sometimes confluent with the terminal. Texture coriaceous; rachis naked; veins numerous, very oblique. Sori in a continuous line along both edges.

Borneo, Banca, Sumatra; Ceylon.

§ 2. EUSCHIZOLOMA.

a. *Fronds dimorphous, simple, forked or ternate.*

*. *Barren fronds cordate-oblong, simple.* (7) *S. cordatum.*

.. *Barren fronds deltoid, 3-partite.* (8) *S. trilobatum.*

a.a. *Fronds uniform, pinnate, rarely simple or 2—3 × pinnate.*

b. *Pinnae simple, spreading or falcate, 1—2 c.M. long.*

(9) *S. Guerinianum.*

b.b. *Pinnae erecto-patent, $3\frac{1}{2}$ —15 c.M. long or longer.*

c. *Pinnae simple, linear or lanceolate, acuminate.*

d. *Fronds approximate, thin-herbaceous.* (10) *S. ensifolium.*

d.d. *Fronds distant, subcoriaceous.* (11) *S. coriaceum.*

c.c. *Pinnae varying from simple to irregularly 2-pinnate. Leaflets semi-orbicular, triangular, trapezoidal or cuneate-obovate, blunt or bluntish.* (12) *S. heterophyllum.*

* (7) **S. cordatum**, *Gaud.*, *Bedd.*, *Handb. Ind. Ferns*, 79; *Ferns Br.I.*, II, tab. CCIC; *Lindsaya cordata*, *Gaud.*, *Hk. Bk.*, *Syn. Fil.*, 111.

Rhizome short-creeping. Stipes 7—20 c.M. long, slender, erect, wiry. Fronds dimorphous. Barren fronds $3\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, $2\frac{1}{2}$ —4 c.M. broad, cordate-oblong, quite entire, fertile ones 7—15 c.M. long, linear, simple, forked or tripartite. Texture coriaceous. Sori in a continuous line along the margin.

Malaya.

(8) **S. trilobatum**, *v. A. v. R.*; *Lindsaya trilobata*, Bk., Journ. of Bot., XXIX, 107; *L. Hosei*, C. Chr., Ind. Fil.

Rhizome short-creeping. Stipes naked, those of the barren fronds slender, $2\frac{1}{2}$ —10 c.M. long. Fronds dimorphous, the barren ones deltoid, 3-partite, with an obovate-cuneate terminal lobe slightly repand in the upper half, and 2 shorter side-lobes produced on the lower side. Texture subcoriaceous; surfaces naked; veins immersed, obscure, flabellate, anastomosing. Fertile fronds with the terminal segment lanceolate. Sori running down both sides of the terminal segment, from the apex nearly to the base; indusium coriaceous, narrow.

Borneo.

(9) **S. Guerinianum**, *Gaud.*, Freyc., Voy. Ur. & Phys., Bot., 380, tab. XVIII; *Lindsaya Gueriniana*, *Gaud.*, Hk. Bk., Syn. Fil., 111.

Rhizome creeping, scaly. Stipes 10—15 c.M. long. Fronds 15—25 c.M. long, oblong-lanceolate, simply pinnate. Pinnae 1—2 c.M. long, ovate or oblong, nearly entire, horizontal or falcate, slightly auricled at the base on the upper side. Texture subcoriaceous. Sori in a continuous line along both margins.

Malaya.

* (10) **S. ensifolium**, *J. Sm.*, Journ. of Bot., III, 414; *Bedd.*, Ferns S.I., tab. XXV; *Lindsaya ensifolia*, *Sw.*, Rac., Flor. Btz., I., 143; *Hk.*, Gard. Ferns, tab. LXII; *Hk. & Grev.*, Ic. Fil., tab. CXI.

Rhizome short-creeping, densely scaly, the scales fibril-like, brown, subadpressed. Stipes 7—25 c.M. long, wiry, flexuose, rather approximate, stramineous or brownish. Fronds 15—30 c.M. long, $3\frac{1}{2}$ —10 c.M. broad, simply pinnate, the apex long, linear, simple or pinnatifid. Pinnae in 2—12 pairs, stalked or sessile, $3\frac{1}{2}$ —15 c.M. long, $2\frac{1}{2}$ —25 m.M. broad, linear or lanceolate, acuminate, \pm erecto-patent, equilateral, the base obliquely truncate to cuneate, sterile pinnae toothed, fertile ones rather narrower, entire. Texture thin, herbaceous. Sori in continuous lines along the margins. — The young plants have the fronds simple or ternate, but sterile.

Tropical Asia, Australia and Polynesia.

* (11) **S. coriaceum**, *v. A. v. R.*

Rhizome creeping, the scales dense, subadpressed, brown, small, fibril-like. Stipes rather distant, naked, glossy, dark-brown or purpureous, \pm 8—15 c.M. long or longer. Fronds simple, linear-lanceolate, 10—20 c.M.

long, 5—15 m.M. broad, narrowed gradually at both ends, forked, or pinnate with \pm 1—5 lateral pinnae on each side similar to the simple fronds, the edges entire, much recurved. Texture firm, subcoriaceous; surfaces naked; rachis like the stipe; veins anastomosing; costa prominent beneath, \pm wavy. Sori long, continuous along the margins; indusium firm. — The description is taken from a specimen determined as *Taenitis blechnoides*, Sw., (Ann. Btz., XX, 128).

Borneo (Hallier No. 1934).

* (12) **S. heterophyllum**, J. Sm., Copel., Polypod. Philipp., 59; Bedd., Ferns S.I., tab. XXVI; *Lindsaya heterophylla*, Dry., Hk. Bk., Syn. Fil., 112; Bedd., l. c., tab. CCVI; (? *L. orbiculata*, Mett., p. p., Copel., Philipp. Journ., I, Suppl., 148).

Rhizome short-creeping, the scales fine, brown. Stipes approximate, erect, 5—20 c.M. long, naked. Fronds 10—50 c.M. long, pinnate and lanceolate or 2-pinnate and deltoid with erecto-patent branches 7—10 c.M. long. Leaflets semiorbicular and fixed by the middle of the straight side with radiate venation and a continuous submarginal indusium around the upper side, or triangular with the indusium along 2 sides, or trapezoidal with the indusium usually continuous along the upper and outer sides, 1—2½ c.M. long, ½—1 c.M. broad, the apex blunt or bluntish, the margin entire or finely toothed. Texture papyraceo-herbaceous; surfaces naked. — A very variable fern.

Trop. Asia, South. China.

31a. SACCOLOMA, Kaulfuss.

Sori roundish or transversally oblong, submarginal, small, numerous, terminal on the veins, placed in submarginal depressions. Indusium slightly convex, covering the depression, attached at the base or the base and sides, open at the outer edge.

Rhizome short or creeping, scaly. Stipes not articulate to the rhizome. Fronds divided; veins free.

Malayan, Polynesian and Neotropical.

Arrangement of the sections.

- § 1. **CYSTODIUM.** Indusium transversally oblong, attached by the broad base, covering a marginal, distinctly cucullate outgrowth. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 113, E.
- § 2. **PARASACCOLOMA.** Indusium cuneate-oblong, the base acute, covering a half-cupshaped submarginal depression. Hk., Spec. Fil., I, tab. LVIII, A, (as *Davallia inequalis*, Kze, var. *minor*).
- § 3. **EUSACCOLOMA.** Indusium transversally oblong, the base broad, covering an intramarginal depression. Diels, l. c. fig. 113, C. (Not in Malaya).

§ 1. **CYSTODIUM.**

* (1) **S. sorbifolium**, *Christ*, Farnkr. d. Erde, 309; *Dicksonia sorbifolia*, *Sm.*, Hk. Bk., Syn. Fil., 52; Hk., Spec. Fil., I, tab. XXV, A; *Cystodium sorbifolium*, *J. Sm.*, Bau. & Hk., Gen. Fil., tab. XCVI; — var. *dentatum*.

Stipes tufted, 50 c.M. or much more long, the base densely clothed with long, fine, silky or woolly, yellowish hairs. Fronds large, 2-pinnate, 60 c.M. or much more long, ovate-lanceolate. Largest pinnae 25—35 c.M. long, ± 8 c.M. broad, stalked, the lowest sometimes smaller than the next in order. Pinnulae numerous, the highest sessile with a broad base, the lowest largest, ± 7—10 m.M. broad, short-stalked, the base truncate or cordate, whether or not auricled on the upper side, the edge sharply serrate towards the apex. Texture subcoriaceous; rachises and under side of the costae and veins more or less hairy; veins simple or forked. Fertile pinnulae bluntly toothed, bearing the sori on the teeth. Sori numerous, 1 to each tooth; indusium semiglobose, outer valve cucullate, formed by the apex of a tooth, inner valve smaller, ± plane.

Var. dentatum: Lower pinnulae deeply toothed.

Malaya.

§ 2. **PARASACCOLOMA.**

* (2) **S. moluccanum**, *Mett.*, Diels, in Engl. & Prantl, Nat. Pfl., Fam., I⁴, 211; *Davallia moluccana*, *Bl.*, Rac., Flor. Btz., I, 136; *D. inequalis*, *Kze.*, var. *minor.*, Hk., Spec. Fil., I, tab. LVIII, A; *D. asperrima*, *Ces.*, Fel.

Becc. Polin., 5 & 6; *Microlepia moluccana*, *Bedd.*, Handb. Ind. Ferns, Suppl., 15; — var. *stenolobum*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1005.

Rhizome creeping or suberect, clothed with large, pale-brown, lanceolate scales. Stipes strong, erect, to 40 c.M. long, \pm naked. Fronds deltoid to lanceolate, 3-pinnate to 4-pinnatifid, 60—120 c.M. long, 20—55 c.M. broad. Lower pinnae linear-lanceolate, to 8—15 c.M. broad, stalked. Pinnulae to 2 c.M. broad. Segments elliptical, the base decurrent, toothed or cut down to the costa into toothed, oblong lobes in the lower part. Texture membranous to subcoriaceous; both surfaces \pm naked. Sori 1—6 to a lobe, small, placed in the teeth near the base; the apices of indusia often produced beyond the margin of the lobes.

Var. *stenolobum*: Pinnulae much narrower than in the type.

Malaya; Polynesia; — var. stenolobum: Luzon.

31b. LEPTOLEPIA, *Mettenius*.

Sori roundish or transversally oblong, submarginal, solitary and terminal on short veinlets, whether or not placed on marginal, recurved outgrowths. Indusium of the same shape, fixed at the base, but by the apex of the veinlet only, otherwise free, the outer edge lacerato-lobate, opening outwardly.

Rhizome creeping and hairy (as far as known). Stipes not articulate to the rhizome. Fronds copiously divided; veins free. *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam., I, fig. 115, A—B.

Australia and Papua.

L. novae guineae, v. A. v. R.; *Davallia Novae Guineae*, *Rst.*, Fed., Rep., V, 36.

Rhizome and stipes unknown. Fronds large, \pm 2 M. long, \pm 1 M. broad, ovate-lanceolate, acuminate, 4-pinnatifid to 4-pinnate. Pinnae on stalks 3—5 c.M. long, subfalcate or erecto-patent, subsymmetrical, lanceolate, acuminate, the anterior side a little broader than the posterior one, the apex serrato-pinnatifid. Pinnulae and tertiary segments of the same shape as the pinnae, the former to 20 c.M. long and 8 c.M. broad, the latter to 5 c.M. long and \pm 1 $\frac{3}{4}$ c.M. broad at the base. Quaternary segments sessile, suberecto-patent, narrowed at the base, decurrent, elongate-ovate, acute, the edge obtusely serrato-dentate, the lowest to 10 m.M. long and 4 m.M. broad. Texture char-

taceo-herbaceous; surfaces naked; colour olive-green, paler beneath; rachis, costae and costulae furnished beneath with scattered hairs; veins simple or forked, lower veins pinnate, the veinlets simple or forked. Sori subglobose, occupying the apex of the higher anterior veinlets, submarginal, placed on subscariose, recurved, blunt or acute lobules; indusium transversally oblong, fastened by the apex of the veinlet only, whitish-yellowish, scariose, forming with the lobule a 2-valved involucre to the sorus, the outer edge dilacerate, the lacinulae piloso-fimbriate.

New Guinea.

32. HUMATA, *Cavanilles.*

Sori roundish or oblong, solitary and terminal on the veins or veinlets, submarginal or somewhat remote from the margin. Indusium \pm of the same shape, fixed by a broad base at the inner side of the sorus, free at the sides, opening outwardly (*).

Rhizome scaly, creeping. Stipes articulate to the rhizome, scattered. Fronds mostly divided; veins free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 112.

Paleotropical.

Arrangement of the sections.

§ 1. **EUHUMATA.** Indusium thick, coriaceous.

§ 2. **LEUCOSTEGIA.** Indusium thin, membranaceous (*Davallia*, p. p., C. Chr., Ind. Fil.).

§ 1. **EUHUMATA.**

a. *Barren fronds simple, entire, crenate or toothed.*

*. *Fertile fronds at best toothed.*

(1) *H. angustata.*

.. *Fertile fronds deeply pinnatifid.*

(2) *H. heterophylla.*

(*) The genera **Oleandra**, **Davallia** & **Microleptia** have the indusia sometimes more or less resembling those of **Humata**.

a.a. Barren fronds varying from pinnatifid to 3-pinnate.

b. Lowest primary segments simple or pinnatifid, not much broader or larger than the higher ones.

§. Rhizome scales adpressed. Stipes \pm 5—10 c.M. long.

(3) *H. Gaimardiana*.

§.§. Rhizome scales spreading. Stipes wanting or to \pm 5 c.M. long.

(4) *H. sessilifolia*.

b.b. Lowest primary segments at least pinnatifid, distinctly broader or larger than the higher ones.

c. Fronds uniform.

d. All the divisions very narrow, almost filiform, at utmost 1 m.M. broad. Fronds under 5 c.M. long. (5) *H. parvula*.

d.d. The divisions not almost filiform, at least 3 m.M. broad. Fully developed fronds 5—20 c.M. long.

e. Lobes of the lowest primary segments entire.

f. Primary segments deltoid-lanceolate.

(6) *H. pinnatifida*.

f.f. Primary segments linear-oblong. (7) *H. pectinata*.

e.e. Lobes of the lowest primary segments crenate or lobed.

(8) *H. repens*.

c.c. Fronds more or less dimorphous; i. e. the fertile segments narrower than the barren ones or the fertile fronds much larger than the barren fronds.

1. Barren fronds small, \pm $2\frac{1}{2}$ —4 c.M. long.

*. Fertile fronds \pm as large as the barren ones, with the segments contracted and sharply toothed. (9) *H. pusilla*.

.. Fertile fronds much larger than the barren ones, with the segments not contracted. (10) *H. lepida*.

1.1. Both barren and fertile fronds 4—20 c.M. long when fully developed.

2. Fertile fronds with the segments not toothed.

(11) *H. neoguineensis*.

2.2. Fertile fronds with the segments toothed, the teeth often very sharp.

3. Lowest pinnae pinnatifid or pinnate in the barren fronds.

4. Secondary segments entire or crenate in the barren fronds. (12) *H. alpina*.

4.4. *Secondary segments bluntly toothed in the barren fronds.* (13) *H. botrychioides*.

3.3. *Lower pinnae 2-pinnate in the barren fronds.*

(14) *H. vestita*.

* (1) ***H. angustata***, *J. Sm.*, Copel., Polypod. Philipp., 50; Bedd., Ferns Br.I., II, tab. CCXXXVII; *H. heterophylla*, *Desv.*, var., C. Chr., Ind. Fil.; *Davallia angustata*, *Wall.*, Hk. Bk., Syn Fil., 88; Hk. & Grev., Ic. Fil., tab. CCXXXI.

Rhizome wide-creeping; scales brown, ovate-lanceolate, acuminate, erectopatient to erect, the base peltate. Fronds sessile or stalked, linear-lanceolate, $7\frac{1}{2}$ —20 c.M. long, $\frac{1}{2}$ —2 c.M. broad, slightly or more distinctly and irregularly crenate to toothed at the margin, sometimes once forked, the stipes varying from short to 6 c.M. long, often slightly winged. Texture coriaceous; surfaces naked. Sori in a row along the edges.

Malaya.

* (2) ***H. heterophylla***, *Desv.* (not *J. Sm.*), Copel., Polypod. Philipp., 50; Bedd., Ferns Br.I., I, tab. C; *Davallia heterophylla*, *J. Sm.*, Rac., Flor. Btz., I, 126; Hk., Fil. Exot., tab. XXVII; Hk. & Grev., tab. CCXXX; — var. *nervosa*.

Rhizome wide-creeping, the scales dense, with an oblong, peltate, adpressed base and a long, linear-subulate, recurved or spreading, shortly ciliate or finely toothed point. Stipes 1— $4\frac{1}{2}$ c.M. long, 1—3 c.M. apart. Fronds dimorphous, the barren ones lanceolate, 6—15 c.M. long, $1\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, the apex acute, the edge entire or slightly crenate, the base more or less suddenly narrowed or rounded and then often decurrent so as to form a long, narrow wing to the stipe. Texture coriaceous, rigid; surfaces naked; veins commonly once or more times forked. Fertile fronds narrower, deeply pinnatifid into blunt, entire or broadly toothed lobes 3—4 m.M. broad. Sori 2—10 to a lobe, often close; indusium transversally oblong.

Var. *nervosa*: Fronds crisped; veins more conspicuous.

Malaya; Polynesia.

* (3) ***H. Gaimardiana***, *J. Sm.*, Copel., Polypod. Philipp., 50; *H. parallela*, *Brack.*, Bedd., Ferns Br.I., I, tab. IC; *Davallia Gaimardiana*, *Pr.*,

Rac., Flor. Btz., I, 127; *D. parallela*, Wall., Hk., Spec. Fil., I, tab, XLII, A; Hk. Bk., Syn. Fil., 89; *D. intramarginalis*, Bl., Hk., l. c., 154.

Rhizome wide-creeping, densely clothed with adpressed, lanceolate, peltate scales. Stipes 1—5 c.M. apart, 5—10 c.M. long, erect, more or less scaly like the rachis. Fronds 10—20 c.M. long, 3—5 c.M. broad, triangular-lanceolate, cut down nearly or quite to the rachis into long, nearly or quite close, parallel, linear-oblong, entire, bluntish or acute segments, the lowest pair often a little shorter and sometimes auricled. Texture coriaceous; surfaces naked; veins forked. Sori oblique, occupying the apices of the anterior veinlets, placed in 1 row on each side of the costulae; indusium with the base slightly excavate.

Malaya; Burmah, Polynesia.

* (4) **H. sessilifolia**, *Mett.* (not *Sm.*), Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 209; *Davallia sessilifolia*, Bl., Rac., Flor. Btz., I, 128; — var. *polypodioides*; *H. polypodioides*, Brack., Wilk., Expl. Exp., XVI, 228, tab. XXXII, fig. 1.

Rhizome wide-creeping, the scales dense, with the base dilated and peltate, the point lanceolate-subulate, spreading. Stipes very short, to 1½ c.M. long. Fronds ovate or deltoid-lanceolate, acuminate, 5—10 c.M. long, 2½—4 c.M. broad, cut down nearly to the costa into linear-oblong, parallel, entire or toothed segments, the lower ones sometimes deeply pinnatifid or auricled at the lower base. Segments ± 3—4 m.M. broad. Texture varying from thin to subcoriaceous; surfaces naked; veins forked. Sori in 1 row on each side of the costulae, occupying the greater part of the space between the costulae and the edges.

Var. polypodioides: Fronds larger, more acute, on stipes 2½—5 c.M. long.

Forms reported from Celebes had the fronds both sessile and stalked on the same rhizome.

Java, Celebes; Fiji.

* (5) **H. parvula**, *Mett.* (not *J. Sm.*), Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 209; *Davallia parvula*, Wall., Hk. Bk., Syn. Fil., 92; Hk. & Grev., Ic. Fil., tab. CXXXVII; C. Chr., Ind. Fil.

Rhizome wide-creeping, branched, the scales lanceolate, peltate near the base, entire or nearly so, ferruginous. Stipes scattered, 1—7½ c.M. long. Fronds 1½—4 c.M. long, to 5 c.M. broad, deltoid-ovate in general outline,

2—5-pinnate, with all the divisions almost filiform and the ultimate segments sharp at the point. Sori placed at the sinuses of the ultimate forks; indusium coriaceous, broader than the segments.

Malacca, Banca, Borneo, Celebes.

(6) **H. pinnatifida**, *Bedd.*, Handb. Ind. Ferns, Suppl., 12; *H. intermedia*, *C. Chr.*, Ind. Fil.; *Davallia pinnatifida*, *Bk.*, Journ. of Linn. Soc., Bot., XXIV, 257.

Rhizome slender, firm, wide-creeping, clothed with close-pressed, ovate, peltate scales, glaucous beneath the scales. Stipes distant, firm, erect, stramineous, 5½—10 c.M. long. Fronds elongate-deltoid, broadest at the base, narrowed gradually towards the apex, 7½—15 c.M. long, 5½—6½ c.M. broad at the base, pinnatifid nearly to the costa into numerous, deltoid-lanceolate, entire, crenate or slightly lobed segments 4—7 m.M. broad, the lowest pair much the broadest, deltoid, lobed on the lower side. Texture rigidly coriaceous; veins simple or forked. Sori small, terminal on the veins; indusium rigid, persistent, broader than long.

Borneo, Malacca.

(7) **H. pectinata**, *Desv.*, Prod., 525; *Davallia pectinata*, *Sm.*, Hk. Bk., Syn. Fil., 89; Hk. & Grev., Ic. Fil., tab. CXXXIX.

Rhizome creeping, scaly. Stipes distant, 5—10 c.M. long, more or less scaly. Fronds 10—20 c.M. long, 5—8 c.M. broad, elongate-deltoid, cut down nearly or quite to the rachis into many, parallel, oblong or linear-oblong segments. Segments approximate, blunt, the higher ones entire, the lower ones inciso-pinnatifid, the lowest ones a little broader and deeply pinnatifid on the lower side, with the lobes sometimes 1—1½ c.M. long. Texture coriaceous. Sori oblique, submarginal; indusium transversally oblong, broader than deep, attached at the central part of the base.

Borneo, New Guinea; Polynesia.

* (8) **H. repens**, *Diels* (not *J. Sm.*), Copel., Polypod. Philipp., 50; *H. pedata*, *J. Sm.*, *Bedd.*, Handb. Ind. Ferns, 48; Ferns S. I., tab. XII; *Davallia pedata*, *Sm.*, Rac., Flor. Btz., I, 128; Hk., Spec. Fil., I, tab. XLV, A; Gard. Ferns, tab. VII; *Adiantum repens*, *L.*, (oldest name).

Rhizome wide-creeping; scales dense, lanceolate, the apex acute, the base peltate, blunt. Stipes distant, 5—10 c.M. long, sparingly scaly. Fronds

5—20 c.M. long, 4—10 c.M. broad, deltoid, or ovate-deltoid, cut down nearly or quite to the rachis into few to several, oblong or linear-oblong segments. Segments rather close, blunt, crenate or toothed, the lowest ones broader; sessile or subsessile with a narrowed base, the lower side the broadest, lobed or pinnatifid, with the lobes crenate, toothed or lobed again, the upper side narrow, toothed or lobed. Texture coriaceous. Sori submarginal, 1 to each crenation or tooth; indusium \pm orbicular, sometimes slightly and transversally oblong, but not much broader than deep.

Trop. Asia & Australia, Japan, South. China, Mascarenes, Seychelles.

* (9) **H. pusilla**, *Carr.* (not *Sm.*), Diels, in Engl. & Prantl, Nat. Pfl.Fam., 1⁴, 209; *Davallia pusilla*, *Mett.*, Hk. Bk., Syn. Fil., 89.

Rhizome creeping, scaly. Stipes to 3 c.M. long. Fronds $2\frac{1}{2}$ —4 c.M. long, $1\frac{1}{2}$ —3 c.M. broad, or the barren fronds not rarely smaller, cordato deltoid in general outline, cut down to a narrowly winged rachis. Upper segments linear-oblong, deeply inciso-dentate in the fully developed fronds, the lower pair deeply inciso-pinnatifid on the lower side, those of the fertile fronds narrower than in the barren ones, with the teeth sharper. Texture coriaceous; veins \pm prominent. Sori small, placed at the base of the small teeth.—Plants suggesting *H. alpina*, *Moore*, but much smaller, with the barren fronds not rarely very small and the segments nearly undivided.

Borneo, Mindanao, New Guinea; Melanesia.

* (10) **H. lepida**, *Moore*, C. Chr., Ind. Fil.; *H. Cumingii*, *Brack.*, Copel., Polypod. Philipp., 51; *Davallia Cumingii*, *Hk.*, Hk. Bk., Syn. Fil., 90; *Hk.*, Spec. Fil., I, tab. XVI, B; *D. longula*, *Kze*, Bot. Zeit., 1848, 215; *D. lepida*, *Pr.*, (oldest name).

Rhizome creeping, scaly. Stipe $7\frac{1}{2}$ —10 c.M. long, both it and the rachis rather scaly. Fronds dimorphous. Barren fronds deltoid-cordate \pm $2\frac{1}{2}$ —3 c.M. each way, cut down nearly to the costa. Upper lobes oblong, blunt, slightly toothed, the lower pair deeply pinnatifid below. Texture coriaceous. Fertile fronds \pm the same shape, but more divided, 10— $12\frac{1}{2}$ c.M. long, 7—8 c.M. broad, pinnate. Upper pinnae toothed to pinnatifid, united at the base by a narrow wing, linear-oblong, lower pinnae cut down nearly or quite to the rachis at the base into toothed lobes or pinnulae. Sori placed in the teeth. — Allied to *H. repens*, *Diels* and like this with the fertile segments not contracted.

Philippines.

FILICES.

(11) **H. neoguineensis**, C. Chr., Ind. Fil.; *Davallia bipinnatifida*, Bk., Kew Bull., 1899, 119.

Rhizome slender, wide-creeping, the scales lanceolate, adpressed, membranaceous, pale-brown. Stipes slender, naked, 7—10 c.M. long. Fronds lanceolate-deltoid, 2-pinnatifid, 12—15 c.M. long, \pm 5 c.M. broad. Pinnae sessile, the lower largest, the barren ones oblong, obtuse, slightly pinnatifid, cuneate at the base, the fertile ones narrower, deeply pinnatifid, with the lobes erecto-patent, the lower lobes oblong, obtuse. Texture coriaceous; surfaces naked like the slender rachis; veins flabellate, conspicuous. Sori solitary on the higher lobes, 2—5 on the lower ones; indusium semiorbicular, firm. *New Guinea*.

* (12) **H. alpina**, Moore (not Sm.), Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 209; *H. repens*, Diels, var., C. Chr., Ind. Fil.; *Davallia alpina*, Bl., Rac., Flor. Btz., I, 127; — var. *montana*.

Like *H. repens*, Diels, but dimorphous. Stipes 1—10 c.M. long. Fronds 4—10 c.M. long, 3—9 c.M. broad. Barren fronds with the central pinnae oblong, crenate, toothed or obliquely pinnatifid and the lower pinnae pinnate or 2-pinnatifid. Fertile fronds brown when dry, the central pinnae linear, simple and the lower pinnae pinnate, the laciniae slightly narrowed, the edges toothed, the teeth commonly sharp, upcurved. Sori occupying the upper base of the teeth.

Var. montana: Fertile fronds more finely divided, blackish when dry, the laciniae much contracted so that the leafy membrane is nearly wanting. Sori occupying the base of the always very sharp, hamate teeth.

Malaya; Polynesia; — var. *montana*: *Java* (Forbes), *Billiton* (Teysmann).

(13) **H. botrychioides**, Brack. (not J. Sm.), Copel., Polypod. Philipp., 51; Wilk., Expl. Exp., XVI, tab. XXXII, fig. 1; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 112, C—D; *Davallia botrychioides*, Bk. (not Brack.), Hk. Bk., Syn., Fil., 90.

Rhizome wide-creeping, scaly. Stipes 5—12½ c.M. long, scaly below. Fronds 6—20 c.M. long, 5—12½ c.M. broad, deltoid in outline; dimorphous, the barren ones with a narrowly winged rachis. Lowest pinnae cut down nearly to the rachis with deep, bluntly toothed segments. Fertile pinnae more finely divided; pinnulae of the lower pinnae cut down to a narrow rachis into narrow, sharply toothed segments. Sori in the sinuses of the teeth.

Philippines, Celebes; Polynesia.

* (14) **H. vestita**, *Moore* (not *J. Sm.*), *Copel.*, *Polypod. Philipp.*, 50; *Davallia vestita*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 126; *Hk. Spec. Fil.*, I, tab. XLI, C; *D. bipinnatifida*, *Bl.*, *Hk.*, l. c., 156; *D. chaerophylla*, *Ces.*, *Fel. Becc. Polin.*, 3; *D. Bélangeri*, *Bory*, *Bél.*, *Voy. Bot.*, II, tab. VII, fig. 1.

Rhizome wide-creeping, the scales dense, linear-lanceolate, membranaceous, peltate at the base. Stipes rather distant, $7\frac{1}{2}$ —20 c.M. long, sparingly covered with long, thin scales especially near the base. Fronds deltoid to lanceolate-ovate, 10—20 c.M. long, 7—10 c.M. broad. Barren fronds with the central pinnae pinnate and the lower pinnae 2-pinnate, the segments oblong, crenate (scarcely toothed), united at the base by a narrow wing. Fertile fronds with the segments narrower, sharper toothed. Texture coriaceous. Sori placed on the teeth. — Near *H. repens*, *Diels*, and *alpina*, *Moore*.

Malaya.

§ 2. LEUCOSTEGIA.

a. *Fronds small, under 20 c.M. long* (*). (15) *H. falcinella*.

a.a. *Fully developed fronds middle-sized, 20—60 c.M. long.*

b. *Sori on the dilated tips of the ultimate segments.* (16) *H. cicutarioides*.

b.b. *Sori at or near the base of the ultimate segments.*

c. *Scales of the rhizome obtusely rounded, adpressed.* (17) *H. pulchra*.

c.c. *Scales of the rhizome acute or acuminate.*

d. *Sori ± immersed, prominent on the upper surface.*

(18) *H. immersa*.

d.d. *Sori not immersed.*

e. *Rhizome scales with the apex not recurved.*

*. *Fronds 2-pinnate.* (19) *H. Hosei*.

.. *Fronds 4—5-pinnate.* (20) *H. hymenophylloides*.

e.e. *Rhizome scales with the apex spreading or recurved.*

f. *Rachis narrowly winged.* (21) *H. Kingii*.

f.f. *Rachis not at all winged.* (22) *H. viscidula*.

a.a.a. *Fronds large, 60—90 c.M. long when fully developed.*

(23) *H. nephrodioides*.

(*) See also under **H. immersa**, *Mett.*

(15) **H. falcinella**, *Copel.*, Polypod. Philipp., 51; *Davallia falcinella*, *Pr.*, Hk. Bk., Syn. Fil., 91; *Pr.*, Rel. Haenk., I, tab. XI, fig. 2; C. Chr., Ind. Fil.

Rhizome wide-creeping, thickly beset with stiff, spreading hairs, ferruginous, turning toward black. Stipes 5—10 c.M. long, naked. Fronds 10—15 c.M. long, $7\frac{1}{2}$ —10 c.M. broad, deltoid, 5—4-pinnatifid; main rachis slightly winged. Lowest pinnae inequilateral. Pinnulae oblong, cut down nearly to the rachis into segments which are again deeply toothed. Ultimate segments of the fertile fronds falcate-mucronate, of the barren fronds broader and not so sharp. Texture subcoriaceous; surfaces naked. Sori 2—6 to a lobe, placed at the base of the teeth. — Ferns with the aspect of Colposoria and the rhizome hairs of *Microlepia*.

Philippines.

(16) **H. cicutarioides**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 210; *Davallia cicutarioides*, *Bk.*, Journ. of Bot., XXVIII, 106; C. Chr., Ind. Fil.

Fronds decomposed. Lower pinnae ovate-lanceolate, 15—20 c.M. long. Ultimate segments linear, 2—4 m.M. long, the sterile ones acute, the fertile ones obtuse. Texture moderately firm; rachises slender, naked, polished, brown; veins 1 to each ultimate segment. Sori minute, solitary on the dilated tips of the ultimate segments; indusium pale-coloured, suborbicular.

New Guinea.

* (17) **H. pulchra**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 209; *Davallia pulchra*, *Don*, Christ, Farnkr. d. Erde, 305; C. Chr., Ind. Fil.; *D. chaerophylla*, *Wall.*, Hk. Bk., Syn. Fil., 92; *Leucostegia pulchra*, *J. Sm.*, *Bedd.*, Handb. Ind. Ferns, 52; *Acrophorus pulcher*, *Moore*, *Bedd.*, Ferns S.I., tab. X.

Rhizome wide-creeping, clothed with broad, obtuse, adpressed scales. Stipes 10—15 c.M. long, naked. Fronds 20—40 c.M. long, 10—30 c.M. broad, lanceolate-deltoid, 5—4-pinnatifid. Lowest pinnulae $2\frac{1}{2}$ —5 c.M. long, $2\frac{1}{2}$ c.M. broad, cut down to a narrowly winged rachis into deeply pinnatifid segments. Ultimate lobes narrow and acute in the fertile, rather broader in the barren fronds. Texture thinly herbaceous, rather flaccid. Sori copious, placed at the base of the teeth of the segments, with they about equal in breadth.

New Guinea, Malacca; Ceylon, Noth. India, Yunnan.

* (18) **H. immersa**, *Mett.* (not *Diels*), *Copel.*, Polypod. Philipp., 51; *Davallia immersa*, *Wall.*, *Rac.*, *Flor. Btz.*, I, 129; *Hk.*, *Fil. Exot.*, tab.

LXXIX; C. Chr., Ind. Fil.; *Leucostegia immersa*, Pr., Bedd., Handb. Ind. Ferns, 51; *Acrophorus immersus*, Moore, Bedd., Ferns S.I., tab. XI; — var. *amplissima*, Christ, Ann. Btz., XV, 90; — var. *minuta*; — var. *nana*, Copel., Philipp. Journ., I, Suppl., 147.

Rhizome wide-creeping, clothed with lanceolate, ferruginous fibrils. Stipes 10—50 c.M. long, firm, erect, naked. Fronds 30—45 c.M. long, 15—25 c.M. broad, triangular, 3-pinnate. Lowest pinnulae lanceolate-triangular, 5—8 c.M. long, with the ultimate divisions rhomboidal, the base, obliquely cuneate, the upper side broader than the lower one, the edge lobed, with the lobes rounded and crenate above. Texture herbaceous or flaccido-coriaceous; surfaces naked. Sori commonly large, immersed, \pm prominent on the upper surface, 1—6 to a leaflet; indusium broader than long, the outer edge semiorbicular, the base straight.

Var. amplissima: Fronds lanceolate, to 75 c.M. long; indusium reniform.

Var. minuta: Fronds small, to 10 c.M. long, to 5 c.M. broad. Stipes slender.

Var. nana: Stipes 5—10 c.M. long, filiform; fronds 5—10 c.M. long, 2-pinnate; pinnae obtuse; pinnulae rounded.

Philippines, Java, Malacca; Br. India; — var. *amplissima*: *Celebes*; — var. *minuta*: *Timor* (Forbes, no. 3522); — var. *nana*: *Luzon*.

(19) **H. Hosei**, v. *A. v. B.*; *Davallia Hosei*, Bk., Journ. of Bot., XXVI, 525; C. Chr., Ind. Fil.

Rhizome creeping, the scales minute, adpressed, linear, brown. Stipes naked, brownish, 30—40 c.M. long. Fronds lanceolate-oblong, 2-pinnate, 45—60 c.M. long, 12—15 c.M. broad. Pinnae distant, lanceolate, erecto-patent, 2—2½ c.M. broad, cut down to the rachis into unequal-sided, lanceolate-oblong, deeply crenate pinnulae \pm 4—5 m.M. broad, with the upper base produced and the lower cut away. Texture firm; surfaces naked; veins indistinct, 1 to each lobule. Sori small, 1 at the base of each lobule; indusium rigid, naked, as broad as long.

Borneo.

* (20) **H. hymenophylloides**, Copel., Polypod. Philipp., 51; *Davallia affinis*, Hk., Rac., Flor. Btz., I, 129; Hk., Spec. Fil., I, tab. LII, B; *D. hymenophylloides*, Kuhn, Ann. Mus. Bot. L.B., IV, 286; (not *Bl.*), Christ, Farnkr. d. Erde, fig. 957; C. Chr., Ind. Fil.; *Acrophorus affinis*, Moore, Bedd.,

Ferns S.I., tab. CCLII; *Leucostegia hymenophylloides*, *Bedd.*, Handb. Ind. Ferns, 54; *Aspidium hymenophylloides*, *Bl.*, (oldest name).

Rhizome wide-creeping, the scales \pm dense, ferruginous, lanceolate, long-acuminate, peltate at the base. Stipes scattered, 10—50 c.M. long. Fronds 50—60 c.M. long, 15—40 c.M. broad, deltoid-lanceolate, 4—5-pinnate. Lowest pinnulae with oblong-rhomboidal divisions, the segments of which are entire or deeply and *finely* inciso-pinnatifid with sharp teeth. Texture herbaceous; surfaces naked; veins fine, blackish, 1 in each ultimate segment. Sori minute, solitary near the base of the segments, \pm 2—6 to each tertiary division, terminal on very short veinlets and therefore often apparently medial on the ultimate veins; indusium broad, semiorbicular or subcordate.

Trop. Asia & Polynesia.

(21) **H. Kingii**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 210, fig. 112, H; *Davallia Kingii*, *Bk.*, Hk., Icon. Pl., XVII, tab. MDCXXII; C. Chr., Ind. Fil.

Rhizome strong, woody, wide-creeping, the scales dense, spreading, subulate, membranaceous, peltate at the base. Stipes erect, straight, naked, \pm 15 c.M. long. Fronds oblong-lanceolate, \pm 45 c.M. long, 20—25 c.M. broad, 3-pinnatifid, sparingly hairy on both surfaces; rachis hairy, narrowly winged. Pinnae lanceolate, the central and lowest longest, subspreading, \pm 5—4 c.M. broad. Pinnulae oblong, adnate, decurrent, deeply pinnatifid, the lobes close, oblong, bluntish or acute. Sori solitary at the base of the lobes; indusium subtriangular-oblong. — Hardly different from *H. viscidula*, *v. A. v. R.* It is much like *Davalia hirsuta*, *v. A. v. R.*, but totally different in the structure of the indusium.

Java.

* (22) **H. viscidula**, *v. A. v. R.*; *Davallia viscidula*, *Mett.*, Rac., Flor. Btz., I, 131; C. Chr., Ind. Fil.

Rhizome wide-creeping, densely scaly, the scales with the base rounded, adpressed, reniform, the apex spreading, long-acuminate to filiform. Fronds 5—4-pinnatifid, \pm 35 c.M. long, to \pm 22 c.M. broad, acuminate, the lower pinnae shorter than the following. Pinnae sessile, oblong-lanceolate, acuminate, to \pm 12 c.M. long. Pinnulae close, to $2\frac{1}{2}$ c.M. long, erecto-patent, lanceolate, bluntish or acute, free and decurrent at the base or more or less confluent, cut down to a \pm winged rachis into toothed or pinnatifid segments to 5 m.M. long and 3 m.M. broad, narrowed at the base, (of the same shape as

the pinnulae). Texture herbaceous; stipes, rachises and surfaces glanduloso-pubescent. Sori near the sinuses of the teeth; indusium \pm cordate-oblong.

Java.

(23) **H. nephrodioides, v. A. v. R.**; *Davallia nephrodioides, Bk.*, Journ. of Linn. Soc., Bot., XXIV, 257; C. Chr., Ind. Fil.

Rhizome wide-creeping, the scales fine, blackish, spreading, hair-like, pel-tate at the base. Stipes stiffly erect, naked, \pm 15 c.M. long. Fronds 60—90 c.M. long, oblong-lanceolate, 20—25 c.M. broad at the middle, 2-pinnate, hairy on the costae (rachises), scaleless. Pinnae subpetiolate, lanceolate, the central ones $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, the lower rather smaller, all except the lowest cuneate-truncate on the lower side at the base. Pinnulae multijugate, oblique, oblong-lanceolate, 3—5 m.M. broad, cut $\frac{1}{2}$ -way down into close, ascending, lanceolate lobes. Veins 1 to each final lobe, with a small medial sorus; indusium naked, subreniform, small.

Borneo.

33. ACROPHORUS, *Presl.*

Sori roundish, submarginal, terminal on a veinlet, which is springing \pm from the middle of the fertile vein. Indusium broadly ovate, attached at the inner side of the sorus, free at the sides, opening outwardly.

Rhizome erect. Stipes tufted, not articulate to the rhizome. Fronds much divided; veins free. Diels, in Engl. & Prantl., Nat. Pfl.Fam., I⁴, 165, 89, D—F.

Malaya, Northern India.

* **A. nodosus, Pr.**, Christ, Farnkr. d. Erde, 285; Bedd., Ferns Br.I., I, tab. XCIII; *A. stipellatus, Moore*, C. Chr., Ind. Fil.; *Davallia nodosa, Hk.*, Rac., Flor. Btz., I, 150; — var. *alpinus*, Christ, Ann. Btz., XV, 89.

Rhizome short, erect, the scales large, ovate or oval, entire, membranaceous. Stipes tufted, 50—100 c.M. long, strong, erect, scaly below. Fronds deltoid-ovate, 60—120 c.M. long, to 80 c.M. broad, 3—4-pinnate. Pinnae subopposite, sessile or short-stalked, the rachis glandular at the base, lower pinnae \pm 15—25 c.M. broad. Pinnulae lanceolate, the rachis glandular

at the base; lower pinnulae \pm 2—5 c.M. broad, with the segments cut down nearly or quite to the rachis into oblong, blunt, entire or toothed lobes \pm 5 m.M. long, \pm 1½ m.M. broad, \pm narrowed at the base. Texture herbaceous; surfaces sparingly hairy; rachises scaly at the base of the divisions. Sori small, solitary and medial in the ultimate lobes.

Var. alpinus: Fronds small, 10—15 c.M. long. Stipes clothed with spreading, acuminate scales.

Malaya; North. India; — var. alpina: Celebes.

34. DAVALLIA, *Smith.*

Sori roundish or oblong, submarginal, solitary and terminal on the veins or veinlets; indusium \pm half-cupshaped or semicylindrical, attached at the base and sides, opening outwardly.

Rhizome scaly, short or creeping, the scales not minute. Stipes articulate to the rhizome. Fronds variously divided; veins free; soriferous veinlets sometimes very short and then the sori appearantly medial on the veins.

Tropical and subtropical in the old world.

Arrangement of the sections.

- § 1. **SCYPHULARIA.** Fronds pinnate with 1—4 linear, undivided pinnae on each side below the terminal one. Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 115, J.
- § 2. **COLPOSORIA.** Fronds at least 2-pinnatifid with small, ultimate segments. Diels, l. c., fig. 115, K.

§ 1. **SCYPHULARIA.**

*. *Fronds 3-foliate.*

(1) *D. triphylla.*

.. *Fronds 5—9-foliate.*

(2) *D. pentaphylla.*

(1) **D. triphylla**, *Hk.*, Hk. Bk., Syn. Fil., 94; Hk., Spec. Fil., I, tab. XLVI, A; Bedd., Ferns Br.I., I, tab. CV.

Rhizome creeping, stout, densely clothed with chaffy scales. Stipes scattered, 5—10 c.M. long, erect, strong. Fronds ternate. Central segment linear, 10—15 c.M. long, 1—2 c.M. broad, obtuse, entire, the very base cuneate, short-stalked, lateral segments similar but shorter, sessile, oblique at the base. Texture coriaceous; veins copious, crowded, parallel, forked, flat. Sori placed in a close row along the margin; indusia semicylindrical, compressed, contiguous so as to form an uninterrupted marginal line along whole the length of the leaflets.

Malacca.

* (2) **D. pentaphylla**, *Bl.*, Rac., Flor. Btz., I, 153; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 113, J; Christ, Farnkr. d. Erde, fig. 947; Hk., Fil. Exot., tab. XXXVII.

Rhizome wide-creeping, the scales dense, rounded, adpressed, peltate at the base, the long, hair-like point spreading. Stipes scattered, strong, erect, 5—10 c.M. long. Fronds with 2—3 pairs of lateral pinnae and a terminal one. Terminal segment 4—15 c.M. long, 1—2 c.M. broad, of the barren frond the broadest and shortest, stalked, linear, bluntish, the base cuneate, the edge more or less crenate or serrulate, lateral segments similar, but commonly shorter, sometimes sessile, the lowest pair sometimes with 1—2 similar, smaller pinnulae at the base. Texture coriaceous; veins close, forked, parallel, flat. Sori close, numerous, in rows along the margin; indusia semicylindrical, often contiguous.

Malaya.

§ 2. COLPOSORIA.

a. Rhizome scales with a small, roundish or oblong, peltate base, the apex suddenly narrowed into a very long, hair-like point. Rachis and surfaces hairy or fibrillose. (Davallodes). (5) *D. hirsuta*.

a.a. Rhizome scales otherwise. Rachis and surfaces naked or \pm scaly. (Eudavallia).

b. Veins with a terminal, calcareous scale on the upper surface of the frond. (4) *D. Korthalsii*.

b.b. Upper surface without calcareous scales.

c. Indusium \pm as broad as long or broader.

- d.** *Indusium* more broad than long, or shortly attached in the lower half of the sides.
- *. *Indusium* half-cupshaped, shortly attached at the sides nearly as in *Humata*. (5) *D. Griffithiana*.
- *.*. *Indusium* broader than deep, attached at the sides throughout. (6) *D. Lobbiana*.
- d.d.** *Indusium* \pm as broad as long, attached at the sides throughout.
- e.** *Indusium* cuneate. (7) *D. stenoloba*.
- e.e.** *Indusium* half-cupshaped or suburceolate, the apex subrotund. (8) *D. pallida*.
- e.e.e.** *Indusium* half-cupshaped, the apex truncate.
- f.** Apex of *indusium* \pm reaching the margin of the segment. Spurious veins more or less distinctly present.
- g.** Rhizome scales crisped and interlaced, short-ciliate. (9) *D. denticulata*.
- g.g.** Rhizome scales flat, entire. (10) *D. epiphylla*.
- f.f.** Apex of *indusium* at some distance from the margin of the segment. No spurious veins. (11) *D. decurrens*.
- e.e.e.e.** *Indusium* \pm orbicular or elliptical, the apex rounded or deltoid-roundish. (12) *D. elata*.
- c.c.** *Indusium* generally longer than broad (*).
- 1.** *Indusium* less than 2 \times as long as broad.
- 2.** Fronds under 50 c.M. long.
- 3.** Rhizome scales nearly black. (13) *D. Lorrainii*.
- 3.3.** Rhizome scales whitish or pale-brown.
- 4.** Edges of the ultimate segments not incised at the inner side of the sori, the teeth therefore not lobe-like. (14) *D. pyxidata*.
- 4.4.** Edges of the ultimate segments incised at the inner side of the sori, which gives the teeth the appearance of very small, soriferous lobes.
- 5.** *Indusium* half-cupshaped to semicylindrical.
- 6.** Soriferous teeth short and broad. (15) *D. bullata*.
- 6.6.** Soriferous teeth long and narrow. (16) *D. fructuosa*.

(*) See also *D. elata*, Spr.

5.5. *Indusium urceolate.*(17) *D. trichomanoides.*3.3.3. *Rhizome scales brown. Rhizome stout, scandent.*(18) *D. Wagneriana.*2.2. *Fully developed fronds more than 50 c.M. long. **(19) *D. divaricata.*1.1. *Indusium $\pm 2 \times$ as long as broad.**α. Rhizome scales strongly adpressed, which gives the rhizome the appearance to be quite naked.* (20) *D. solida.**α.α. Rhizome scales not strongly adpressed.**β. Indusium rostrate.* (21) *D. embolostegia.**β.β. Indusium not rostrate.**γ. Soriferous segments without horns, but denticulate at the apex.* (22) *D. Veitchii.**γ.γ. Soriferous segments with 1—2 distinct horns.**δ. Tips of the ultimate segments rather hair-like. Stipes winged.* (23) *D. brevipes.**δ.δ. Tips of the ultimate segments not hair-like. Stipes not winged.* (24) *D. dissecta.*1.1.1. *Indusium $\pm 3—4 \times$ as long as broad.* (25) *D. fijiensis.*

* (3) ***D. hirsuta***, v. ***A. v. R.***, Hort. Bog.; *D. ciliata*, Hk., Hk. Bk., Syn. Fil., 98; Hk., Spec. Fil., I, tab. LX, A; *Microlepia ciliata*, Copel., Polypod. Philipp., 55; *M. hirsuta*, Pr., Epim., 97; C. Chr., Ind. Fil.; *Davallodes grammatosorum*, *hirsutum* & *gymnocarpum*, Copel., Philipp. Journ., III, 33—34; *Leucostegia hirsuta*, J. Sm., (oldest name).

Rhizome firm, creeping, scaly, the scales with a bluntish, roundish or oblong, peltate base and a suddenly very long-acuminate, hair-like, erectopate or spreading apex $\pm 1—1\frac{1}{2}$ c.M. long. Stipes scattered, 5—10 c.M. long, pubescent, firm, erect. Fronds $\pm 20—60$ c.M. long, $\pm 7\frac{1}{2}—25$ c.M. broad, lanceolate, 2—5-pinnatifid. Pinnae spreading, lanceolate, the central ones the longest, $1\frac{1}{2}—4$ c.M. broad, deeply pinnatifid with oblong or linear-oblong, entire to pinnatifid, blunt or acute, close segments. Texture thin, flaccid; rachis and under surface and occasionally also the upper surface hairy or fibrillose especially on the costae and veins. Sori 2—12 to a segment, very small, semicupuliform or semicylindrical, placed near the centre of the teeth at the base, whether or not ciliate. — Copeland distinguishes 5 forms:

Davallodes grammatosorum, Copel, l. c., tab. VI: Indusium narrowly semi-cylindrical, naked; lower pinnae hardly reduced.

Davallodes hirsutum, Copel.: Indusium semicupuliform; lower pinnae moderately reduced.

Davallodes gymnocarpum, Copel., l. c., tab. V: Indusium broader than long, attached by the base (as in *Humata*): lower pinnae gradually much reduced.

Philippines, Celebes, Borneo.

(4) **D. Korthalsii**, *Mett.*, Linn., XXXVI, 142.

Rhizome probably creeping. Fronds \pm 45 c.M. long, ovate, acuminate, 4-pinnatifid; main rachis stramineous, canaliculate at the upper side. Pinnae petiolate, trapezoidal or oblong-lanceolate, the apex curved and caudato-acuminate. Pinnulae short-stalked, trapezio-ovate or oblong, the apex short, narrowed, obtuse. Ultimate divisions trapezio-oblong, obtuse, pinnatifid at the base. Lobes spatulate-cuneate, with the upper edge crenate. Texture thinly chartaceous, firm; veins thickened at the apex, covered with a calcareous scale on the upper surface of the frond. Sori unknown.

Sumatra, Singapore.

* (5) **D. Griffithiana**, *Hk.*, Hk. Bk., Syn. Fil., 96; Hk., Spec. Fil., I, tab. XLIX, B.

Rhizome stout, the scales dense, pale-brown or whitish, linear. Stipes erect, wiry, 10—20 c.M. long. Fronds 20—50 c.M. long, 10—20 c.M. broad, deltoid, 5—4-pinnatifid. Pinnulae of the lower pinnae lanceolate-deltoid, 5—7½ c.M. long, 2½ c.M. or more broad. Lower segments toothed in the barren fronds, cut down nearly to the rachis in the fertile ones. Texture coriaceous. Sori large; indusium half-cupshaped, \pm as long as broad, very shortly attached on the sides nearly as in *Humata*, the apex whether or not reaching the margin.

Malacca; North. India, China, Formosa.

(6) **D. Lobbiana**, *Moore*, Hk. Bk., Syn. Fil., 94.

Fronds 30—45 c.M. or more long, 15—25 c.M. broad, lanceolate-deltoid, 2—5-pinnatifid; main rachis slightly winged above. Lower pinnae 10—15 c.M. long, \pm 5 c.M. broad, lanceolate-deltoid, acuminate, cut down to the rachis in the lower part into linear- or deltoid-oblong, blunt segments, only the lowest of which are deeply toothed. Sori copious, placed obliquely in 2 long rows in the pinnulae near the edge; indusium broader than deep.

Borneo.

(7) **D. stenoloba**, *Bk.*, Becc., Mal., III, 55; *D. corniculata*, *Ces.* (not *Moore*), Fel. Becc. Polin., 5.

Rhizome wide-creeping, stout, the scales minute, lanceolate, castaneous. Stipes slender, brown, naked, \pm 50 c.M. long. Fronds lanceolate-deltoid, 5-pinnatifid, \pm 45 c.M. long. Pinnae sessile, acuminate, the lowest the longest, lanceolate-deltoid, the highest lanceolate. Lower pinnulae lanceolate, sessile, deeply pectinato-pinnatifid. Lobes distant, erecto-patent, inequilateral, lanceolate. Texture firm; surfaces naked; rachis sparingly scaly; veins 1 to each lobe. Sori small, subterminal; indusium firm, cuneate, as long as broad.

Amboyna.

* (8) **D. pallida**, *Mett.*, Hk. Bk., Syn. Fil., 469; Hk., Icon. Pl., XVII, tab. MDCXXIV; *D. Mooreana*, *Masters*, Gard. Chron., 1869, 964, with figure; *D. Beccariana*, *Ces.*, Fel. di Bor., 14, tab. III, fig. 6—6c.

Rhizome thick, wide-creeping, the scales lanceolate, dark-brown. Stipes 50—45 c.M. long, naked, stramineous. Fronds 60—90 c.M. long, subdeltoid, 4-pinnatifid. Pinnae deltoid, lowest largest, 15—20 c.M. broad, produced on the lower side. Pinnulae and tertiary divisions deltoid, stalked, obliquely cuneate at the base; ultimate lobes obovate-cuneate, blunt, $2\frac{1}{2}$ —4 m.M. broad, bearing the sorus on the upper side at the base. Texture firm-membranaceous; surfaces naked; veinlets immersed. Sori immersed, prominent on the upper surface; indusium half-cupshaped or suburceolate, submarginal, the apex subrotund.

Borneo, Mindanao, New Guinea; Aneitum, Samoa.

* (9) **D. denticulata**, *Mett.*, C. Chr., Ind. Fil.; *D. elegans*, *Sw.*, Rac., Flor. Blz., I, 159; Christ, Farnkr. d. Erde, fig. 949; Bedd. Ferns S.I., tab. XVIII; Hk., Spec. Fil., I, tab. XLIII, B; *Adiantum denticulatum*, *Burm.*, (oldest name).

Rhizome stout, creeping; scales dense, linear-lanceolate, ferruginous, the base \pm dilated, peltate, the point long, linear-subulate, crisped, interlaced, short-ciliate, giving the rhizome a woolly appearance. Stipes firm, naked, 10—50 c.M. long. Fronds 50—60 c.M. long, 15—40 c.M. broad, deltoid to ovate-lanceolate, 5—4-pinnatifid; main rachis slightly winged towards the apex. Pinnulae of the lower pinnae \pm $2\frac{1}{2}$ c.M. broad, deltoid-lanceolate, cut down to the rachis in the lower part with oblong-deltoid, more or less deeply toothed segments, which are obliquely cuneate at the base. Texture coriaceous; surfaces naked; venation prominent, irregular, with spurious veins be-

tween the true ones. Sori in the teeth of the \pm narrowed and more deeply toothed fertile segments, several to a segment, with 1—2 sharp horns projecting beyond them; indusium half-cupshaped, \pm as broad as long, with the apex reaching the margin of the frond.

Trop. Asia, Australia & Polynesia, Madagascar, Fernando-Po.

* (10) **D. epiphylla**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 137; *D. elegans*, *Sw.*, *var. pulchra* (not *bidentata*), *Hk.*, *Spec. Fil.*, I, tab. XLIII, A; *D. corniculata*, *Moore*, *C. Chr.*, *Ind. Fil.*

Rhizome wide-creeping; scales linear-lanceolate, the base peltate, the edge entire. Stipes distant, 10—50 c.M. long, erect firm. Fronds 20—60 c.M. long, 15—40 c.M. broad, deltoid to broad-lanceolate, 5—5-pinnatifid; main rachis hardly at all winged. Pinnulae of the lowest pinnae lanceolate, 1—4 c.M. long, 2—10 m.M. broad. Segments narrow, sharply toothed, mucronate. Texture coriaceous; veins not immersed, 1—3 carried into each tooth, with spurious veins between the true ones. Sori small, with the sharp mucro of the tooth extending beyond; indusium half-cupshaped, semi-orbicular, \pm as long as broad. — Scarcely different from *D. denticulata*, *Mett.*

Malaya.

* (11) **D. decurrens**, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 95; *Hk.*, *Spec. Fil.*, I, tab. XLIV, B.

Rhizome stout, creeping, densely fibrillose. Stipes 10—15 c.M. long, stout, erect. Fronds 30—65 c.M. long, 15—45 c.M. broad, deltoid, 5-pinnatifid; main rachis hardly at all winged at the apex. Pinnulae of the lower pinnae lanceolate-acuminate, 7 $\frac{1}{2}$ —10 c.M. long, 2—3 c.M. broad, cut down throughout within a short distance of the costulae with broadly toothed, linear-oblong segments. Texture subcoriaceous. Sori 1 to each tooth, the blunt teeth slightly projecting beyond them; indusium half-cupshaped, as long as broad, not reaching the margin.

Sumata, Celebes, Philippines.

* (12) **D. elata**, *Spr.* (not *Sw.*), *Hk.*, *Spec. Fil.*, I, 166, tab. LV, A; *D. denticulata*, *Mett.*, *var.*, *C. Chr.*, *Ind. Fil.*

Stipes firm, naked. Fronds \pm 40 c.M. long or longer, deltoid to lanceolate-triangular, 3—4-pinnate. Pinnae erecto-patent, stalked. Ultimate divisions ovate-lanceolate, the edge crenate, inciso-serrate or pinnatifid with entire

or crenate, oblong lobes. Texture coriaceous; main rachis slightly winged towards the apex, secondary and tertiary rachises narrowly winged; surfaces more or less distinctly striated or marked with lines or spurious veins between the true veins. Sori in the teeth of the mostly narrower fertile segments, several to a segment, with 1—2 horns projecting beyond them; indusium \pm orbicular or elliptical, the both ends rounded or roundish-deltoid, the apex reaching the margin of the segment.

Tropical Asia, Polynesia, Australia.

(15) **D. Lorrainii**, *Hance*, Hk. Bk., Syn. Fil. 469; Bedd., Ferns S.Br.I., Suppl., tab. CCCLI.

Rhizome thick, clothed with linear-subulate, nearly black, densely grey-ciliated scales. Stipes $7\frac{1}{2}$ —10 c.M. long, naked, brownish. Fronds 15—50 c.M. long, deltoid, 4-pinnatifid. Pinnae stalked, deltoid, lowest largest, produced on the lower side, their rachises winged to base. Pinnulae and segments subsessile, crowded, deltoid, \pm reduced on the lower side; final lobes ligulate, $\frac{1}{2}$ —1 m.M. broad, with sorus at base of inner side. Texture subcoriaceous, surfaces naked; barren lobes 1-veined. Indusium subcylindrical.

Malacca.

(14) **D. pyxidata**, *Cav.*, Hk. Bk., Syn. Fil., 96; Hk., Spec. Fil., I, tab. LV, C; Bau. & Hk., Gen., Fil., tab. XXVII.

Rhizome stout, creeping, densely clothed with pale-brown, linear scales. Stipes erect, strong, 10—15 c.M. long. Fronds 20—45 c.M. long, 15—25 c.M. broad, deltoid, 3—4-pinnatifid. Pinnulae of the lower pinnae lanceolate, 5—8 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, with deltoid or oblong segments, the lowest of which are cut down nearly to the rachis. Texture coriaceous. Sori in the teeth, with a broad space outside them, which projects like a horn beyond them; indusium half-cupshaped.

Kei Islands; Australia.

* (15) **D. bullata**, *Wall.*, Hk. Bk., Syn. Fil., 97; Hk., Spec. Fil., I, tab. L, B; Christ, Farnkr. d. Erde, fig. 955; Bedd., Fern S.I., tab. XVII.

Rhizome creeping, stout; the scales light-brown or whitish, the base round, peltate, the apex long, filiform, hair-pointed, short-ciliate, \pm recurved. Stipes strong, $7\frac{1}{2}$ — $12\frac{1}{2}$ c.M. long, erect. Fronds 15—50 c.M. long, 10—20 c.M. broad, \pm deltoid, 3—4-pinnatifid. Pinnulae of the lower pinnae lanceolate, 5—8 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, with deeply inciso-pinnatifid, oblong rhom-

boidal segments. Texture coriaceous; surfaces naked. Sori occupying the greater part of the teeth in which they are placed, with usually a large horn or 1—2 teeth projecting beyond; indusium deeply half-cupshaped, more long than broad, the apex \pm reaching the margin.

Trop. Asia, China, Japan.

* (16) **D. fructuosa**, *Christ*, Warb. Mons., I, 86.

Scales of the rhizome filiform, pale-brown, long, spreading. Stipes naked, stramineous, \pm 15 c.M. long. Fronds deltoid, \pm 25 c.M. long, \pm 18—24 c.M. broad, 4-pinnatifid. Pinnae 10—12, the lower ones ovate-acuminate, the rachis narrowly winged. Pinnulae deltoid-ovate, spreading, pinnate, with the divisions pinnatifid to the rachis; segments narrow, linear, bearing the sori on the \pm dilated apices. Texture coriaceous. Indusium half-cupshaped to semicylindrical, opaque, slightly emarginate, narrowly winged by the edge of the segments, with a short tooth beyond. — Near *D. fijiensis*, *Hk.*

Java.

(17) **D. trichomanoides** *Bl.*, *Hk. Bk.*, *Syn. Fil.*, 95; *D. stenomera*, *Kze.*, *Bot. Zeit.*, 1848, 216.

Rhizome wide-creeping, the scales linear, squarrose, pale-brown. Stipes 5—10 c.M. long, grey-stramineous, naked. Fronds deltoid, 4-pinnatifid, 15—25 c.M. long, acuminate. Pinnae short-stalked, lanceolate, lowest largest, deltoid-lanceolate, 5—5 c.M. broad, rachis winged. Pinnulae close, lanceolate. Ultimate segments (teeth) cuneate-oblong when barren, spatulate when fertile and forming a narrow wing on each side of the sorus. Texture \pm coriaceous; surfaces naked. Sori terminal on the lobe-like teeth; indusium rather deeper than broad, urceolate, with mostly a horn beyond on both sides.

Malaya.

(18) **D. Wagneriana**, *Copel.*, *Polypod. Philipp.*, 54.

Rhizome stout, scandent, the scales dense, lanceolate, acuminate, adpressed, brown. Stipes stout, scaly near the base. Fronds 20—25 c.M. long, 10—15 c.M. broad, 5-pinnatifid at the base only, the fertile narrower than the sterile ones. Lowest pinnae not larger than the succeeding, with a few, distinct, deeply toothed, equal pinnulae at the base of each, followed by broadly linear, acutely toothed segments. Texture coriaceous; rachis winged, naked like the surfaces. Sori usually a little longer than broad, rarely reaching the margin.

Mindanao.

* (19) **D. divaricata**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 138; *D. polyantha*, *Hk.*, *Spec. Fil.*, I, tab. LIX, A; *Bedd.*, *Ferns Br.I.*, I, tab. CVII.

Rhizome stout, creeping, the scales linear-lanceolate, dense, ferruginous or pale-brown, with the base truncate, not peltate, the edge entire or beset with short, spreading teeth or ciliae, the apex acuminate or acute, not adpressed, deciduous. Stipes firm, erect, 15—50 c.M. long, naked. Fronds 60—90 c.M. long, 3-pinnatifid to 4-pinnate. Lower pinnae often 50 c.M. long by 15 c.M. broad. Ultimate segments linear-oblong, toothed. Texture coriaceous; surfaces naked. Sori numerous, 1 to each tooth of the ultimate segments; indusium half-cupshaped, often $\pm 1\frac{1}{2} \times$ as long as broad, the apex at some distance ($\pm \frac{1}{2}$ —1 m.M.) from the margin of the segment.

Trop. Asia.

* (20) **D. solida**, *Sw.*, *Rac.*, *Flor. Btz.*, I, 157; *Hk.*, *Spec. Fil.*, I, tab. XLII; *Hk.*, *Fil. Exot.*, tab. LVII; *Bedd.*, *Ferns Br.I.*, I, tab. CIV; *D. elegans*, *Hedw.*, *Fil. Gen. & Spec.*, 1803; — var. *caudata*; *D. caudata*, *Cav.*, *Descr.*, 1802, 279; — var. *Lindleyi*; *D. Lindleyi*, *Hk.*, l. c., 165, tab. LVIII, B; — var. *ornata*; *D. ornata*, *Wall.*, *List*, No. 246.

Rhizome stout, wide-creeping; scales adpressed, black, white when young, the base roundish, peltate, persistent, the apex finely acuminate, the margin beset with spreading, cobwebby, deciduous, adpressed, interlaced fibres altogether forming a dense, flat coat to the rhizome. Stipes distant, strong, erect, 10—20 c.M. long, naked. Fronds ovate-deltoid, 15—60 c.M. long, 15—40 c.M. broad, 5-pinnatifid to -pinnate, the apex with a moderately broad, undivided centre. Ultimate segments ovate-rhomboidal, $\pm 5 \times$ as long as broad, deeply toothed, narrower and sharper in fertile fronds. Texture coriaceous; surfaces naked. Sori in the teeth, sometimes with 1—2 horns beyond; indusium semicylindrical, $\pm 2 \times$ as long as broad, the apex nearly or quite reaching the edge of the frond.

Var. caudata: Ultimate segments acuminate to caudate.

Var. Lindleyi: Pinnulae more divided, with narrower segments.

Var. ornata: Ultimate segments broad, slightly cut.

Malaya; Polynesia, Queensland.

(21) **D. embolostegia**, *Copel.*, *Philipp. Journ.*, I, Suppl., 147, tab. III.

Rhizome creeping, densely clothed with ferruginous, brunnescent, linear scales. Stipes distant, 50—50 c.M. long, naked, glossy. Fronds deltoid,

FILICES.

60—100 c.M. long, acuminate, 4—5-pinnatifid. Pinnae and lower pinnulae long-acuminate, deltoid-ovate, the higher ones narrower. Ultimate segments lanceolate or oblong, serrate or incised. Texture hardly coriaceous; surfaces \pm naked; penultimate rachises winged; no spurious veins. Sori flanked mostly by 1—2 teeth; indusium semicylindrical, conspicuously rostrate. — Fronds resembling *D. solida*, Sw. & *divaricata*, Bl.

Luzon, Mindoro.

(22) ***D. Veitchii*, Bl.**, Journ. of Bot., XVII, 39.

Rhizome wide-creeping, the scales small, shining, linear, acuminate, castaneous. Stipes naked, erect, slender, castaneous, 50—45 c.M. long. Fronds oblong-deltoid, decomposed, 50—45 c.M. long, \pm 15 c.M. broad. Pinnae distant, deltoid, stalked, erecto-patent, many of the lower ones subequal, 10—15 c.M. long by half as broad. Pinnulae and tertiary divisions subpetiolate, rhomboidal, the base cuneately cut away on the lower side. Ultimate lobes subremote, ligulate or rather broader towards the truncate apex, \pm 1—5½ m.M. long, \pm ½ m.M. broad, with only a single central vein, the apex denticulate. Texture subcoriaceous; surfaces naked. Sori small, oblong, usually immersed in the tips of the segments.

Borneo.

(23) ***D. brevipes*, Copel.**, Philipp. Journ., I, Suppl., 147, tab. II.

Rhizome creeping, densely clothed with fuscous, lanceolate, \pm adpressed scales, which are whitish at the apex and the edge. Stipes remote, \pm 1¾ c.M. long, naked, narrowly winged. Fronds deltoid, \pm 10 c.M. long, \pm 6 c.M. broad, the rachis furcato-alate on the upper side. Pinnae ascending, \pm straight, narrowly ovate, the lowest largest, broadly ovate, falcate, slightly deflexed. Pinnulae minute, cuneate, cut down to the costa into narrow, long-subulate, rather hair-like segments. Texture coriaceous; surfaces naked; no spurious veins. Sori on the ultimate segments, flanked by a horn on each side; indusium semicylindrical, marginal.

Mindanao.

(24) ***D. dissecta*, J. Sm.**, Rac., Flor. Btz., I, 159; Moore, in Gardn. Chron., 1855, 469, with figure.

Rhizome stout, wide-creeping, the scales brown or pale-brown, with the base roundish, peltate, adpressed, the apex long-acuminate, recurved, spreading, ciliate, at length deciduous, giving the rhizome a woolly appearance when

young. Stipes 5—15 c.M. long, naked. Fronds $12\frac{1}{2}$ —45 c.M. long, 10—30 c.M. broad, deltoid to lanceolate-triangular, 3—4-pinnatifid. Pinnae lanceolate, the lower deltoid, produced on the lower side. Pinnulae lanceolate-deltoid, subsessile, obliquely cuneate at the base. Segments oblong, cuneate at the base, deeply pinnatifid with erecto-patent, ligulate, simple or 2-fid lobes 5—5 m.M. long, 1—2 m.M. broad, those of the fertile fronds oblong, $\frac{1}{2}$ m.M. broad at the base, with a suddenly dilated, cuneate, soriferous apex. Texture \pm coriaceous; surfaces naked. Sori numerous, 1 to each ultimate lobe, commonly with 1—2 horns beyond; indusium nearly 2 \times as long as broad, the apex \pm reaching the margin of the segment.

Java, Sumatra.

(25) **D. fijiensis** (*fejensis*), **Hk.**, Hk. Bk., Syn. Fil., 97; Hk., Spec. Fil., I, tab. LV, D; *D. solida*, Sw., var., C. Chr., Ind. Fil.

Rhizome creeping, densely fibrillose. Stipes 15—25 c.M. long, erect, strong. Fronds 30—45 c.M. long, 15—30 c.M. broad, deltoid, 4-pinnatifid. Pinnulae of the lower pinnae deltoid-lanceolate, the lobes of the tertiary segments cut down nearly to the rachis into narrow, linear laciniae. Texture coriaceous. Sori semicylindrical, on the dilated apices of the laciniae, without horns; indusium long, 3—4 \times as long as broad.

New Guinea; Fiji.

35. MICROLEPIA, *Presl.*

Sori roundish or oblong, submarginal, solitary and terminal on the veins. Indusium half-cupshaped or semicylindrical, attached at the base and sides, open at the outer edge. Receptacle distinct.

Rhizome creeping, hairy or fibrillose. Stipes 2-seriate, adherent to the rhizome. Fronds variously divided; ultimate free divisions rhomboidal-oblong or linear-triangular, broadest \pm at the base, the upper base rather broadly and the lower rather narrowly cuneate; veins free, simple or branched. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 116, A; Bedd., Handb. Ind. Ferns, fig. 53 (*).

Pantropical.

(*) Copeland thinks that **Microlepia** would better be abandoned as a genus and regarded as a section of **Dennstaedtia** (Philipp. Journ., II, 126).

- a.** *Fronds simply pinnate or at best 2-pinnatifid.*
- ‡. *Rhizome erect. Fronds rather small, naked.* (1) *M. firmula.*
- ‡‡. *Rhizome creeping. Fronds rather large, hairy beneath.* (2) *M. marginalis.*
- a.a.** *Fronds at least 2-pinnate at the base.*
- b.** *Texture membranaceous or herbaceous.*
- c.** *Indusium naked.*
- ×. *Fronds small, ± 25 c.M. broad.* (3) *M. scaberula.*
- ×.×. *Fronds large, ± 90 c.M. broad.* (4) *M. manilensis.*
- c.c.** *Indusium hispid, hairy or bristly.*
- d.** *Receptacle rather small, not exerted.*
- e.** *Ultimate lobes entire or crenato-lobate. The hairs of the under surface bulbous at the base.* (5) *M. trichosticha.*
- e.e.** *Ultimate segments toothed or pinnatifid. The hairs of the under surface not bulbous at the base.* (6) *M. speluncae.*
- d.d.** *Receptacle prominent, exerted.* (7) *M. exserta.*
- b.b.** *Fronds more or less coriaceous.*
- 1.** *Surfaces naked.*
- ×. *Small. Pinnulae crenato-lobate.* (8) *M. philippinensis.*
- ×.×. *Large. Pinnulae deeply pinnatifid.* (9) *M. platyphylla.*
- 1.1.** *At least the under surface hairy or strigose.*
- 2.** *Pinnulae crenate or broadly toothed.* (10) *M. strigosa.*
- 2.2.** *Pinnulae (at least the lower ones) ± deeply pinnatifid.*
- 3.** *Tertiary segments bluntly serrate.* (11) *M. Kurzii.*
- 3.3.** *Tertiary segments entire or broadly crenate.*
- 4.** *Stipes naked.* (12) *M. proxima.*
- 4.4.** *Stipes pubescent.* (13) *M. trapeziformis.*
- 2.2.2.** *Pinnulae (at least the lower ones) pinnate.* (14) *M. hirta.*

(1) **M. firmula**, **C. Chr.**, Ind. Fil.; *Davallia firmula*, **Bk.**, Ann. of Bot., VIII, 125.

Stipes tufted, 20—25 c.M. long, slender, firm, naked above, scaly at the base, stramineous, the scales lanceolate, membranaceous, ferruginous. Fronds deltoid, 20—25 c.M. each way, simply pinnate. Pinnae sessile, ∞-jugate,

linear, often entire, with the base cuneate, the lower ones 10—15 c.M. long, $\pm 7\frac{1}{2}$ m.M. broad. Texture firm; surfaces naked like the stramineous rachis; veins not close, erecto-patent, deeply forked, not reaching the margin. Sori small, terminal on the veins; indusium semicupuliform, naked, persistent.

Sumatra.

(2) **M. marginalis**, *Bedd.* (not *Hance*), *Christ, Farnkr. d. Erde*, 507; *M. scabra*, *J. Sm., Bedd., Ferns Br.I., I, tab. CII*; *M. marginata*, *C. Chr., Ind. Fil.*; *Davallia marginalis*, *Bk., Hk. Bk., Syn. Fil., 98*; *D. villosa*, *Wall., Hk., Spec. Fil., I, tab. XLVIII, A.*

Rhizome creeping, villose. Stipes 50—60 c.M. long, erect, strong. Fronds 45—75 c.M. long, 20—55 c.M. broad, pinnate. Pinnae linear, distant, alternate, slightly stalked, straight or subfalcate, $\pm 2\frac{1}{2}$ c.M. broad at the base, narrowed gradually towards the apex, the edge cut $\pm \frac{1}{3}$ — $\frac{2}{3}$ -way down into numerous, close, deltoid-oblong, crenato-dentate lobes, the upper base broadly and the lower rather narrowly cuneate. Texture subcoriaceous; rachis and stipe downy, the latter at length glabrous; under surface hairy. Sori submarginal, 2—8 to a lobe, 1 to a serrature; indusium broadly half-cupshaped, densely hairy.

New Guinea; Japan, China, Ceylon.

(5) **M. scaberula**, *Mett., Linn., XXVI, 148.*

Fronds ± 30 c.M. long, ± 25 c.M. broad, deltoid, acuminate, sub-5-pinnatisect. Pinnae alternate, not close, spreading, short-stalked, acuminate. Pinnulae approximate, spreading, sessile, trapezio-oblong-lanceolate, unequal-sided, the upper base truncate, auricled, the lower cuneate. Segments oblong, obtuse, pinnatifid, incised or crenate with a few teeth at the apex. Texture thinly chartaceous; costae and veins beneath covered with subadpressed, reddish-brown, small, sharp bristles; rachis short-hairy; veins prominent beneath. Sori submarginal, small; indusium naked.

New Guinea.

(4) **M. manilensis**, *Pr., Tent. Pterid., 125*; *Davallia manilensis*, *Hk., Ann. Bot., XV, 93, tab. XIV, fig. 9, a—b*; *D. Sarasinorum*, *Christ, Fil. Saras., IV, 7.*

Fronds large, to ± 2 M. long, deltoid, 2-pinnate to 5-pinnatifid. Pinnae 40—50 c.M. long, oblong, 15—18 c.M. broad at the base, the apex long-caudate. Pinnulae numerous, 25—50 on each side below the pinnatifid apex

of the pinnae, alternate, the lower ones 8—10 c.M. long, \pm 3 c.M. broad at the base, caudate, deeply pinnatifid, with the upper base truncate, auricled, the lower cuneate, the auricle sometimes free; higher pinnulae close, cut $\frac{1}{2}$ -way down to the costa or more into numerous, close, rhomboidal, blunt, crenate lobes. Lobes to 1 c.M. long, with 4—8 crenations on each side. Texture firm-herbaceous; rachises and surfaces covered with short, brown hairs. Sori numerous, large, immersed, slightly intramarginal below the teeth or crenations; indusium grey.

Java, Celebes, Philippines.

(5) **M. trichosticha**, **J. Sm.**, Journ. of Bot., III, 416; *Davallia trichosticha*, **Hk.**, Spec. Fil., I, 185; *D. Speluncae*, **Bk.**, var. *trichosticha*, **Hk.** **Bk.**, Syn. Fil., 470.

Fronds large, 2—5-pinnate. Pinnae \pm 50 c.M. long, the rachis winged above. Ultimate segments sessile, obliquely cuneate at the base, lanceolate-oblong, acuminate, deeply pinnatifid, with the lowest anterior lobe the largest. Lobes ovate, obtuse, nearly entire or crenato-lobate. Texture submembranaceous; upper surface slightly hairy, under surface densely and minutely pubescent, the hairs bulbous at the base, almost concealing the veins. Sori small, rather distant from the margin; indusium small, semicupuliform, very downy.

Malaya; Samoa

* (6) **M. speluncae**, **Moore**, Christ, Farnkr. d. Erde, 308; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 116, A—C; *Davallia Speluncae*, **Bk.**, Rac., Flor. Btz., I, 155; *D. polypodioides*, **Don**, var. *subglabra* & *pubescens*, **Hk.**, Spec. Fil., I, 182; *D. flaccida*, **R. Br.**, Prod. N.H., 157; — var. *immersa*.

Rhizome creeping, more or less densely clothed with pale or brownish hairs or fibres. Stipes strong, 20—100 c.M. long, stramineous, brownish or purpleous, minutely hairy or squamulose. Fronds 40—200 c.M. long, 20—100 c.M. broad, triangular or ovate-lanceolate, 5—5-pinnatifid. Lower pinnae ovate-lanceolate. Pinnulae lanceolate, cut down to a winged rachis into toothed or pinnatifid lobes, the lowest lobe on the upper side the largest; lobes and teeth \pm rounded. Texture herbaceous, membranaceous or flaccid; rachis and surfaces more or less hairy or setose, or with few or many, long, glistening, flaccid, scale-like hairs; veins conspicuous. Sori at the base of the teeth or lobes, 2—20 to a segment; indusium half-cupshaped, hispid or hairy, rarely glabrous.

Var. immersa: Sori immersed, distinctly prominent on the upper surface.

Tropical and subtropical.

(7) **M. exserta**, *Mett.*, Linn., XXXVI, 148; *Davallia Speluncae*, *Bk.*, var. *exserta*, *Hk. Bk.*, Syn. Fil., 470.

Fronds large, 4-pinnatisect. Pinnae 50 c.M. long or longer, acuminate. Pinnulae short-stalked, 5—8 c.M. long, trapezio-oblong, acuminate, or the apex narrow and obtuse. Ultimate divisions trapezio-oblong, obtuse, unequal-sided, the anterior base the broadest. Lobes contiguous, oblong, oblique, obtuse, the higher ones crenate, the lower pinnatifid. Texture membranaceous; upper surface naked or nearly so, under surface pubescent; main rachis clothed beneath which a dense, pale-brown tomentum. Sori submarginal, subapproximate, 1 or more to a lobe, placed near the sinuses; indusium membranaceous, purveyed with some few bristles; receptacle prominent, exserted.

? *Malaya; Fiji.*

(8) **M. philippinensis**, *Copel.*, Polypod. Philipp., 56; *Davallia philippinensis*, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 27.

Rhizome thickly clothed with narrow, reddish-brown scales. Stipes \pm 15 c.M. long, dark-brown, with a few scales below like those of the rhizome, naked above. Fronds deltoid, 7—25 c.M. long, nearly as broad; rachis naked. Lower pinnae deltoid, pinnate, the higher ones simple, those at the apex confluent. Pinnulae linear, crenato-lobate, the lobes extending $\frac{1}{2}$ -way to the costa, lowest pinnulae on the lower side much larger than the other ones. Texture coriaceous; surfaces naked; veins ascending. Indusium intramarginal, broader than deep.

Philippines.

(9) **M. platyphylla**, *J. Sm.*, Christ, Farnkr. d. Erde, 507; *Bedd.*, Ferns S.I., tab. XIII; *Davallia platyphylla*, *Don*, *Hk. Bk.*, Syn. Fil., 99; *D. lonchitidea*, *Wall.*, *Hk.*, Spec. Fil., I, tab. XLVI, B; *Fil. Exot.*, tab. XIX.

Rhizome stout. Stipes firm, erect, 60—90 c.M. long. Fronds 90—120 c.M. long, 5-pinnatifid. Lower pinnae 50—40 c.M. long, 15—25 c.M. broad. Pinnulae distant, linear, the higher ones adnate, the lower narrowed gradually from the base towards the apex, the edge cut $\pm \frac{1}{3}$ — $\frac{2}{3}$ -way down to the costa into many, bluntish, toothed, oblong-deltoid lobes, the upper base broadly, the lower rather narrowly cuneate. Texture subcoriaceous; surfaces naked.

Sori 2—12 to a lobe, 1 to a crenation, a short distance from the edge; indusium naked.

Philippines; Ceylon, North. India.

* (10) **M. strigosa**, *Pr.*, Christ, Farnkr. d. Erde, 508; Bedd., Ferns S.I., tab. CCLV; *Davallia strigosa*, (*Sw.*) *Kze*, *Rac.*, *Flor. Btz.*, I, 135.

Rhizome creeping, hairy. Stipe erect, strong, 15—40 c.M. long, both it and the rachis pubescent throughout. Fronds ovate-lanceolate, acuminate, 30—100 c.M. long, 15—50 c.M. broad, 2-pinnate. Pinnae short-stalked, erecto-patent, linear-lanceolate, acuminate, straight or falcate, the lowest longest, 10—20 c.M. long, \pm 2 c.M. broad, cut down to the rachis with numerous close pinnulae. Pinnulae rhomboidal, obliquely cuneate at the base, the lowest largest, oblong, broadly and rather bluntly toothed or crenated. Texture subcoriaceous; rachises and veins beneath more or less copiously hairy; veins beneath prominently raised. Sori small, 2—12 to a pinnula, placed near the bases of the crenations at a short distance from the sinuses; indusium half-cupshaped, hairy.

Trop. Asia & Polynesia, Japan, China.

(11) **M. Kurzii**, *Bedd.*, *Hand. Ind. Ferns*, 66; *Davalia Kurzii*, *Clarke*, *Trans. Linn. Soc., Bot.*, 2^d Series, I, 446.

Fronds large, 5-pinnatifid. Lower pinnae 30—40 c.M. long, 15—25 c.M. broad, lanceolate, with linear-lanceolate pinnulae, which are cut down to the rachis (which is not wavy, or only slightly so towards the apex) below into triangular, subobtusate or lanceolate-caudate lobes, which are bluntly serrate. Texture more or less coriaceous; under surface pubescent even when adult; venation moderately prominent. Indusium finely ciliated round the margin. — Near *M. platyphylla*, *J. Sm.*

Malacca; Burmah.

* (12) **M. proxima**, *Pr.*, *C. Chr.*, *Ind. Fil.*; *Davalia proxima*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 154.

Stipes 30 c.M. or more long, naked. Fronds 100 c.M. or more long, to 80 c.M. broad, 2-pinnate; rachis rough-hairy. Pinnae stalked, to 40 c.M. long, to 10 c.M. broad, lanceolate, the lower ones shorter than the following, the stalks 1—1½ c.M. long. Pinnulae short-stalked, lanceolate, inequilateral, to 6 c.M. long, to 2 c.M. broad, toothed at the apex, the base deeply pinnatifid. Segments rounded, entire, the lower to 15 m.M. long, to 6 m.M. broad,

dark-green above, light-green below. Texture subcoriaceous; upper surface short-haired on the veins, the under one covered with pale hairs. Sori on a short distance from the edge; indusium very small, semiorbicular, covering only the base of the sorus when mature, slightly hairy.

Malaya.

(13) **M. trapeziformis**, *Kuhn*, C. Chr., Ind. Fil.; *M. rhomboidea*, *Pr.*, Tent. Pterid., 125; *M. polypodioides*, *Bedd.*, Ferns S.I., tab. XV; *Davallia rhomboidea*, *Wall.*, Bot. Zeit., 1850, 158; *D. strigosa*, *Sw.*, var. *rhomboidea*, *Hk. Bk.*, Syn. Fil., 99; *D. polypodioides*, *Don*, var. *hispida* & *rhomboidea*, *Hk.*, Spec. Fil., I, 182; *D. trapeziformis*, *Rb.*, (oldest name).

Near *M. strigosa*, *Pr.*, but rather larger in all parts. Rachis downy or hispid beneath. Lower pinnulae lanceolate-deltoid, cut down nearly to the rachis into oblong, entire or irregularly inciso-lobate or again pinnatifid, sub-oblique, very obtuse lobes. Sori rather large, usually solitary on the entire lobes, several on the pinnatifid ones, near the sinuses, within the margin; indusium half-cupshaped, naked or hispid.

Trop. Asia.

* (14) **M. hirta**, *Pr.*, Tent. Pterid., 125; *Bedd.*, Ferns S.I., tab. CCLVI; *Davallia hirta*, *Klf.*, *Hk. Bk.*, Syn. Fil., 100.

Stipes strong, 50—60 c.M. long, deciduously short-hairy. Fronds 45—180 c.M. long, 50—60 c.M. broad, ovate-deltoid, subcaudate, 5-pinnate. Lower pinnae erecto-patent, 7—10 c.M. broad, triangular-lanceolate. Pinnulae triangular-lanceolate; tertiary segments rhomboidal-oblong, the lowest largest, blunt, broadly toothed or deeply pinnatifid, the base obliquely cuneate, slightly decurrent. Texture coriaceous; rachises and especially the under surface densely pubescent. Sori 2—12 to a tertiary segment, 1 or more to each ultimate tooth or lobe; indusium \pm hairy or covered by the hairs of the veins. — Stature and habit of *M. speluncae*, *Moore*, but firmer.

Trop. Asia; Hawaii.

36. TAPEINIDIUM, *Christensen*.

Sori roundish, terminal on the veins, submarginal, single or in \pm confluent pairs. Indusium broadly half-cupshaped, attached at the base and sides, open at the outer edge. Receptacle indistinct.

Rhizome creeping, minutely scaly or fibrillose. Stipes 2-seriate, \pm distant or approximate, adherent to the rhizome. Fronds variously divided; leaflets narrow, linear-lanceolate or divided into narrow, oblong or linear segments; veins free, simple or forked. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 116, J & H. (*Wibelia*).

Malaya, Polynesia.

a. Fronds at best 2-pinnate.

b. Fronds small, under 10 c.M. long. (1) *T. oligophlebium*.

b.b. Fronds larger, more than 10 c.M. long.

c. Pinnae slightly crenate or toothed. (2) *T. pinnatum*.

c.c. Pinnae deeply toothed, pinnatifid or pinnate. (3) *T. gracile*.

a.a. Fronds at least deeply 3-pinnatifid at the base.

*. Fronds rather small, 3-pinnatifid or 3-pinnate, coriaceous.

(4) *T. amboinense*.

.. Fronds rather large, 4-pinnatifid, herbaceous.

(5) *T. Denhami*.

* (1) ***T. oligophlebium***, *C. Chr.*, Ind. Fil.; *Davallia oligophlebia*, Bk., Journ. of Bot., XXVI, 523; *Wibelia oligophlebia*, Christ, Ann. Btz., XX, 154.

Rhizome slender, short-creeping, the scales minute, adpressed, linear, brown. Stipes slender, naked, deciduously scaly near the base, \pm 1 $\frac{1}{2}$ —7 $\frac{1}{2}$ c.M. long. Fronds triangular, \pm 5—7 $\frac{1}{2}$ c.M. long, 4—9 c.M. broad at the base, pinnate, the lower pinnae \pm spreading, pinnate again, \pm 2 c.M. broad at the base. Ultimate segments oblong to linear, 5—20 m.M. long, 1 $\frac{1}{2}$ —2 $\frac{1}{2}$ m.M. broad, the base cuneate, the edge, crenate or bluntly toothed to lobed, the point narrowed, bluntish. Texture subcoriaceous; rachis and surfaces naked; veins 1 to each tooth or lobe. Sori terminal on the crenations, 1 to each; indusium firm naked, suborbicular, small. — The description is taken from specimens gathered by Hallier, determined by Christ. — Baker's description: »Fronds pinnate, 3 $\frac{1}{2}$ —5 c.M. long, 8—9 m.M. broad, with the pinnae rhomboid, obtuse, 3-lobed,» is probably taken from a young specimen.

Borneo.

* (2) ***T. pinnatum***, *C. Chr.*, Ind. Fil.; *Davallia pinnata*, Cav., Rac., Flor. Btz., I, 153; Christ, Farnkr. d. Erde, fig. 967: *D. longipinnula* &

intramarginalis, Ces., Fel. Becc. Polin., 7 & 8; *D. flagellifera*, Wall., Hk. & Grev., Ic. Fil., tab. CCLXXXIII; *Microlepia pinnata*, J. Sm. (not Cav.), Copel., Polypod. Philipp., 55; Bedd., Ferns S.I., tab. XIV; *Wibelia pinnata*, Fée (not Bernh.), Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 116, J & H.

Rhizome creeping, densely clothed with subulate-lanceolate, fibrillose scales. Stipes erect, strong, 10—30 c.M. long, scaly towards the base. Fronds oblong, 15—40 c.M. long, 7—20 c.M. broad, simply pinnate with 8—20 short-stalked, erecto-patent pinnae on each side below the terminal one. Pinnae linear, 4—20 c.M. long, 3—10 m.M. broad, acuminate, the base cuneate, the edge slightly crenate or toothed. Texture coriaceous; rachis and surfaces naked. Sori 1—2 to each tooth, more broad than long, sometimes confluent.

Malaya; Polynesia.

* (3) **T. gracile**, v. A. v. R.; *T. pinnatum*, C. Chr., var., C. Chr., Ind. Fil.; *Davallia gracilis*, Bl., Rac., Flor. Btz., I, 154; *D. Luzonica*, Hk., Spec. Fil., I, tab. LX, B, fig. 2, 3 & 5; *Microlepia pinnata*, J. Sm., var. *gracilis*, Copel., Polypod. Philipp., 55; *M. pinnata*, J. Sm., var. *Luzonica*, Bedd., Handb. Ind. Ferns, Suppl., 15; *Wibelia gracilis*, Christ, Ann. Btz., XX, 154.

Rhizome creeping, densely clothed with small, brown, subulate-lanceolate, fibrillose scales. Stipes \pm 5—10 m.M. apart, 10—15 c.M. long, densely scaly at the base. Fronds oblong-lanceolate, 20—50 c.M. long, to 17 c.M. broad. Texture coriaceous; rachis and surfaces naked. Sori 1—2 to each tooth, the pairs often \pm confluent. — Two forms.

Forma grandidentata: Fronds simply pinnate. Pinnae linear, \pm 5—15 m.M. broad, deeply toothed or subpinnatifid, the teeth or lobes broad, obliquely deltoid, reaching down nearly to the costa.

Forma dissecta: Fronds pinnate. Pinnae \pm 2—5 c.M. broad, deeply pinnatifid nearly to the costa or pinnate again, the segments linear or oblong, entire or obliquely toothed, the base adnate; decurrent or cuneate.

Both forms may be found on the same rhizome.

Malaya.

* (4) **T. amboinense**, C. Chr., Ind. Fil.; *Davallia amboinensis*, Hk., Hk. Bk., Syn. Fil., 470; Hk., Spec. Fil., I, tab. LVI, C; *D. moluccana*, Rb., Wall., List, No. 2219; *Wibelia amboinensis*, Kuhn, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 216.

Rhizome creeping, the scales lanceolate-subulate, fibrillose, brown. Stipes

seriate, \pm 5—10 m.M. apart, 15—45 c.M. long, firm, erect, \pm scaly at the base. Fronds lanceolate-ovate, 15—35 c.M. long, deeply 3-pinnatifid or 3-pinnate at least at the base. Pinnae erecto-patent, the higher ones pinnate, the segments entire, toothed or subpinnatifid, lowest pinnae largest, with the lower pinnulae deeply pinnatifid or pinnate, those on the lower side often much produced. All the teeth and segments erecto-patent or ascending. Texture coriaceous; rachis and surfaces naked. Sori near the base on the inner side of the ultimate segments or teeth, minute, intramarginal.

Moluccas; Samoa.

(5) **T. Denhami**, *C. Chr.*, Ind. Fil.; *Davallia Denhami*, *Hk.*, *Hk. Bk.*, Syn. Fil., 100; *Hk.*, 2^d Cent. of Ferns, tab. XLVII; *Wibelia Denhami*, *Kuhn*, *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam. I⁴, 217.

Rhizome creeping, the scales subulate-fibrillose. Stipes slender, erect, 10—20 c.M. long. Fronds 30—60 c.M. long, 15—30 c.M. broad, 4-pinnatifid. Pinnae erecto-patent. Pinnulae of the lower pinnae erecto-patent, lanceolate, $5\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, \pm $2\frac{1}{2}$ c.M. broad, cut down to the rachis into oblong, toothed segments, the lower of which \pm $\frac{1}{2}$ c.M. long, \pm $\frac{1}{4}$ c.M. broad. Texture herbaceous; surfaces naked. Sori numerous, intramarginal, very small; indusium obpyriform, the outer edge rounded.

New Guinea; Polynesia.

TRIBE IV. PTERIDEAE.

37. ADIANTUM, *Linné.*

Sori roundish, obreniform, transversally oblong or linear, terminal on the veins, the sporangia springing from the veins only (*Eu-adiantum*), or placed both on the veins and the spaces between them (*Adiantellum*). Indusium of the same shape, marginal, consisting of the modified, reflexed margin, bearing the sori on its inner surface.

Rhizome creeping or erect. Stipes often black and polished, not articulate to the rhizome. Fronds erect or climbing (not in malayan material), simple or commonly variously divided; leaflets inclined to be dimidiate, but often cuneato-flabellate, the outer edge not rarely more or less deeply incised, the laciniae generally finely toothed; veins free or sometimes anastomosing. Ferns mostly known and cultivated under the name of »chevelure” or »maiden-hair”. Hk. Bk., Syn. Fil., tab. II, fig. 21; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 150 (*).

Tropical and subtropical; some species also in the temperate zones.

Arrangement of the sections and groups.

§ 1. **EU-ADIANTUM** (incl. *Adiantellum*). Veins free.

- a. **Reniformiae.** Fronds simple.
- b. **Oligosoriae.** Fronds pinnately divided. Leaflets not cuneato-flabellate. Sori linear, in continuous or interrupted lines.
- c. **Polysoriae.** Fronds pinnately divided. Leaflets not cuneato-flabellate. Sori transversally oblong, or sometimes obreniform.
- d. **Cuneatae.** Fronds pinnately divided. Leaflets cuneato-flabellate, ± oblique, the petiole near the centre at the base. Sori obreniform, or sometimes transversally oblong.
- e. **Pedatae.** Fronds pedato-dichotomous, i.e. forked, with the secondary branchings springing from the upper side of the 2 primary ones.

§ 2. **HEWARDIA.** Veins anastomosing. (Not in Malaya).

(*) With view on the many beautiful species of this genus, which are much beloved as ornamental plants, I inserted in this work the imported species (not the varieties got by cultivation), which I know are found in our indian gardens.

Key to the malayan groups. (§ 1. *Eu-adiantum*).

- a.* Fronds simple. § 1. a. **Reniformiae.**
- a.a.* Fronds divided.
- b.* Fronds forked, the primary branches pinnately branched on the upper (inner) side only. § 1. e. **Pedatae.**
- b.b.* Fronds varying from simply to more copiously pinnate.
- c.* Sori long, linear, continuous or interrupted. Leaflets dimidiate or unequal-sided. § 1. b. **Oligosoriae.**
- c.c.* Sori short, transversally oblong or obreniform.
- d.* Leaflets dimidiate or unequal-sided. § 1. c. **Polysoriae.**
- d.d.* Leaflets obliquely cuneato-flabellate, sometimes tending towards orbicular, or trapeziform- or rhomboidal-dimidiate. § 1. d. **Cuneatae.**

 § 1. EU-ADIANTUM. a. **Reniformiae.**

(1) **A. Parishii**, *Hk.*, Hk. Bk., Syn. Fil., 114; Hk., Spec. Fil., III, tab. CXLII, A; Fil. Exot., tab. LI; Bedd., Ferns Br.I., I, tab. XVI.

Stipes tufted, slender, naked, dark-brown, polished, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long. Fronds 1— $2\frac{1}{2}$ c.M. each way, suborbicular, slightly undulate, cuneate at the base. Texture papyraceo-herbaceous; veins radiant, not prominent. Sori few, placed in the crenations of the fronds.

Malacca; Burmah.

 § 1. EU-ADIANTUM. b. **Oligosoriae.**

*. *Fronds simply pinnate. Leaflets (pinnae) sessile or subsessile, rather opposite.*

(2) **A. macrophyllum.**

.. *Fronds once or more times pinnate. Leaflets (pinnae or pinnulae) distinctly stalked, rather alternate.*

(3) **A. platyphyllum.**

* (2) **A. macrophyllum**, Sw., Hk. Bk., Syn. Fil., 121; Hk., Fil. Exot., tab. LV; Hk. & Grev., Ic. Fil., tab. CXXXII; — var. *striatum*, Hort. Bog.

Stipes 15—30 c.M. long, firm, erect, polished, naked, blackish. Fronds 20—40 c.M. long, 10—20 c.M. broad, simply pinnate with \pm 4—6 pairs of \pm opposite lateral pinnae below the terminal one. Lower pinnae (leaflets) of the barren fronds 7—10 c.M. long, \pm 5 c.M. broad, so broad at the base that the opposite ones frequently overlap; all the pinnae sessile or the lower ones very short-stalked, inequilateral, but not dimidiate, the margin rather distinctly lobed; fertile pinnae narrower than the barren ones. Texture flaccido-papyraceous. Sori linear, in long-continuous or interrupted marginal lines. — Occasionally this well-known species becomes 2-pinnate in the lower part. The young fronds are often beautifully tinged with red.

Var. striatum: Beautifully variegated with long or short, broad or narrow, white stripes placed irregularly between or along the veins.

Much cultivated. — Orig. loc.: Trop. America.

(3) **A. platyphyllum**, Sw., C. Chr., Ind. Fil.; *A. Seemanni*, Hk., Hk. Bk., Syn. Fil., 121; Hk., Spec. Fil., II, tab. LXXXI, A.

Stipes 15—25 c.M. long, erect, blackish, polished. Fronds 15—50 c.M. long, 1—2 \times pinnate. Leaflets 7—10 c.M. long, $3\frac{1}{2}$ —5 c.M. broad, ovate, acuminate, rather unequal-sided, but not dimidiate, the barren ones finely serrate, one side usually cordate at the base, the other obliquely truncate; petioles of the lowest leaflets nearly $2\frac{1}{2}$ c.M. long. Texture subcoriaceous; veins prominent; rachis naked, polished. Sori in long, continuous, marginal lines, often accompanied by roundish or oblong ones. — Resembles *A. peruvianum*, Kl. in habit.

In cultivation. — Orig. loc.: Centr. America, Brasilia, Bolivia.

§ 1. EU-ADIANTUM. c. **Polysoriae.**

a. *Fronds 1—2 \times pinnate.*

b. *Sori few, typically 1—2 to a leaflet.*

*. *Fronds simply pinnate. Indusium narrow.* (4) *A. gomphophyllum.*

.. *Fronds 2-pinnate. Indusium broad.* (5) *A. Hosei.*

b.b. *Sori many, typically 3 or more to a leaflet.*

c. *Surfaces pilose or strigose at least on the veins.*

§. *Texture thinly herbaceous or membranaceous.* (6) *A. diaphanum.*

§.§. *Texture chartaceous or coriaceous.*

*. *Rhizome scales bristle-like. Lobes of the leaflets ovate or transversally oblong.* (7) *A. Zollingeri*.

.. *Rhizome scales lanceolate. Lobes of the leaflets oblong or linear-oblong.* (8) *A. caudatum*.

c.c. *Surfaces naked or nearly so.*

d. *Rachis not winged.*

e. *Fronds unbranched, often proliferous at the apex.*

f. *Pinnae (leaflets) rather deeply lobed.*

g. *Rachis naked.* (9a) *A. rhizophorum*.

g.g. *Rachis rather long-hairy.* (9b) *A. ciliatum*.

f.f. *Pinnae (leaflets) entire or rather shallowly lobed.*

‡. *Fronds 1—2¹/₂ c.M. broad.*

(10a) *A. Spencerianum*.

‡.‡. *Fronds 3—8 c.M. broad.* (10b) *A. lunulatum*.

e.e. *Fronds mostly branched, not proliferous at the apex* (1).

*. *Leaflets dimidiate, trapeziform.* (11) *A. trapeziforme*.

.. *Leaflets unequal-sided, rather obliquely ovate.*

(12) *A. peruvianum*.

d.d. *Rachis winged.*

(13) *A. Mettenii*.

a.a. *Fronds 3—4-pinnate at least at the base* (2):

1. *Rachises pubescent or tomentose at least on the upper side.*

‡. *Rhizome wide-creeping; leaflets bluntish at the apex.*

(14) *A. pulchellum*.

‡.‡. *Rhizome short-creeping; leaflets bluntly rounded at the apex.*

(15a) *A. aculeolatum*.

(15b) *A. mindanaoense*.

1.1. *Rachises naked, but often scabrous.*

2. *Leaflets small, 1—2 c.M. long, 3—5 m.M. broad.* (16) *A. formosum*.

2.2. *Leaflets larger, 2—3 c.M. long, ± 10 m.M. broad.*

(17) *A. polyphyllum*.

(4) **A. gomphophyllum**, *Bl.*, *Ann. of Bot.*, VIII, 124.

Stipes tufted, erect, straight, black-brown, nearly naked. Fronds ligulate, 5—10 c.M. long, 2—2¹/₂ c.M. broad, simply pinnate. Leaflets 5—6-jugate, 7—10 m.M. each way, short-stalked, deltoid, the sides entire, straight, the

(1) See also **A. diaphanum**, *Bl.*, var. **affine** & **A. aculeolatum**, *v. A. v. R.*

(2) See also **A. peruvianum**, *Kl.* & **trapeziforme**, *L.*

apex lobed. Lobes rounded, crenulate. Surfaces naked; rachis slender, black-brown, naked; veins flabellate. Sori often 2 on a pinna, terminal on the lobes; indusium narrow, obreniform, naked, persistent.

Malacca.

(5) **A. Hosei**, *Bk.*, Journ. of Bot., XXVI, 524.

Stipes slender, castaneous, \pm 50 c.M. long, sparingly covered with small, reflexed, linear, brown scales towards the base. Fronds 2-pinnate, rhomboidal in outline. Pinnae lanceolate, 10—25 c.M. long, 2—2½ c.M. broad. Leaflets (pinnulae) close, numerous, rhomboidal, dimidiate, obtuse, short-stalked, the central ones 2—2½ c.M. long, 6—9 m.M. broad, the barren ones minutely crenate. Texture moderately firm; surfaces naked like the castaneous rachis. Sori 1—2, placed on the upper margin of the leaflets, to 4—7 m.M. long; indusium broad, firm. — Pendulous plants.

Borneo.

* (6) **A. diaphanum**, *Bl.*, Rac., Flor. Btz., I, 145; *Hk.*, Spec. Fil., II, tab. LXXX, C; — var. affine; *A. affine*, *Hk.*, l. c., 52.

Rhizome short, erect, densely clothed with small, lanceolate scales. Stipes tufted, 6—20 c.M. long, slender, erect, blackish, polished, scaly at the very base. Fronds 10—20 c.M. long, simply pinnate, linear-lanceolate, or with 1—5 branches varying from short to nearly as long as the terminal one. Leaflets short-stalked, 1—2 c.M. long, ½—1 c.M. broad, unilateral or sometimes slightly inequilateral, ovate to rhomboidal, the lower line straight or upcurved, the upper one \pm parallel to the lower, crenate or slightly lobed like the rounded outer edge. Texture very thin; rachis naked; upper surface purveyed with few, scattered, brown, subulate bristles especially near the lower edge, the under surface clothed with numerous small, adpressed hairs; veins terminating in the ultimate crenations or serratures. Sori obreniform, placed in nearly circular hollows on the upper and often also the outer edge; capsules springing from the veins and parenchyma. — In habit this shows an approximation towards the pedate group when branched.

Var. affine: Surfaces naked except the subulate bristles on the upper surface, olive-green when dry.

Trop. Asia, Polynesia, Australia, New Zealand.

(7) **A. Zollingeri**, *Mett.*, Ann. Mus. Bot. L. B., IV, 280.

Rhizome short, clothed like the base of the stipes with reddish-brown,

bristle-like scales. Stipes dense, \pm 17—18 c.M. long, ebeneous and glossy like the rachis. Fronds \pm 45 c.M. long, linear, simply pinnate, proliferous, narrowed gradually towards the apex. Leaflets subsessile, spreading, to 2 c.M. long, 5—8 m.M. broad, the upper base obliquely truncate or cuneate, the outer edge obtuse. Lobes ovate or transversally oblong, the lower ones 2-fid, the barren ones obtusely denticulate, the fertile ones retuse. Texture rigidly chartaceous; rachis hairy above; veins prominent, strigose. Sori solitary on the lobes; indusium coriaceous, transversally oblong.

Java; Angola.

* (8) **A. caudatum**, *L.*, *Rac., Flor. Btz., I, 145; Bedd., Ferns S.I., tab. II.*

Rhizome short-creeping or erect, clothed with brown, lanceolate scales. Stipes tufted or approximate, 5—15 c.M. long, wiry, dark-brown, clothed with a brown tomentum. Fronds simply pinnate, 15—50 c.M. long, the apex often nudate, elongate, caudate, rooting at the point. Leaflets dimidiate, \pm sessile with a cuneate base, 1—2 c.M. long, half as broad, the lower edge straight and entire, the upper and outer more or less broadly cut, often deeply and repeatedly, the outer edge bluntly rounded, lowest leaflets not rarely sub-flabellate. Lobes linear-oblong or oblong, with the apex rounded or truncate, entire or serrulate. Texture coriaceous; rachis and surfaces more or less densely pilose. Sori roundish or transversally oblong, terminal on the lobes; sporangia springing from the veins only.

Trop. Asia & Africa; New Hebrides.

(9a) **A. rhizophorum**, *Sw., Hk., Spec. Fil., II, 12, tab. LXXX, A.*

Near *A. caudatum*, *L.* Fronds elongate-lanceolate, simply pinnate, often tapering into a long rooting and proliferous apex bare of pinnae. Leaflets alternate, short-stalked, semiovate, the upper base truncate, parallel to the rachis, the upper edge lobed, the lobes retuse, soriferous, shining above. Texture chartaceo-coriaceous; rachis ebeneous or reddish-brown like the slightly chaffy stipe, naked, glossy; surfaces \pm naked; veins numerous, compact, fine. Indusium obreniform, thick, coriaceous.

Timor; South. Africa.

* (9b) **A. ciliatum**, *Bl., Enum., 215; A. caudatum, L., var., C. Chr., Ind. Fil.; A. caudatum, L., var. β Hk., Spec. Fil., II, 14.*

Near *A. caudatum*, *L.* Fronds elongate-lanceolate, simply pinnate, often

tapering into a long, rooting and proliferous apex bare of pinnae. Leaflets sessile or subsessile, dimidiate, the base cuneate, the inner edge \pm parallel to the rachis, the upper and outer edge laciniate, the laciniae linear or oblong, \pm 2-fid, the apices denticulate, frequently ciliate. Texture firm, rachis hirsute with moderately long, spreading hairs; surfaces naked or clothed with some scattered hairs. Sori terminal on the lobes; indusium lunulate to obreniform.

Trop. Asia & Africa.

(10a) **A. Spencerianum**, *Copel.*, Philipp. Journ., I, Suppl., 154, tab. XI.

Rhizome very short, scaly. Stipes approximate, 2—15 c.M. long, purple-black, glossy, the base clothed with, small, thin, deciduous scales. Fronds 7—17 c.M. long, or longer and rooting, 1—2½ c.M. broad, simply pinnate. Leaflets approximate, nearly sessile, sometimes falcate, the base obliquely cuneate, the inner edge excised, the upper edge entire or rarely slightly incised, the apex acute. Texture papyraceous; surfaces naked; veins inconspicuous. Sori continuous or interrupted.

Luzon.

* (10b) **A. lunulatum**, *Burm.*, Rac., Flor. Btz., I, 144; Hk. & Grev., Ic. Fil., tab. CIV; Bedd., Ferns S.I., tab. I; *A. philippense*, *L.*, Copel., Poly-pod. Philipp., 95.

Stipes 5—20 c.M. long, tufted, wiry, naked, polished, dark-brown. Fronds 15—30 c.M. long, 3—8 c.M. broad, simply pinnate, often elongated and rooting at the apex. Leaflets 1½—4 c.M. long, to 1—2½ c.M. broad, dimidiate or subdimidiate, the base broadly cuneate or truncate (upper and lower base in a continuous, straight or curved line), the lower base in a line or oblique with the petiole, the upper and outer edge rounded, more or less lobed or nearly entire; petioles of the lower leaflets spreading, to 1—1½ c.M. long. Lobes rounded, ½—5 m.M. deep. Texture papyraceo-herbaceous; both surfaces naked. Sori occupying the apices of the lobes, in continuous rows along the edge, sometimes confluent; capsules on the veins only.

Trop. Africa, Asia, Australia & Polynesia, Centr. America.

* (11) **A. trapeziforme**, *L.*, Hk. Bk., Syn. Fil., 118; *A. pendactylon*, *L. & F.*, Hk. & Grev., Ic. Fil., tab. XCVIII.

Stipes 15—40 c.M. long, firm, erect, naked, blackish, polished. Fronds 2—5-pinnate with a terminal branch 10—20 c.M. long, 5—8 c.M. broad and 2—4 large ones on each side, the lowest of which are often branched again. Leaflets 5—6 c.M. long, 1—3 c.M. broad, dimidiate, the apex \pm acute, the upper and lower edges parallel, the lower sometimes decurved, the outer edge oblique, the inner deflexed, the upper and outer edge more or less deeply or shallowly lobed, the lobes blunt; lowest leaflets on stalks $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long. Texture papyraceo-herbaceous; rachis and both surfaces naked; veins terminating in the ultimate lobes or their crenations or serratures when present. Sori numerous, terminal on the lobes, approximate, transversally oblong; capsules on the veins only, but densely crowded.

Much cultivated. — Orig. loc.: Trop. America.

* (12) **A. peruvianum**, *Hk.*, Hk. Bk., Syn. Fil., 116; Hk. Spec. Fil., II, tab. LXXXI, C.

Stipes 15—25 c.M. long or longer, erect, naked, polished, nearly black, deciduously scaly at the base. Fronds either simply pinnate or 2-pinnate with 1—3 branches at the base or even with some of the latter slightly branched again. Leaflets large, 4—7 c.M. long, 3—5 c.M. broad, unequally ovate or subtrapezoidal, the base broadly cuneate or subtruncate, the edge slightly lobed, the apex varying from bluntish to acuminate; lowest leaflets on stalks 1—2 c.M. long. Texture papyraceous; veins copious, flabellate, terminating in the ultimate lobes or their crenations or serratures when present; rachis and surfaces naked. Sori many or numerous along the upper and outer edge; capsules springing from the veins only. — Leaflets mostly resembling those of *A. platyphyllum*, *Sw.*

In cultivation. — Orig. loc.: Peru.

* (15). **A. Mettenii**, *Kuhn*, Hk. Bk., Syn. Fil., 472; *A. lunulatum*, *Burm.*, var. *Mettenii*, *Bedd.*, Handb. Ind. Ferns, 83; Ferns S.Br.I., Suppl., tab. CCCLIV; *A. alatum*, *Copel.*, Polypod. Philipp., 95.

Rhizome erect, short. Stipes \pm 8—12 c.M. long, blackish-brown, glossy, slightly scaly at the base, more or less narrowly winged throughout like the rachis and the stalks of the pinnae. Fronds 10—50 c.M. long, \pm 4—5 c.M. broad, simply pinnate. Leaflets more or less spreading, the lower often subdeflexed, 1— $1\frac{1}{2}$ c.M. broad; the lower side entire, nearly straight, the upper and outer ones rounded, \pm irregularly lobed, the inner one subparallel to the rachis. Texture papyraceous; surfaces naked; veins conspicuous. Sori trans-

versally oblong, terminal on the lobes, 5—10 to a pinna, straight or subcurved; capsules on the veins only.

Philippines; Trop. India, West. Africa.

* (14) **A. pulchellum**, *Bl.*, Rac., Flor. Btz., I, 146; *A. Lobbianum*, *Hk.*, Spec. Fil., II, tab. LXXXVI, C.

Rhizome wide-creeping, the scales dense, linear-lanceolate, brown. Stipes remote, 15—60 c.M. long, polished, blackish, naked, scaly towards the base, the scales linear-subulate, spreading, crisped, leaving the stipes rough when they fall. Fronds 20—40 c.M. each way, 2—5-pinnate with a terminal branch 15—25 c.M. long, 2—4 c.M. broad and a few lateral ones on each side, the lower ones branched again with 2—4 erecto-patent secondary branches. Leaflets dimidiate or slightly inequilateral, close, the largest to ± 2 c.M. long, to ± 1 c.M. broad, the lower edge upcurved, the upper one nearly straight, the apex bluntish, but not rounded. Texture coriaceous; rachises hirsute-tomentose on the upper side; surfaces naked; veins terminating in the ultimate serratures. Sori obreniform, small, numerous, placed on the upper and often also on the outer edge in slightly rounded hollows; capsules springing from the veins and parenchyma.

Java.

* (15a) **A. aculeolatum**, *v. A. v. R.*, Hort. Bog.; (? *A. mindanaense*, *Copel.*).

Rhizome short, the scales narrow, dense, brown, narrowed gradually from the base towards the acuminate point. Stipes approximate, ± 30 —40 c.M. long, polished, dark-purple-brown or nearly black, with the lower $\frac{1}{3}$ covered with irregularly scattered, spreading, crisped, brown scales, which leave the stipes rough (subaculeate) when they fall. Fronds ± 20 —35 c.M. long, 15—30 c.M. broad, 2—5-pinnate with a terminal branch 12—15 c.M. long, ± 3 c.M. broad, and 1—4 lateral branches on each side. Lowest branches largest, erecto-patent, generally branched again at the base, with the secondary branch on the lower side often the longest. Leaflets dimidiate, often close and imbricate, the largest ± 2 c.M. long, ± 1 c.M. broad, erecto-patent, the lower line upcurved, the upper edge \pm straight, the apex rounded, denticulate, the upper edge slightly lobed and denticulate, the base cuneate, the lowest leaflets of each branch \pm flabellate. Texture coriaceous; rachises purple-blackish, rather hirsute-tomentose on the upper side; surfaces naked; veins terminating in the ultimate serratures. Sori transversally oblong or

nearly obreniform, placed in shallow or semiorbicular hollows along the upper edge, solitary on the lobes; sporangia springing from the veins as well as from parenchyma. — The description is taken from a specimen cultivated in the Buitenzorg gardens.

Amboyna.

(15b) **A. mindanaoense**, *Copel.*, Philipp. Journ., I, Suppl., 154, tab. X.

Rhizome short-creeping. Stipes approximate, 25—40 c.M. long, clothed towards the base with glossy, linear-subulate, fuscous scales, otherwise naked, black-fuscous. Fronds deltoid-ovate, 25—40 c.M. long, 20—30 c.M. broad, acuminate. 3-pinnate. Leaflets short-stalked, dimidiate, $\pm 1\frac{1}{2}$ — $1\frac{3}{4}$ c.M. long, $\pm \frac{3}{4}$ c.M. broad, the outer edge obtuse, denticulate, the upper one slightly lobed, the base truncate on the upper side. Texture papyraceous; rachis densely furnished on the upper side with minute hairs; surfaces naked. Sori ± 6 to a leaflet, solitary on the lobes, suborbicular-reniform, the sporangia springing from the veins and parenchyma.

Mindanao.

* (16) **A. formosum**, *R. Br.*, Hk. Bk., Syn. Fil., 119; Hk., Spec. Fil., II, tab. LXXXVI, B.

Stipes 50—50 c.M. long, erect, scabrous, strong. Fronds 45—75 c.M. long, 3—4 \times pinnate. Lowest branches 30—40 c.M. long, erecto-patent, 15—25 c.M. broad, deltoid. Leaflets short-stalked, 1—2 c.M. long, 5—5 m.M. broad, dimidiate, numerous, rhomboidal, the lower edge straight, erecto-patent, the inner one parallel to the rachis, the upper and outer edge lobed and toothed, the teeth often sharp. Texture firm; rachis scabrous or hirsute towards the apex; veins prominent above, terminating in the ultimate serratures. Sori numerous, transversally oblong, slightly obreniform, in slightly rounded hollows of the upper and outer edge; capsules springing from the veins and parenchyma.

In cultivation. — *Orig. loc.: Australia, New Zealand.*

* (17) **A. polyphyllum**, *Willd.*, Hk. Bk., Syn. Fil., 118; Hk., Gard. Ferns, tab. XII; *A. cardiochlaena*, *Kze*, Hk., Spec. Fil., II, tab. LXXXIII.

Stipes 50—45 c.M. long, strong, erect, blackish, polished, deciduously scaly, rather scabrous. Fronds 60—120 c.M. long, 50—60 c.M. broad, 3-pinnate, with a long, terminal branch and several lateral branches on each side,

the lower of which are branched again below a long terminal branch. Secondary branches \pm 8 c.M. long or longer. Leaflets dimidiate, rhomboidal (almost a parallelogram), sessile, numerous, close, 2—3 c.M. long, \pm 1 c.M. broad, the upper and lower sides parallel, the apex bluntly rounded, the upper and outer edge crenate, or lobed and serrate. Texture firm-membranaceous or subcoriaceous; rachises and surfaces naked; veins terminating in the ultimate serratures. Sori numerous, placed in shallow hollows at the apices of the lobes or crenations, 1 to each lobe, along the upper and sometimes also the outer edge; capsules springing from the veins only.

Much cultivated. — *Orig. loc.:* Venezuela, Peru.

§ 1. EU-ADIANTUM. d. **Cuneatae.**

- a.** *Ultimate segments (leaflets) articulate at the base; ultimate petioles distinctly nodose at the apex.* (18) *A. tenerum.*
- a.a.** *Ultimate segments (leaflets) not articulate at the base; ultimate petioles not nodose at the apex.*
- b.** *Leaflets with the outer edge entire or shallowly lobed.*
- c.** *Sori rather numerous, typically 3—6, sometimes 2 to a leaflet.*
- d.** *Leaflets often dimidiate or trapezoidal, the terminal ones cuneate.*
(19a) *A. neoguineense.*
(19b) *A. opacum.*
- d.d.** *Leaflets cuneato-flabellate to suborbicular.*
- e.** *Sori in rather half-rounded hollows; sporangia springing from the veins and parenchyma.* (20) *A. aethiopicum.*
- e.e.** *Sori in shallow hollows; sporangia springing from the veins only.* (21) *A. suborbiculare.*
- c.c.** *Sori few, typically 1—2, sometimes 3 to a leaflet.*
(22) *A. celebicum.*
- b.b.** *Leaflets with the outer edge rather deeply lobed.*
- §.** *Lower leaflets of the pinnae overlapping the main rachis, adpressed to the latter.*
- ‡.** *Pinnae spreading.* (23) *A. concinnum.*
- ‡.‡.** *Pinnae erecto-patent.* (24) *A. tinctum.*
- §.§.** *Lower leaflets not overlapping the main rachis.*

*. *Sori roundish-obreniform. Veins terminating in the angles between the ultimate serratures of the barren leaflets.*

(25) *A. cuneatum.*

.. *Sori transversally oblong. Veins terminating in the tops of the ultimate serratures of the barren leaflets.*

(26) *A. capillus veneris.*

* (18) **A. tenerum**, *Sw.*, Hk. Bk., Syn. Fil., 124; *A. nobile*, *Fisch.*, hort.

Stipes to 50 c.M. long, erect, naked, blackish, glossy. Fronds 25—90 c.M. long, 20—45 c.M. broad, deltoid, 3—4-pinnate. Leaflets cuneate or tending toward rhomboidal-dimidiolate in shape, the largest $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, 1—2 c.M. broad, articulate to the nodose apex of the petioles, the upper and outer edge rounded or somewhat angular, broadly and often rather deeply lobed. Texture thin: rachis glossy and naked like the surfaces; veins terminating in the ultimate serratures. Sori placed in roundish or transversally oblong patches in slightly rounded, shallow hollows, terminal on the lobes, the sporangia springing from the veins only.

In cultivation. — *Orig. loc.:* Bermuda, West. India, Mexico, Juan-Fernandez.

(19a) **A. neoguineense**, *Moore*, Ann. of Bot., V, 208.

Stipes naked, castaneous, 15—20 c.M. long. Fronds deltoid, 3-pinnate, naked, above 50 c.M. long and broad; rachises very slender, naked, nearly black. Terminal leaflets cuneate, lateral trapezoidal, $1-1\frac{1}{2}$ c.M. long, cuneate. Sori orbicular, 3—4 to a leaflet, placed in deep sinuses. — Intermediate between *A. tenerum*, *Sw.* and *A. aethiopicum*, *L.*

New Guinea.

(19b) **A. opacum**, *Copel.*, Philipp. Journ., I, Suppl., 255, tab. III.

Rhizome short, clothed with narrow, castaneous scales. Stipes approximate, slender, \pm 20 c.M. long, the base furnished with long, scattered, deciduous hairs, otherwise pubescent like the lower part of the rachis. Fronds 16—20 c.M. long, ovate, 3-pinnate. Leaflets deciduous (? articulate, the stalks persistent), often dimidiolate, \pm 8 m.M. broad, $1-1\frac{1}{4}$ c.M. long, the margin narrowly cartilagineous, the upper and outer edge rounded, slightly lobed with 2—4 shallow lobes. Texture papyraceous, opaque; surfaces naked, pale-

coloured. Sori transversally oblong, 1 to each lobe, the sporangia nearly restricted to the veins.

Palawan.

* (20) **A. aethiopicum, L.**, Hk. Bk., Syn. Fil., 125; Hk., Spec. Fil., II, tab. LXXVII; Bedd., Ferns S.I., tab. V.

Stipes 15—25 c.M. long, erect, naked, dark-brown or blackish, polished. Fronds 15—45 c.M. long, 10—25 c.M. broad, \pm deltoid in outline, 3—4-pinnate. Leaflets suborbicular, broadly flabellate, rather equilateral, the base straight, rounded or broadly cuneate, the outer edge rounded, broadly and shallowly lobed. Texture thinly herbaceous to papyraceous; rachis and surfaces naked; veins terminating in the ultimate crenations or serratures. Sori in several roundish or transversally oblong patches in rather half-rounded hollows; capsules springing from the veins and parenchyma.

Banca, New Guinea; Australia, New Zealand; South. & Trop. Africa, ? Spain.

* (21) **A. suborbiculare, v. A. v. R.**

Rhizome short, densely clothed with linear-subulate, entire, bright-reddish-brown scales. Stipes approximate, to 20 c.M. long, erect, blackish, polished like the rachises, scaly at the very base. Fronds \pm 15—20 c.M. each way, 3-pinnate with several alternate, erecto-patent lateral branches below the terminal one, the lowest largest, branched again in the lower part. Leaflets \pm 1 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, obliquely flabellate or suborbicular, stalked, the base cuneate, truncate or rounded, the outer edge rounded, often irregularly and shallowly lobed. Texture papyraceous, firm, subpellucid; surfaces naked like the rachises and the stipes; veins terminating in the serratures. Sori 2—6 to a leaflet, roundish or transversally oblong, placed in indistinct or shallow hollows of the outer edge, the sporangia springing from the veins only.

Sumatra (Burck).

* (22) **A. celebicum, Christ**, Ann. Btz., XV, 105, tab. XV, fig. 15, a—c.

Stipes wiry, black, polished, shorter than the fronds. Fronds 20—50 c.M. long, 7—15 c.M. broad, 3-pinnate at least at the base; rachises black, naked, polished, erecto-patent. Leaflets 1—2 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, the base more or less broadly cuneate, sometimes rounded, whether or not

oblique, shallowly and broadly lobed with 2—5 subentire lobes. Texture membranaceous-herbaceous; surfaces naked, opaque, often bluish. Sori obreniform, 1—2, rarely 5 to a leaflet, placed in orbicular hollows along the outer edge at the tops of the sinuses; indusium adpressed, persistent.

Celebes.

* (23) **A. concinnum**, *H. B. K.*, Hk. Bk., Syn. Fil., 123.

Stipes 10—25 c.M. long, blackish, naked, polished. Fronds 50—50 c.M. long, 15—30 c.M. broad, ovate or ovate-deltoid, 3-pinnate. Branches many, spreading, alternate, the lowest 10—15 c.M. long, 3—7½ c.M. broad at the base. Leaflets 6—20 m.M. across, broadly flabellate, the outer edge irregularly rounded, deeply lobed, the lobes again crenate, the base broadly cuneate, truncate or subcordate. Texture papyraceo-herbaceous; main rachis naked, glossy, flexuose, overlapped by the large, adpressed lower leaflets; surfaces naked; veins terminating in the angles of the sinuses between the ultimate serratures. Sori numerous, obreniform, placed in round or semiorbicular hollows, the sporangia springing from the veins and parenchyma.

In cultivation. — *Orig. loc.: Trop. America.*

* (24) **A. tinctum**, *Moore*, Hk. Bk., Syn. Fil., 474; *A. Wagneri*, *Mett.*, Hk. Bk., l. c. 473; *A. decorum*, *Moore*, *Gardn. Chron.* 1869, 582.

Stipes 10—20 c.M. long, erect, naked, blackish, glossy. Fronds ± deltoid, 15—45 c.M. long, 2—4-pinnate. Lower primary and secondary branches erecto-patent, stalked, deltoid. Terminal leaflets obliquely cuneate-flabellate, the lateral ones subrhomboidal, ¾—1½ c.M. long, the lower edge straight, the inner one ± touching the rachis or even wrapped over it, upper and outer edge more or less distinctly lobed, the lobes blunt, lowest leaflets equilateral, broadly flabellate, imbricated over the main rachis and adpressed to it. Surfaces naked; texture papyraceous; rachis glossy, naked, slightly zigzag; veins terminating in the tops of the sinuses between the ultimate serratures. Sori round, obreniform, in ± circular hollows of the upper and outer edge; capsules on and between the veins. — Intermediate between *A. concinnum*, *H. B. K.* and *A. cuneatum*, *L. & F.*

In cultivation. — *Orig. loc.: Costa Rica, Peru.*

* (25) **A. cuneatum**, *L. & F.*, Hk. Bk., Syn. Fil., 124; Hk. & Grev., *Ic. Fil.*, tab. XXX.

Stipes 15—25 c.M. long, slender, naked, blackish, polished. Fronds 20—40

c.M. long, 15—25 c.M. broad, deltoid, 3—4-pinnate. Lower branches 5—8 c.M. broad. Leaflets numerous, $\frac{1}{2}$ —2 c.M. broad, the base cuneate, the outer edge obliquely rounded, deeply lobed, the lobes whether or not lobed again. Texture thin; rachis and surfaces naked; veins terminating in the angles of the sinuses between the ultimate serratures. Sori \pm 4—6 to a leaflet, obreniform, placed in distinct semiorbicular or circular hollows, the sporangia springing from the veins and parenchyma. — A variable species with many garden varieties.

In cultivation. — *Orig. loc.:* *Brasilia, Peru, Argentina.*

* (26) **A. capillus veneris**, *L.*, *Hk. Bk., Syn. Fil., 125*; *Bedd., Ferns S.I., tab. IV*; *Hk., Spec. Fil., II, tab. LXXIV, B*; *Bau. & Hk., Gen. Fil., tab. LXVI, B.*

Stipes slender, 10—25 c.M. long, naked, polished, blackish. Fronds 2-pinnate at least at the base, 20—40 c.M. long, deltoid-oblong. Branches alternate. Leaflets to 3 c.M. long, 1—2 $\frac{1}{2}$ c.M. broad, the base cuneate, the outer edge obliquely rounded, deeply lobed from the circumference in the direction of the base, the lobes again bluntly crenate and slightly lobed and whether or not serrulate. Texture pellucido-herbaceous; rachises and surfaces naked; veins terminating in the ultimate crenations or serratures. Sori roundish or transversally oblong, slightly curved, placed in slightly rounded, shallow hollows, the sporangia springing from the veins only. — Variable, with several garden varieties.

Paleotropical, Temperate Asia & Africa, South. Europe.

§ 1. EU-ADIANTUM. e. **Pedatae.**

§. *Surfaces hispid.* (27) *A. hispidulum.*

§.§. *Surfaces naked.*

*. *Leaflets imbricated over the rachises.* (28) *A. stenochlamys.*

.. *Leaflets not or hardly imbricated over the rachises.* (29) *A. flabellulatum.*

* (27) **A. hispidulum**, *Sw., Rac., Flor. Btz., I, 145*; *Bedd., Ferns S.I., tab. III.*

Rhizome short-creeping or erect, scaly. Stipes strong, erect, polished, dark-brown, 15—35 c.M. long, scabrous and rough-hairy. Fronds irregularly

broadly ovate, 12—15 c.M. each way, pedato-subdichotomous with the main divisions flabellately branched. Central branches 15—25 c.M. long, 1—2½ c.M. broad. Leaflets ½—2¼ c.M. long, 5—10 m.M. broad, dimidiate, subrhomboidal, rather broader on the side nearest the stipe, the outer edge bluntly rounded or oblique, upper and outer margin often crenate, finely toothed; lower leaflets slightly stalked. Texture subcoriaceous; rachises and both surfaces hispid. Sori roundish, contiguous round the upper and outer edge; indusium obreniform, hairy.

Paleotropical.

(28) **A. stenochlamys**, *Bk.*, Ann. of Bot., V, 209.

Stipes long, naked, wiry, nearly black. Fronds pedate, deltoid, 3-pinnate, ± 15—25 c.M. each way. Terminal leaflets cuneate, 1—2 c.M. broad, shallowly lobed, crenate round the upper edge; lateral leaflets subrhomboidal, short-stalked, little longer than broad, the upper and outer edge shallowly lobed and crenate, the inner edge inflexed so that it is imbricated over the rachis. Texture firm; surfaces naked. Sori 6—8 to a leaflet; indusium firm, narrow.

Borneo.

* (29) **A. flabellulatum**, *L.*, Rac., Flor. Btz., I, 146; Bedd., Ferns S.I., tab. CCXVIII; *A. amoenum*, *Hk. & Grev.*, Ic. Fil., tab. CIII.

Rhizome erect, short, clothed with narrow, lanceolate scales. Stipes erect, strong, polished, dark, naked, to 25 c.M. long. Fronds irregularly broadly ovate, to ± 20 c.M. or more long and to ± 15 c.M. or more broad, pedato-subdichotomous and the branches once or twice branched again. Central branches to 20 c.M. long, to 2½ c.M. broad. Leaflets to ± 10 m.M. broad and to ± 7½ m.M. deep, flabellato-cuneate to dimidiate, the base truncate, rounded or cuneate the upper and outer margin rounded, entire or slightly or finely toothed. Texture coriaceous, rachis naked or tomentose above, surfaces naked. Sori in several transversally oblong patches round the upper and outer edge, contiguous, often confluent.

Trop. Asia, China, Japan.

38. HYPOLEPIS, *Bernhardi*.

Sori round, subterminal on the veins on a thickened receptacle, sub-

.marginal, solitary at the tops of the sinuses of the ultimate segments, protected by reflexed, membranaceous, roundish lobes, too small to cover them permanently.

Rhizome whether or not creeping. Stipes not articulate to the rhizome. Fronds variously divided, herbaceous, more or less hairy; veins free. *Hk. Bk.*, *Syn. Fil.*, tab. II, fig. 24; *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 147.

Pantropical.

Arrangement of the sections.

§ 1. **ASPIDOTIS.** Rhizome not creeping; fronds small, tufted. (Not in Malaya).

§ 2. **EUHYPOLEPIS.** Rhizome wide-creeping; fronds large, scattered.

§ 2. **EUHYPOLEPIS.**

§. Fronds 2—3-pinnate, under 50 c.M. long. (1) *H. alpina.*

§.§. Fronds at least 4-pinnatifid, more than 50 c.M. long.

*. Surfaces naked except a few hairs on the midribs. (2) *H. punctata.*

.. Surfaces sparingly glanduloso-pubescent. (5) *H. tenuifolia.*

* (1) **H. alpina**, *Hk.*, *Rac.*, *Flor. Btz.*, I, 149; *Cheilanthes alpina*, *Bl.*, *Enum.*, 158.

Rhizome branched, densely clothed with reddish-brown, spreading, narrow fibrils. Stipes 5—20 c.M. long, densely clothed with spreading hairs. Fronds triangular-lanceolate, less than 50 c.M. long, 15—30 c.M. broad, 2—3-pinnate. Ultimate segments oblong, blunt, not close, 5—10 m.M. long, 2½—5 m.M. broad at the base, adnate to the rachis, very slightly decurrent, often united at the base by a narrow wing ± ¼ m.M. broad, the edges crenate or toothed. Texture herbaceous, but firm; rachises and surfaces hairy. Sori 1—8 to each segment.

Java.

(2) **H. punctata**, *Bedd.*, *Handb. Ind. Ferns, Suppl.*, 19; (? *Phegopteris punctata*, *Mell.*; *Dryopteris punctata*, *C. Chr.*, *Ind. Fil.*).

Rhizome large, wide-creeping. Fronds 120—560 c.M. long, 4-pinnatifid; rachises strigose. Pinnae ± 60 c.M. long, more than 50 c.M. broad at the

base, on petioles \pm 5 c.M. long. Pinnulae lanceolate, 15—20 c.M. long. Tertiary segments \pm 3 $\frac{1}{2}$ c.M. long, \pm 1 c.M. broad, cut down to a narrowly winged rachis into 10—12 oblong, obtuse segments on each side, which are bluntly lobed. Texture herbaceous; surfaces naked; tertiary rachises and mid-ribs with a few, longish, white hairs on both sides. Sori in the axils of the 1—2 lower lobes on each side of the segments; indusium distinct, membranaceous.

Malacca.

* (3) **H. tenuifolia**, *Bernh.*, Rac., Flor. Btz, I, 148; Hk., Spec. Fil., II, tab. LXXXIX, C & XC, A; *H. pallida*, *Hk.*, l. c., 64; *Cheilanthes pallida*, *Bl.*, Enum., 159.

Rhizome densely clothed with short, spreading fibrils. Stipes 20—100 c.M. long, puberulous like the rhizome but less so. Fronds 50—500 c.M. long, 4—5-pinnate. Lower pinnae ovate-lanceolate, 20—100 c.M. long. Ultimate segments linear-oblong, bluntish, free or joining at the base by a very narrow wing, the edges toothed or shallowly pinnatifid. Texture herbaceous; surfaces sparingly glandular-pubescent, the rachises more so. Sori in the ultimate sinuses.

Malaya; Australia, New Zealand, Polynesia, China.

39. CHEILANTHES, *Swartz.*

Sori roundish or oblong, terminal on the veins, at first small and distinct, afterwards more or less laterally confluent into a submarginal line. Indusium semiorbicular or transversally oblong, contiguous or confluent, consisting of the wholly or partially recurved, modified margin of the frond.

Rhizome ascending or creeping, mostly short, scaly. Stipes dark-coloured, not articulate to the rhizome. Fronds divided; veins free, the tips thickened. Hk. Bk., Syn. Fil., tab. III, fig. 25; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I^a, fig. 116.

Arrangement of the sections.

§ 1. **PHYSAPTERIS.** Ultimate segments very small, roundish, \pm 1—2 m.M. each way. Diels, l. c., fig. 116, D.

§ 2. **ALEURITOPTERIS.** Ultimate segments not very small. Fronds coated beneath with a white or yellow powder.

§ 3. **EUCHEILANTHES.** Like § 2, but the fronds not powdery beneath.

§ 1. **PHYSAPTERIS.**

(1) **Ch. Boltoni, Copel.**, Polypod. Philipp., 95.

Stipes densely tufted, $1\frac{1}{2}$ —3 c.M. long, brown-black, sparsely covered with hairs of the same colour. Fronds linear-oblongate, to \pm 16 c.M. long, $1\frac{1}{2}$ c.M. broad, 2-pinnate, the broadest part $\frac{1}{3}$ of the way from the apex. Pinnae almost sessile, remote, the largest to \pm 8 m.M. long, 6 m.M. broad. Pinnulae in 1—5 pairs, ovate, minute, crenulate, plane when moist, curling up when dry. Texture very thin, but firm when dry; rachis blackish-brown, nearly naked; surfaces naked. Indusium interrupted.

Philippines.

§ 2. **ALEURITOPTERIS.**

a. Primary segments not reaching down to the rachis, *i. e.* joined at the base.

(2) *Ch. argentea.*

a.a. At least the lowest primary segments reaching down quite to the rachis, *i. e.* free from the next higher ones.

b. Fronds 2-pinnatifid.

(3) *Ch. rufa.*

b.b. Fronds 3-pinnatifid.

(4) *Ch. farinosa.*

* (2) **Ch. argentea, Kze** (not *Hk.*), *Hk. Bk.*, *Syn. Fil.*, 142; *Bedd.*, *Ferns Br.I.*, I, tab. CXLIII; *Pteris argentea, Gmel.*, (oldest name).

Stipes densely tufted, 7—15 c.M. long, polished, dark-brown, scaly at the base especially when young. Fronds 7—10 c.M. long, 5—8 c.M. broad, 3-pinnatifid at the base, the primary segments not reaching down quite to the rachis. Lowest segments the largest with the lowest basipetal lobe the largest. Costae polished like the stipe; upper surface naked, the under one covered with white, waxy powder. Sori very small, fusing; indusium crenate or fimbriate.

Siberia to Malaya.

FILICES.

(5) **Ch. rufa**, *Don* (not *Desv.*), *Hk. Bk., Syn. Fil.*, 141; *Hk., Spec. Fil.*, II, tab. IC, A; *Bedd., Ferns Br.I.*, I, tab. CXLIV.

Stipes tufted, 2½—5 c.M. long, densely clothed with rusty-brown, woolly tomentum like the rachis. Fronds 8—25 c.M. long, 4—8 c.M. broad, ovate-lanceolate, 2-pinnatifid. Pinnae opposite, the lower ones with 2—5 c.M. space between them, oblong, much shorter than the central ones, with the lowest segments on the lower side produced, the higher ones linear-oblong, regularly lobed or wavy. Texture herbaceous or flaccido-coriaceous; surfaces somewhat tomentose, under surface more or less coated with white powder. Sori roundish, small, numerous; indusium brown, ragged or ciliate.

Philippines; North. India.

* (4) **Ch. farinosa**, *Kuf.*, *Rac., Flor. Btz.*, I, 147; *Bedd., Ferns S.I.*, tab. CXCI; *Hk. & Grev., Ic. Fil.*, tab. CXXXIV; *Pteris farinosa*, *Forsk.*, (oldest name); — var. *chrysophylla*; *Ch. chrysophylla*, *Hk., Spec. Fil.*, II, 113; *Ch. argentea*, *Kze, var. chrysophylla*, *Hk. Bk., Syn. Fil.*, 142; *Hk., Fil. Exot.*, tab. XCV.

Stipes densely tufted, 10—30 c.M. long, polished, dark-brown, clothed with scattered, deciduous, linear-lanceolate scales. Fronds 5—30 c.M. long, 4—15 c.M. broad, 3-pinnatifid. Pinnae in opposite pairs, sessile, the lower ones the largest, remote, reaching to the rachis, with the segments on the lower side longer than the others and deeply lobed, the higher ones confluent at the base so as to form a deeply pinnatifid, acuminate terminal pinna. Texture sub-coriaceous; rachis and costae polished, naked, dark-brown; under surface covered with white, waxy powder. Indusia brown, scariose, rounded, contiguous or confluent and then waved or lobed, the margin entire or toothed.

Var. chrysophylla: Powder yellow.

Trop. Asia, Africa, America, Fiji, China, Japan.

§ 3. EUCHEILANTHES.

1. *Rachis naked.*

2. *Fronds 2-pinnatifid.*

(5) *Ch. varians.*

2.2. *Fronds at least 3-pinnatifid.*

3. *Indusium transversally oblong, narrow.*

(6) *Ch. tenuifolia.*

3.3. *Indusium roundish-obreniform, broad.*

(7) *Ch. multifida.*

1.1. *Rachis tomentose.*

- *. *Fronde 2-pinnatifid, herbaceous.* (8) *Ch. fragilis.*
 .. *Fronde 3-pinnatifid, subcoriaceous.* (9) *Ch. hirta.*

* (5) **Ch. varians**, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 136; *Hk.*, *Spec. Fil.*, II, tab. CIII, A; *Bedd.*, *Ferns S.I.*, tab. CLXXXIX; *Ch. Belangeri*, *C. Chr.*, *Ind. Fil.*; *Pteris varians*, *Wall.*, *List*, No. 86.

Stipes densely tufted, 5—15 c.M. long, dark-brown, polished, rather slender and brittle, fibrillose below. Fronds 15—25 c.M. long, 3½—5 c.M. broad, lanceolate, 2-pinnatifid. Pinnae in numerous pairs, the lower ones opposite, often 3½—5 c.M. apart, 2½—3½ c.M. long, ± 2 c.M. broad, deltoid-acuminate, cut down to the rachis in the lower part into oblong or linear, oblong, entire or slightly toothed lobes. Texture herbaceous; rachis polished like the stipe; both surfaces naked. Sori copious, continuous; indusium narrow, membranaceous.

Trop. Asia.

* (6) **Ch. tenuifolia** *Sw.*, *Rac.*, *Flor. Btz.*, I, 148; *Hk.*, *Spec. Fil.*, II, tab. LXXXVII, C; *Bedd.*, *Ferns S.I.*, tab. CLXXXVIII; *Ch. hispidula*, *Kze.*, *Bot. Zeit.*, 1848, 212; *Trichomanes tenuifolium*, *Burm.*, (oldest name).

Stipes not tufted, 10—20 c.M. long, wiry, flexuose, naked, dark-brown, polished, slightly fibrillose towards the base. Fronds 10—20 c.M. long, 7—10 c.M. broad, deltoid to lanceolate, 3-pinnatifid. Pinnae in numerous, opposite, erecto-patent pairs, the lowest 5—10 c.M. long, 2—5 c.M. broad, deltoid. Pinnulae on the lower side the largest, the lowest ± 2½ c.M. long, cut down to the rachis into oblong, entire or sinuato-pinnatifid segments. Texture subcoriaceous; rachis polished like the stipe; both surfaces naked. Sori roundish or subcontinuous; indusium brownish, narrow, transversally oblong, ± toothed.

Trop. Asia, Australia, Polynesia, Tasmania, New Zealand.

(7) **Ch. multifida**, *Sw.*, *Hk. Bk.*, *Syn. Fil.*, 138; *Hk.*, *Spec. Fil.*, II, tab. C, B; *Gard. Ferns*, tab. XXXIX.

Stipes tufted, 7—25 c.M. long, strong, erect, dark-chestnut-brown, polished, slightly scaly towards the base. Fronds 7—30 c.M. long, 5—20 c.M. broad, ovate-lanceolate or deltoid, 3—4-pinnatifid. Lower pinnulae opposite, remote, erecto-patent, deltoid, 5—15 c.M. long. Ultimate divisions linear-oblong, deeply lobed and the margin in the fertile fronds much recurved. Texture subco-

riaceous; rachis dark-chestnut-brown, naked and polished; surfaces naked. Sori small, roundish, slightly confluent; indusium roundish-obreniform.

Java; South. Africa, St. Helena.

(8) **Ch. fragilis**, *Hk.*, Hk. Bk., Syn. Fil. 135; Hk., Fil. Exot., tab. XCVI; Bedd., Ferns Br.I., II, tab. CCXXIII.

Stipes densely tufted, not more than 2—5 c.M. long, strong, tomentose and fibrillose, erect. Fronds 15—30 c.M. long, 5—10 c.M. broad, ovate-lanceolate, 2-pinnatifid. Pinnae numerous, alternate, the lowest 2—5 c.M. apart, 2—5 c.M. long, 1—1½ c.M. broad, oblong-lanceolate, cut down to the rachis below into numerous linear-oblong, blunt, sinuato-crenate lobes. Texture herbaceous; rachises bright-brown, tomentose, but the surfaces naked except on the midribs beneath. Sori small, roundish or subcontinuous; indusium pale-brown, membranaceous.

Malacca; Moulmein, Yunnan.

(9) **Ch. hirta**, *Sw.*, Hk. Bk., Syn. Fil., 136; Hk., Spec. Fil., II, tab. CI, B; *Ch. parviloba*, *Sw.*, Syn., 128.

Stipes tufted, 5—10 c.M. long, strong, erect, densely coated with spreading, bright-reddish-brown, woolly hairs. Fronds 10—30 c.M. long, 5—15 c.M. broad, ovate-lanceolate, 3-pinnatifid. Pinnae opposite, spreading from the main rachis at right angles, lanceolate, 2½—7½ c.M. long, cut down to the rachis into numerous oblong pinnulae, which are scarcely more than ½ c.M. long and again inciso-pinnatifid. Texture subcoriaceous; rachis and both surfaces more or less tomentose; margin of the segments much incurved. Sori copious. — *Ch. parviloba*, *Sw.* has the rachis and surfaces less tomentose, the pinnae shorter and the segments and apex slightly divided and convolute.

Java; Réunion, South. Africa.

40. PLAGIOGYRIA, *Mettenius*.

Sori oblong, on the somewhat thickened ends of the mostly branched veins, soon becoming confluent and quite covering the space between the costae and the edges of the narrowed segments at maturity. Sporangia long-stalked, with an oblique, complete annulus. Indusium long, linear, continuous, consisting of the more or less modified, reflexed or recurved margin of the pinnae.

Rhizome erect. Stipes clustered, adherent to the rhizome, the base abruptly dilated, bearing on the outer side generally ± 2 —12 spongy tubercles or projections. Fronds simply pinnate, dimorphous; barren pinnae linear-lanceolate, fertile ones much contracted, linear. Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 149.

Eastern Asia, Tropical America.

a. *Pinnae of barren fronds adnate to the rachis by a broad base.*

*. *Fronds ± 20 —25 c.M. long, ± 6 —7½ c.M. broad, submembranaceous.* (1) *P. stenoptera.*

.. *Fronds ± 30 —45 c.M. long, ± 10 —15 c.M. broad, subcoriaceous.* (2) *P. adnata.*

a.a. *At least the lower pinnae of barren fronds not adnate to the rachis.*

b. *Tubercles wanting on the stipes at the base.* (3) *P. Christii.*

b.b. *Tubercles present.*

c. *Pinnae of barren fronds glaucous or white-powdery beneath.*

(4) *P. glauca.*

c.c. *Pinnae of barren fronds not glaucous, nor powdery beneath.*

d. *Pinnae of barren fronds \pm truncate at the base.*

(5) *P. pycnophylla.*

d.d. *Pinnae of barren fronds narrowed at the base.*

e. *Barren pinnae serrate throughout.* (6) *P. euphlebia.*

e.e. *Barren pinnae serrate towards the apex only, otherwise entire.* (7) *P. tuberculata.*

(1) **P. stenoptera**, **Diels**, Copel., Philipp. Journ., I, Suppl., 154; *Blechnum stenopterum*, Hance, Journ. of Bot., XXI, 268; *Lomaria concinna*, Bk., Journ. of Bot., XXIII, 105; *L. stenoptera*, Bk., Ann. of Bot., V, 220.

Rhizome short, erect. Stipes tufted, greenish, nearly naked, 10—25 c.M. long, basal tubercles present, the fertile fronds on the longest stipes. Barren fronds oblong-lanceolate, 20—25 c.M. long, 6—7½ c.M. broad. Pinnae ± 20 on a side, lanceolate, contiguous, adnate by a dilated base, ± 7 ½ m.M. broad, acute, minutely denticulate. Texture nearly membranaceous; surfaces naked; veins distinct, simple or forked. Fertile fronds with the pinnae less numerous, remote, linear.

Luzon; Formosa, China.

* (2) **P. adnata**, *Bedd.*, Handb. Ind. Ferns, 127; Ferns Br.I., I, tab. LI; *Lomaria adnata*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 162; *Hk.*, *Spec. Fil.*, III, tab. CXLVII.

Rhizome short, stout. Stipes 15—45 c.M. long, firm, erect, naked, basal tubercles present. Fronds ovate-lanceolate, 30—45 c.M. long, 10—15 c.M. broad. Pinnae linear, falcate, the largest $\pm 7\frac{1}{2}$ c.M. long, to $1\frac{1}{2}$ c.M. broad, the lower ones narrowed at the base below, dilated at the base above, 1— $2\frac{1}{2}$ c.M. apart, the apex narrowed gradually and finely toothed. Texture subcoriaceous; rachis naked; veins conspicuous. Fertile fronds similar, but the pinnae more distant and narrowly linear.

Java, Sumatra, Philippines; Japan, Yunnan, Khasia.

(3) **P. Christii**, *Copel.*, *Philipp. Journ.*, I, Suppl., 153.

Rhizome short, erect. Stipes approximate, 20—30 c.M. long, naked, castaneous, the base thickened, *without* tubercles. Fronds pinnate, 40—60 c.M. long. Barren fronds with ± 13 pinnae on each side. Pinnae nearly horizontal, ± 13 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, acuminate, serrulate, the lower ones stalked, the higher ones adnate. Texture herbaceous; surfaces naked; veins simple or forked. Fertile fronds with 12—16 pinnae, the pinnae linear, ± 10 c.M. long, 2—3 m.M. broad, stalked, or the higher ones adnate. — Fronds with the aspect of *Stenochlaena*, and the absence of the basal projections of the stipes helps to conceal the real affinity. However, the oblique annulus and the insertation of the sori establish the place beyond doubt (Copel.).

Mindanao.

* (4) **P. glauca**, *Mett.*, *Christ, Farnkr. d. Erde*, 173; *Bedd.*, *Ferns Br.I.*, I, tab. XC; *Lomaria glauca*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 162; — var. *philippinensis*, *Christ, Bull. Herb. Boiss.*, VI, 150.

Rhizome erect. Stipes erect, naked, ± 30 c.M. or more long, basal tubercles present. Fronds ovate, 30—150 c.M. long, 15—30 c.M. broad, pinnate, the apex pinnatifid, narrowed. Pinnae contiguous, linear, $1\frac{1}{4}$ —2 c.M. broad, sessile or the lower short-stalked, the apex acuminate, the edge sharply serrulate, but more distinctly towards the point, the base rounded or narrowed, often provided with a large gland beneath. Texture coriaceous; veins fine, copious, compact, simple or forked; under surface glaucous or white-powdery. Fertile fronds with the pinnae distant and narrowly linear.

Var. philippinensis: Pinnae of barren fronds scarcely acuminate,

minutely serrulate, separated by as much as their own breadth; terminal pinna like the lateral ones.

Malaya; North. India, Yunnan; — var. philippinensis: Luzon.

* (5) **P. pycnophylla**, *Mett.*, Christ, Farnkr. d. Erde, 175; Bedd., Ferns Br.I., I, tab. LII; *Lomaria pycnophylla*, Kze, Rac., Flor. Btz., I, 161; Hk., Spec. Fil., III, tab. CXLVIII; — var. *remota*, Christ, Bull. Herb. Boiss., VI, 150; — var. *mixta*, Copel., Philipp. Journ., I, Suppl., 154.

Rhizome stout, erect, woody. Stipes erect, naked, 15—40 c.M. long, basal tubercles present. Fronds ovate-lanceolate, \pm 60—150 c.M. long, 15—30 c.M. broad, the apex pinnatifid, narrowed. Pinnae numerous, linear, spreading or nearly so, contiguous, 1—1½ c.M. broad, the apex acuminate or subcaudate, the edge finely serrulate, but more strongly towards the point, the base truncate or subtruncate, mostly purveyed beneath with 1—2 conspicuous, prominent glands. Texture coriaceous; veins numerous, close, simple or forked. Fertile pinnae distant and narrowly linear.

Var. remota: Pinnae separated by \pm their own breadth.

Var. mixta: Fertile pinnae often dilated and sterile at the apex.

Malacca, Java, Borneo; Yunnan, North.India; — var. remota: Philippines; — var. mixta: Luzon.

(6) **P. euphlebia**, *Mett.*, Bedd., Handb. Ind. Ferns, 129; Ferns Br.I., II, tab. CLXV; *Lomaria euphlebia*, Kze, Hk. Bk., Syn. Fil., 183; Hk., 2^d Cent. of Ferns, tab. LXXXIX; *L. articulata*, Muell., Fragm., V, 174.

Rhizome stout, woody. Stipes 30 c.M. or more long, naked, erect, basal tubercles present. Fronds ovate-lanceolate, 30—60 c.M. long, 15—30 c.M. broad. Pinnae 2½ c.M. or more apart at the base, erecto-patent, linear, serrulate, 10—15 c.M. long, 1—2 c.M. broad, narrowed at the base on both sides and at the toothed apex. Texture subcoriaceous; veins prominent. Fertile fronds similar, but the pinnae more distant and narrowly linear.

New Guinea, Malacca; China, Japan.

(7) **P. tuberculata**, *Copel.*, Philipp. Journ., I, Suppl., 153.

Rhizome erect, short. Stipes close, 20—40 c.M. long, the fertile fronds on the longest stipes, the base bearing \pm 12 tubercles. Barren fronds \pm 60 c.M. long, 20—25 c.M. broad, the apex abruptly acuminate, the rachis quadrangular, naked, glossy, reddish-brown, bearing glands on the under side

at the base of the pinnae. Pinnae 30—40-jugate, nearly horizontal, the largest $\pm 1\frac{1}{4}$ c.M. broad, the base acute, the edge entire, the apex acuminate, serrate; lower pinnae short-stalked, higher adnate, highest confluent. Texture papyraceous; surfaces naked; veins conspicuous, simple or forked. Fertile fronds narrower, the pinnae spreading, ± 11 c.M. long, ± 2 —3 m.M. broad.

Luzon.

41. PELLAEA, *Link.*

Sori intramarginal, *either* dot-like to oblong, solitary and terminal on the veins, but soon running into a submarginal line *or* linear, placed on a submarginal receptacle uniting the vein tips. Indusia roundish, contiguous or linear, consisting of the reflexed, modified margin of the frond.

Rhizome more or less creeping. Stipes seriate, not articulate to the rhizome, with a single, 2—4-archous vascular strand (∇ , Υ , \times). Fronds divided; leaflets mostly \pm stalked, more or less distinctly articulate, not adnate at the base; veins free or sometimes anastomosing, the apices not or hardly thickened. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 142.

Trop. Asia, Australia, Polynesia, America, Africa.

Arrangement of the sections.

- § 1. **PLATYLOMA.** Sori terminal on the veins, free, but soon running into a submarginal line. Veins always free. Diels, l. c., fig. 142, B.
- § 2. **PTERIDELLA.** Sori linear, on a submarginal receptacle uniting the vein tips. Veins free in malayan material. Diels, l. c., fig. 142, F.

§ 1. **PLATYLOMA.**

✧. *Fronds simply pinnate.*

×. *Stipes scattered. Lower pinnae not or hardly reduced.* (1) *P. falcata.*

×.×. *Stipes tufted. Lower pinnae much reduced.* (2) *P. Zippelii:*

✧.✧. *Fronds more compoundly divided.* (3) *P. pteroides.*

* (1) **P. falcata**, *Fée*, Hk. Bk., Syn. Fil., 151; *Platyloma falcatum*, *J. Sm.*, Bedd., Ferns S.I., tab. XXII; *Pteris falcata*, *Burm.*, (oldest name).

Rhizome wide-creeping, densely scaly; scales linear-lanceolate, with a broad or narrow black line in the centre, the margin brown. Stipes $7\frac{1}{2}$ —15 c.M. long, erect, more or less densely clothed with short (narrow) and longer (linear-lanceolate), ferruginous scales. Fronds 15—45 c.M. long, $2\frac{1}{2}$ —5 c.M. broad, linear-oblong to lanceolate, simply pinnate. Pinnae more or less numerous, nearly sessile, 1 — $2\frac{1}{2}$ c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, lanceolate or oblong, blunt or mucronate, often slightly falcate, cuneate or dilated and cordate at the base. Texture coriaceous; rachis densely scaly and tomentose, both surfaces nearly naked. Sori in broad marginal lines, soon hiding the narrow indusium.

Trop. Asia; Australia, Tasmania, New Zealand, New Caledonia.

(2) **P. Zippelii**, *Bk.*, Hk. Bk., Syn. Fil., 477; *Allosurus Zippelii*, *Miq.*, Ann. Mus. Bot. L. B., IV, 98.

Stipes short, tufted, scaly. Fronds lanceolate, \pm 45 c.M. long, \pm 7 c.M. broad, simply pinnate. Pinnae very numerous, opposite, patent, short-stalked, linear-lanceolate, obtuse, straight or slightly falcate, the central ones \pm $5\frac{1}{2}$ c.M. long, with the base \pm 5 m.M. broad, the lower ones much reduced. Texture firm, subcoriaceous; rachis clothed with subulate scales; upper surface naked, lower scaly on the midribs; veins simple or 2-fid, tumid at the tip.

New Guinea.

(3) **P. pteroides**, *Prantl*, Engl. Jahrb., III, 420; *Diels*, in Engl. & *Prantl*, Nat. Pfl.Fam., I⁴, fig. 142, G—H; *Cheilanthes pteroides*, *Sw.*, Hk. Bk., Syn. Fil., 131; Hk., Spec. Fil., II, tab. CI, A; *Adiantum pteroides*, *L.*, (oldest name).

Stipes 15—50 c.M. long, strong, erect, polished, dark-brown, naked. Fronds 20—40 c.M. long, 15—25 c.M. broad, deltoid, 5-pinnate; upper part simply pinnate, lower with several opposite pairs of wiry, erecto-patent branches growing gradually larger down, the lowest often again branched. Ultimate segments \pm 1 — $2\frac{1}{4}$ c.M. long, \pm $\frac{1}{2}$ —1 c.M. broad, oblong, entire, broadly rounded at both ends, sessile. Texture subcoriaceous; rachis polished; both surfaces naked. Sori small, roundish, distinct but contiguous; indusia oblong, contiguous, numerous.

? *Java; South. & West. Africa.*

§ 2. PTERIDELLA.

* (4) **P. timorensis**, *v. A. v. R.*

Rhizome short, the scales long, linear-subulate, brown, eroso-denticulate. Stipes approximate, 12—20 c.M. long, blackish, polished, fibrilloso-tomentose on the upper (inner) side, deciduously scaly at the base. Fronds lanceolate-oblong, 15—25 c.M. long, 8—12 c.M. broad, simply pinnate. Pinnae \pm 10—15 on each side, stalked, \pm spreading, \pm 1—1½ c.M. broad, the lower ones longer or smaller than the others, lanceolate, subacute, the edge recurved when dry, crenate throughout when barren, the base rounded or slightly cordate. Texture coriaceous; rachis and stalks of the pinnae blackish, polished, tomentose on the upper side; surfaces naked; midribs conspicuous, black in the lower half; veins rather hidden. Sori on a marginal receptacle uniting the vein-tips, soon hiding the indusium. — Probably closely allied to the african *P. Doniana*, *Hk.*, (Spec. Fil., II, tab. CXXV, A).

Timor (Forbes, No. 5918).

42a. **DORYOPTERIS**, *J. Smith.*

Sori roundish or oblong, *either* solitary and terminal on the veins, but soon running into a submarginal line, *or* linear and placed on a submarginal receptacle uniting the vein tips. Indusium linear, continuous, consisting of the reflexed, modified margin of the frond.

Rhizome short, the scales costate. Stipes black or dark-coloured, polished, not articulate to the rhizome, with a single, 2—4-archous vascular strand (∇ , Υ , \times). Fronds simple or divided, if divided the segments adnate or joined, not articulate at the base; veins free or anastomosing, the tips not or hardly thickened. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 143.

Pantropical.

Arrangement of the sections.

- § 1. **DORYOPTERIDASTRUM**. Sori solitary and terminal on the veins, but soon running into a submarginal line. Veins free, or anastomosing near or towards the margin. Diels, l. c., fig. 142, B (sub Pellaea).
- § 2. **EUDORYOPTERIS**. Sori linear, on a submarginal receptacle uniting the vein tips. Veins anastomosing copiously. Diels, l. c., fig. 142, F (sub Pellaea).

§ 1. DORYOPTERIDASTRUM.

* (1) **D. concolor**, *Kuhn*, Copel., Polypod. Philipp., 94; *Pellaea concolor*, *Bk.*, Christ, Farnkr. d. Erde, 157; *P. geraniaefolia*, *Fée*, Hk. Bk., Syn. Fil., 146; *Pteris geraniaefolia*, *Raddi*, Rac., Flor. Btz., I, 146; Hk., Icon. Pl., X, tab. CMXV; *Bedd.*, Ferns S.I., tab. XXXVII; *Pt. concolor*, *L. & F.*, (oldest name).

Rhizome short-creeping, clothed with short, lanceolate scales. Stipes numerous, subtufted, 5—20 c.M. long, naked, erect, blackish, scaly at the base. Fronds ovate-deltoid, 5—10 c.M. each way, cut down nearly or rarely quite to the rachis into 3—5 pinnae on each side, the lowest pair the largest, with the segments on the lower side produced and deeply lobed with linear-oblong lobes, the segments of the fertile fronds not always deeply lobed. Texture coriaceous; lateral veins once forked; rachis and costae dark-coloured and polished. Sori in broad, marginal lines.

Pantropical.

§ 2. EUDORYOPTERIS.

(2) **D. ludens**, *J. Sm.* Copel., Polypod. Philipp., 94; *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 145, C; *Pellaea ludens*, *Prantl*, Engl. Jahrb., III, 419; *Pteris ludens*, *Wall.*, Hk. Bk., Syn. Fil., 166; *Litobrochia pedata*, *Bedd.*, Ferns Br.I., I, tab. XXVI; *L. ludens*, *Bedd.*, l. c., tab. XXVII.

Rhizome creeping, the scales adpressed, linear, brown, pale at the margin. Stipes polished, blackish, naked, sometimes with a few scales and often with dusky, subtomentose pubescence at the base. Barren fronds on stipes 7—10 c.M. long, triangular or ovate-lanceolate, deeply cordate or auriculate, sometimes with 2—3 pairs of lobes, acuminate, with an entire or crisped margin. Texture coriaceous, rather flaccid; costa polished, surfaces naked; veins hidden, anastomosing copiously. Fertile fronds on stipes to 30 c.M. long, 10—20 c.M. each way, cut down into 5—7 lanceolate or linear-lanceolate lobes, the lowest of which are deeply forked, with the lower segment deflexed. Sori in narrow marginal lines.

Malacca, Philippines; Br. India.

42b. SCHIZOSTEGE, Hillebrand.

Sori roundish, submarginal, on terminal T-shaped expansions of the veins or veinlets or uniting the apices of the fork branches of forked veins. Indusium of the same shape, remarkably broad. Spores tetrahedral.

Rhizome erect. Stipes not articulate to the rhizome, with a single U-shaped vascular strand. Fronds divided; veins free or joining.

Hawaii, Mindanao.

×. *Pinnae pinnatifid* $\frac{3}{4}$ -way down to the costa, the segments entire.

(1) *S. pachysora*.

×.×. *Pinnae pinnatifid* nearly or quite to the costa, the segments crenate or lobed.

(2) *S. calocarpa*.

(1) ***S. pachysora*, Copel.**, Philipp. Journ., I, Suppl., 155.

Rhizome short, erect, thick, crowned with fuscous, lanceolate scales. Stipes tufted, \pm 50 c.M. long, yellow-fuscous, clothed with scales of the same colour. Fronds \pm 40 c.M. long, \pm 20 c.M. broad, 2-pinnatifid. Lowest pinnae somewhat deflexed, \pm 10 c.M. long, sessile, forked at the base on the lower side, the branches subacute, cut $\frac{3}{4}$ -way down to the costa into entire, obtuse lobes 5—10 m.M. long, 5—7 m.M. broad, central pinnae similarly pinnatifid, decurrent, nearly confluent, the higher ones gradually reduced. Texture carnosu-subcoriaceous; surfaces naked; veins hidden, joining in 1 row of costal areolae, rarely forming a second row of areolae in the segments. Sori solitary on the vein tips, free or confluent.

Mindanao.

(2) ***S. calocarpa*, Copel.**, Philipp. Journ., I, Suppl., 155, tab. XII.

Like *S. pachysora*, Copel., but: Fronds 35—40 c.M. long, \pm 20 c.M. broad, acute, 2-pinnate or sub-2-pinnate. Lower pinnae oblique, deltoid, the following lanceolate, 10 c.M. long, \pm 3 c.M. broad, acuminate, cut down nearly or quite to the costa. Secondary segments \pm 15 m.M. long, \pm 4 m.M. broad, decurrent, crenate or cut $\frac{1}{2}$ -way down to the costula. Texture carnosu-subcoriaceous; main rachis cinereous, marginate; veins free. Sori not confluent.

Mindanao.

43. HISTIOPTERIS, *J. Smith.*

Sori linear, submarginal, on a receptacle connecting the vein tips, commonly continuous around the sinuses. Indusium linear, consisting of the reflexed, modified margin. Spores 2-lateral.

Rhizome wide-creeping; stipes polished, stramineous to purplish, not articulate to the rhizome. Fronds at least 2-pinnate; lowest pinnulae shorter than the next in order, stipule-like, close to the main rachis; veins free or anastomosing. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 155, F.

Pantropical and farther south.

(1) **H. montana**, *Copel.*, Polypod. Philipp., 104.

Rhizome wide-creeping, beset with minute, brown scales. Stipes firm, 10—20 c.M. high, erect, black-brown. Fronds deltoid, 15—30 c.M. long, often broader than long, 2—3-pinnatifid. Lowest pinnae somewhat larger than the next following ones, pinnate. Pinnulae sinuate or pinnatifid, the lowest close to the rachis. Lobes often deltoid-ovate, obtuse, entire. Texture coriaceous; veins prominent above, anastomosing. Indusium broad; margin everywhere rolled in, giving the segments a lomarioid appearance.

Philippines.

* (2) **H. incisa**, *J. Sm.* (not *Agardh.*), *Copel.*, Polypod. Philipp. 104; Diels, in Engl. & Prantl, Nat Pfl.Fam., I⁴, fig. 155, F; *Pteris incisa*, *Thb.*, *Rac.*, *Flor. Btz.*, I, 159; *Pt. Vespertilionis*, *Lab.*, *Nov. Holl.*, II, tab. CCXLV; *Litobrochia incisa*, *Pr.*, *Bedd.*, *Handb. Ind. Ferns*, 120; — var. *aurita*; *Pt. aurita*, *Bl.*, *Enum.*, 215; *L. aurita*, *Pr.*, *Bedd.*, *Fern S.I.*, tab. CCXXI; — var. *integri-folia*, *Bedd.*, *Handb. Ind. Ferns*, *Suppl.*, 25.

Rhizome wide-creeping, clothed with narrow, linear-lanceolate scales. Stipes distant, stout, erect, sometimes slightly muricate towards the base, 8—150 c.M. long. Fronds 10—600 c.M. long, or even longer (!), 2—3-pinnate at the base, often scandent or twining in the largest forms, with long, spreading, rigid branches. Central pinnae opposite, with numerous, lobed or pinnatifid pinnulae $\pm 2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, ± 8 —20 m.M. broad, the lowest pinnulae shorter, adpressed to the rachis, reduced in size, stipule-like, lowest pinnae more compound. Texture herbaceous to coriaceous; surfaces naked or the under one glaucous; veins free or anastomosing.

Var. aurita: Stipule-like pinnulae quite entire.

Var. integrifolia: Pinnae petiolate; pinnulae sub-petiolate, long, linear, acuminate, entire, obliquely cordate at the base, the basal pair not auricle-like, distant from the rachis. Sori continuous almost to the finely acuminate apex.

Pantropical and farther south.

44. ONYCHIUM, *Haulfuss.*

Sori linear, submarginal, on a continuous receptacle connecting the vein tips. Indusium consisting of the reflexed, modified margin, broad enough to reach \pm the costa. Spores tetrahedral.

Rhizome creeping. Stipes pale-coloured or brown, with 1—2 vascular strands, not articulate to the rhizome. Fronds finely divided, fertile and sterile divisions \pm unlike; veins free. Hk. Bk., Syn. Fil., tab. III, fig. 27.

Tropical and extra-tropical Asia, Africa & America.

§. *Sporangia yellow.*

*. *Fronds 10—15 c.M. long. Indusium whitish.* (1) *O. tenue.*

.. *Fronds 15—60 c.M. long. Indusium yellow.* (2) *O. siliculosum.*

§.§. *Sporangia brown.* (3) *O. japonicum.*

(1) ***O. tenue***, *Christ*, Bull. Herb. Boiss., 2^e Série, I, 451.

Rhizome short, erect. Stipes densely tufted, \pm 5 c.M. long, slender. Fronds 10—15 c.M. long, narrowly ovate. Pinnae and pinnulae erect; ultimate segments narrow, cuneate and 5-lobate when sterile, lanceolate and acuminate when fertile. Sori pale-yellow; indusium whitish.

Java, New Guinea, Philippines.

* (2) ***O. siliculosum***, *C. Chr.*, Ind. Fil.; *O. auratum*, *Klf.*, Rac., Flor. Btz., I, 165; *Bedd.*, Ferns S.I., tab. XXX; *Cryptogramme aurata*, *Prantl*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 279; *Pteris chrysocarpa (chrysosperma)*, *Hk. & Grev.*, Ic. Fil., tab. CVII; *Pt. siliculosa*, *Desv.*, (oldest name).

Rhizome short, creeping, clothed with subulate-lanceolate, brown, spreading scales. Stipes 10—35 c.M. long, stout, erect, naked, stramineous or pale-

brown. Fronds 15—60 c.M. long, 7—20 c.M. broad, ovate or lanceolate-deltoid, 3-pinnate to 4-pinnatifid. Lower pinnae subdeltoid, erecto-patent. Pinnulae and segments numerous; ultimate divisions dimorphous, the sterile ones often obovate-cuneate, 3-fid at the apex, not more than 5—10 m.M. long and $\frac{1}{2}$ —1 m.M. broad, the fertile ones pod-like, $\frac{1}{2}$ —2 $\frac{1}{2}$ c.M. long, 1—2 m.M. broad. Texture coriaceous; rachis and both surfaces naked. Indusium membranaceous and like the sori long, linear or linear-oblong, of a rich golden-yellow colour.

Trop. Asia.

* (5) **O. japonicum**, *Hæ.*, *Rac.*, *Flor. Btz.*, I, 164; *O. lucidum*, *Spr.*, *Bedd.*, *Ferns Br.I.*, I, tab. XXI; *Cryptogramme japonica*, *Prantl*, *Diels*, in *Engl. & Prantl*, *Nat. Pfl.Fam.*, I⁴, 279; *Trichomanes japonicum*, *Thb.*, (oldest name).

Rhizome creeping, clothed with narrow, lanceolate scales. Stipes stout, erect, naked, glossy, 10—30 c.M. long, stramineous or pale-brown. Fronds ovate-lanceolate, 20—35 c.M. long, 7—20 c.M. broad, 5-pinnate or 4-pinnatifid. Lower pinnae lanceolate-deltoid, erecto-patent. Pinnulae and segments numerous, the ultimate divisions uniform or nearly so, 3-lobate, copious, the lobes mucronate, 4—10 m.M. long, 1—2 $\frac{1}{2}$ m.M. broad. Texture herbaceous to subcoriaceous; rachis and both surfaces naked. Sori short, oblong; indusium pale or brown, membranaceous; capsules brown or pale-brown.

Java, Timor, Philippines; Japan, China, North. India.

45a. PTERIS, *Linné.*

Sori linear, submarginal, on a receptacle uniting the vein tips, occupying both (posterior and anterior) sides of the ultimate segments, generally interrupted at the tops of the sinuses between the ultimate lobes. Indusium of the same shape, consisting of the reflexed, modified margin of the frond. Spores globose to tetrahedral.

Rhizome creeping or erect, scaly. Stipes not articulate to the rhizome, traversed by a single, mostly ∇ - or \cup -shaped vascular strand. Fronds commonly variously divided, sometimes variegated, i. e. the upper surface provided with broad, whitish or pale-coloured bands running down the costae; veins free or anastomosing. *Hk. Bk.*, *Syn. Fil.*, tab. III, fig. 31, a—c; *Diels*, in *Engl. & Prantl*, *Nat. Pfl.Fam.* I⁴, fig. 154.

Arrangement of the sections and groups.§ 1. **EUPTERIS.** Veins free.

- a. **Pinnatifidae.** Fronds pinnatifid throughout, at best pinnate at the very base.
- b. **Pinnatae.** Fronds 3-foliolate or pinnate, the lowest pinnae simple, or forked with 2—3 undivided segments in the barren fronds.
- c. **Compositae.** Fronds 3-foliolate or pinnate, the lowest pinnae at least pinnatifid in the barren fronds.
- *. **Bipinnatifidae.** Fronds not 3- or 5-partite; lowest pinnae not much longer than the next higher ones and similarly pinnatifid or pinnate, with the lowest segments on the lower side not rarely elongate and divided again.
- *.*. **Partitae.** Fronds 3- or 5-partite; lateral branches nearly as long as the central one and similarly 2-pinnatifid, with the lowest segments on the lower side not rarely elongate and divided again.

§ 2. **CAMPTERIA.** Veins slightly anastomosing, forming 1 single row of costal areolae on each side, otherwise free.§ 5. **LITOBROCHIA.** Veins anastomosing more copiously.*Key to the sections and groups.*

- a. Veins free. § 1. **EUPTERIS.**
- b. Fronds pinnatifid, at best pinnate at the very base.
 - § 1, a. **Pinnatifidae.**
 - b.b. Fronds 3-foliolate, pinnate or more compoundly divided.
 - c. Lower primary segments simple, or forked at the base with 2—3 undivided secondary segments in the barren fronds.
 - § 1, b. **Pinnatae.**
 - c.c. Lower primary segments at least pinnatifid in the barren fronds
 - § 1, c. **Compositae.**
 - d. Fronds not 3- or 5-partite, i. e. the lowest pinnae not much longer than and similarly pinnatifid or pinnate as the next higher ones, but not rarely forked at the base with similarly divided branches. § 1, c*. **Bipinnatifidae.**
 - d.d. Fronds 3- or 5-partite, i. e. the fronds divided into 3 or

5 branches clustered at the apex of the stipe, the lateral branches often nearly as long and similarly 2-pinnatifid as the central one, not rarely forked near the base with more or less divided secondary branches. § 1, c**. **Partitae.**

a.a. Veins free except that the central ribs (costulae) of the ultimate segments are more or less connected at the very base by arching veins forming 1 single row of costal areolae on each side. § 2. **CAMPTERIA.**

a.a.a. Veins anastomosing more copiously than in **a.a.** § 3. **LITOBROCHIA.**

§ 1. **EUPTERIS.** a. **Pinnatifidae.**

* (1) **Pt. reducta**, *Bk.*, Journ. of Bot., XVIII, 211; *Pt. biauwaita*, *L.*, var., C. Chr., Ind. Fil.

Stipes tufted, 15—25 c.M. long, erect, naked, the base often dark-coloured and slightly scaly. Fronds 25—55 c.M. long, 5—7 c.M. broad, linear-lanceolate, cut down in the upper part nearly, in the lower nearly or quite to the costa into 50—40 pairs of crowded, linear lobes below the caudate apex. Lobes ± close or slightly imbricate, rather spreading, subfalcate, 6—9 m.M. broad, the apex bluntly rounded, entire or slightly crenate, the edges parallel, entire, the lower lobes hardly shorter than the others. Texture firm; costules and veins prominent, the latter simple or forked: surfaces naked. Sori not reaching the apex of the lobes, often (but not always) extending to the top of the sinuses; indusium narrow.

Sumatra.

§ 1. **EUPTERIS.** b. **Pinnatae.**

a. *Fronds dimorphous. Lowest barren pinnae simple, lowest fertile ones often contracted, mostly forked or pinnatifid on the upper side only.* (2) *Pt. rangiferina.*

a.a. *Fronds uniform. Lowest pinnae simple or 2—3-furcate, the fertile ones sometimes contracted, not divided as in a.*

b. *Pinnae cordate, truncate or rounded at the base.*

‡. *Veins close; lower pinnae ± sessile.*

*. *Lowest pinnae ± largest. Rachis reddish or purpureous, deciduously villose.* (3) *Pt. orientalis.*

- *.*. *Central or highest pinnae largest. Rachis yellowish-green, naked or slightly scaly.* (4a) *Pt. longifolia.*
- ✧.✧. *Veins distant; lower pinnae stalked.* (4b) *Pt. platysora.*
- b.b.** *Pinnae cuneate at the base.*
- c.** *Rachis pubescent.* (5) *Pt. opaca.*
- c.c.** *Rachis naked.*
- d.** *Largest lateral pinnae of fully developed fronds more than 27 $\frac{1}{2}$ c.M. long.*
- ×. *Fronds middle-sized. Lowest pinnae 2—3-furcate.* (6) *Pt. Treubii.*
- ×.×. *Fronds large. Lowest pinnae simple.* (7) *Pt. moluccana.*
- d.d.** *Largest lateral pinnae under 27 $\frac{1}{2}$ c.M. long.*
- e.** *Barren fronds with 3—5 pinnae clustered at the apex of the stipe.* (8) *Pt. stenophylla.*
- e.e.** *Barren fronds with the pinnae not clustered at the apex of the stipe.*
- f.** *Barren pinnae entire or nearly so, at best serrate or toothed towards the apex.*
- g.** *Lowest pinnae simple or 2-furcate.* (9) *Pt. pellucida.*
- g.g.** *Lowest pinnae 3-lobate or 3-furcate.* (10) *Pt. papuana.*
- f.f.** *Barren pinnae more copiously serrate throughout.*
- ✧. *Stipes stramineous or pale-brown.* (11) *Pt. cretica.*
- ✧.✧. *Stipes black.* (12) *Pt. melanocaulon.*

* (2) ***Pt. rangiferina***, *Pr.*, Ann. Mus. Bot. L. B., IV, 95; *Pt. Dalhousiae*, *Rac.* (not *Hk.*), Flor. Btz., I, 155; *Pt. laevigata*, *Zipp.*, MS. in Herb. Bog.

Rhizome erect, thick, densely clothed with brownish, tomentose hairs. Stipes 50—150 c.M. long, erect, naked, the outer side convex, the inner 5-sulcate, the base reddish-brown, the rest stramineous. Fronds dimorphous. Barren fronds lanceolate-oblong, 45—250 c.M. long, pinnate with \pm 8—12 pairs of stalked, erecto-patent pinnae below the terminal one. Terminal and lateral pinnae similar, lanceolate, 15—50 c.M. long, 2—4 c.M. broad, the

point acuminate, the sides \pm parallel, the base broadly cuneate, often slightly unequal-sided, the edge entire, bordered by a continuous, narrow, \pm pellucid line, the stalks 1—40 m.M. long. Texture coriaceo-chartaceous when dry; rachis and both surfaces naked; upper surface glossy, lower pale, opaque; veins numerous, simple or forked, the tips dilated and often confluent. Fertile fronds with the pinnae linear, acuminate, 5—8 m.M. broad when simple, the base cuneate; higher pinnae simple, the lower with a long, terminal segment and 1—5 distant, linear lobes near the base on the upper side, the lower side undivided or rarely lobed like the upper. Indusium membranaceous. — Besides both barren and fertile fronds there are semi-fertile ones, which have the pinnae broad or narrow, simple or forked at the base with stalked segments, the sori often interrupted into many shorter, oblong ones. — *Pt. Dalhousiae*, *Hk.* is quite another plant.

Java.

* (5) ***Pt. orientalis*, v. *A. v. B.***

Stipe and rachis reddish or purpureous, deciduously villose, at length naked. Fronds \pm 40—55 c.M. long, \pm 50 c.M. broad in the broadest part. Pinnae in \pm 20—50 pairs below the terminal one, simple, linear, erectopate, sessile, close or imbricate, the lowest largest, or a little smaller than the next higher ones, 15—25 c.M. long, 1—2 c.M. broad near the base, narrowed gradually towards the apex, the point somewhat bluntish, the margin entire or serrate in the barren pinnae, often crispate, the base rounded, the highest pinnae the smallest, 5—15 c.M. long. Texture coriaceo-membranaceous, pellucid; costae prominent and rounded beneath; veins fine, numerous, close, not very oblique, not rarely nearly horizontal, deciduously and sparingly pilose on both sides, at length naked. Sori long; indusium narrow. — The description is taken from specimens determined as *Blechnum orientale*, *L.* probably by the resemblance with that species.

Banda Islands (Treub).

* (4a) ***Pt. longifolia*, *L.***, *Rac.*, *Flor. Btz.*, I, 151; *Bedd.*, *Ferns S.I.*, tab. XXXIII; — var. *diversifolia*, *Rac.*, l. c.; *Pt. diversifolia*, *Sw.*, *Syn.*, 96, 288.

Rhizome short, the scales yellowish, linear-lanceolate. Stipes clustered, 10—50 c.M. long, erect, pale-green, more or less clothed towards the base with pale scales. Fronds 10—100 c.M. long, 5—20 c.M. broad, simply pinnate, oblong or lanceolate. Pinnae sessile, linear, the central ones the longest,

acuminate, entire or serrulate when barren, \pm 5—12 m.M. broad, truncate or cordate, sometimes slightly auricled at the base; lower pinnae gradually dwarfed. Texture subcoriaceous; rachis naked or slightly scaly; surfaces naked; veins close, mostly once forked. Indusium narrow, membranaceous.

Var. diversifolia: Fronds narrowed gradually towards the base; the lowest pinnae reduced, roundish or oblong, the highest the longest. United to the type by intermediate forms.

Tropical and subtropical, everywhere.

(4b) **Pt. platysora**, *Bk.*, Journ. of Bot., XVIII, 211.

Stipes 60—90 c.M. long, naked, stramineous upwards, castaneous towards the base. Fronds 45—60 c.M. long, \pm 15 c.M. broad, oblong-lanceolate, simply pinnate. Pinnae \pm 25, simple, linear, erecto-patent, the lower distinctly stalked, 12—15 c.M. long, 6—9 m.M. broad, entire, 5—8 c.M. apart, the upper sessile, the terminal one like the others. Texture moderately firm; surfaces naked; veins distant, simple or forked. Sori continuous, not reaching the apex of the pinnae; indusium broad and distinct, but hidden when the fruit is fully developed. — Near *Pt. longifolia*, *L.*

Sumatra.

(5) **Pt. opaca**, *J. Sm.*, Hk. Bk., Syn. Fil., 154; Hk., Spec. Fil., II, tab. CXIV, A.

Stipes strong, erect, pale-coloured, finely pubescent like the rachis. Fronds 60—200 c.M. long, oblong. Pinnae in numerous \pm opposite pairs, linear, cuneate at the base, the edge quite entire towards the point, the largest \pm 50 c.M. long, \pm 1—1½ c.M. broad. Texture coriaceous; surfaces naked except on the midribs beneath; veins sunk in the frond and only visible as faint striations. Indusium narrow, brownish.

Celebes, Batjan, Philippines.

* (6) **Pt. Treubii**, *v. A. v. R.*

Stipes \pm 20 c.M. long, brownish and naked like the rachis. Fronds 50—45 c.M. long, pinnate. Terminal pinna linear, 20—30 c.M. long, \pm 5 m.M. broad, the base gradually narrowed, the edge entire, the apex acuminate, spinuloso-denticulate. Lateral pinnae 3—4 on each side, erecto-patent, similar to the terminal one, the highest ones simple or forked at the base, the lowest the longest, to 40 c.M. long, \pm 5 m.M. broad, stalked, 2—5-partite, the stalks \pm 1½—2 c.M. long; segments of the divided pinnae as long,

sometimes forked again. Texture moderately firm, but thin; surfaces naked; veins obscure, spreading, simple or forked. Sori very long, not reaching the serrate apex of the pinnae; indusium narrow. — Barren fronds not seen.

Ceram (Treub).

(7) **Pt. moluccana**, *Bl.*, Hk. Bk., Syn. Fil., 154; Hk., Spec. Fil., II, tab. CXII, B; *Pt. indica*, *Christ* (not *Lam.*), Farnkr. d. Erde, 165.

Stipes strong, erect, naked, polished, dark-brown. Fronds 60—100 c.M. long, oblong. Pinnae in numerous, nearly opposite pairs, cuneate at the base, spinuloso-serrate towards the apex, the largest 30—45 c.M. long, 1—3 c.M. broad. Texture subcoriaceous; rachis naked; both surfaces shining; veins conspicuous, very close, \pm 20—25 to a c.M., usually simple. Indusium narrow, membranaceous.

Malaya; West. Polynesia.

* (8) **Pt. stenophylla**, *Wall.*, Hk. & Grev., Ic. Fil., tab. CXXX; *Pt. cretica*, *L.*, var. *stenophylla*, Hk. Bk., Syn. Fil., 154; *Pt. pellucida*, *Pr.*, var. *stenophylla*, *Bedd.*, Handb. Ind. Ferns, 107.

Near *Pt. pellucida*, *Pr.*, but the fronds with 3—5 pinnae clustered at the apex of the stipe, the central pinna sometimes with a pair of basal segments. Fertile fronds often on the longest stipes, with the pinnae longer and narrower than in the barren ones.

Malacca, Philippines; North. India.

* (9) **Pt. pellucida**, *Pr.*, *Rac.*, Flor. Btz., I, 152; *Bedd.*, Ferns S.I., tab. XXXVIII; *Pt. venusta*, *Kze.*, Bot. Zeit., 1848, 195; *Pt. venulosa*, *Bl.*, Enum., 209.

Rhizome erect or short-creeping, the scales linear-lanceolate, blackish. Stipes 20—50 c.M. long, erect, naked, glossy, stramineous or brownish. Fronds 15—60 c.M. long, ovate-lanceolate or deltoid-ovate, pinnate. Pinnae in 3—6 opposite or alternate pairs below the sometimes very long terminal one, erect-patent, 10—20 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, linear-lanceolate, the base cuneate, the edge entire or repand to undulate, the apex acuminate, entire, serrate or toothed, the point \pm bluntish; higher pinnae \pm decurrent, free or joining with the terminal one, the lowest \pm stalked, simple or forked at the base, the fertile ones whether or not contracted, not rarely $\frac{1}{2}$ \times as broad as the barren ones. Texture papyraceous to coriaceous, pellucid; veins

visible, fine, simple or forked, close, \pm 10—15 to a c.M. Sori not reaching the apex. — The young plants have the fronds ternate or quinate.

Malaya; North. India.

(10) **Pt. papuana**, *Ces.*, Fel. Becc. Polin., 3 & 7.

Near *Pt. pellucida*, *Pr.*, but much finer. Pinnae more approximate, about $\frac{1}{3}$ narrower and $\frac{1}{2}$ shorter, entire, the highest ones simple, the following slightly 2-lobed, the lower 3-furcate or 3-lobed, petiolate. Sori long, reaching the apex of the pinnae except on the upper side.

New Guinea.

* (11) **Pt. cretica**, *L.*, Hk. Bk., Syn. Fil., 154; Bedd. Ferns S.I., tab. XXXIX.

Stipes 15—50 or even to 60 c.M. long, erect, wiry, naked, stramineous or pale-brown, polished. Fronds 15—40 c.M. long, 15—25 c.M. broad, ovate, sometimes subflabellate. Pinnae usually in 2—6 pairs below the terminal one, the upper ones sometimes more or less decurrent, sessile, ascending, the longest linear-lanceolate, 7—20 c.M. long, $\frac{1}{4}$ —2 c.M. broad, the sterile ones not seldom considerably the broadest, acuminate, narrowed towards the base, spinuloso-serrate at the edge, the lower ones stalked, often cleft down nearly or quite to the base into 2—3 linear segments. Texture coriaceous, but not thick; rachis and surfaces naked; veins fine, close, parallel, simple or forked. Indusium narrow, pale, membranaceous. — Exceedingly variable.

Tropical and extra-tropical all round the world.

* (12) **Pt. melanocaulon**, *Fée*, Copel., Polypod. Philipp., 100; *Pt. cretica*, *L.*, var. *melanocaulon*, Hk. Bk., Syn. Fil., 154; C. Chr., Ind. Fil.; *Pt. Treacheriana*, *Bk.*, Journ. of Bot., XVII, 65.

Rhizome erect, short. Stipes tufted, slender, naked, blackish, polished, fragile, \pm 10—30 c.M. long. Fronds ovate-lanceolate, simply pinnate, 15—36 c.M. long. Pinnae stalked, linear, arcuate, suberect, 5—25 c.M. long, 5—5 m.M. broad, narrowed towards the base, serrate or toothed in the barren fronds, crenate and sterile towards the apex in the fertile fronds, the higher ones \pm decurrent, the lowest often with 2—3 fork branches at the base. Texture rather firm; rachis and surfaces naked, (\pm glaucous?); veins generally once forked, sometimes scarcely visible. Indusium mostly broad.

Philippine and Sulu Islands.

§ 1. EUPTERIS. c. **Compositae**. * **Bipinnatifidae**.

- a.** *Pinnae with a long, undivided terminal segment and mostly few, rarely pectinate lateral ones near the base.*
- b.** *Fronde dimorphous.* (15) *Pt. ensiformis*.
- b.b.** *Fronde uniform or nearly so.*
- c.** *Main rachis winged by the decurrent bases of the pinnae.*
(14) *Pt. serrulata*.
- c.c.** *Main rachis at best winged towards the apex.*
- d.** *Pinnae divided on the lower side only.*
- ***. *Lowest pinnae with 2 large segments on the lower side.*
(15) *Pt. sumatrana*.
- *.*.** *Lowest pinnae with more than 2, not very large segments on the lower side.* (16) *semipinnata*.
- d.d.** *At least the lower pinnae divided on both sides.*
- e.** *Pinnae generally with 1—6 segments on each side below the long terminal one.*
- f.** *Lower pinnae with the lateral segments short, close or approximate.*
- g.** *Secondary barren segments serrate throughout.*
- h.** *Segments somewhat distant, decurrent.*
(17) *Pt. dispar*.
- h.h.** *Segments close, not decurrent.*
(18) *Pt. distans*.
- g.g.** *Secondary barren segments entire or at best spinulose serrate at or towards the apex.*
(19) *Pt. heteromorpha*.
- f.t.** *Lower pinnae with the segments long, distant.*
(20) *Pt. Dalhousiae*.
- e.e.** *Pinnae with numerous lateral segments on each side below an often very long terminal one.*
(21) *Pt. Beccariana*.
- a.a.** *Pinnae pectinato-pinnatisect throughout, with a rather short, rarely subcaudate point and ± many lateral segments on each side.*
- 1.** *Lateral pinnae few, 1—2 on each side below the terminal one.*
- ***. *Veins indistinct.* (22a) *Pt. Grevilleana*.
- *.*.** *Veins distinct.* (22b) *Pt. pluricaudata*.
- 1.1.** *Lateral pinnae more numerous, at least 3 on each side.*
- 2.** *Fronde mostly under 125 c.M. long.*

3. *Pinnae not pinnatifid quite to the base, the lowest segments distinctly decurrent.*
 ×. Veins 2-furcate (23) *Pt. longipinnula.*
 ×.×. Veins 4-furcate. (24) *Pt. torricelliana.*
- 3.3. *Pinnae pinnatisect quite to the base, the lowest segments not or hardly decurrent.*
4. *At best the lowest pinnae forked at the base.*
5. *Stipes and rachises green, stramineous or pale-brown.*
6. *Ultimate segments entire or slightly serrate, herbaceous, papyraceous or coriaceous.*
7. *Upper surface not bluish.*
8. *Ultimate segments joined at the base.*
 (25a) *Pt. quadriaurita.*
- 8.8. *Ultimate segments generally free at the base.*
 (25b) *Pt. parviloba.*
- 7.7. *Upper surface bluish.* (25c) *Pt. caesia.*
- 6.6. *Ultimate segments appresso-serrate, membranaceous.*
 (26) *Pt. amoena.*
- 5.5. *Stipes and rachises black or dark-brown.*
- ✧. *Fronds rather small; ultimate segments erecto-patent, not conspicuously dilated at the base.*
 (27) *Pt. furcans.*
- ✧.✧. *Fronds rather large; ultimate segments horizontally spreading, conspicuously dilated at the base.*
 (28) *Pt. decussata.*
- 4.4. *Several lower pinnae forked at the base.* (29) *Pt. Walkeri.*
- 2.2. *Fully developed fronds more than 125 c.M. long.* (50) *Pt. excelsa.*

* (13) ***Pt. ensiformis***, *Burm.*, *Rac.*, *Flor. Btz.*, I, 154; *Pt. crenata*, *Sw.*, *Hk.*, *Spec. Fil.*, II, tab. CXXVII, A; *Bedd.*, *Ferns S.I.*, tab. XXXV; — var. *Victoriae*, *Bk.*, *Gard. Gron.*, III⁷, 576; *Pt. Victoriae*, *hort.*; — var. *permixta*, *Christ*, *Philipp. Journ.*, II, 173.

Rhizome creeping, sparingly clothed with deltoid-lanceolate, slightly adpressed scales. Stipes 5—50 c.M. long, those of the barren fronds the shortest, slender, naked, erect, stramineous. Fronds 10—50 c.M. long, the barren ones the shortest, ± 5—15 c.M. broad, with a long terminal pinna and 2—7 pairs of lateral ones. Pinnae of the barren fronds slightly compound, the

lower ones cut down to the rachis below into 1—3 pairs of oblong, sharply toothed pinnulae, which are often $1\frac{1}{2}$ c.M. broad, the higher ones simple, decurrent, often joined. Texture herbaceous to subcoriaceous; rachis and surfaces naked; veins not very conspicuous, often 1—2 \times forked. Pinnae of fertile fronds long, linear, narrow, entire or toothed at the apex, erect, simple, forked or pinnate at the base with linear segments. — Exceedingly variable.

Var. Victoriae: Fronds with broad, whitish bands running down the costae.

Var. permixta: Apex of fertile fronds elongated into a linear, caudiform appendage much exceeding the plant, furnished with irregular, short, obtuse lobes.

Malaya; Trop. Australia, Polynesia, China, North. India.

* (14) **Pt. serrulata, L. f.**, Hk. Bk., Syn. Fil., 155; *Pt. multifida, Poir.*, C. Chr., Ind. Fil.

Stipes 15—25 c.M. long, erect, wiry, naked, pale or brownish. Fronds 20—45 c.M. long, 15—25 c.M. broad, ovate, pinnate, the main rachis winged by the decurrent and confluent bases of the pinnae, the wing growing gradually narrower downwards, sometimes wanting between the 2 lowest pinnae, $3\frac{1}{2}$ —5 m.M. broad at the top of the rachis. Pinnae 4 or more on each side, opposite, the upper ones simple, often 10—15 c.M. long, 3—7 m.M. broad, acute or acuminate, with the edges \pm parallel, serrate in the barren region, the lower ones with several (rarely 1 or few), long, linear, erecto-patent pinnulae on each side. Texture scarcely coriaceous; rachis and both surfaces naked; veins distinct, simple or forked. Indusium slightly intramarginal, narrow, membranaceous.

In cultivation. — *Orig. loc.: China, Japan.*

(15) **Pt. sumatrana, Bl.**, Journ. of Bot., XIX, 367.

Stipes naked, castaneous like the rachis, \pm 15 c.M. long. Fronds obovate, 20—25 c.M. long, pinnate. Pinnae \pm 5—6-jugate below the simple terminal one; upper lateral pinnae decurrent to the next lower pair, simple, linear, 7—10 c.M. long, 8—12 m.M. broad, the lowest ones with 2 large segments on the lower side, the apices strongly inciso-crenate. Texture subcoriaceous; surfaces naked; veins obscure, immersed, fine, simple or forked. — Near *Pt. serrulata, L. f.*

Sumatra.

* (16) **Pt. semipinnata**, *L.*, *Rac.*, *Flor. Btz.*, I, 153; *Hk.*, *Gard. Ferns*, tab. LIX; *Bedd.*, *Ferns S.I.*, tab. XXXIV; *Pt. dimidiata*, *Willd.*, *Spec.*, V, 581; — var. *inaequalis*; *Pt. inaequalis*, *Bk.*, *Journ. of Bot.*, XIII, 199.

Rhizome creeping, clothed with blackish-brown, linear-lanceolate, finely acuminate scales. Stipes 20—50 c.M. long, strong, erect, naked, stramineous, reddish or bright-brown. Fronds 15—45 c.M. long, 10—25 c.M. broad, ovate-lanceolate, the upper part cut down nearly or quite to the rachis into close, entire, linear or oblong, \pm falcate or straight segments, the lower $\frac{2}{3}$ part with 4—8 pairs of opposite, distant pinnae with a \pm long entire point and a broad, undivided, leafy membrane on the upper side, but the lower side with several linear or oblong lobes or pinnulae 2—5 c.M. long, with the base \pm decurrent; barren segments finely serrate. Texture subcoriaceous; rachis and surfaces naked; veins simple or forked. Indusium membranaceous.

Var. inaequalis: Two times as large as the type.

Trop. Asia, China, Japan; — var. *inaequalis*: *Celebes; Hongkong*.

(17) **Pt. dispar**, *Kze*, *Copel.*, *Polypod. Philipp.*, 101; *Pt. semipinnata*, *L.*, *var.*, *C. Chr.*, *Ind. Fil.*

Like *Pt. semipinnata*, *L.*, but the pinnae more or less pinnate on the upper side also.

Philippines; China, Japan.

(18) **Pt. distans**, *J. Sm.*, *Hk. Bk.*, *Syn. Fil.*, 156; *Hk.*, *Spec. Fil.*, II, tab. CXXIV, B.

Stipes about 15 c.M. long, slender, erect, naked, pale. Fronds about 50 c.M. long, 10—15 c.M. broad, terminated by a pinna with a long, narrow, sharply serrate, linear apex, below which are 12—20 pairs of closely placed short lobes to $1\frac{1}{2}$ c.M. long and $\frac{1}{3}$ c.M. broad, the lower $\frac{2}{3}$ part of the fronds with 6—9 pairs of opposite, distantly placed pinnae with a long, linear apex, and at the base a few, short, oblong, close, spreading pinnulae. Texture coriaceous; rachis scabrous; veins conspicuous, simple or forked; barren lobes sharply and finely spinuloso-serrate. Indusium membranaceous.

Philippines.

* (19) **Pt. heteromorpha**, *Fée*, *Hk. Bk.*, *Syn. Fil.*, 156; *Hk.*, *Spec. Fil.*, II, tab. CXXVII, B; *Bedd.*, *Ferns Br.I.*, I, tab. LXXXIX; *Pt. cretica*, *L.*, *var. heteromorpha*, *Bedd.*, *Handb. Ind. Ferns*, 106.

Stipes 15—45 c.M. long, naked, erect, pale-coloured. Fronds 30—65 c.M. long, 15—25 c.M. broad, ovate-lanceolate, pinnate. Terminal pinna with the lower half sinuate and furnished with a few short, linear lobes. Lateral pinnae usually in 4—6 opposite pairs, erecto-patent, the higher ones linear-lanceolate, $7\frac{1}{2}$ —15 c.M. long, \pm 1 c.M. broad, simple or here and there with 1—2 lobes or ears near the base, lower pinnae 5—10 c.M. apart, 10—50 c.M. long, cut down nearly to the costa with \pm 4—6 oblong or linear-oblong, \pm spreading lobes on each side near the base below the often very long terminal segment, the lateral lobes varying from very short to \pm $2\frac{1}{2}$ c.M. long, $\frac{1}{2}$ —1 c.M. broad, the margin entire or sinuato-subcrenate, the apex provided with 1—5 minute spines. Texture subcoriaceous; rachis and both surfaces naked; the margin rather slightly crisped; veins conspicuous, usually once forked. Indusium narrow, membranaceous. — Very variable; the lowest pinnae sometimes with 1—12 lateral lobes on each side. The javan plant found by Mr. Buysman has the lobes of the terminal pinna in \pm 9 pairs, linear, to 4 c.M. long.

Java (Pasuruan), Celebes, Philippines; Burmah.

* (20) **Pt. Dalhousiae**, *Hk.*, Hk. Bk., Syn. Fil., 157; Hk., Spec. Fil., II, 170, tab. CXXI, A.

Rhizome short. Stipes \pm clustered, strong, erect, reddish-brown and naked like the rachis, polished, \pm 30—40 c.M. long. Fronds oblong, 50—60 c.M. long, the apex deeply pinnatifid with \pm 5 erecto-patent, distant, linear, acuminate segments below the long terminal one, \pm slightly joined by their suddenly dilated, decurrent bases; below this \pm 4—5 distant, erecto-patent, upcurved, \pm stalked pinnae on each side. Higher pinnae with a long, undivided, acuminate terminal segment, and with the lower side provided with 1—4 lateral segments similar to those of the apex of the frond, the upper side undivided, (\pm as in *Pt. semipinnata*, *L.*, but the segments distant), central and lower pinnae similar, but with 4—5 segments on the lower and a few similar ones on the upper side near the base (\pm as in *Pt. dispar*, *Kze*), lower pinnae 20—50 c.M. long, with the lowest segment on the lower side simple, or divided like the upper pinnae. Ultimate segments linear, upcurved, acuminate, the longest 10—15 c.M. long, 5—8 m.M. broad in both barren and fertile fronds, the edges finely crenate in the barren region. Texture firm; rachis polished; surfaces naked; veins distinct, simple or forked, the apices not thickened.

Java, Sumatra, Penang.

* (21) **Pt. Beccariana**, *C. Chr.*, Ind Fil., *Pt. concinna*, *Bk.* (not *Hew.*), *Becc.*, *Mal.*, III, 57; *Pt. longipinnula*, *Ces.*, *Fel. Becc. Polin.*, 5.

Stipes tufted, black-brown, naked, 15—50 c.M. long. Fronds deltoid, 2-pinnate, 15—25 c.M. long, naked; rachis black-brown, naked. Pinnae opposite, paucijugate, lanceolate, often conspicuously caudate, the lowest largest, forked at the base. Segments linear, contiguous, multijugate, entire, arcuate-ascending, adnate. Sori not reaching the apices of the pinnulae; indusium naked, broad, persistent, firm. — Near *Pt. Griffithiana*, *Hk.*, (*Spec. Fil.*, II, tab. CXXIII, A). — If this plant is the same as found at the same locality (Andai) by *Teysmann*, then the segments are very irregular, varying from 1—50 m.M. (mostly \pm 20 m.M.) long, joined at the base, the edges broadly crenate, the sori lateral on the fully developed segments, but terminal on the reduced ones.

New Guinea.

* (22a) **Pt. Grevilleana**, *Wall.*, *Trans. Linn. Soc., Bot.*, 2nd Series, I, 466, tab. LIV; *Pt. quadriaurita*, *Retz*, *var. digitata*, *Bk.*, *Journ. of Bot.*, XVII, 40; — *var. ornata*, *Hort. Bog.*

Stipes tufted, to 16 c.M. long, firm, erect, naked. Fronds pinnate with 1—2 pinnae on each side below the terminal one, the lowest pair forked at the base on the lower side in the fully developed plants. Pinnae and fork branches of the lowest pinnae lanceolate, acuminate, pectinato-pinnatifid, to 15 c.M. long, to 8 c.M. broad, the apex crenato-serrate. Lobes linear-oblong, falcate, often broader upwards, erecto-patent, to 4 c.M. long, to 1 c.M. broad, not rarely shorter and narrower in the fertile than in the barren fronds, crenate throughout or towards the rounded or bluntish apex only. Texture coriaceous; surfaces naked; veins very obscure, intermixed with numerous, fine, spurious veins. Sori not reaching the apex of the lobes.

Var. ornata: Fronds provided with broad, whitish or pale-grey-green bands running down the centre of the pinnae.

Malacca, Borneo, Philippines; Siam, North. India.

(22b) **Pt. pluricaudata**, *Copel.*, *Philipp. Journ.*, I, *Suppl.*, 156; *Pt. Blumeana*, *Ag.*, *var. pluricaudata*, *Christ*, *Philipp. Journ.*, II, 175.

Rhizome very short, crowned with lanceolate-setaceous, cinnamon-coloured scales. Stipes close, 10—25 c.M. long, slender, erect, naked, rubro-stramineous, sparingly scaly at the base. Fronds orbicular-ovate, \pm 10—15 c.M. long, 5—5-foliate. Pinnae cut down nearly or quite to the costa, the lowest ones

forked. Ultimate segments approximate, linear, $2\frac{1}{2}$ —5 m.M. broad, obtuse, serrulate, the terminal one narrowed, the fertile ones hardly contracted. Texture papyraceous; veins conspicuous, albo-setaceous above; the rest naked. Sori not reaching the apex of the segments. — Probably a local derivative of *Pt. quadriaurita*, *Retz* (Copel.).

Mindanao.

* (23) ***Pt. longipinnula***, *Wall.*, Hk. Bk., Syn. Fil., 158; Hk., Spec. Fil., II, tab. CXXXIV, A; Bedd., Ferns S.I., tab. XLIII; *Pt. quadriaurita*, *Retz*, var. *longipinnula*, Rac., MS. in Herb. Bog., *Pt. umbraculifera*, *Mett.*, Ann. Mus. Bot. L. B., IV, 97; — var. *sumatrana*.

Stipes erect, naked, 60—90 c.M. long, yellowish-green, rarely brown. Fronds 30—90 c.M. long, 50—45 c.M. broad, the terminal pinna to 30 c.M. long, 5—8 c.M. broad, regularly pectinato-pinnatifid with numerous nearly spreading or falcate, entire, crenate or serrate, blunt, linear, close or \pm remote lobes on each side, which are to 5 c.M. long and reach down nearly to the costa, the terminal lobe \pm caudiform. Lateral pinnae several on each side, similar to the terminal one, the lowest \pm 5 c.M. apart, whether or not forked at the base. Texture subcoriaceous; rachis and both surfaces naked; costae not rarely spinulose above; veins once forked. Sori reaching nearly to the apex of the segments. — Hardly different from *Pt. quadriaurita*, *Retz*, but the stalks of the pinnae \pm shortly winged by the decurrent bases of the lowest lobes, the lowest anterior lobe generally shorter than the opposite posterior one.

Var. *sumatrana*: Pinnae irregularly pinnatifid, the lobes different in length, varying from $\frac{1}{2}$ —5 c.M. deep, the edges crenato-serrate.

Malaya; Samoa, Japan, China, Burmah, North. India; — var. *sumatrana*: *Sumatra* (Hagen).

(24) ***Pt. torricelliana***, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., Suppl., 45.

Like *Pt. longipinnula*, *Wall.*, but the sterile pinnae ovate-oblong, narrowed at the base; pinnulae lanceolate-acuminate, very approximate, the veins 4-furcate.

New Guinea.

* (25a) ***Pt. quadriaurita***, *Retz*, Rac., Flor. Btz., I, 156; Hk., Spec. Fil., II, tab. CXXXIV, B; Bedd., Ferns S.I., tab. XXXI; *Pt. biaurita*, *L.*, var., C. Chr., Ind. Fil.; — var. *setigera*, Copel., Polypod. Philipp., 102; *Pt.*

asperula, J. Sm., Christ, Bull., Herb. Boiss., VI, 148; *Pt. Blumeana*, Ag., var. *asperula*, Christ, Philipp. Journ., II, 172; — var. *Blumeana*, Bedd., Handb. Ind. Ferns, 112; *Pt. Blumeana*, Ag., Rec., 22; — var. *argentea*, Bedd., l. c., 111; ? *Pt. argyrea*, Moore; — var. *normalis*; *Pt. normalis*, Don, Bl., Enum., 211; — var. *albido-maculata*; *Pt. normalis*, Don, var. *B.*, Bl., l. c., 211; — var. *nemoralis*; *Pt. nemoralis*, Willd., Bl., l. c., 210; Bau. & Hk., Gen. Fil., tab. LXIV; — var. *trachyphylla*; *Pt. trachyphylla*, Kze, Bot. Zeit., 1848, 196; — var. *Whitfordi*; *Pt. Whitfordi*, Copel., Philipp. Journ., I, Suppl., 255.

Stipes 15—60 c.M. long, erect, strong, naked, or scaly at the base, smooth, green, stramineous or brown. Fronds 15—100 c.M. long, 10—50 c.M. broad or broader, with a terminal pinna and 5—11 lateral ones on each side. Pinnae erecto-patent, pinnatifid $\frac{1}{3}$ -way down or more to the costa, 5—50 c.M. long, the lowest ones forked at the base on the lower side with 1—2 similar, but smaller, regularly pinnatifid segments. Lobes several to numerous, rather close, parallel, pectinate, linear or linear-oblong, rather erecto-patent or falcate, entire or serrate, blunt or acute, $\pm 1-2\frac{1}{2}$ c.M. long, $\pm 2\frac{1}{2}-7\frac{1}{2}$ m.M. broad, the lowest not or hardly decurrent at the base. Texture herbaceous to subcoriaceous; rachis and surfaces naked, smooth, rachis and costae often canaliculate above and purveyed there with distant, spine-like bristles; veins conspicuous, simple or often forked, the lowest posterior ones springing from the base of the costulae (main veins). Sori often continuous along the whole margin of the lobes. — A very variable species.

Var. setigera: Costulae spinulose like the costae.

Var. Blumeana: Very large; lobes 50—60 to a pinna, not falcate, nearly free at the base, equally wide throughout or broader at the apex, the apex rounded, barren, entire or crenulate.

Var. argentea: Fronds provided with broad, whitish bands running down the centre of the pinnae. Lobes not retuse at the apex.

Var. normalis: Lobes subtruncate, retuse at the apex; lowest posterior veins often springing from the costae at some distance from the costulae.

Var. albido-maculata: Like var. *normalis*, but the fronds variegated like var. *argentea*.

Var. nemoralis: Lowest posterior veins springing from the costae as in var. *normalis*, sometimes uniting with the opposite anterior lowest veins and then forming here and there a single costal areola.

Var. trachyphylla: Stipe, rachis and costae more or less scabrous.

Var. Whitfordi: Lobes conspicuously erecto-patent, nearly close.

All warm countries, the varieties scattered here and there.

(25b) **Pt. parviloba**, *Christ*, Bull. Sci. Fr. & Belg., XXVIII, 264, tab. XII.

Near *Pt. quadriaurita*, *Retz*, but small. Stipes slender, rufo-stramineous, scaberulous. Fronds broadly deltoid, the lowest pinnae produced, sharply deflexed. Ultimate segments close, linear-oblong, short, rather free. Texture rigidly papyraceous; colour pale-ochraceous-green; upper surface often rough with rigid setae; veins distinct beneath. Sori and indusia generally very slender. — This form is often found strongly reduced in and size resembling *Pt. Grevilleana*, *Wall.*

Luzon, Palawan.

(25c) **Pt. caesia**, *Copel.*, Philipp. Journ., I, Suppl., 156.

Rhizome short, erect. Stipes tufted, stramineous or brownish, slightly puberulous like the rachis, the base densely clothed with fuscous, filiform scales. Fronds 20—55 c.M. long, 10—18 c.M. broad. Pinnae \pm 1 c.M. broad, cut down nearly or quite to the costa, the terminal one the longest, the basal ones forked. Ultimate segments \pm 1½ m.M. broad, entire, obtuse, subauriculate, \pm overlapping at the base, the sinuses closed. Texture papyraceous; upper surface bluish, spinulose on the costae and veins, under one pale-green. Sori narrow. — Near *Pt. quadriaurita*, *Retz*, distinct enough to constitute a species. *Christ* points out that this species is near *Pt. furcans*, *Bk.* (*Copel.*)

Mindanao.

(26) **Pt. amoena**, *Bl.*, Enum., 210; *Hk. Bk.*, Syn. Fil., 479.

Stipes naked. Fronds 2-pinnatifid. Pinnae opposite, short-stalked, lanceolate, narrowed at both ends, pinnatifid, the lowest ones 2-partite. Segments subfalcate-oblong, somewhat acute, the lower ones obtuse, the edges appressate-serrate. Texture membranaceous; rachis and surfaces naked. — *Baker* considers this plant to be a form of *Pt. semipinnata*, *L.* See the information in *Hk.*, Spec. Fil., II, 205.

Java.

* (27) **Pt. furcans**, *Bk.*, Journ. of Bot., XXVI, 324.

Stipes naked, slender, blackish-purple or castaneous, glossy, 12—50 c.M. long. Fronds 15—20 c.M. long. Pinnae erecto-patent, lanceolate, simply pinnate, 2—2½ c.M. broad, lowest largest, \pm 15 c.M. long, forked at the base in the fully developed plants. Ultimate segments linear, entire, obtuse, adnate to the rachis, free or sometimes slightly joined, \pm 2—5 m.M. broad.

Texture moderately firm; surfaces naked; rachises dark-coloured; veins copious. Sori nearly reaching the apex of the segments; indusium narrow, naked.

Borneo.

* (28) **Pt. decussata**, *J. Sm.*, Journ. of Bot., III, 405; *Pt. patens*, *Hk.*, Hk. Bk., Syn. Fil., 165; *Hk.*, Spec. Fil., II, tab. CXXXVII; *Bedd.*, Ferns S.I., tab. CCV.

Stipes tufted, stout, scaly at the base, glossy, rather dark-brown. Fronds large, 90—120 c.M. long, 60 c.M. broad or broader. Terminal pinna 15—25 c.M. long, $3\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad, with several, \pm horizontally spreading, linear lobes on each side reaching down nearly or quite to the costa, dilated suddenly on both sides at the base, the barren ones slightly serrate or crenate. Lateral pinnae numerous, similar to the terminal on but longer, sometimes 45 c.M. long, to 5 c.M. broad, the lowest forked. Texture thinly coriaceous; rachis and both surfaces naked; veins oblique, the lowest sometimes anastomosing (*Christ*, *Farnkr. d. Erde*, 168), commonly once forked. Sori continued nearly to the end of the segments. — I am sceptical to the anastomosing of the veins. A plant gathered in Borneo by *Hallier* and determined by *Christ* as *Pt. patens*, *Hk.*, (*Ann. Btz.*, XX, 150) has the fronds smaller, the veins quite free and the costae purveyed with 2 interrupted, narrow wings on the upper side. (See *Bedd.*, *Handb. Ind. Ferns*, 114, line 3 from bottom).

Malacca, Borneo, Celebes, Philippines; Ceylon, Samoa, Society-Islands

(29) **Pt. Walkeri**, *Bk.*, Journ. of Bot., XXVI, 524.

Stipes naked, castaneous, 15—20 c.M. long. Fronds rhomboidal, irregularly 5-pinnate, \pm 30 c.M. long. Pinnae usually caudate, terminal and higher ones simply pinnate, several lower forked at the base. Ultimate segments obtuse, at utmost \pm 2 c.M. long, \pm 2 m.M. broad. Texture moderately firm; surfaces naked; rachises castaneous; veins copiously pinnate in the ultimate segments; veinlets indistinct. Sori occupying nearly the whole margin of the segments; indusium narrow, naked.

Borneo.

* (30) **Pt. excelsa**, *Gaud.*, *Rac.*, *Flor. Btz.*, I, 155; *Hk.*, *Spec. Fil.*, II, tab. CXXXVI; *Bedd.*, *Ferns Br.I.*, II, tab. CCXVIII.

Rhizome erect, short, densely clothed with narrow, lanceolate, brown scales. Stipes tufted, naked, erect, stout, reddish, greenish or pale-brown, to

± 170 c.M. long. Fronds 150—200 c.M. long. Terminal pinna 30 c.M. or more long, 7 c.M. or more broad, with numerous closely placed, linear, ± spreading, straight or falcate segments on each side, which are sometimes 5 c.M. long, 1 c.M. broad, dilated at the base and slightly serrate in the barren part. Lateral pinnae several, similar to the terminal one, the lowest distant, to 60 c.M. long and to 20 c.M. broad, whether or not forked at the base. Texture subcoriaceous; rachis channelled and sparingly bristly above; both surfaces naked; veins once forked. Sori not reaching the apex of the segments.

Malaya; North. India, Fiji, Hawaii.

§ 1. EUPTERIS. c. **Compositae.** **. **Partitae.**

- *. *Fronds 3-partite.* (31) *Pt. longipes.*
 ***. *Fronds 5-partite.* (32) *Pt. radicans.*

* (31) **Pt. longipes**, *Don*, *Rac., Flor. Btz., I, 154*; *Pt. pellucens*, *Agardh.*, *Bedd., Ferns S.I., tab. XXXII*; *Pt. Zollingeri*, *Mett., Ann. Mus. Bot. L. B., IV, 97.*

Rhizome erect, densely clothed with lanceolate scales. Stipes 30—120 c.M. long, erect, naked, stramineous or pale-brown. Central branch with a terminal pinnula about 7—15 c.M. long, 1—2½ c.M. broad, cut down nearly to the costa into numerous erecto-patent, linear-oblong lobes on each side about ½—1½ c.M. long, 2—5 m.M. broad, bluntly toothed towards the point when barren; below this numerous lateral, pinnate or pinnatifid pinnulae on each side. Lateral branches similar to the central one. Texture herbaceous; rachises and both surfaces naked or the secondary rachises slightly scaly above; veins not prominent, forked. Sori not reaching the apex of the segments.

Trop. Asia.

(32) **Pt. radicans**, *Christ*, *Ann. Btz. XV, 110.*

Stipes about 100 c.M. long, reddish-yellow. Fronds ± 150 c.M. long, deltoid, with a central branch and 2 lateral branches on each side, which are half as long as the central one. Branches pinnate; pinnulae distant, ± 15 on each side of the central branch and ± 6 on each side of the lateral

branches, \pm 5 c.M. long and 2 c.M. broad, cut down nearly or quite to the costa into ligulate, blunt, slightly crenate segments. Texture thinly herbaceous; both surfaces naked; veins inconspicuous, forked. Sori not reaching the base and the apex of the segments; indusium narrow. — Branches proliferous at the apex.

Celebes.

Species of an indefinite systematical place ().*

(35) **Pt. ligulata**, *Gaud.*, Freyc., Voy. Ur. Phys., Bot., 385; *Pt. quadriaurita*, *Retz*, var. *Lauterbachii*, Christ, in Schum. & Laut., Flor. deut. Sch.geb., 154, p. p.

Stipes tufted, black, glossy, \pm 50—45 c.M. long. Fronds 3-partite. Branches whether or not branched, deeply pinnatifid or pinnate, the apex long-acuminate, the base auricled. Lobes unequal, linear, entire, sinuate, toothed or lobed, the lower sometimes distinct, free, stalked, auricled. Sori irregular, lateral on the fully developed lobes or terminal on the reduced ones. — The scales (of the rhizome?) capillary, reddish, thin, diaphanous.

Borneo, Moluccas, New Guinea.

(34) **Pt. mixta**, *Christ*, Schum. & Laut., Flor. deut. Sch.geb., Suppl., 45; *Pt. quadriaurita*, *Retz*, var. *Lauterbachii*, Christ, p. p.

Rachis blackish-castaneous. Fronds 2-pinnatifid, nearly 3-partite. Pinnae deeply pinnatifid, the apex caudato-elongate, subentire or irregularly incised; lobes pectinate, linear-acuminate. Sori thick, ochraceous; indusium nearly wanting. — Fertile fronds slightly suggesting *Pt. ligulata*, *Gaud.* but smaller.

New Guinea.

(35) **Pt. philippinensis**, *Fée*, Mém., VIII, 114; X, 17, tab. XXXIII, fig. 2.

Fronds ovate-lanceolate, 2-pinnate at the base. Pinnae stalked, lanceolate, the apex caudiform. Segments oblong, toothed. Sori reaching neither the base nor the apex of the segments.

Philippines.

(*) Having not seen the 3 following species and being not able to judge their systematical place from the descriptions alone, I placed them here.

§ 2. CAMPTERIA.

- a.* Fronds pinnately divided, not 3-partite; lowest pinnae not much longer than the next higher ones (*).
- b.* Fronds small or middle-sized, under 100 c.M. long. (36) *Pt. biaurita*.
- b.b.* Fronds large, more than 100 c.M. long. (37) *Pt. geminata*.
- a.a.* Fronds 3-partite; branches similarly 2-pinnatifid, the lateral ones sometimes nearly as long as the central one, whether or not forked at the base on the lower side, and if so the secondary branches not rarely forked again.
- *. Texture herbaceous; barren ultimate segments flat at the edge. (38) *Pt. Wallichiana*.
- *.*. Texture coriaceous; barren ultimate segments revolute at the edge. (59) *Pt. heterophlebia*.

* (36) ***Pt. biaurita*, L.**, *Rac.*, *Flor Btz.*, I, 157; *Pt. quadriaurita*, *Retz.*, *var. biaurita*, *Christ*, *Ann. Btz.*, XV, 110; *Campteria biaurita*, *Hk.*, *Bedd.*, *Ferns S.I.*, tab. XLIV; *Bau. & Hk.*, *Gen. Fil.*, tab. LXV, A.

Rhizome erect, short, densely clothed with short, deltoid-lanceolate scales. Stipes 50—70 c.M. long, strong, erect, naked, or scaly at the base, the scales very fugacious. Fronds to 90 c.M. long, with a terminal pinna 7—30 c.M. long, 2—5 c.M. broad, cut down nearly to the costa into numerous spreading or erecto-patent, linear, straight or falcate, entire lobes on each side, which are to $\pm \frac{1}{2}$ c.M. broad. Lateral pinnae few or several on each side, similar to the terminal one, the lower ones often forked at the base. Texture herbaceous to subcoriaceous; rachis and surfaces naked, under surface sometimes glaucous; veins usually forked. Sori commonly continued to the apex. — Differs from *Pt. quadriaurita*, *Retz* only in the costulae being united at the base by an arching vein.

Pantropical.

(37) ***Pt. geminata*, Wall.**, *List*, No. 2180; *Pt. maxima*, *Bk.*, *Hk.*, *Bk.*, *Syn. Fil.*, 480; *Pt. Kleiniana*, *Christ* (not *Pr.*), *Copel.*, *Polypod. Philipp.*, 103; *Campteria Kleiniana*, *Pr.*, *Bedd.*, *Handb. Ind. Ferns*, 116; *Pr.*, *Tent. Pterid.*, tab. V, fig. 19; *C. anamallayensis*, *Bedd.*, *Ferns S.I.*, tab. XLV.

(*) See also ***Pt. quadriaurita*, Retz**, *var. nemoralis* & ***Pt. decussata*, J. Sm.**

Large, 120—180 c.M. high. Fronds deltoid-ovate, pinnate. Pinnae subsessile, lanceolate, long-acuminate, deeply pinnatifid, the lowest to \pm 50 c.M. long, with 1—2 pinnatifid branches on the lower side. Segments erecto-patent, falcate-oblong, sterile ones broadly crenate, 5—4 c.M. long, 7—10 m.M. broad, fertile ones crenate at their barren apices. Texture subcoriaceous; rachis and surfaces naked; veins anastomosing in costal arches. Sori never reaching the base or apex of the segments. — Scarcely more than a variety of *Pt. biaurita*, *L.*
? *Java, Philippines; Japan, China, South. India.*

* (38) **Pt. Wallichiana**, *Agardh.*, *Rac.*, *Flor. Btz.*, I, 157; *Campteria Wallichiana*, *Moore*, *Ferns Br.I.*, I, tab. XXV; II, tab. CCXVII.

Rhizome erect, clothed with roundish, entire or dentate, soft scales. Stipes 50—180 c.M. long, strong, naked, glossy, green or brown. Fronds 5-partite. Central branch mostly 25—60 c.M. long, 15—50 c.M. broad, with numerous linear-lanceolate, sessile, opposite pinnulae on each side below the terminal one. Pinnulae 7—15 c.M. long, to 5 c.M. broad, cut down nearly to the costa into numerous flattened, contiguous, linear-oblong, rounded, finely toothed or crenate lobes. Lateral branches similar to the central one and nearly as long, forked near the base, the lower fork branch sometimes forked again. Texture herbaceous; rachises and both surfaces naked; veins not conspicuous, commonly once forked. Sori continuous along nearly the whole length of the edge, or occupying the lower half of the lobes only.

Java, Celebes, Philippines; Samoa, North. India.

* (39) **Pt. heterophlebia**, *Kze.*, *Bot. Zeit.*, 1848, 196; *Campteria heterophlebia*, *Moore*, *Ind.*, XLIII.

Whole plant to 1 $\frac{1}{4}$ M. high. Stipes yellowish. Fronds 5-partite; branches to 60 c.M. long, to 25 c.M. broad. Pinnulae approximate, deeply pinnatifid, 10—15 c.M. long, to 4 c.M. broad, short-stalked, spreading, elongate-lanceolate, subacuminate. Lobes falcate, oblong, with a broad base, the apex bluntish, the edge crenato-serrulate, revolute. Texture coriaceous; surfaces naked; rachises, costae and costulae subterete, the upper side flat, reddish-yellow-brown, pubescent; main veins united at the base by arching veins, the other veins free, forked, some of them forming here and there a single costular areola in the lobes.

Java.

§ 3. LITOBROCHIA.

1. *Fronds simply pinnate; pinnae entire.* (40a) Pt. Warburgii.
 (40b) Pt. finisterrae.
- 1.1. *Fronds pinnate; pinnae pinnatifid, the lowest ones not much longer than the next higher ones, whether or not forked at the base.*
2. *Indusium inconspicuous.* (41) Pt. microdictyon.
- 2.2. *Indusium distinct.*
3. *Fronds to \pm 60 c.M. long; lowest pinnae not forked at the base.* (42) Pt. Junghuhnii.
- 3.3. *Fully developed fronds more than 60 c.M. long; lowest pinnae forked at the base.* (43) Pt. Milneana.
- 1.1.1. *Fronds 3-partite; branches similarly 2-pinnatifid, the lateral ones sometimes nearly as long as the central one, whether or not forked at the base on the lower side, and if so the secondary branches not rarely forked again (*).* (44) Pt. tripartita.

(40a) **Pt. Warburgii**, *Christ*, Warb., Mons., I, 70.

Stipes 45 c.M. or more long, canaliculate in the higher part, naked, stramineous. Fronds large, ovate, probably \pm 100 c.M. long or longer, \pm 40 c.M. broad, simply pinnate. Pinnae distant, \pm 4 c.M. apart, ligulate-linear, as long, entire, long-acuminate, adnate at the base. The lower ones free, not reduced, the higher decurrent so as to form a broad wing to the rachis, the longest \pm 24 c.M. long, \pm 2 c.M. broad. Texture herbaceous or membranaceous; surfaces naked; costae prominent; no main veins; areolae numerous, hexagonal-oblong, in 5—4 rows; no free veinlets. Sori reaching nearly the apex. — Sterile fronds smaller, with the apex pinnatifid, the pinnae to \pm 3½ c.M. broad.

Moluccas, New Guinea.

(40b) **Pt. finisterrae**, *Rst.*, Fed., Rep., V, 57.

Rhizome and stipes unknown. Fronds elongate-ovate, to 60 c.M. long or longer, to 50 c.M. broad, pinnate at the base, pinnatifid towards the apex. Lowest pinnae 5—10 c.M. apart, sometimes more approximate, short-stalked, forked at the base on the lower side, the 1—3 following pinnae decurrent,

(*) See also **Pt. heterophlebia**, *Kze.*

with the upper base free and truncate or joined with the next higher pinnae, the higher pinnae joined at the base by a wing $\frac{3}{4}$ — $1\frac{1}{2}$ c.M. broad. Lower and central pinnae to 25 c.M. long, to 5 c.M. broad, the fertile ones narrower than the barren, entire, linear, or slightly dilated towards the acuminate apex, the higher ones rather suddenly abbreviated, $\pm \frac{1}{3} \times$ shorter, the terminal one nearly as long as or longer than the highest ones. Texture membranaceous, pellucid; colour grey-green; surfaces naked; rachis and costae dark-purple beneath, yellowish above; costulae and veins distinct; areolae roundish- or elongate-subhexagonal. Sori continuous along the sides, interrupted at the base and apex of the segments; indusium dark-purple, rigidly membranaceous.

New Guinea.

(41) ? **Pt. microdictyon**, *Hk.* (not *Fée*), *Spec. Fil.*, II, 228; *Litobrochia microdictyon*, *Fée*, *Gen.*, 138.

Stipes reddish. Fronds tall, ovate; rachis subtrigonus, channelled. Pinnae short-stalked, alternate, acuminate, pinnatifid. Segments lanceolate, falcate, slightly dilated at the base, obtuse, crenate. Indusium narrow, inconspicuous, receptaculiform; the margin thick and black.

Philippines.

* (42) **Pt. Junghuhnii**, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 170; *Pt. tripartita*, *Sw.*, var. *Junghuhnii*, *Hk.*, *Spec. Fil.*, II, 226; *Campteria Junghuhnii*, *Reunw.*, MS.

Stipes firm, erect, polished, nearly stramineous. Fronds to 60 c.M. long, with several or numerous pinnae on each side, which are cut down $\frac{2}{3}$ — $\frac{3}{4}$ of the way to the costa into linear-oblong, falcate, entire lobes $\pm \frac{1}{2}$ c.M. broad at the base; lowest pinnae simple (i. e. not forked at the base), to 25 c.M. long. Texture herbaceous to subcoriaceous; rachis and both surfaces naked; veins fine, anastomosing principally in costal arches (as in *Campteria*) and in the space between the sinuses and the costae, sometimes also in the lobes. Sori reaching nearly to the apex of the lobes. — Habit of *Pt. quadriaurita*, *Retz* & *biaurita*, *L.*

Java, Banca.

(45) **Pt. Milneana**, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 170; *Pt. tripartita*, *Sw.*, var. *Milneana*, *Hk.*, *Spec. Fil.*, II, 226, tab. CXXXVIII, B.

Stipes strong, erect, polished, yellowish-brown. Fronds 60—90 c.M. long, with numerous pinnae on each side cut down throughout nearly to the costa

into linear-oblong, falcate lobes 1—2 c.M. long, \pm 5 m.M. broad, lowest pinnae 15—25 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, with a single, similar, but smaller pinnula at the base on the lower side. Texture herbaceous; rachis and surfaces naked; veins anastomosing copiously. Sori not reaching the subentire apices. — The form found in Batjan has the veins only slightly anastomosing.

Batjan; Polynesia, Trop. Australia.

* (44) **Pt. tripartia**, Sw., Hk., Spec. Fil., II, 225 (excl. β & γ); *Pt. marginata*, Bory, Rac., Flor. Btz., I, 158; *Litobrochia marginata*, Pr., Bedd., Handb. Ind. Ferns, 122; *L. tripartita*, Pr., Bedd., Ferns S.I., tab. CCXX.

Stipes 50—150 c.M. long, strong, erect, naked, polished, stramineous or brownish. Fronds deltoid, \pm 50—150 c.M. each way. Central branch with a terminal pinnula 7—25 c.M. long, 8—25 m.M. broad, cut down nearly to the costa into numerous closely placed, linear, straight or falcate, spreading or erecto-patent lobes on each side, which are entire or slightly toothed when barren; below this many closely placed, similar lateral pinnulae on each side. Lateral branches similar to the central one, but a little smaller, unbranched or branched again at the base on the lower side. Texture commonly herbaceous; rachises and surfaces naked; veins fine, anastomosing principally in arches against the midribs and costulae. Sori falling rather short of the apex of the segments.

Paleotropical.

Species of an indefinite systematical place.

(45) **Pt. mertensioides**, Willd., Spec., V, 594.

Fronds 60 c.M. or more long, 2-pinnate. Lowest pinnae 15—17 $\frac{1}{2}$ c.M. long, highest 7 $\frac{1}{2}$ c.M. long. Pinnulae linear-lanceolate, sessile, subacute, entire, subconfluent, 1—1 $\frac{1}{2}$ c.M. long. Texture thinly membranaceous. Sori narrow. — Pinnae resembling the leafy branches of *Mertensia*.

Java, Borneo, Amboyna.

45b. HEMIPTERIS, Rosenstock.

Like *Pteris*, but the sori occupying only the posterior edge of the ultimate segments, the anterior side destitute of sori. Fed., Rep., V, 38.

Papuan.

H. Wernerii, Rst., Fed., Rep., V, 38.

Rhizome and stipes unknown. Fronds (ex sched.) pedate, with the primary segments (of the primary branches?) elongate-ovate, 50 c.M. long or longer, \pm 30 c.M. broad, pinnate like the terminal segments. Secondary segments multijugate, alternate, subcontiguous, linear-lanceolate, erecto-patent, deeply pinnatifid, narrowed towards the serrate apex, the point entire, the largest ones \pm 25 c.M. long, \pm 3 c.M. broad, the higher ones suberect and smaller than the lower ones. Lobes multijugate, erecto-patent, not very distant, joined at the base by a wing \pm 1 m.M. broad, linear, $1\frac{1}{2}$ c.M. long, $\frac{1}{2}$ c.M. broad, the apex obtuse, the margin serrate-dentate, the teeth slightly crenate. Texture membranaceo-herbaceous; surfaces naked; costae and costulae sparingly clothed with minute, linear or irregularly dilacerate scales, flattened above and provided there with whitish-membranaceous, subinterrupted, unequal wings (the internodes narrow at the base and broad at the apex); veins anastomosing, forming principally costal and costular areolae, the branches free or slightly anastomosing. Sori 4—5 on each side of the lobes, occupying the posterior side of the ultimate lacinulae (i. e. the teeth) from the base to the apex; indusium grey-brown.

New Guinea.

46. PAESIA, St. Hilaire.

Sori linear, submarginal, uniting the apices of several veins. Indusium linear, consisting of the reflexed, modified margin of the frond, often 2-valved, the inner valve attached at the inner side of the sorus. Spores bilateral.

Rhizome creeping. Stipes distant, not articulate to the rhizome, purveyed with a single, \pm semicylindrical vascular strand. Fronds compound; veins free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 156, B—E.

America, Polynesia, Malaya.

* (1) **P. radula, C. Chr.,** Ind. Fil.; *Pteris Radula, Bk.,* Journ. of Bot., XVIII, 211.

Rhizome wide-creeping. Stipes distant, brown, naked, 15—50 c.M. long, rough with minute, raised points like the flexuose rachis. Fronds lanceolate, 5-pinnatifid or 5-pinnate, 20—40 c.M. long, 7—10 c.M. broad. Lower pinnac erecto-patent, often cut away on the lower side at the base, lanceolate-deltoid.

Pinnulae unequal-sided, \pm deltoid, \pm 2 c.M. long, \pm 1 c.M. broad, cut down nearly to the costa into small, oblong-cuneate, entire or bluntly toothed segments. Upper pinnae rather lanceolate, with nearly simple secondary divisions. Texture firm; surfaces naked; veins free, immersed, obscure. Sori reaching from the base nearly to the apex of the segments; indusium narrow.

Sumatra.

(2) **P. rugulosa**, *Kuhn*, Copel., Polypod. Philipp., 105; *Pteris rugulosa*, *Lab.*, Hk. Bk., Syn. Fil., 480; ? *Pt. divaricatissima*, *Bk.*, Hk. Bk., l. c., 163.

Rhizome wide-creeping. Stipes and rachises slender, glutinoso-pubescent, canstaneous. Fronds ample, 4-pinnatifid, rhomboidal. Pinnae close, lanceolate, short-stalked, erecto-patent, to 50 c.M. or more long. Pinnulae lanceolate, close, patent, nearly sessile, 1—2 c.M. broad. Segments lanceolate, cuneate-truncate on lower side at base, lower deeply pinnatifid with oblique, blunt lobes. Texture cartilagineo-coriaceous; rachis zigzag; surfaces glutinose and minutely scaly, especially below; edges of fertile segments revolute, the sori reaching to the midribs.

Philippines; New Caledonia, Tahiti.

47. PTERIDIUM, *Gleditsch.*

Sori linear, on a submarginal strand uniting the vein tips. Indusium linear, consisting of the reflexed, modified margin, often 2-valved with the inner valve attached at the inner side of the sorus.

Rhizome creeping. Stipes distant, seriate, not articulate to the rhizome, purveyed with many (\pm 12—24), irregularly placed vascular strands. Fronds at least 2-pinnate; veins free. Hk. Bk., Syn. Fil., tab. III, fig. 31, d—f (as *Pteris*, § *Paesia*); Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 156, A.

Cosmopolitan.

* **Pt. aquilinum**, *Kuhn*, Copel., Polypod. Philipp., 104; *Pteris aquilina*, *L.*, Rac., Flor. Btz., I, 156; Bedd. Ferns S.I., tab. XLII; — var. *lanuginosum*, Hk., Spec. Fil., II, 196; *Pteris lanuginosa*, *Spr.*, Nova Acta, X, 251, tab. XVI, fig. 1—2; — var. *esculentum*, Hk., l. c., 197; *Pteris esculenta*, *Forst.*, Pl. escul., 74.

Rhizome stout, wide-creeping, scaly. Stipes distant, 50—100 c.M. long, strong, erect, naked, stramineous or brownish. Fronds deltoid to broad-lanceolate, 60—200 c.M. long or longer, 30—60 c.M. or more broad. Uppermost pinnae simple, the next pinnatifid, the following pinnate, the lowest largest, 2—5-pinnatifid. Ultimate segments linear-oblong, 1—2½ c.M. long, 3—5 m.M. broad, numerous, close. Texture coriaceous; rachis and surfaces naked; veins close, conspicuous, often 1—2 × forked.

Var. lanuginosum: Fronds pubescent or silky-tomentose beneath; pinnulae more regularly pinnatifid.

Var. esculentum: Rhizome fibrillose or hairy; ultimate segments more distant, connected by their suddenly dilated, decurrent bases.

Cosmopolitan, chiefly northern; — var. lanuginosum: chiefly tropical; — var. esculentum: Borneo, New Guinea; Australia.

48. LOMARIA, *Willdenow.*

Sori linear, continuous, on a broad receptacle, parallel to the margin, at length filling up the whole space between the midrib and the margin. Indusium linear, consisting of the modified, reflexed margin of the frond (*).

Rhizome or caudex erect. Stipes not articulate, purveyed with 2 stout and 1 to several finer vascular strands placed altogether in a circle, the stout ones abreast at the inner (upper) side. Fronds simple to 2-pinnatifid, dimorphous, the fertile ones with the pinnae narrowly linear; veins free, but united by the receptacle in the fertile segments. Hk. Bk., Syn. Fil., tab. III, fig. 53; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 151 (as *Blechnum*, § *Lomaria*).

Tropical and extra-tropical.

- a.* Barren fronds simple, pinnatifid or pinnate; lobes or pinnae suddenly dilated and decurrent at the base, adnate to the rachis. (1) *L. Patersoni.*
- a.a.* Barren fronds pinnate; pinnae slightly dilated, but not decurrent at the base, adnate to the rachis. (2) *L. vulcanica.*
- a.a.a.* Barren fronds pinnate; lower pinnae truncate or rounded at the base, not adnate to the rachis.

(*) See also under **Stenochlaena.**

b. Barren fronds small, 10—15 c.M. long.

*. Stipes tufted. Pinnae crenate, auricled at the base on both sides.

(5) *L. egenolfioides*.

.. Stipes scattered. Pinnae finely serrate, not auricled at the base.

(4) *L. areolaris*.

b.b. Barren fronds larger, 30—200 c.M. long.

c. Stipes naked, or scaly at the base.

d. Pinnae 5—7½ c.M. long, 5—7½ m.M. broad.

(5) *L. punctulata*.

d.d. Pinnae 7½—30 c.M. long, 10—25 m.M. broad. (6) *L. capensis*.

c.c. Stipes densely scaly.

(7) *L. vestita*.

a.a.a.a. Barren fronds 2-pinnatifid.

(8) *L. Fraseri*.

* (1) **L. Patersoni**, *Spr.*, Hk. Bk., Syn. Fil., 174; Hk., Fil. Exot., tab. II; *L. elongata*, *Bl.*, var. *Cumingiana*, *Hk.*, Spec. Fil., III, tab. CXLIII; *Blechnum Patersoni*, *Mett.*, Christ, Farnkr. d. Erde, 176, p. p.; C. Chr., Ind. Fil.; *Stegania Patersoni*, *R. Br.*, (oldest name); — var. *elongata*; *L. elongata*, *Bl.*, Rac., Flor. Btz., I, 160.; *B. Patersoni*, *Mett.*, var. *majus*, Copel., Philipp. Journ., I, Suppl., 152.

Rhizome short-creeping or erect, clothed with ovate-lanceolate, brown scales. Stipes 5—8 c.M. long, erect, rather scaly below. Barren fronds simple, ± 30 c.M. long, under 2½ c.M. broad, broadest ± ⅓ of the distance from the top, narrowed gradually downwards, the point acuminate, the margin cartilagineous, wavy. Texture coriaceous: veins inconspicuous. Fertile fronds linear, ± 5 m.M. broad, simple, or branched with 1 or more segments on 1 or both sides. Sori occupying the whole space between the midrib and the edge.

Var. elongata: Sterile and fertile fronds deeply pinnatifid, 60 c.M. or more long, the barren ones cut down into ± 6—10 segments on each side, which are ± 15—25 c.M. long, ± 2½ c.M. broad, suddenly dilated and decurrent, and free or confluent at the base; fertile pinnae often numerous, erecto-patent, as long as the barren ones, but narrowly linear, mostly connected at the base by a very narrow wing.

Malaya; South. India, Fiji, New Zealand, Australia, Tasmania.

* (2) **L. vulcanica**, *Bl.*, Rac., Flor. Btz., I, 161; Hk., Icon. Pl., X, tab. CMLXIX; *Blechnum vulcanicum*, *Kuhn*, Farnkr. d. Erde, 179; C. Chr., Ind. Fil.

Rhizome short, erect, clothed with long, linear-lanceolate or subulate

scales. Stipes 10—30 c.M. long, pale, erect, paleaceous below. Fronds 15—45 c.M. long, 7—15 c.M. broad, the barren ones ovate-lanceolate, narrowed at the base or not. Pinnae numerous, close, lanceolate, subfalcate, the point acute or bluntish, 5—10 c.M. long, $\frac{1}{2}$ —2 c.M. broad, the base not decurrent, slightly dilated, lowest pinnae deflexed. Texture coriaceous; veins fine, prominent beneath; surfaces and edges often hairy. Fertile pinnae linear, very narrow, distant, suddenly dilated at the base, $3\frac{1}{2}$ —10 c.M. long.

Java, Celebes & Luzon to Polynesia, Queensland, Tasmania, New-Zeeland.

(5) **L. egenolfoides**, *Bk.*, Kew Bull., 1894, 7; *Blechnum egenolfoides*, *C. Chr.*, Ind. Fil.

Rhizome erect. Stipes densely tufted, $5\frac{1}{2}$ —15 c.M. long, the fertile fronds on the longest stipes. Fronds simply pinnate, the barren ones 12—15 c.M. long, 5—4 c.M. broad at the middle, the stipes slender, brown. Pinnae linear, obtuse, crenate, the base auricled on both sides, lower pinnae deflexed, gradually reduced. Texture subcoriaceous; surfaces naked; veins distant, erecto-patent, immersed, obscure, simple, the lower forked. Fertile fronds $2-2\frac{1}{2}$ c.M. broad, with the pinnae narrower, more distant.

Borneo.

(4) **L. areolaris**, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 28; *Blechnum areolare*, *Copel.*, Polypod. Philipp., 90; *C. Chr.*, Ind. Fil.; *Stenochlaena areolaris*, *Copel.*, Philipp. Journ., II, 406.

Rhizome long, slender, naked, twisted. Stipes of barren fronds \pm 15 c.M. long. Barren fronds ovate, 10—15 c.M. long, 7—10 c.M. broad, pinnate. Pinnae 7—9, sessile, linear-lanceolate, 1—2 c.M. broad, acute, finely serrate, the base truncate or rounded. Texture coriaceous; veins forming a series of costal arches with free, simple or forked veins beyond; surfaces naked; costae and veins slightly chaffy below. Fertile fronds on stipes \pm 30 c.M. long, with the pinnae narrowly linear, $5\frac{1}{2}$ —5 c.M. long, rather distant. Indusium narrow.

Philippines.

(5) **L. punctulata**, *Hae*, Hk. Bk., Syn. Fil., 179; *Blechnum punctulatum*, *Sw.*, Christ, Farnkr. d. Erde, 178; *C. Chr.*, Ind. Fil.

Rhizome stout, densely paleaceous at the crown. Stipes 7—15 c.M. long, strong, erect. Barren fronds oblong-lanceolate, 50—60 c.M. long, 10—15 c.M. broad. Pinnae very numerous, contiguous, linear, subfalcate, $5-7\frac{1}{2}$

c.M. long, $\frac{1}{2}$ — $\frac{2}{3}$ c.M. broad, rounded or cordate and auricled at the base, the lower ones deflexed and the lowest reduced to auricles. Texture coriaceous; veins inconspicuous; rachis stout, erect, naked, stramineous. Fertile fronds similar, but the pinnae narrowed, often not more than $2\frac{1}{2}$ m.M. broad. Indusium not rarely \pm intramarginal.

Java; South. Africa.

* (6) **L. capensis**, *Willd.*, Spec., V, 291; *L. procera*, *Spr.*, Hk. Bk., Syn. Fil., 179, p. p.; Hk., Icon. Pl., II, tab. CXXVII—CXXVIII; Gard. Ferns, tab. LIII; *Blechnum capense*, *Schl.*, Christ, Farnkr. d. Erde, 178; C. Chr., Ind. Fil.; *Osmunda capensis*, *L.*, (oldest name).

Rhizome stout, erect, woody, clothed with large, ovate or lanceolate, acuminate, ferruginous scales. Stipes 15—50 c.M. long, stout, erect, scaly below. Barren fronds 50—120 c.M. long, 15—60 c.M. broad. Pinnae numerous, linear, $7\frac{1}{2}$ —30 c.M. long, 1— $2\frac{1}{2}$ c.M. broad, narrowed gradually from the base to the apex, the base unequally cordate, sometimes auricled, the margin serrate, the apex acuminate, lower pinnae sessile or slightly stalked, higher adnate to the rachis, the terminal one free, often very long. Texture coriaceous; rachis stout, slightly scaly, the scales not large; costae \pm chaffy beneath; veins fine and parallel. Fertile fronds with the pinnae narrowly linear, distant, 10—15 c.M. long, to $\frac{1}{2}$ — $\frac{2}{3}$ c.M. broad. Indusium broad, membranaceous, occasionally ciliate, sometimes intramarginal.

Malaya; Polynesia, Trop. America, South. Africa.

* (7) **L. vestita**, *Bl.*, Rac., Flor. Btz., I, 161; *L. procera*, *Spr.*, Hk. Bk., Syn. Fil., 179, p. p.; *Blechnum vestitum*, *Kuhn*, Ann. Mus. Bot. L. B., IV, 284; C. Chr., Ind. Fil.

Rhizome erect, forming a short, thick caudex, densely clothed with broad, brown, acuminate scales. Stipes \pm 25 c.M. long, thick, densely clothed throughout with large, acuminate, brown or pale-brown scales. Fronds dimorphous. Barren fronds pinnate, 50—200 c.M. long. Pinnae numerous, linear, 12—24 c.M. long, 1— $1\frac{1}{2}$ c.M. broad at the base, acuminate, the edge serrate, the base truncate or rounded. Texture coriaceous; rachis scaly like the stipe; costae \pm chaffy beneath; veins numerous, fine, parallel. Fertile fronds with the pinnae of the same length, erecto-patent, 4—5 m.M. broad.

Malaya.

(8) **L. Fraseri**, *Cunn.*, Hk. Bk., Syn. Fil., 182; Icon. Pl., II, tab.

CLXXXV; *Blechnum Fraseri*, Luerss., Christ, Farnkr. d. Erde, 180; C. Chr., Ind. Fil.; — var. *philippinesis*, Christ, Bull. Herb. Boiss., VI, 149, tab. II.

Caudex elongate, stout, suberect, densely clothed with linear, dark-brown scales at the crown. Stipes 10—15 c.M. long, strong, erect, scaly. Fronds ovate-acuminate, 2-pinnatifid, 30—45 c.M. long, 10—15 c.M. broad. Pinnæ linear-lanceolate, the lower ones 5—7 c.M. long, 1—2 c.M. broad, cut down very nearly to the costa into linear-oblong, mucronate, slightly toothed segments, their bases decurrent into a pinnatifid wing to the main rachis, the lobes triangular. Texture herbaceous; veins fine, simple or forked. Fertile fronds similar in size and cutting.

Var. philippinensis: Caudex subarborescent, to 1½ M. high, the scales lanceolate-ovate to subulate. Stipes 5—4 c.M. long. Fronds 20—25 c.M. broad, long-acuminate, narrowed at the base. Indusium narrow, evanishing.

New Zealand; — var. *philippinensis:* *Philippines.*

TRIBE V. BLECHNEAE.

49. BLECHNUM, *Linné.*

Sori long, linear, 1 on each side, \pm close to the costa, continuous or sometimes interrupted. Indusium of the same shape, attached at the outer side of the sorus, opening towards the costa.

Caudex mostly erect, sometimes subarborescent. Stipes tufted, not articulate to the caudex. Fronds simple or commonly divided; veins free, but those of the fertile fronds united near the base by a costal strand occupied by the sori. Hk. Bk., Syn. Fil., tab. IV, fig. 54; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 150.

Tropical and subtropical.

Arrangement of the sections.

- § 1. **EUBLECHNUM.** Fronds erect.
§ 2. **SALPICHLAENA.** Fronds twining. (Not in Malaya).

§ 1. **EUBLECHNUM.**

- 1.** *Fronds simple.* (1) *B. Treubii.*
1.1. *Fronds pinnatifid.*
* *Stipes naked.* (2) *B. nitidum.*
** *Stipes scaly and muricate in the lower part.* (3) *B. cartilagineum.*
1.1.1. *Fronds pinnate.*
2. *Pinnae entire.*
3. *Pinnae narrowed at the base.* (4) *B. Finlaysonianum.*
3.3. *Pinnae rounded, truncate or cordate at the base.*
4. *Pinnae auricled at the base.* (5) *B. Blumei.*
4.4. *Pinnae not auricled.* (6) *B. orientale.*
2.2. *Pinnae serrate or finely incised.*
* *Pinnae narrow at the base.* (7) *B. serrulatum.*
** *Pinnae broad at the base.* (8) *B. egregium.*

* (1) **B. Treubii**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 15.

Small. Rhizome short, erect or nearly so, slightly scaly, copiously stoloniferous so as to form a dense mass of low plants. Stipes tufted, clothed with few, scattered scales, but more copiously towards the base, slender, erect, 2—5 c.M. long. Fronds simple, lanceolate, 4—9 c.M. long, 8—12 m.M. broad, narrowed gradually towards both ends, the apex acuminate, but the point bluntish, the margin entire or slightly irregular, the very base not rarely purveyed with 1—2 minute auricles. Texture subcoriaceous; surfaces naked; veins not very close, some of them sometimes anastomosing, mostly forked, sometimes simple or 3-furcate, the apex thickened, not reaching the margin. Sori long, continuous or interrupted, close to the costa, not reaching the base and apex of the fronds. — Very near the Brazilian and Panaman *B. lanceola*, Sw. (Hk. Bk., Syn. Fil., 185; Hk., Icon. Pl., X, tab. CMLXX, right-hand figure), but distinguished from Baker's description and Hooker's drawing by the tufted stipes, the smaller fronds, the mostly once forked and sometimes anastomosing veins. — It bears the name of Prof. Dr. M. Treub, Director of the Agricultural Department in Netherlands India.

Java (Mount Salak).

(2) **B. nitidum**, *Pr.*, Hk. Bk., Syn. Fil., 184; Hk., Spec. Fil., III, tab. CLV; *B. elongatum*, *Gaud.*, Freyc., Voy. Ur. Phys., Bot., 595; C. Chr., Ind. Fil.

Stipes stout, erect, naked, 7—10 c.M. long. Fronds oblong-lanceolate, 50 c.M. or more long, 10—15 c.M. broad. Segments numerous, erecto-patent, subfalcate, linear, 7—10 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, narrowed gradually towards the point, dilated and connected at the base, the edge undulato-dentate; lower segments linear, acute and not much smaller than the rest. Texture coriaceous, rachis and both surfaces naked; veins fine. Sori in a continuous line on each side close to the midrib; indusium conspicuous.

Philippines; Polynesia, Brasilia.

(3) **B. cartilagineum**, *Sw.*, Hk. Bk., Syn. Fil., 184; *B. nitidum*, *Pr.*, var. *contracta*, Hk., Spec. Fil., III, 44, tab. CLVI.

Rhizome oblique, densely clothed at the crown with blackish, fibrillose scales. Stipes strong, erect, 10—15 c.M. long, scaly and muricate in the lower part. Fronds ovate-oblong, 50—60 c.M. long, 15—25 c.M. broad. Segments numerous, erecto-patent, linear, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, narrowed gradually towards the apex, the margin finely toothed, dilated and connected at the base, the lower ones not much smaller than the others. Texture coriaceous; rachis

and both surfaces naked; veins fine. Sori in broad lines close to the midribs.

Philippines; Mishmee, Australia,

* (4) **B. Finlaysonianum**, *Wall.*, Hk. Bk., Syn. Fil., 187; Hk. & Grev., Ic. Fil., tab. CCXXV; Bedd., Ferns Br.I., II, tab. CCIL.

Rhizome short. Stipes erect, brown, polished, 30 c.M. long or longer. Fronds 90—120 c.M. long or longer, with a large terminal pinna and numerous erecto-patent lateral ones on each side. Largest pinnae 20—30 c.M. long, 2½ c.M. broad or broader, narrowed at both ends, the apex acuminate, the base ± obliquely cuneate, the margin entire; upper pinnae decurrent at the base, lower reduced to mere auricles. Texture coriaceous; both surfaces naked. Sori in a continuous line on each side close to the midrib.

Malacca, Borneo, New Guinea.

(5) **B. Blumei**, *Moore*, Ind., 195; *Lomaria auriculata*, *Bl.* (not *Desv.*), Enum., 202.

Stipes subpaleaceous. Fronds pinnate. Pinnae sessile, approximate, naked. Sterile pinnae linear-lanceolate, obtuse; upper base truncate, auriculate, lower subcordate; fertile pinnae linear, truncate, auriculate.

Java.

* (6) **B. orientale**, *L.*, Rac., Flor. Btz., I, 165; Hk., Fil. Exot., tab. LXXVII; Bedd., Ferns S.I., tab. XXIX; — var. *undulatum*, Hk., Spec. Fil., III, 52.

Caudex stout, erect, densely clothed with linear, acuminate, glossy, brown scales. Stipes 10—40 c.M. long, erect, scaly at the base. Fronds 25—200 c.M. long, 10—40 c.M. broad, linear-lanceolate or ovate, with very numerous, nearly contiguous pinnae on each side. Pinnae sessile, the apex acuminate, the upper ones decurrent at the base, the lower ones with the base truncate, rounded or cordate, 5—20 c.M. long, ½—2 c.M. broad, the lowest ones reduced to mere auricles or glands. Texture subcoriaceous; rachis and both surfaces naked; veins fine and close. Sori in a long continuous lines close to the costae.

Var. undulatum: Pinnae transversally undulate, the apex finely acuminate.

Trop. Asia, Australia & Polynesia.

* (7) **B. serrulatum**, *Rich.*, Hk. Bk., Syn. Fil., 186; *B. striatum*, *R. Br.*, Hk., Spec. Fil., III, tab. CLIX.

Caudex elongate, stout, ascending. Stipes 15—30 c.M. long, strong, erect, smooth, nearly naked. Fronds oblong-acuminate, 50—60 c.M. long, 15—25 c.M. broad, with 12—24 pairs of quite distinct, articulate, linear-oblong pinnae on each side, the largest of which are 10—15 c.M. long, 1—2 c.M. broad, narrowed gradually towards the point, and downwards to a narrow base, the margin finely incised. Texture coriaceous; rachis rigid, naked; veins fine and close, not conspicuous. Fertile pinnae narrower. Sori in continuous lines close to the midribs.

Trop. America, Asia & Australia, New Caledonia.

(8) **B. egregium**, *Copel.*, Polypod. Philipp., 90.

Stipes clustered in form of a nest, 10—20 c.M. long, stout, brown, black below and densely clothed with long, black scales. Fronds \pm 100 c.M. long, \pm 30 c.M. broad, pinnate, naked, moderately dimorphous. Pinnae acuminate, serrate, the higher ones confluent, the central ones attached to the rachis by a broad base but free, the lowest reduced into distant auricles; fertile pinnae coriaceous, \pm 15 c.M. long, $1\frac{1}{2}$ —2 c.M. broad at the base, contracted gradually until the sorus occupies very nearly the whole breadth; barren pinnae thinner but broader. Indusium coriaceous, persistent.

Philippines.

50. SADLERIA, *Kaulfuss.*

Sori linear, close to the costa on both sides, continuous or interrupted. Indusium of the same shape as the sorus, attached at its outer side, opening inwardly.

Caudex arborescent. Stipes tufted, not articulate to the caudex. Fronds large, 2-pinnate; veins united near the base by transversal strands, which are occupied by the sori in the fertile segments. Hk. Bk., Syn. Fil., tab. IV, fig. 55; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 132, A—B.

* **S. cyatheoides**, *Hkf.*, Hk. Bk., Syn. Fil., 187; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 132, A—C.

Stipes strong, erect, 15—45 c.M. long, densely clothed with long, linear scales at the base, the rest naked. Fronds 100—250 c.M. long, 2-pinnate.

Pinnae erecto-patent or spreading, 20—40 c.M. long, 1—3 c.M. broad, cut down into very numerous pinnulae. Pinnulae linear, subfalcate, acute or bluntish, sessile with a dilated base, to 4 m.M. broad, those of the upper pinnae connected at the base, the lower ones free. Texture coriaceous; veins inconspicuous; rachis and both surfaces naked. Sori placed on elevated receptacles.

Sumatra; Hawaii.

51. DOODIA, R. Brown.

Sori oblong, straight or slightly recurved, superficial, placed in 1 or more rows on each side, parallel to the costa. Indusium of the same shape as the sorus, attached at the outer side, opening inwardly.

Rhizome \pm erect. Stipes not articulate to the rhizome. Fronds pinnatifid or pinnate; veins forming 1 or more series of arches between the midrib and edge, on which the sori are placed. Hk. Bk., Syn. Fil., tab. IV, fig. 57; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 154, C—D.

* *Fronds pinnatifid, at best pinnate in the lower half, the 1—3 lowest pairs of segments occasionally quite free, rounded and subauriculate at the base.*

(1) *D. dives.*

** *Fronds pinnatifid in the upper, pinnate in the lower half, with some of the lower pairs of pinnae cordate or distinctly auricled, or even hastate at the base.*

(2) *D. media.*

* (1) **D. dives**, *Hae*, Rac., Flor. Btz., I, 164; Kze, Farnkr., II, tab. CV; Bedd., Ferns S.I., tab. CCXXII.

Rhizome \pm erect, densely clothed with dark-brown, lanceolate, acuminate scales. Stipes slender, erect, naked, blackish like the rachis, scaly at the base, 10—30 c.M. long. Barren fronds ovate-lanceolate, to \pm 50 c.M. long, 7—13 c.M. broad, deeply pinnatifid throughout, at best pinnate in the lower half. Segments several to many, linear or linear-oblong, \pm blunt, subundulate, serrate, dilated at the base, \pm 1 c.M. broad, the higher ones confluent, the lower ones confluent or contiguous and occasionally the 1—5 lowest pairs quite free, rounded and subauriculate at the base. Texture coriaceous; surfaces naked. Fertile fronds often longer, with narrower segments; seg-

ments linear, the central ones $\pm 2\frac{1}{2}$ c.M. apart, connected by a decurrent wing, the lowest pairs free. Sori oblong, straight or recurved, placed in 2 irregular rows on each side.

Java; Ceylon.

* (2) **D. media**, *R. Br.*, Hk. Bk., Syn. Fil., 190; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 154, C—D.

Near *D. dives*, *Kze*, but the fronds mostly longer, ± 30 — 45 c.M. long, 4 — 20 c.M. broad, lanceolate or subtriangular-lanceolate, deeply pinnatifid in the upper, but pinnate in the lower half. Segments linear, bluntish or acute, toothed or serrate, the higher ones confluent, the central ones confluent or contiguous and some of the lowest pairs quite free and cordate or distinctly auricled, or even hastate at the base. Texture more or less coriaceous; surfaces naked. Sori oblong, straight or recurved, \pm distant, in 1 — 2 rows on each side, the inner rows not rarely at some distance from the midrib.

Sumbawa, Timor; Australia, New Zealand, Polynesia.

52. WOODWARDIA, *Smith.*

Sori oblong or linear-oblong, in 1 row on each side, parallel and nearly close to the costa, immersed in cavities occupying the costal areolae. Indusium of the same shape as the sorus, attached at the outer side, covering the cavity like a lid, opening inwardly.

Rhizome erect. Stipes not articulate to the rhizome. Fronds pinnate or 2-pinnatifid; veins anastomosing to form 1 or more rows of areolae between the midrib and the edge. Hk. Bk., Syn. Fil., tab. IV, fig. 36; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 134, A—B.

Warmer regions, chiefly of the northern hemisphere.

Arrangement of the sections.

- § 1. **EUWOODWARDIA**. Fronds uniform. Veins anastomosing between the sori and the margin.
- § 2. **ANCHISTEA**. Fronds uniform. Veins free between the sori and the margin. (Not in Malaya).
- § 3. **LORINSERIA**. Fronds dimorphous. Veins anastomosing as in § 1. (Not in Malaya).
-

§ 1. EUWOODWARDIA.

* **W. radicans**, *Sm.*, Hk. Bk., Syn. Fil., 188; Bedd., Ferns Br.I., I, tab. LXXXVIII; Bau. & Hk., Gen. Fil., tab. XVII; *W. auriculata*, *Bl.*, Rac., Flor. Btz., I, 165; *Blechnum radicans*, *L.*, (oldest name); — var. *prolifera*, Copel., Philipp. Journ., III, 55; *W. prolifera*, *Hk. & Arn.*, Beech. Voy., 275, tab. LVI.

Rhizome erect. Stipes strong, erect, naked, scaly at the base. Fronds 1—2 M. long, 50—70 c.M. broad, 2-pinnatifid. Pinnae linear-lanceolate, erecto-patent, the lowest to 40 c.M. long, on stalks 0—5 m.M. long, cut down into linear, subfalcate or nearly straight, acute, serrulate lobes 1—4 c.M. long, the lower lobes often shorter. Texture coriaceous; rachis and under surface naked or covered with long or short, narrow, brown scales or hairs; veins anastomosing, forming 1 row of arches on each side of the costulae (mid-ribs of the lobes), and 1 row of areolae outside the fruit, the rest free. (*W. auriculata*, *Bl.*: Pinnae auricled at the base).

Var. prolifera: Fronds small, membranaceous, proliferous.

Sumatra, Java, Philippines; Formosa, Japan, China, North. India, South. Europe, Atlant. Islands; — var prolifera; Batan Island (Philippines).

TRIBE VI. ASPLENIEAE

53. DIPLAZIUM, Swartz.

Sori linear or linear-oblong, straight or slightly curved, springing from the sides of the veins, \pm oblique to the costa, the lower veins of each group mostly, the higher ones rarely bearing 2 sori, i.e. with a sorus on each side. Indusia single or double (according to the sori), the single ones opening towards the main veins or the central veins of the ultimate segments, the double ones in opposite directions.

Rhizome creeping or erect, the scales finely cancellate, with the areolae mostly dark-coloured, i. e. the cells with the side walls thin and the lumina mostly dark-coloured (Sadeb., in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 40, A). Stipes not articulate to the rhizome, with 2 free, \pm flattened fibro-vascular bundles, which mostly unite towards the lamina (or higher) into a single diarchous one ($\setminus /$, \vee , \sphericalangle , \square , \cup). Fronds simple or divided; veins free or anastomosing. Hk. Bk., Syn. Fil., tab. IV, fig. 38, f—g.

Arrangement of the sections.

- § 1. **EUDIPLAZIUM.** Veins free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 121, A—D.
- § 2. **ANISOGONIUM.** Veins anastomosing. Bedd., Handb. Ind. Ferns, fig. 92—94.

§ 1. EUDIPLAZIUM.

- a.** Fronds simple, the margin entire, crenate or toothed. (1) *D. subserratum.*
- a.a.** Fronds pinnatifid, at best pinnate at the very base.
- §. Fronds herbaceous. Rhizome creeping.
- *. Fronds pinnatifid throughout. Stipes approximate. (2a) *D. Merrilli.*
- *.*. Fronds pinnate at the very base. Stipes rather remote.
- (2b) *D. zeylanicum.*
- §.§. Fronds subcoriaceous. Rhizome erect or short-creeping. Stipes \pm tufted.
- (3) *D. porphyrorachis.*
- a.a.a.** Fronds pinnate or 2-pinnatifid, the lower pinnae sometimes pinnate at the base.
- b.** Pinnae entire, crenate, serrate, toothed or slightly lobed, the lowest sometimes more deeply lobed.

- c.** *Texture ± herbaceous to papyraceous. (*)*
- d.** *Under surface densely scaly on the veins. (4) D. chlorophyllum.*
- d.d.** *Under surface not densely scaly on the veins.*
- e.** *Pinnae truncate, rounded or broadly cuneate at the base.*
- §.** *Stipes ± scaly.*
- ***. *Pinnae ± 6 on each side, not hastate at the base. (5a) D. inconspicuum.*
- *.*.** *Pinnae ± 8—9 on each side, hastate at the base. (5b) D. Weinlandii.*
- §.§.** *Stipes naked, or scaly at the base.*
- ***. *Main veins simple or forked. (6) D. camarinum.*
- *.*.** *Main veins pinnate. (7) D. silvaticum.*
- e.e.** *Pinnae more narrowly cuneate at the base, or the lower base obliquely cut away & narrower than the upper.*
- f.** *Stipe and rachis ± scaly or fibrillose, the scales or fibres persistent or deciduous.*
- g.** *Fronds narrow, ± 3 c.M. broad. Pinnae prominently auricled at the base above. (8) D. acrotis.*
- g.g.** *Fronds broader, ± 3½—5 c.M. broad. Pinnae not auricled at the base. (9) D. biseriale.*
- f.f.** *Stipe and rachis naked. (10) D. aequibasale.*
- c.c.** *Texture ± chartaceous to coriaceous.*
- d.** *Rachis pubescent, at least on the upper (inner) side.*
- §.** *Pinnae paucijugate, rather broad.*
- ×**. *Fronds small. (11a) D. barisanicum.*
- ×.×**. *Fronds middle-sized. (11b) D. tabacinum.*
- §.§.** *Pinnae multijugate, rather narrow. (12) D. confertum.*
- d.d.** *Rachis ± naked, or sometimes slightly pubescent.*
- e.** *Pinnae unequal-sided at the base, i.e. the upper base ± broadly, the lower ± narrowly cuneate.*
- ***. *Stipes stramineous. Fronds pale-green. (13) D. pallidum.*
- *.*.** *Stipes bluish-glaucous. Fronds saturate-green. (14) D. vacillans.*

(*) See also **D. bulbiferum**, Brack., **Williamsi**, Copel. & **Grammitoides**, Pr.

e.e. Pinnae rather equal-sided at the base.

f. Pinnae narrowed at the base.

g. Sori reaching \pm the margin.

h. Pinnae in \pm 1—8 pairs.

(15) *D. bantamense*.

h.h. Pinnae in \pm 13—25 pairs.

(16) *D. xiphophyllum*.

g.g. Sori distant from the margin.

(17) *D. palauanense*.

f.f. Pinnae broad at the base. (*)

a. Fronds narrow, under $7\frac{1}{2}$ c.M. broad.

(18) *D. larutense*.

a.a. Fronds broader, more than $7\frac{1}{2}$ c.M. broad.

β. Lateral pinnae \pm 3—5 on each side below the pinnatifid terminal one.

(19) *D. cultratum*.

β.β. Lateral pinnae several to many below the pinnatifid terminal one.

γ. Largest pinnae $1-2\frac{1}{2}$ c.M. broad, auricled or hastate at the base.

δ. Pinnae rather close, broadly cuneate to truncate at the base.

(20) *D. crenato-serratum*.

δ.δ. Pinnae rather distant, cordate to truncate at the base. (21) *D. Hosei*.

γ.γ. Largest pinnae 2—4 c.M. broad, not auricled. (22) *D. Lobbianum*.

b.b. Pinnae more deeply lobed often to $\frac{1}{2}-\frac{3}{4}$ -way down or more to the costa, sometimes pinnate at the base.

1. Texture \pm herbaceous to papyraceous.

2. Lower pinnae deeply lobed or pinnatifid.

3. Rachis naked.

4. One or both surfaces slightly hairy. (25) *D. japonicum*.

4.4. Surfaces naked or nearly so.

5. Pinnae rather few. (24) *D. shepherdoides*.

5.5. Pinnae rather many.

6. Fronds rather narrow. (25a) *D. Williamsi*.

(*) See also under *D. bantamense*, Bl.

6.6. *Fronds rather broad.* (25b) *D. acuminatum.*

3.3. *Rachis more or less scaly or hairy.*

§. *Pinnae acute at the base.* (26) *D. petiolare.*

§.§. *Pinnae broadly cuneate to truncate at the base.*

‡. *Pinnae 2—10 c.M. long.*

+. *Stipes scaly or fibrillose throughout.*

×. *Pinnae ± 2—3½ c.M. long, ½—1 c.M. broad.* (27a) *D. grammitoides.*

×.×. *Pinnae to ± 10 c.M. long, ± 2 c.M. broad.* (27b) *D. Woodii.*

+.+. *Stipes villose or tomentose, at least on the upper (inner side), sometimes scaly at the base.*

×. *Surfaces villose or short-hairy.*

(28a) *D. Petersenii.*

×.×. *Surfaces naked, except on the costulae beneath.* (28b) *D. Burchardi.*

‡.‡. *Pinnae 10—20 c.M. long in the fully developed fronds.* (29a) *D. sorzogonense.*

(29b) *D. brachysoroides.*

2.2. *Lower pinnae pinnate, at least at the base.*

α. *Stipes muricate.* (30) *D. Prescottianum.*

α.α. *Stipes not muricate.*

β. *Rachis ± naked.*

γ. *Lower pinnae reduced.*

(31a) *D. brachypodium.*

γ.γ. *Lower pinnae not reduced.* (31b) *C. Bolsteri.*

β.β. *Rachis scaly.*

(32) *D. crinitum.*

1.1. *Texture rather coriaceous.*

‡. *Rachis hairy or tomentose.* (33) *D. tomentosum.*

‡.‡. *Rachis naked or ± scaly.*

+. *Lateral pinnae commonly ± 6—10-jugate, lobed.*

(34) *D. bulbiferum.*

+.+. *Lateral pinnae commonly ± 10—20-jugate, pinnatifid.*

!. *Fronds ± 6—15 c.M. broad.* (35a) *D. Whitfordi.*

!:. *Fronds ± 20—35 c.M. broad.*

(35b) *D. speciosum.*

a.a.a.a. Fronds copiously 2-pinnate to 3-pinnatifid (*).

b. Pinnulae lobed at best $\frac{1}{2}$ -way down to the costa.

c. Rachis naked or glabrescent.

d. Fronds herbaceous or membranaceous.

+. Lobes sharply toothed. (36a) *D. cyatheifolium*.

+.+. Lobes obscurely denticulate. (36b) *D. davaoense*.

d.d. Fronds papyraceous to coriaceous.

e. Stipes smooth. (57a) *D. latifolium*.

e.e. Stipes at length sparingly muricate (37b) *D. dolichosorum*.

c.c. Rachis tomentose or scaly.

×. Pinnulae entire to serrate; main rachis scaly.

(38) *D. manilense*.

×.×. Pinnulae broadly and shallowly lobed; main rachis tomentose.

(39) *D. vestitum*.

b.b. Pinnulae pinnatifid at least $\frac{1}{2}$ -way down to the costa.

1. Fully developed fronds under 50 c.M. long. (40) *D. deltoideum*.

1.1. Fully developed fronds more than 50 c.M. long.

2. Main rachis smooth, unarmed.

3. Sori reaching nearly to the edge.

+. Texture thinly coriaceous. (41a) *D. Meyenianum*.

+.+. Texture nearly woody. (41b) *D. atratum*.

3.3. Sori not reaching nearly to the edge.

4. Rachises not ebeneous.

5. Pinnae 30—45 c.M. long; pinnulae $\pm 2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad. (42) *D. amplissimum*.

5.5. Pinnae 20—30 c.M. long; pinnulae ± 1 —2 c.M. broad.

6. Pinnae paucijugate. (43a) *D. oligosorum*.

6.6. Pinnae multijugate.

(43b) *D. polypodioides*.

4.4. Rachises ebeneous. (44) *D. ebeneum*.

2.2. Main rachis \pm asperous, muricate or aculeate.

‡. Costae and costulae not paleaceo-asperous.

+. Ultimate lobes \pm rounded. (45a) *D. asperum*.

+.+. Ultimate lobes truncate. (45b) *D. fructuosum*.

(*) See also *D. crinitum*, C. Chr.

✧✧. *Costae and costulae paleaceo-asperous.*

(46) *D. prionophyllum.*

a.a.a.a. *Fronde 3-pinnate to 4-pinnatifid.*

✧. *Tertiary divisions crenato-pinnatifid at the base.* (47) *D. ceramicum.*

✧✧. *Tertiary divisions deeply pinnatifid.* (48) *D. Huttoni.*

* (1) **D. subserratum**, *Moore*, *Bedd.*, *Handb. Ind. Ferns*, 174: *Ferns Br.I.*, II, tab. CCLXXXIX; *Asplenium subserratum*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 220; *Hk.*, *Spec. Fil.*, III, tab. CLXIV, B.

Rhizome short, erect, (after *Bedd.* creeping & elongated). Stipes firm, erect, naked, $2\frac{1}{2}$ —20 c.M. long. Fronds 20—50 c.M. long, 1—4 c.M. broad, narrowed gradually towards both ends or rounded at the base, the apex acuminate, toothed or bluntly crenate. Texture herbaceous or firm-membranaceous; surfaces naked; costa slender, firm, prominent, scaly beneath; main veins pinnate with few veins. Sori linear, occupying $\pm \frac{2}{3}$ of the space between the costa and the edge.

Java, Malacca.

(2a) **D. Merrilli**, *Copel.*, *Philipp. Journ.*, II, 128, tab. II, A.

Rhizome creeping. Stipes clustered, 2—5 c.M. long, densely clothed with narrow, horizontal, black-brown scales. Fronds 15—25 c.M. long, narrowed towards both ends, acuminate, deeply pinnatifid, the sinuses narrow or the lobes often imbricated. Lobes oblong, nearly horizontal, the central ones the largest, ± 1 c.M. broad, the apex rounded, the edge sometimes denticulate. Texture herbaceous; costa densely scaly beneath, but slightly so above; veins forked, minutely scaly beneath; upper surface black-greenish, lower olive-coloured. Sori linear, not reaching the costulae.

Mindoro.

* (2b) **D. zeylanicum**, *Moore*, *C. Chr.*, *Ind. Fil.*; *D. Ceylanicum*, *Christ*, *Frankr. d Erde*, 216, fig. 679; *Asplenium zeylanicum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 250; *Hk.*, *2nd Cent. of Ferns*, tab. XVI.

Rhizome creeping, slightly covered with lanceolate, brownish scales. Stipes erect, firm, scaly, $\frac{1}{2}$ —3 c.M. apart, to 10—20 c.M. long. Fronds 15—30 c.M. long, $2\frac{1}{2}$ —5 c.M. broad, lanceolate, the apex acuminate, the upper $\frac{1}{4}$ — $\frac{1}{3}$ slightly lobed, the lower $\frac{2}{3}$ — $\frac{3}{4}$ more or less deeply lobed to pinnatifid, the base cut down to the rachis into 1—4 sessile or subsessile pinnae, with

the lowest segments shorter than the next in order. Lobes and pinnae bluntly rounded, linear-oblong, to $\pm \frac{1}{2}$ —1 c.M. broad. Texture herbaceous; veins forked. Sori reaching nearly from the midribs (costulae) to the edge.

Java (escaped from gardens?); Ceylon.

* (3) **D. porphyrorachis**, *Diels*, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 225; *Asplenium porphyrorachis*, *Bk.*, Journ. of Bot., XVII, 40; *Hk.*, Icon. Pl., XVII, tab. MDCL; *A. zeylanicum*, *Ces.*, Fel. di Bor., 21; *A. lomariaceum*, *Christ*, Fil. Saras., III, 229; *Polypodium subserratum*, *Hk.*, *Hk. Bk.*, Syn. Fil., 525.

Rhizome erect or short-creeping, clothed with lanceolate-acuminate, dark-brown scales at the extremity. Stipes \pm tufted, firm, slender, 5—50 c.M. long, scaly at the base, black-hairy or scaly above, the hairs or scales deciduous. Fronds 15—40 c.M. long, $2\frac{1}{2}$ — $6\frac{1}{2}$ c.M. broad, lanceolate, acuminate, the apex lobed or toothed, the lower $\frac{2}{3}$ — $\frac{5}{6}$ deeply pinnatifid into many or numerous, close, spreading, subfalcate, linear-oblong, blunt, slightly crenate, serrate or toothed segments 8—15 m.M. broad, which have not seldom the sides entire and only the apex serrate. Lower segments gradually growing smaller, the 1—4 lowest free and deflexed. Texture subcoriaceous; rachis and under surface slightly covered with scattered scales; veins distinct, mostly once forked. Sori oblique, parallel, reaching \pm the edge.

Malacca, Borneo, Celebes.

(4) **D. chlorophyllum**, *Bedd.*, Handb. Ind. Ferns, Suppl., 39; *Asplenium chlorophyllum*, *Bk.*, Journ. of Bot., XXIII, 104.

Stipes tufted, dull-grey-green, pubescent like the rachis, \pm 15 c.M. long. Fronds oblong-lanceolate, simply pinnate, \pm 30 c.M. long, 10—13 c.M. broad. Pinnae \pm 20 on each side, close, lanceolate, sessile, acute, crenulate towards the apex, conspicuously auricled on the upper side at the base, the central ones 1— $1\frac{1}{2}$ c.M. broad, the lowest strongly deflexed. Texture moderately firm; upper surface minutely paleaceous, under surface densely clothed with minute, linear-subulate scales on the veins; veins distinct, erecto-patent, deeply forked. Sori not quite reaching midrib and margin; indusium crisped, narrow, moderately firm, pale-brown.

Penang; Formosa.

(5a) **D. inconspicuum**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1000.

Rhizome suberect, black. Stipes few, \pm 20 c.M. long, the base black, clothed with subulate, black-brown scales, the rest rufo-stramineous like the

rachis, sparingly scaly towards the frond. Fronds oblong, \pm 20 c.M. long, \pm 11 c.M. broad, hardly narrowed at the base. Terminal pinna \pm 11 c.M. long, acuminate, lobed at the base; lateral pinnae \pm 6 on each side, oblong, nearly sessile or short-stalked, rather distant, spreading, \pm 5 c.M. long, \pm 1½ c.M. broad, the base subtruncate on the upper, rounded on the lower side, the edges densely 2-crenulate, the apex obtuse. Texture herbaceous; rachis and surfaces subglabrous; veins dense, forked, subpinnate in the terminal pinna. Sori distant, unequal, linear, not reaching the margin; indusium thin, inconspicuous, greyish.

Luzon.

(5b) **D. Weinlandii**, *Christ*, Bull. Herb. Boiss., 2^e Série, I, 452.

Rhizome oblique, black, decumbent or creeping. Stipes subfasciculate, erect, subflexuose, slender, 10—16 c.M. long, the base black, the rest green, clothed like the rhizome with few, distant, black, subulate scales. Fronds elongate-lanceolate, \pm 15 c.M. long, 4—5 c.M. broad at the base, pinnate. Pinnae 8—9 on each side, distant, the lower ones short-stalked, obtuse, the base hastate, (the auricles obtuse, rarely deeply incised), the margin broadly, but not deeply incised and crenate, the teeth acute; terminal pinna \pm 6 c.M. long, pinnatifid. Texture herbaceous; surfaces naked; veins oblique, 7—8 on each side, forked or sometimes 3-partite. Sori reaching from the midrib to the margin, narrow; indusium thin, grey.

New Guinea.

(6) **D. camarinum**, *Copel.* (not *Bk.*), Polypod. Philipp., 73; *Asplenium camarinum*, *Bk.*, Hk. Bk., Syn. Fil., 230.

Stipe and rachis dull-brown, naked. Fronds oblong, simply pinnate, 20—45 c.M. long, 15—25 c.M. broad. Pinnae in 8—12 pairs, distant, stalked, linear-ligulate, acute, crenate, 10—15 c.M. long, \pm 2 c.M. broad, rather reduced on the lower side, rounded on both sides at the base; lowest pinnae not reduced. Texture membranaceous; surfaces naked; colour dull-green; veins distinct, in close, little-ascending pairs, posterior forked, anterior simple. Sori reaching from the costa to the edge; indusium persistent.

South. Camarines.

* (7) **D. silvaticum**, *Sw.*, Christ, Farnkr. d. Erde, 217; Bedd., Ferns S.I., tab. CLXI (small); *D. acuminatum*, *Pr.*, Tent. Pterid., 113; *Asplenium sylvaticum*, *Pr.* (not *Mett.*), Rac., Flor. Btz., I, 223; *Callipteris sylvatica*, *Bory*, (oldest name).

Rhizome erect or decumbent, slightly clothed at the apex with lanceolate scales. Stipes 10—30 c.M. long, firm, erect, naked, yellowish to brownish, scaly at the base. Fronds \pm 20—60 c.M. long, \pm 10—30 c.M. broad, ovate-to triangular-lanceolate, with many or numerous spreading pinnae. Largest pinnae 5—15 c.M. long, 1—2 $\frac{1}{2}$ c.M. broad, acuminate, or the upper ones blunt or bluntish, the edges crenate and broadly, mostly shallowly lobed (commonly to 2 $\frac{1}{2}$ m.M. deep), the base rounded, broadly cuneate or truncate, whether or not auricled on the upper side; lower pinnae stalked, upper sessile; apex of the frond deeply pinnatifid. Texture thin, herbaceous; veins pinnate in the lobes. Sori long, linear.

Pantropical.

(8) **D. acrotis**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1000.

Rhizome oblique, short. Stipes tufted, numerous, 3—4 c.M. long, slender, pale-castaneous like the rachis, crinite with subulate, black-brown scales. Fronds caespitose, 12—14 c.M. long, \pm 3 c.M. broad, pinnate, the rachis slender, nearly filiform, not winged, sparingly furfuraceous. Pinnae \pm 14 on each side below the pinnatifid, acuminate apex, \pm horizontal, stalked, regular, oblong, \pm 1 $\frac{1}{2}$ c.M. long, \pm 8 m.M. broad, subfalcate, the base cuneate on the lower, truncate and prominently and acutely auricled on the upper side, the edge coarsely toothed with 5—6 teeth on each side, the apex bluntish or acute; lower pinnae becoming shorter, higher sessile. Texture herbaceous; veins 1 to each tooth, simple, the lowest anterior one pinnate in the auricle. Sori oblong, oblique, regular, reaching the midrib and the edge, often double.

Luzon.

(9) **D. biseriale**, *C. Chr.*, Ind. Fil.; *Asplenium biseriale*, *Bk.*, Trans. Linn. Soc., Bot., 2nd Series, IV, 252.

Rhizome short-creeping. Stipes erect, 3 $\frac{1}{2}$ —5 c.M. long, clothed above with lanceolate, spreading, brown scales. Fronds oblong-lanceolate, simply pinnate, naked, 10—15 c.M. long, 3 $\frac{1}{2}$ —5 c.M. broad; rachis covered with small scales. Pinnae multijugate, sessile, lanceolate, the higher ones entire, the lower \pm 5 m.M. broad, deeply crenate, with the lower base reduced; veins obscure, erecto-patent. Sori in 1 row on each side of the costae, linear-oblong, often diplazioid; indusium broad, membranaceous, naked, persistent.

Borneo.

(10) **D. aequibasale**, *C. Chr.*, Ind. Fil.; *Asplenium aequibasale*, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 225.

Rhizome erect. Stipes tufted, 7—10 c.M. long, with few, black, lanceolate, spreading scales near the base. Fronds oblong, 15—20 c.M. long, 10—15 c.M. broad, simply pinnate. Pinnae alternate, 9—11, adnate, distant, lanceolate, acuminate, crenulate, the base narrowed and equal on both sides, the middle 1—1½ c.M. broad. Texture rather thin; surfaces naked; veins conspicuous, erecto-patent, simple or forked. Sori medial on the veins, parallel; indusium narrow, naked, persistent.

Borneo.

(11a) **D. barisanicum**, *C. Chr.*, Ind. Fil.; *Asplenium barisanicum*, *Bk.*, Ann. of Bot., VIII, 126.

Stipes ± 15 c.M. long, slender, green, naked, clothed near the base with linear, brown scales. Fronds deltoid-oblong, 15—20 c.M. long, 12—15 c.M. broad, simply pinnate. Terminal pinna pinnatifid; lateral pinnae paucijugate, lanceolate-oblong, 3—4 c.M. broad, deeply crenate, short-stalked, the lower ones not or hardly shorter. Both surfaces naked; rachis pubescent above; main veins pinnate; veins erecto-patent, simple. Sori occupying the lower portion of the veins; indusium membranaceous, naked, persistent.

Sumatra.

(11b) **D. tabacinum**, *Copel.*, Philipp. Journ., I, Suppl., 149, tab. VI.

Rhizome erect, clothed with brown, lanceolate-subulate scales. Stipes 30—40 c.M. long, canaliculate, the lower part sparingly scaly, the upper part short-pubescent like the rachis and the stalks of the pinnae. Fronds 30—40 c.M. long, ovate, pinnate. Pinnae 2—4 on each side below the terminal one, simple, lanceolate-oblong, the base cuneate-truncate, the edge serrate towards the falcato-acuminate apex; lowest pinnae largest, ± 15 c.M. long, ± 5 c.M. broad, the terminal one simple or with a pair of basal lobes. Texture subcoriaceous; surfaces naked; veins 3—4-furcate. Sori linear, reaching nearly the costae and the margin.

Mindanao.

(12) **D. confertum**, *C. Chr.*, Ind. Fil.; *Asplenium confertum*, *Bk.*, Ann. of Bot., VIII, 125.

Stipes ± 15 c.M. long, scaly at the base, slender, green, the scales

lanceolate, firm, castaneous. Fronds lanceolate-oblong, 15—25 c.M. long, 6½—7½ c.M. broad, pinnate. Pinnae multijugate, 15—18 m.M. broad, close, sessile, obtuse, crenate, the upper base auricled; lower pinnae not much shorter, deflexed. Texture firm; surfaces naked; rachis pubescent; veins obscure, erecto-patent, forked. Sori linear, reaching from the costae to the margin; indusium naked, narrow, persistent.

Sumatra.

* (13) **D. pallidum**, *Moore* (not *Bl.*), *Copel.*, *Polypod. Philipp.*, 75; *Bedd.*, *Ferns Br.I.*, II, tab. CXCVI; *Asplenium pallidum*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 222.

Rhizome short, creeping, clothed with black, lanceolate scales. Stipes 15—30 c.M. long, firm, erect, stramineous, naked or scaly at the dark-brown base. Fronds 30—100 c.M. long, with numerous nearly horizontal, straight or falcate pinnae on each side. Pinnae 7—15 c.M. long, short-stalked, linear-lanceolate, the apex acuminate, 1—2 c.M. broad, the edge often sharply toothed, the upper base truncate, often rounded, the lower cuneate, slightly rounded. Texture coriaceous; veins once or twice forked. Sori in regular lines, reaching from the midribs to the edge or occupying the lower half of the veins.

Malaya; Queensland.

(14) **D. vacillans**, *C. Chr.*, *Ind. Fil.*; *Asplenium vacillans*, *Kze*, *Bot. Zeit.*, 1848, 172.

Rhizome short-creeping, clothed with black scales. Stipes 40—45 c.M. long, bluish-pruinose, the base blackish, scaly like the rhizome. Fronds oblong, ± 50 c.M. long, 15—20 c.M. broad, acuminate. Pinnae ± stalked, spreading, rather distant, linear-oblong, falcate, the apex narrowed, the edge regularly and broadly serrate, the upper base truncate, the lower subcuneate. Texture thinly coriaceous; surfaces naked; costae prominent; veins forked, the higher simple, the lowest once or more times forked, oblique. Sori rather distant, rarely diplazioid, commonly reaching from the midribs to the margin.

Java.

* (15) **D. bantamense**, *Bl.*, *Christ*, *Farnkr. d. Erde*, 216; *D. fraxinifolium*, *Bedd.*, *Ferns Br.I.*, I, tab. LXIX; *Asplenium fraxinifolium*, *Wall.*, *Hk.*, 2nd Cent. of *Ferns*, tab. XIX; *A. bantamense*, *Bk.*, *Rac.*, *Flor. Btz.*, I, 221; — var. *alternifolium*; *A. alternifolium*, *Mett.*, *Ann. Mus. Bot. L. B.*, II, 237; *D. alternifolium*, *Bl.*, *Hk.*, *Fil. Exot.*, tab. XVII.

Rhizome short, stout, clothed with brown, lanceolate scales. Stipes tufted, 15—30 c.M. long, erect, firm, stramineous, scaly at the base. Fronds \pm 20—100 c.M. long, with a large terminal and 1—8 pairs of \pm stalked, lateral pinnae. Pinnae 10—20 c.M. long, \pm 2—5 c.M. broad, narrowed at both ends, the point acuminate, the edge serrate or nearly entire; higher pinnae sessile. Texture coriaceous but rather thin; surfaces naked; veins \pm 3 \times forked. Sori narrow, irregular, the longest reaching nearly from the midrib to the edge.

Var. alternifolium: Stipes green, passing gradually into brown at the base. Pinnae ovate, broad, $7\frac{1}{2}$ —10 c.M. long, the base rounded or cordate, the edge entire, obscurely sinuate or subcrenato-serrate towards the apex. Texture subcarnosocoriaceous.

Trop. Asia; — var. *alternifolium*: *Java*.

(16) **D. xiphophyllum**, *C. Chr.*, *Ind. Fil.*; *Asplenium xiphophyllum*, *Bk.*, *Journ. of Bot.*, XVII, 40.

Rhizome suberect. Stipes close, to 30 c.M. long, the base densely clothed with large, brown, linear scales. Fronds oblong-deltoid or deltoid, simply pinnate, 30—45 c.M. long, with a terminal pinna similar to the lateral ones. Pinnae 13—25, subsessile, lanceolate, 20—25 c.M. long, 2—4 c.M. broad, toothed and much acuminate at the point, cuneate at the base. Texture subcoriaceous; surfaces naked; veins in pinnate groups of 3—4. Sori reaching from the midrib to the edge; indusium narrow, naked, persistent.

Borneo.

(17) **D. palauanense**, *Copel.*, *Polypod. Philipp.*, 72.

Stipes firm, 30—40 c.M. long, the base black and covered with black scales, the higher part stramineous. Fronds 30—60 c.M. long, pinnate. Pinnae alternate, distant, \pm 6-jugate, the largest \pm 30 c.M. long, $2\frac{1}{2}$ —5 c.M. broad, acute, obscurely serrate, the higher ones sessile, the lower narrowed at the base into a short stipe, both sides nearly equal. Texture coriaceous; rachis naked; surfaces naked; veins 2—3 \times forked. Sori linear, distant from the margin.

Philippines.

(18) **D. larutense**, *Bedd.*, *Handb. Ind. Ferns, Suppl.*, 38.

Rhizome erect. Stipes \pm 30 c.M. long, firm, erect, naked. Fronds narrow, linear-lanceolate, \pm 30 c.M. long, 4—5 c.M. broad, broadest at the base, gra-

dually tapering upwards, pinnate, pinnatifid towards the apex; rachis deeply channelled above. Pinnae numerous, horizontal, oblong, with a broad base and rounded apex, ± 1 c.M. broad, the margin slightly crenate, the upper base slightly lobed. Texture coriaceous; veins simple or the lowest forked. Lowest sori sometimes double, the others always single, reaching from the midrib to the margin.

Malacca.

(19) **D. cultratum**, *Pr.*, Copel., Polypod. Philipp., 73; *Asplenium cultratum*, *Hk.* (not *Mett.*), *Hk. Bk.*, Syn. Fil., 231.

Stipes tufted, 10—15 c.M. long, grey, naked. Fronds 15—25 c.M. long, 7—10 c.M. broad, the apex pinnatifid, the lower $\frac{2}{3}$ pinnate with 3—5 pinnae on each side. Lower pinnae distinctly stalked, ± 5 c.M., long, ± 2 c.M. broad, the point acute, rather falcate, the edge nearly entire, both sides narrowed suddenly at the base, the upper base auricled. Texture subcoriaceous; veins twice forked; sori linear, slender, reaching the edge but not the midrib.

Luzon.

* (20) **D. crenato-serratum**, *Moore* (not *Bl.*), *Christ*, *Farnkr. d. Erde*, 217; *D. porrectum*, *Pr.*, *Bedd.*, *Handb. Ind. Ferns*, 176; *Ferns Br.I.*, II, tab. CCXLV; *Asplenium porrectum*, *Wall.*, *Hk. Bk.*, Syn. Fil., 230; *A. crenato-serratum*, *Bl.*, (oldest name).

Stipes tufted, to 50 c.M. long or longer, brownish. Fronds 20—45 c.M. long, 10—25 c.M. broad, deltoid-oblong, with several to numerous horizontal pinnae on each side. Pinnae linear-lanceolate, nearly straight, acute or bluntish, the edge varying from slightly crenate to shallowly lobed or broadly toothed, the base \pm equally and broadly cuneate to truncate and auricled on the upper side; lower pinnae $\pm 1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, short-stalked, bluntly lobed, longer than the next in order, sometimes shorter or deflexed; highest pinnae erecto-patent, sessile, adnate to the rachis. Texture more or less distinctly coriaceous; rachis naked or slightly pubescent; surfaces naked; main veins pinnate; veins simple or forked. Sori in regular, parallel rows, extending from the midrib to the edge.

Malaya.

* (21) **D. Hosei**, *Christ*, *Bull. Soc. Bot. Belg.*, XXXIII², 93.

Stipes long, slightly rough like the rachis. Fronds ± 30 —45 c.M. long, ± 14 —18 c.M. broad, triangular, pinnatifid at the apex. Pinnae ± 12 —20-

jugate, spreading, the highest ones the shortest, the lower ones the largest, short-stalked, broadly lanceolate, ± 3 c.M. apart, ± 2 c.M. broad, cordate to truncate at the base, the upper base auriculate to hastate, the apex acute to acuminate, the edge regularly but slightly crenate to serrate. Texture firm, chartaceous or subcoriaceous; surfaces naked; costae distinct; veins 2—5-jugate, inconspicuous. Sori ± 25 on each side of the costa, ± 1 c.M. long, only double near the margin.

Borneo.

* (22) **D. Lobbianum**, *Moore*, C. Chr., Ind. Fil.; *Asplenium Lobbianum*, *Hk.*, Spec. Fil., III, 244; 2nd Cent. of Ferns, tab. XVII.

Rhizome suberect, clothed with black, subulate scales. Stipes approximate or subtufted, 15—50 c.M. long, scaly at the base. Fronds oblong-ovate, 20—45 c.M. long, with a pinnatifid apex and to 20 or more pinnae on each side. Pinnae subhorizontal, linear-oblong or lanceolate-oblong, $7\frac{1}{2}$ —15 c.M. long, 2—4 c.M. broad, stalked, subfalcate, acuminate, the edge more or less serrate, the base \pm truncate and subauriculate. Texture thinly coriaceous; surfaces naked; veins 2—3 \times forked. Sori not reaching the margin or the midrib.

Java, Philippines, New Guinea.

* (25) **D. japonicum**, *Bedd.*, Christ, Farnkr. d. Erde, 218; *D. decussatum*, *Bedd.*, Ferns Br.I., II, tab. CCXCII; *Asplenium japonicum*, *Thb.*, Hk. Bk., Syn. Fil., 234.

Rhizome slender, wide-creeping. Stipes 15—30 c.M. long, stramineous or brownish, slightly scaly towards the base. Fronds 20—35 c.M. long, 10—15 c.M. broad, ovate-lanceolate to subdeltoid. Terminal pinna pinnatifid, narrowed gradually towards the point; lateral pinnae 8—10 on each side, rather distant, to $2\frac{1}{2}$ c.M. broad, short-acuminate, cut down in the lower part $\frac{2}{3}$ of the way to the costa, the base broadly cuneate or subtruncate. Lobes oblong, rounded or bluntish, slightly toothed, approximate. Texture herbaceous; surfaces slightly hairy; rachis \pm naked; veins 4—6 on each side in the lower lobes. Sori short, not reaching the margin.

Trop. Asia, China, Japan.

(24) **D. shepherdioides**, *C. Chr.*, Ind. Fil.; *Asplenium shepherdioides*, *Bk.*, Ann. of Bot., VIII, 126.

Stipes 20—25 c.M. long, slender, green, naked. Fronds oblong-lanceolate, 30 c.M. and more long, 2-pinnatifid; surfaces naked; rachis slender, green,

naked. Pinnae lanceolate, acuminate, 7—13 c.M. long, $2\frac{1}{2}$ —3 c.M. broad, paucijugate, cut down $\frac{1}{2}$ -way to the costa, the lower base cuneate; higher pinnae sessile, lower short-stalked, not reduced. Lobes oblong, 5— $6\frac{1}{2}$ m.M. broad, entire. Main veins pinnate; veins 4—6-jugate, erecto-patent, simple. Sori not reaching the apices of the veins; indusium narrow, naked, persistent.

Sumatra.

(25a) **D. Williamsi**, Copel., Philipp. Journ., I, Suppl., 150, tab. VII.

Rhizome erect, short, the scales subulate, blackish-fuscous. Stipes 4—10 c.M. long, erect, the base black, scaly, the upper part green, naked. Fronds 10—20 c.M. long, lanceolate, acuminate, pinnate, the rachis slender, green, naked. Pinnae in ± 10 pairs below the pinnatifid apex, alternate, serrate or slightly lobed, obtuse or acute, the upper base truncate-auriculate, the lower cuneate; lowest pinnae distant, short-stalked, ± 2 c.M. long, deltoid-lanceolate, deeply pinnatifid into oblong, obtuse, denticulate lobes. Texture papraceous; surfaces naked; veins subconspicuous, 2—4-furcate. Sori elongate, often double. — Fruiting fronds are found less than 5 c.M. long including the stipe, with the lowest pinnae less incised. In size and shape this species resembles *D. grammitoides*, Pr.

Mindanao.

(25b) **D. acuminatum**, Bl., Christ, Ann. Bot., XII, 220; *D. Christii*, C. Chr., Ind. Fil.; *Asplenium acuminatum*, Wall., List, No. 205.

Stipes naked, yellowish, ± 55 c.M. long. Fronds oblong-deltoid, ± 55 c.M. long, ± 25 —30 c.M. broad. Pinnae ± 20 on each side, the lowest largest, ± 2 c.M. broad, distinctly stalked, the base slightly rounded, the edge cut $\pm \frac{1}{2}$ -way down to the costa, the apex long, toothed. Segments close, broadly ligulate, blunt, truncate, ± 5 m.M. broad, entire or slightly denticulate. Texture herbaceous; surfaces naked. Sori ± 5 on each side of the costulae, not reaching the latter, linear; indusium very narrow, inconspicuous.

Malaya.

(26) **D. petiolare**, Pr., Copel., Polypod. Philipp., 74.

Stipes brown, smooth. Fronds about 50 c.M. long, pinnate, with a pinnatifid apex. Pinnae ± 12 c.M. long, the lowest a little shorter, but longer

stalked, linear, acute, pinnatifid $\frac{1}{2}$ -way down to the costa, the base equilateral, acute. Texture herbaceous; colour bright-green above, paler beneath; surfaces naked; rachis densely beset with short, spreading hairs; veins free, all bearing sori.

Philippines.

* (27a) **D. grammitoides, Pr.**, Copel, Polypod. Philipp., 73; *Asplenium grammitoides, Hk.*, Hk. Bk., Syn. Fil., 491; Hk., Icon. Pl., X, tab. CMXIII.

Rhizome creeping, the scales brown, lanceolate, acuminate. Stipes seriate, approximate, slender, 6—15 c.M. long, deciduously scaly like the rachis, the scales fibril-like, spreading, crisped. Fronds linear-lanceolate, 10—25 c.M. long, 5—6 c.M. broad, the upper $\frac{1}{3}$ — $\frac{1}{2}$ deeply pinnatifid, toothed to serrate towards the acuminate point, the lower part pinnate with 6—10 erecto-patent pinnae on each side. Central pinnae largest, subsessile, lanceolate, subfalcate, blunt or acute, 2—3 $\frac{1}{2}$ c.M. long, $\frac{1}{2}$ —1 c.M. broad, the edge toothed or lobed and crenate, the base broadly cuneate, distinctly auricled on the upper side; lower pinnae shorter, remote, blunt, \pm stalked, the edge distinctly lobed with short, blunt, subentire or crenate lobes, the lowest anterior lobe the largest. Texture herbaceous but firm; surfaces sparingly setulose; rachis winged in the upper part; veins oblique, once or more times forked. Sori moderately large, the 1—3 lower of each pinna often double. — Fruiting young specimens often small and with the lowest pinnae less incised. The description is taken from specimens gathered by Raciborski in Central Java and determined as *Asplenium pusillum, Bl.*

Malaya.

(27b) **D. Woodii, Copel.**, Philipp. Journ., II, 129.

Rhizome erect. Stipes close, 20—50 c.M. long, straight, the base grey-brown and densely clothed with brown, falcate scales, the upper part stramineous-brown and covered like the rachis with few, short, narrow, castaneous, whether or not deflexed, distant scales. Fronds 40—50 c.M. long, hardly 20 c.M. broad, acuminate, 2-pinnatifid, the rachis stramineous-brown. Pinnae \pm 12—17 on each side, the central ones the largest, horizontal, short-stalked, \pm 2 c.M. broad, acuminate, deeply pinnatifid, the base truncate, the rachis castaneous. Lobes oblong, \pm 5 m.M. broad, subfalcate, serrulate, obliquely truncate, separated by sinuses \pm 1 m.M. broad. Texture herbaceous; surfaces naked, upper black-green, lower olivaceous; veins simple 5—6-jugate.

Sori 2—3 m.M. long, oblique, rarely diplazioid, distant from the costulae, not reaching the margin.

Mindoro.

* (28a) **D. Petersenii** *Christ*, Bull. Ac. Int. Bot., 1902, 245; *D. Lasiopteris*, *Kze.*, Linn., XVII, 568; *Bedd.*, Ferns S.I., tab. CLX; *Asplenium lasiopteris*, *Mett.*, Rac., Flor. Btz., I, 226; *A. tomentosum*, *Mett.*, Farng., VI, Asplen., No. 209; *A. Petersenii*, *Kze.* (oldest name).

Rhizome creeping, clothed with short, broad, \pm adpressed scales. Stipes 15—30 c.M. long, erect, villose, scaly at the base, 1—3 c.M. apart. Fronds 15—45 c.M. long, 6—20 c.M. broad, lanceolate, with a toothed to deeply pinnatifid terminal pinna. Lateral pinnae \pm 8—15 on each side, \pm horizontal, straight or falcate, bluntish or acute to acuminate, close or \pm distant, the upper ones sessile, the lowest subsessile, sometimes deflexed and shorter than the next in order, the longest $\frac{3}{4}$ —3 c.M. broad, cut \pm $\frac{1}{2}$ -way down to nearly to the costa, the base broadly cuneate, nearly truncate, or sometimes narrowed in the lower pinnae, the edges of the lobes \pm crenate. Texture herbaceous; rachis and at least the under surface villose or paleaceous; main veins pinnate in the lobes. Sori short.

Java, New Guinea; China, Br. India.

(28b) **D. Burchardi**, *Rst.*, Fed., Rep., IV, 294.

Stipes slender, to 15 c.M. long or longer, the inner side deeply sulcate, the groove tomentose, the rest naked; rachis slightly hirsute. Fronds deltoid-lanceolate, long-acuminate, to 27 c.M. long, 16 c.M. broad at the base, pinnate. Pinnae to 20-jugate, subcontiguous, spreading, falcate; lowest pinnae largest, to 8 c.M. long, $2\frac{1}{2}$ c.M. broad, ovate-elongate, short-acuminate, pinnatifid nearly to the costa, the segments linear, moderately acute, suberect, somewhat falcate, the lowest very abbreviate, the central the largest, \pm $1\frac{1}{4}$ c.M. long, 4 m.M. broad, the highest rather suddenly diminished into the denticulate apex; next following pinnae similar to the basal ones in the fully developed plants, but similar to the central pinnae in the fronds of the younger plants; central pinnae short-stalked, subfalcate, \pm equally truncate at the base on both sides, auricled on the upper side, linear-lanceolate, short-acuminate, pinnatifid \pm $\frac{1}{2}$ -way down to the costa, the segments linear-lanceolate, falcate, the lowest anterior elongate, the lowest posterior slightly reduced or nearly as long as the others, the medial ones \pm 5 m.M. long, 3 m.M. broad at the base, the higher ones gradually tapering into the crenate apex, the point

entire; higher pinnae sessile, obliquely serrate at the margin, otherwise similar to the medial ones; highest pinnae gradually passing into the lobes of the terminal pinna. Texture chartaceo-herbaceous; colour blackish when dry; surfaces naked; costulae pinnate, slightly hirsute beneath; veins simple, to 8-jugate in the lower, to 5-jugate in the central pinnae. Sori copious, straight, parallel, reaching \pm from the costulae to the edge; indusium firm, shortly fimbriate, pilose when young. — Near *D. sorzogonense*, *Pr.*

Sumatra.

* (29a) ***D. sorzogonense***, *Pr.*, Copel. Polypod. Philipp., 74; Bedd., Ferns Br.I., II, tab. CCXLVI; *Asplenium sorzogonense*, *Pr.*, Hk. Bk., Syn. Fil., 236; — var. major, Bedd., Handb. Ind. Ferns, Suppl., 40.

Stipes tufted, 10—15 c.M. long, densely fibrillose below. Fronds 50—65 c.M. long, 20—55 c.M. broad, with numerous pinnae on each side, the lower ones sessile, 10—18 c.M. long, \pm 2½ c.M. broad, \pm truncate at the base, cut down regularly throughout $\frac{2}{3}$ of the way down to the costa into spreading, blunt, subentire lobes \pm 4—5 m.M. broad. Texture herbaceous, rachis slightly fibrillose; main veins pinnate in the lobes; veins simple. Sori in regular rows, reaching from the main veins to the edge.

Var. major: Stipes \pm 75 c.M. long. Fronds 120—180 c.M. long. Pinnae \pm 50 c.M. long, 5—7½ c.M. broad, the 3 lower pairs much deflexed, cut down nearly to the costa into narrow, slightly crenated segments. Surfaces naked; veins generally forked. — Perhaps a distinct species.

Philippines, Malacca; Himalaya; — var. major: *Malacca.*

(29b) ***D. brachysoroides***, *Copel.*, Philipp. Journ., II, 127, tab. I, A.

Stipes 20—40 c.M. long, black in the lower, dirty-brown in the upper part, sparingly clothed like the rachis with narrow, spreading, flexuose scales becoming smaller towards the apex of the frond. Fronds 50—60 c.M. long, \pm 30 c.M. broad, acuminate, hardly 2-pinnate, the apex pinnatifid. Central pinnae largest, horizontal, acuminate, deeply pinnatifid, the base truncate or subtruncate, with a short, black stalk; lowest pinnae slightly remote and reduced, sometimes deflexed. Lobes oblong, subacute, the largest \pm 3 c.M. long, \pm 1 c.M. broad, sharply serrate towards the apex, connected at the base by a narrow wing, but some of them sometimes free, the lowest lobes of the lowest pinnae reduced. Texture papyraceous; upper surface black-

green, lower olive-green; veins forked. Sori small, oblong, oblique, \pm costular, hardly 2 m.M. long, the lowest often double, longer.

Mindoro.

(50) **D. Prescottianum**, *Moore*, C. Chr., Ind. Fil.; *Asplenium Prescottianum*, *Wall.*, *Hk.*, *Spec. Fil.*, III, 251.

Stipes \pm 55 c.M. long, stramineous, muricate. Fronds to 80 c.M. long, ovate-lanceolate, with several to numerous spreading pinnae. Lower pinnae 10 c.M. apart, upper 4 c.M. apart, all petiolate, on stalks to 1 c.M. long except the highest ones, which are sessile; lower pinnae pinnatifid $\frac{2}{3}$ on the way down into blunt lobes and the lowest cut down quite to the costa at the base, (the lower segments 2 c.M. long, linear-oblong, blunt at the apex), the base suddenly narrowed or rounded on both sides. Texture thin-herbaceous; rachis naked, smooth or muricate; main veins fine, pinnate in the lobes with the short sori on the branches.

Malaya.

(31a) **D. brachypodium**, *Copel.*, *Polypod. Philipp.*, 75; *Asplenium brachypodium*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 490; (? *D. grammitoides*, *Pr*, C. Chr., Ind. Fil., 669).

Stipes tufted, slender, 1—2 $\frac{1}{2}$ c.M. long, with a few, minute, linear scales. Fronds oblanceolate, acuminate, 10—15 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad, 2-pinnate. Lower pinnae lanceolate-deltoid, \pm $\frac{1}{2}$ c.M. broad, distinctly stalked, square on upper side, cuneate-truncate on lower, cut down to the rachis below; many lower pinnae distant, reduced. Lowest anterior pinnulae much the largest, cuneate-oblong, dentate, 2—4 m.M. broad, the others lanceolate, entire. Texture moderately firm; rachis naked; main veins pinnate in the lower lobes. Sori medial, linear.

Philippines.

(31b) **D. Bolsteri**, *Copel.*, *Philipp. Journ.*, I, *Suppl.*, 254, tab. II.

Rhizome ascending. Stipes approximate, 5—10 c.M. long, slender, the base black, clothed with black-castaneous scales, the upper part dirty-green and nearly naked like the rachis. Fronds 20—25 c.M. long, 3 $\frac{1}{2}$ —5 c.M. broad, acuminate, the terminal pinna long, pinnatifid at the base. Lateral pinnae many on each side, falcate, erecto-patent, stalked, dentato-lobate, the base subtruncate, conspicuously and deeply auricled on the upper, obliquely cut away on the lower side, the apex serrate; lower pinnae deflexed, pinna-

tifid, pinnate at the base (the auricle free, serrate or crenate); higher pinnae toothed. Texture papyraceous; surfaces naked; veins distinct, forked, pinnate in the auricles. Sori very oblique; indusium narrow, castaneous-fulvous.

Mindanao.

(52) **D. crinitum**, *C. Chr.*, Ind. Fil.; *Asplenium crinitum*, *Bl.*, Journ. of Linn. Soc., Bot., XXIV, 258; *A. sorzogonense*, *Pr.*, var. *majus*, *Hk.*, Spec. Fil., III, 252.

Stipes strong, long, clothed like the rachis with lanceolate, castaneous, crisped scales. Fronds oblong-lanceolate, 2-pinnatifid or 2-pinnate. Pinnae lanceolate or the lower ones lanceolate-oblong, ± 15 c.M. long, with the largest segments ± 5 c.M. long, $1-1\frac{1}{2}$ c.M. broad. Ultimate lobes parallel, oblong, obtuse, 5-9 m.M. broad. Texture membranaceous; main veins pinnate; lower veins forked, oblique. Sori confined to the inner half of the segments, neither very close nor so regular.

Borneo.

* (55) **D. tomentosum**, *Bl.*, Bedd., Handb. Ind. Ferns, 179; Ferns Br.I., II, tab. CXCv; *Asplenium tomentosum*, *Hk.* (not *Mett.*), Rac., Flor. Btz., I, 226.

Rhizome short, the scales dense, lanceolate-acuminate, black or dark-brown. Stipes tufted, dark-brown, fibrillose above, scaly towards the base, $\pm 9-50$ c.M. long. Fronds lanceolate, acuminate, 15-30 c.M. long; rachis dark-brown, fibrillose or villose; the apex deeply pinnatifid with broadly serrate or crenate, blunt, oblong lobes. Pinnae $\pm 15-20$ on each side, the highest oblong, erecto-patent, blunt, sessile with a broad base, slightly auricled above, the edge crenato-serrate; central pinnae sessile or subsessile, spreading, $\pm 3-3\frac{1}{2}$ c.M. long, 5-8 m.M. broad, blunt or acute, the upper base truncate-auriculate, the lower base truncate, rounded or broadly cuneate, the edge distinctly toothed; lowest pinnae stalked, deflexed, lanceolate, $\pm 4-5$ c.M. long, blunt or acuminate, $\pm 1\frac{1}{2}$ c.M. broad at the centre, the edge cut down $\frac{1}{2}-\frac{3}{4}$ on the way to the costa, the segments oblong, blunt, erecto-patent, serrate or crenate, the lowest anterior segment (auricle) free, deflexed. Texture coriaceous; surfaces naked; costae and veins distinct, fibrillose beneath; main veins pinnate in the lobes, $1-2 \times$ forked in the teeth of the central pinnae; veins often \pm hairy. Sori rather few, slightly recurved, reaching \pm the midrib and the margin.

Malaya.

(34) **D. bulbiferum**, *Brack.*, Expl. Exp., XVI, 141, tab. XVIII, fig. 1; *D. Brackenridgei*, *Diels*, Schum. & Laut., Flor. deut. Sch.geb., 124; *Asplenium Brackenridgei*, *Bk.*, Hk. Bk., Syn. Fil., 254.

Rhizome thick, oblique. Stipes tufted, 15—25 c.M. long, the base black, clothed with black, subulate scales, the rest slender, slightly setose with scattered, black bristles. Fronds 20—45 c.M. long, 8—25 c.M. broad, ovate-oblong, acuminate, pinnate, gemmiferous and sometimes viviparous. Pinnae \pm 6—16 on each side below the pinnatifid apex, more or less distinctly stalked, \pm horizontal, acute, to \pm 2 c.M. broad, crenate at the base on the lower side, the upper base truncate, auriculate, the edge coarsely crenatolobate, the lobules short, to 3 m.M. long, 3—5 m.M. broad, decumbent, subacute, tooth-like; lowest pinnae shorter, deflexed. Texture herbaceous; surfaces and rachis naked; colour blackish-green above, paler beneath; veins pinnate in the lobules. Sori on the anterior veinlets, often double, to 7 m.M. long; indusium thin, decoloured, diaphanous. — [Texture subcoriaceous; pinnae cut $\frac{1}{3}$ -way down to the costa. (Bk.)].

Luzon, Mindanao; Polynesia.

(35a) **D. Whitfordi**, *Copel.*, Philipp. Journ., I, Suppl., 150.

Rhizome short, horizontally creeping. Stipes approximate, 10—20 c.M. long, the lower part asperous, the base black, clothed with scattered, brown, caducous scales, the upper part smooth, naked, greenish or stramineous. Fronds 20—35 c.M. long, 6—15 c.M. broad, 2-pinnatifid, the rachis naked. Pinnae 10—20-jugate, alternate, the central ones the largest, \pm spreading, broadly lanceolate, acute, cut down nearly to costa; lower pinnae rather distant, somewhat shorter, deflexed. Lobes oblong, 3—4 m.M. broad, serrate towards the apex. Texture chartaceous; surfaces naked; veins rarely forked. Sori 4—5 on each side of the costulae, occupying the lower part of the veins.

Luzon.

* (35b) **D. speciosum**, *Bl.*, Copel., Polypod. Philipp., 74; Bedd., Ferns Br.I., II, tab. CCXC; *Asplenium speciosum*, *Mett.*, Rac., Flor. Btz., I, 224; (? *D. sorzogonense*, *Pr.*, C. Chr., Ind. Fil.).

Rhizome or caudex erect or suberect, clothed with lanceolate scales. Stipes tufted, 50—50 c.M. long, firm, erect, stramineous or brownish, the base covered with narrow, lanceolate scales. Fronds 50—150 c.M. long, 20—35 c.M. broad, with 10—20 pairs of pinnae below the pinnatifid apex, the lower ones often stalked, the stalks 1—4 m.M. long. Largest pinnae to

17 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, straight or subfalcate, horizontal or nearly erecto-patent, the apex acuminate, the edges pinnatifid to deeply pinnatipartite, the lowest segments generally the longest, \pm spreading and all blunt, entire, crenate or toothed. Texture subcoriaceous; rachis naked or \pm scaly; veins pinnate in the lobes. Sori reaching nearly or quite to the edge.

Malaya.

(36a) **D. cyatheifolium**, *Pr.*, Copel., Polypod. Philipp., 75; *Asplenium cyatheifolium*, *Rich.* (not *Bory*), Hk. Bk., Syn. Fil., 238.

Caudex erect, subarborescent. Stipes erect, firm, dark-brown, nearly naked. Fronds 45—60 c.M. long, 30—45 c.M. broad. Lower pinnae to 20—25 c.M. long, 7—8 c.M. broad, cut down to the rachis in the lower $\frac{2}{3}$ into numerous pinnulae on each side, the upper ones inciso-crenate, the lower ones $3\frac{1}{2}$ —4 c.M. long, \pm 1— $1\frac{1}{2}$ c.M. broad, cut down $\frac{1}{3}$ of the distance to the costa into oblong, falcate, sharply toothed lobes. Texture herbaceous; rachis dark-brown, naked like the stipe; main veins pinnate with 3—4 veins on each side in the lower lobes. Sori falling short of the edge.

Philippines, New Guinea; Polynesia.

(36b) **D. davaoense**, *Copel.*, Philipp. Journ., I, Suppl., 151.

Stipes tufted, \pm 70 c.M. long, conspicuously sulcate, nearly naked, the lower part asperulous. Fronds ovate, 1— $1\frac{1}{2}$ M. long, the lower part 2-pinnate, the higher part simply pinnate, the terminal part pinnatifid, the apex acuminate, serrate. Pinnae few, spreading, \pm 50 c.M. long, 20—25 c.M. broad, the lower ones hardly reduced. Pinnulae short-stalked, $2\frac{1}{2}$ —3 c.M. broad, the base truncate, the edge cut \pm $\frac{1}{3}$ -way down to the costa, the apex acuminate. Lobes falcate, truncate, obscurely denticulate. Texture membranaceous; surfaces naked; veins \pm 6 on each side of the main veins, arcuate, remote, simple. Sori linear. — Near *D. cyatheifolium*, *Pr.* & *latifolium*, *Moore*.

Mindanao.

* (37a) **D. latifolium**, *Moore* (not *Don*), Christ, Farnkr. d. Erde, 220; *D. maximum*, *C. Chr.*, Ind. Fil.; *D. dilatatum*, *Bl.*, Bedd., Ferns S.I., tab. CLXII (not quite correct); *Asplenium latifolium*, *Don*, Rac., Flor. Btz., I, 227.

Caudex erect, short or subarborescent, clothed with linear-lanceolate, dark-brown scales. Stipes strong, erect, 30—90 c.M. long, livid, smooth, scaly at the base. Fronds subdeltoid, 90—200 c.M. long, with \pm 12 pinnae on each side, the largest to \pm 50 c.M. or more long. Pinnulae numerous, the

largest to 18 c.M. long, to ± 3 c.M. broad at the base, linear-lanceolate, the point acute or acuminate, the edge lobed, and crenate or toothed, the base truncate or rounded. Texture subcoriaceous; rachis naked; main veins pinnate in the lobes with 5—8 veins on each side. Sori linear, the lowest much the longest, not reaching the edge. — A variable plant.

Trop. Asia, Polynesia & Australia, China.

(57b) **D. dolichosorum**, *Copel.*, Philipp. Journ., I, Suppl., 151.

Rhizome strong, clothed with long, black-brown scales. Stipes close, ± 35 c.M. long, clothed with fuscous, linear, contorted scales, the upper part subglabrescent, at length sparingly muricate. Fronds ovate, $1-1\frac{1}{2}$ M. long, 2-pinnate, the apex pinnatifid, the main rachis glabrescent, sometimes asperulous. Pinnae remote, the largest ± 45 c.M. long, ± 20 c.M. broad, the lowest shorter. Pinnulae short-stalked, ± 2 c.M. broad, subfalcate, acuminate, the base truncate, the edges lobed $\pm \frac{1}{3}$ -way down in the lower part, the apex serrate. Lobes truncate, denticulate. Texture papyraceous; surfaces naked; veins ± 6 on each side of the main veins, the lower ones curved. Sori linear.

Mindanao.

(58) **D. manilense**, *C. Chr.*, Ind. Fil.; *Asplenium serrulatum*, *Pr.*, Rel. Haenk., I, 46; *A. manilense*, *Spr.*, Syst., IV, 89.

Fronds ovate, ± 45 c.M. long, 2-pinnate; main rachis tetragonal, clothed with small, ovate, fuscous, membranaceous scales. Pinnae distant, alternate, petiolate, pinnate, the apex pinnatifid, acuminate; central pinnae $\pm 12\frac{1}{2}$ c.M. long, the rachises scaly like the main one. Pinnulae spreading, sessile, lanceolate-oblong, $2\frac{1}{2}-4$ c.M. long, $1-1\frac{1}{4}$ c.M. broad, acuminate, the base truncate on both sides, the highest ones entire, the central ones serrate with the serratures 2-denticulate, obtuse, the lowest largest, serrulate. Texture membranaceous; surfaces naked; veins simply branched. Sori long, narrow, double; indusium diaphanous.

Philippines.

(59) **D. vestitum**, *Pr.*, Copel., Polypod. Philipp., 75; *Asplenium vestitum*, *Hk.* (not *Pr.*), *Hk. Bk.*, Syn. Fil., 239; *Hk.*, 2nd Cent. of Ferns, tab. XLVI.

Rhizome erect. Stipes stout, erect, brownish, tomentose, sometimes muricate. Fronds 60—90 c.M. long, 20—45 c.M. broad, with numerous pinnae on each side, the lower ones 10—25 c.M. long, to 10 c.M. broad, with numerous, distinct, subsessile pinnulae, which are $\pm 2\frac{1}{2}$ c.M. broad, blunt, bluntly lobed to a depth of 3—6 m.M., the base narrowed suddenly or even cordate.

Texture thinly herbaceous; rachis chaffy like the stipe; main veins pinnate in the lobes; veins 4—5 on each side. Sori not reaching the margin:

Philippines.

(40) **D. deltoideum**, *Pr.*, Copel., Polypod. Philipp., 76; *Asplenium deltoideum*, *Pr.*, Hk. Bk., Syn. Fil., 257; *Pr.*, Rel. Haenk., I, tab. VII, fig. 2.

Rhizome erect, black-scaly. Stipes 15—20 c.M. or more long, black and black-scaly at the base, naked and green above. Fronds 30—40 c.M. long, 20—25 c.M. broad, deltoid, with a few distant pinnae on each side. Lowest pinnae largest, ± 6 c.M. broad, bearing numerous sessile pinnulae. Pinnulae obtuse, $\pm 1-1\frac{1}{2}$ c.M. broad, cut $\pm \frac{2}{3}$ of the way down to the costa into blunt lobes. Texture herbaceous; surfaces naked; rachises naked, black or green; main veins pinnate in the lower lobes. Sori reaching the margin.

Celebes, Philippines.

(41a) **D. Meyenianum**, *Pr.*, Copel., Polypod. Philipp., 76; *Asplenium Meyenianum*, *Mett.*, Hk. Bk., Syn. Fil., 258.

Fronds probably ± 1 M. long, ovate-triangular. Largest pinnae 30—40 c.M. long, 20—25 c.M. broad, stalked, alternate, oblong-lanceolate, acuminate. Lower pinnulae $2\frac{1}{2}$ —5 c.M. broad, stalked or sessile, linear-oblong, acuminate, the base subcordate, the edge pinnatifid $\pm \frac{3}{4}$ -way down to the costa; higher pinnulae decurrent with a narrow wing or confluent at the base. Lobes subopposite, ovate-oblong, subfalcate, obtuse, crenate or serrulate, to $\pm \frac{3}{4}$ c.M. broad. Texture thinly coriaceous; surfaces glabrous; rachis brown, channelled above; veins pinnate, the higher forked. Sori copious, reaching from the costulae nearly to the edge.

Philippines.

(41b) **D. atratum**, *Christ*, Philipp. Journ., II, 163.

Stipes 25 c.M. or more long, sulcate, ebeneous or black-purple, the base thick, densely clothed on the inner side with long, subulate, black scales, the rest nearly naked or minutely furfuraceous. Fronds broadly deltoid, 5-pinnatifid, acute, 60 c.M. or more long, the base 45 c.M. or more broad. Pinnae distant, ± 20 on each side, on stalks $\pm 2-3$ c.M. long, the lowest not reduced, ± 10 c.M. broad, horizontally spreading, long-acuminate, narrowed at the base, the higher sessile. Pinnulae ± 20 below the incised apex, more than 1 c.M. apart, horizontally spreading, the lowest stalked, broadly lanceolate, acuminate, cut down $\frac{1}{2}$ -way to the costa, but deeply pinnatifid at

the base. Lobes ± 15 on each side, ± 5 m.M. long, porrect, triangular-falcate, acute, the edges serrulate, often reflexed. Texture very rigidly coriaceous, nearly woody; surfaces naked; colour black-brown; veins $\pm 4-7$ -jugate, oblique, stout, simple. Sori reaching from the costulae nearly to the edge, very oblique, convex, black-brown, linear, simple; indusium coriaceous, brown, persistent.

Palawan.

(42) **D. amplissimum**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 227; *Asplenium amplissimum*, *Bk.*, Ann. of Bot., V, 510.

Fronds ample, 3-pinnatifid. Pinnae oblong-lanceolate, 50—45 c.M. long. Pinnulae lanceolate, lower short-stalked, 7—10 c.M. long, $2\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, cut down to a narrow wing into oblong-lanceolate, crenate tertiary segments. Texture moderately firm; surfaces naked; rachises brown-stramineous, naked or sparsely fibrillose; veins 6—7-jugate, mostly forked. Sori oblong, reaching from the midrib $\frac{1}{2}$ -way to the edge.

Amboyna.

(43a) **D. oligosorum**, *Copel.*, Philipp. Journ., II, 128.

Rhizome erect. Stipes close, ± 40 c.M. long, the base castaneous, swollen, aculeate, slightly scaly, the rest smooth and yellowish-brown like the rachises. Fronds 50—60 c.M. long, 40—50 c.M. broad, acuminate, 3-pinnatifid. Lateral pinnae opposite, ± 3 -jugate, rather equal, ± 10 c.M. apart, stalked, horizontal, acuminate. Pinnulae subsessile, ± 7 c.M. long, ± 2 c.M. broad, caudato-acuminate, the base truncate, the edge cut down nearly to the costa into-oblong, obtuse, subfalcate, entire or serrulate segments 4—5 m.M. broad. Texture thinly coriaceous; surfaces naked, the upper black-green, the lower paler; veins distant, often simple. Sori minute, nearer the edge than the costulae.

Mindoro.

* (43b) **D. polypodioides**, *Bl.*, Christ, Farnkr. d. Erde, 219; *Bedd.*, Ferns Br.I., II, tab. CCXCIII; *Asplenium polypodioides*, *Mett.*, Rac., Flor. Btz., I, 227, p. p.

Caudex erect, strong, subarborescent, densely clothed at the crown with long, brown scales. Stipes densely tufted, stout, 50—100 c.M. or more long. Fronds 90—120 c.M. long, $\pm 50-60$ c.M. broad, 2-pinnate, with $\pm 8-9$ distant, stalked, divided pinnae on each side below the simple higher ones.

Pinnulae 10—15 on each side, approximate, sessile, sharply acuminate, 5—10 c.M. long, \pm 1—2 c.M. broad at the suddenly narrowed base, the edge cut $\frac{1}{2}$ — $\frac{3}{4}$ -way down to the costa into oblong, entire, crenate or serrate, bluntish, approximate or close lobes. Texture herbaceous; surfaces naked or nearly so; rachis green, \pm naked; main veins pinnate with 5—7 simple or forked veins on each side. Sori copious, the lower commonly double, commencing at the main vein (costula), generally falling short of the edge.

Malaya; Trop. Australia, North. India.

(44) **D. ebeneum**, **J. Sm.**, Copel., Polypod. Philipp., 76; (? *D. asperum*, *Bl.*, C. Chr., Ind. Fil.).

Differs from *D. polypodioides*, *Bl.* by its slender, naked, ebeneous rachises, and the ultimate lobes, with but 4 distinct veins in each.

Philippines.

* (45a) **D. asperum**, **Bl.**, Christ, Farnkr. d. Erde, 219; *D. polypodioides*, *Bedd.*, Ferns S.I., tab. CLXIII; *Asplenium polypodioides*, *Mett.*, Rac., Flor. Btz., I, 227, p. p.; *A. asperum*, *Hk.* (not *Mett.*), *Hk.*, Spec. Fil., III, 258.

Near *D. polypodioides*, *Bl.*, but the stipes, main and partial rachises asperous and more or less scaly. Pinnulae and lobes \pm spreading, the lobes rounded or bluntish, not truncate. Texture rather coriaceous; veins 8—12 on each side, commonly simple. Sori short, commonly confined to the lower half of the veins.

Malacca, Java, Mindoro; North. India.

(45b) **D. fructuosum**, **Copel.**, Philipp. Journ., I, Suppl., 150, tab. VIII.

Near *D. polypodioides*, *Bl.* Rhizome erect. Stipes \pm 1 M. long, deeply sulcate, the lower part clothed with long, linear, black-brown scales leaving the stipes spinulose when they fall, the upper part glabrescent. Fronds $1\frac{1}{2}$ —2 M. long, \pm 1 M. broad, the rachis muricate. Pinnae \pm 15 c.M. broad, nearly spreading, the lower ones hardly reduced, the rachis smooth. Pinnulae subsessile, acuminate, \pm spreading, \pm $1\frac{1}{2}$ c.M. broad near the base, the edge hardly cut $\frac{1}{2}$ -way down to the costa into entire, falcate, truncate lobes. Texture herbaceous; surfaces naked; veins \pm 6 on each side of the main veins. Sori often filling up the whole under surface.

Mindanao.

(46) **D. prionophyllum**, **Kze.**, Bot. Zeit., 1848, 194.

Fronds large. Pinnae petiolate, falcate, oblong, acuminate, the largest 30—45 c.M. long. Pinnulae short-stalked, spreading, oblong, acuminate, the base cordato-truncate, the edge pinnatifid. Lobes falcate, ovate, bluntish, irregularly inciso-serrulate, the margin revolute. Texture subcoriaceous; surfaces glabrescent; rachis partially aculeolate; costae and costulae paleaceo-asperous, ± flexuose. Sori numerous, at length confluent.

Java.

(47) **D. ceramicum**, *C. Chr.*, Ind. Fil.; *Asplenium ceramicum*, *Miq.*, Ann. Mus. Bot. L. B., IV, 168.

Main rachis covered with few, deciduous scales; secondary rachises dilated at the base, blackish. Fronds oblong, membranaceous, large, 3-pinnate or 4-pinnatifid. Pinnae and pinnulae alternate. Pinnae short-stalked, lanceolate, acuminate, the lower ones 35—40 c.M. long. Pinnulae 10—15 c.M. long, short-stalked, long-acuminate. Tertiary divisions short-stalked, slightly falcate, the apex obtuse, serrulate, the lower part crenato-pinnatifid, the base broadly cuneate. Veins 2-fid or the lower ones forked or pinnate, the higher branches sori-ferous. Sori linear, medial on the veins; indusium membranaceous, entire, brownish.

Ceram.

(48) **D. Huttoni**, *C. Chr.*, Ind. Fil.; *Asplenium Huttoni*, *Bk.*, Ann. of Bot., V, 311.

Fronds large, deltoid, decomposed. Pinnae oblong-lanceolate, 50 c.M. long or longer. Pinnulae lanceolate. Tertiary segments obtuse, 4—5 m.M. broad, deeply pinnatifid. Lobes oblong, obtuse. Texture moderately firm; surfaces naked; main veins (costulae) pinnate in the lobes; veins erecto-patent, simple. Sori medial, short; indusium firm, naked, persistent.

Malaya.

§ 2. ANISOGONIUM.

a. Fronds simple, or with 1—6 lateral pinnae on each side.

b. Simple frond or terminal pinna cordate at the base.

c. Fronds simple, or with 1—2 pairs of lateral pinnae.

(49) **D. cordifolium**.

c.c. Fronds pinnate; pinnae 3—5-jugate.

(50) **D. Forbesii**.

b.b. Simple frond or terminal pinna cuneate or rounded at the base.

*. Stipes scaly throughout. Fronds with 0—3 lateral pinnae on each side. (51) *D. alismifolium*.

.. Stipes scaly at the base. Fronds with 1—6 lateral pinnae on each side. (52) *D. fraxinifolium*.

a.a. Fronds pinnate with more than 6 lateral pinnae on each side.

S. Pinnae entire.

*. Pinnae membranaceous, narrowly cuneate at the base. (53) *D. angustatum*.

.. Pinnae subcoriaceous, rounded at the base. (54) *D. pariens*.

S.S. Pinnae toothed, wavy or slightly lobed. (55) *D. proliferum*.

a.a.a. Fronds at least 2-pinnate (*).

1. Largest pinnae to 25 c.M. long.

2. Pinnulae rounded at the base. (56) *D. Smithianum*.

2.2. Pinnulae truncate at the base. (57) *D. spinulosum*.

1.1. Largest pinnae \pm 30 c.M. long or longer in the fully developed fronds. (58) *D. esculentum*.

* (49) **D. cordifolium**, *Bl.*, Christ. Farnkr. d. Erde, 221; Hk., Icon. Pl., X, tab. CMXXXVI; *Asplenium cordifolium*, *Mett.*, Hk. Bk., Syn. Fil., 245; *A. integrifolium*, *Mett.*, Rac., Flor. Btz., I, 230, p.p.; *Callipteris cordifolia*, *Copel.* (not *J. Sm.*), Polypod. Philipp., 70; *Anisogonium cordifolium*, *Bedd.*, Ferns Br. I., II, tab. CCCXXXI.

Rhizome erect, densely clothed with brown, lanceolate scales. Stipes clustered, \pm 20 c.M. long, firm, erect, scaly below. Fronds \pm 25—35 c.M. long, \pm 10 c.M. broad, undivided, acuminate, the edge entire or subrepand, the base cordate. Below this there are sometimes 1—2 pairs of lateral pinnae, which are \pm erecto-patent, similar to the simple frond but smaller and narrower, the lowest to \pm 15 c.M. long and \pm 5—7 c.M. broad. Texture coriaceous; surfaces naked; veins in groups of 3—4, anastomosing in the outer half of the pagina. Sori reaching from the midrib \pm to the edge.

Africa to Polynesia.

* (50) **D. Forbesii**, *C. Chr.*, Ind. Fil.; *Asplenium Forbesii*, *Bk.*, Ann. of Bot., V, 312.

(*) See also **D. proliferum**, *Thouars*, var. **robustum**.

Stipes naked, 12—15 c.M. long. Fronds deltoid, simply pinnate, \pm 30 c.M. long. Pinnae \pm 3—5-jugate, lanceolate, acuminate, entire, sessile, \pm 15 c.M. long, 5—6 $\frac{1}{2}$ c.M. broad, cordate at the base. Texture moderately firm; surfaces naked; veins 3—5-jugate, anastomosing towards the margin. Sori linear.

Java.

(51) **D. alismifolium**, *v. A. v. R.*; *D. Cumingii*, *C. Chr.*, Ind. Fil.; *Asplenium alismaefolium*, *Hk.*, Hk. Bk., Syn. Fil., 243; *Callipteris alismaefolia*, *J. Sm.*, Copel., Polypod. Philipp., 70; *Anisogonium alismaefolium*, *Christ*, Fil. Saras., III, 253; *Oxygonium alismaefolium*, *J. Sm.*, (oldest name).

Stipes 5—15 c.M. long, firm, erect, scaly throughout. Fronds varying in shape from simple, oblong-lanceolate, 15—25 c.M. long, 5—8 c.M. broad, entire, acuminate, to ternate or pinnate with a large terminal and to 3 pairs of lateral pinnae, each like the entire frond of the simple form. Texture coriaceous; veins \pm 5 to a group in the inner half, anastomosing copiously, with hexagonal areolae towards the edge.

Celebes, Philippines.

* (52) **D. fraxinifolium**, *Pr.*, Rel. Haenk., I, 49; *D. elegans*, *Hk.* Icon. Pl., X, tab. CMXXXIX—CMXL; *D. lineolatum*, *Bl.* Enum., 191; *Asplenium lineolatum*, *Mett.*, Hk. Bk., Syn. Fil., 243; *Callipteris elegans*, *J. Sm.*, Copel., Polypod. Philipp., 70; *Anisogonium lineolatum*, *Bedd.*, Ferns Br.I., II, tab. CCCXXX; *A. elegans*, *Pr.*, Epim., 95.

Rhizome erect, the scales brown, lanceolate. Stipes erect, firm, grey, 15—25 c.M. long, scaly below. Fronds occasionally simple, usually with a terminal pinna and 3—6 pairs of lateral ones, which are 15—30 c.M. long, 4—5 c.M. or more broad, entire, or serrate towards the often suddenly acuminate apex, the base subcuneate or rounded. Texture coriaceous; rachis naked; veins 4—8 to a group, uniting slightly towards the margin. — A small form with bluish-green, succulent fronds and 1—3 lateral pinnae was reported from Celebes (*Christ*, Ann. Btz.; XV, 122).

Malaya; Japan.

(53) **D. angustatum**, *Amann*, Nat. Tijdschr. Ned. Ind., XXIII, 406.

Stipes \pm terete, smooth, reddish-brown, glossy. Fronds \pm ovate, 30—45 c.M. long (incl. the stipe), pinnate. Pinnae alternate, the higher ones sessile, decurrent, the lower short-stalked, lanceolate or oblong, acuminate, the base

narrowly cuneate, the edge entire, \pm revolute. Texture membranaceous. Sori reaching from the midrib to the edge, parallel, dichotomous as in *D. cordifolium*, *Bl.*

Banca.

(54) ***D. pariens***, *C. Chr.*, *Ind. Fil.*; *Callipteris pariens*, *Copel.*, *Polypod. Philipp.*, 70.

Rhizome strong, erect, black, clothed with lanceolate, dark-brown scales. Stipes canaliculate, \pm 55 c.M. long, blackish and slightly scaly at the base, above green and clothed with minute fibrils like the rachis. Fronds \pm 50 c.M. long, \pm 20 c.M. broad, pinnate. Terminal pinna ovate, acuminate, entire, rounded at the base; lateral pinnae \pm 11 on each side, similar to the terminal one, those of the lower half of the frond larger, \pm 10 c.M. long, \pm 4 c.M. broad, short-stalked. Texture subcoriaceous; surfaces naked; veins 1—2 \times forked, anastomosing, indistinct in the marginal $\frac{1}{3}$. Sori linear, straight, occupying the lower $\frac{2}{3}$ of the veins.

Philippines.

* (55) ***D. proliferum***, *Thouars*, *Flor. Trist. d'Ac.*, 35; *D. decussatum*, *Christ*, *Farnkr. d. Erde*, 221; *Callipteris prolifera*, *Bory*, *Copel.*, *Polypod. Philipp.*, 70; *Asplenium decussatum*, *Sw.*, *Rac.*, *Flor. Btz.*, I, 251; *A. proliferum*, *Lam.*, (oldest name); — var. *robustum*; *Digrammaria robusta*, *Fée*, *Gen.*, 218; — var. *accedens*; *Dipl. accedens*, *Bl.*, *Enum.*, 192; *A. spinulosum*, *Mett.*, *Farnng.*, VI, *Aspl.*, 172 (not *Ann. Mus. Bot. L. B.*, II, 257).

Rhizome erect (or creeping after *Christ*), the scales long, lanceolate. Stipes 30—60 c.M. long, strong, erect, scaly and slightly muricate at the base. Fronds large, 60—250 c.M. long, the apex pinnatifid, and below this many to numerous sessile pinnae. Pinnae 8—30 c.M. long, 2—6 c.M. broad, the lower ones often stalked, the base truncate or subtruncate, the apex acute or acuminate, the edge toothed, wavy or slightly lobed, the lobes blunt, \pm toothed. Texture herbaceous to subcoriaceous; main veins uniting \pm $\frac{1}{3}$ or less of the distance from the costa to the margin. Sori very regularly running along the veins, reaching \pm the edge, often copiously double.

Var. robustum: Pinnae 30—45 c.M. long, deeply lobed in the upper part, in the lower cut down to the costa.

Var. accedens. Stipes muricate or spinulose throughout.

Africa to Polynesia and Australia.

(56) **D. Smithianum**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 228; *Asplenium Smithianum*, *Bk.*, Hk. Bk., Syn. Fil., 245; *Anisogonium Smithianum*, *Bedd.*, Handb. Ind. Ferns, 192; *Callipteris Smithiana*, *Bedd.*, Ferns Br.I., II, tab. CCCXXXII.

Rhizome oblique, decumbent. Stipes thick, \pm 50 c.M. long, slightly scaly below, furfuraceous throughout and rather densely muricate. Fronds 45—60 c.M. long, by nearly as broad, with 4—5 pinnated pinnae and 4—5 that are only pinnatifid below the point on each side, the lowest 7—8 c.M. apart, \pm 20—25 c.M. long, \pm 10 c.M. broad. Pinnulae sessile, 5—8 c.M. long, $1\frac{1}{4}$ —2 c.M. broad, the edge subentire, the point slightly toothed, acuminate, the base rounded on both sides. Texture subcoriaceous; both surfaces and rachis naked, the latter muricate in the lower part; veins in groups of 2—3 on a side, the groups usually joining half-way across to the edge.

New Guinea, Batjan, Celebes, Luzon; Ceylon.

(57) **D. spinulosum**, *Bl.*, Enum., 193; *Asplenium spinulosum*, *Mett.*, Ann. Mus. Bot. L. B., II, 257 (not *Farng.*, VI, *Aspl.*, 172); *A. heteropteron*, *Mett.*, *Farng.* VI, *Aspl.*, 174.

Stipes \pm 15 c.M. or more long, scaly and tomentose-chaffy, the scales membranaceous, blackish-brown, lanceolate, acuminate, thinly ciliate, deciduous. Fronds ovate, the base slightly narrowed. Pinnae stalked, \pm 20—25 c.M. long, ovate or lanceolate-oblong, the lower ones distant, pinnate, the higher pinnatifid. Pinnulae approximate, the lower ones \pm 7—8 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, subsessile, oblong-acuminate or slightly obtuse, the edge truncato-crenate, the base truncate; higher pinnulae adnate to the rachis, sometimes decurrent, the highest confluent. Lobes of the highest pinnae oblong, \pm bluntish. Texture membranaceous, rigid, opaque; lowest veins anastomosing; rachis deciduously scaly. Sori 2—5 on each side, the lowest acroscopical double; indusium narrow, membranaceous. — (Stipe and rachis aculeate after *Blume*, l. c.).

Moluccas, New Guinea.

* (58) **D. esculentum**, *Sw.*, *Christ*, *Farnkr. d. Erde*, 222; *Asplenium esculentum*, *Pr.*, *Rac.*, *Flor. Btz.*, I, 251; *Callipteris esculenta*, *J. Sm.*, *Copel.*, *Polypod. Philipp.*, 71; *Bedd.*, *Ferns*, S. I., tab. CLXIV; *Anisogonium esculentum*, *Pr.*, *Bedd.*, *Handb. Ind. Ferns*, 192; *Gymnogramme edulis*, *Ces.*, *Fel. di Bor.*, 28; *Hemionitis esculenta*, *Retz.* (oldest name).

Caudex erect, to 15 c.M. high, densely clothed with narrow, lanceolate scales. Stipes 15—60 c.M. long, strong, erect, tufted. Fronds 60—180 c.M. long or longer, 2-pinnate, deltoid or broadly lanceolate. Lower pinnae 30—45 c.M. long, 15—20 c.M. broad. Pinnulae 7—15 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, short-stalked, the point acuminate, the edge crenate to shallowly lobed, the lobes entire or serrate, the base truncate and often auricled on both sides. Texture subcoriaceous; rachis often pubescent; main veins distinct, copiously pinnate with 6—10 veins on each side in each lobe, which are beginning to unite a short distance from the midrib, the higher ones often free. Sori copious, occupying nearly all the veins, reaching \pm the margin. — Lowest pinnae sometimes simple.

Trop. Asia & Polynesia.

54. ATHYRIUM, *Roth.*

Characters of Diplazium, except that the sori and indusia are oblong, often strongly recurved, sometimes horseshoe-shaped or \pm reniform and then sometimes hardly different from those of Lastraea, the lower of the same group rarely, the higher never double. Fronds at least simply pinnate; veins free. Hk. Bk., Syn. Fil., tab. IV, fig. 38, e; Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 120.

a. Fronds simply pinnate. Pinnae entire or slightly lobed, at best the lower ones pinnatifid.

§. Rachis winged throughout. (1a) *A. nanum.*

§.§. Rachis not winged, or only winged in the upper half.

‡. Fronds herbaceous, \pm $7\frac{1}{2}$ —15 c.M. long.

*. Fronds \pm 1—2 c.M. broad. (1b) *A. pusillum.*

.. Fronds \pm 2— $2\frac{1}{2}$ c.M. broad. (1c) *A. Toppingianum.*

‡.‡. Fronds coriaceous, \pm 15—50 c.M. long. (2) *A. gedeanum.*

a.a. Fronds more copiously 2-pinnatifid to 3-pinnate.

b. Pinnae pinnatifid, sometimes pinnate at the base.

‡. Lower pinnae not reduced.

*. Fronds small, to \pm 9 c.M. broad.

×. Indusium reniform. (3a) *A. anisopterum.*

(3b) *A. benguetense.*

- ×.×. *Indusium not reniform.* (3c) *A. Copelandi*.
 .. *Fully developed fronds larger, more than 10 c.M. broad.*
 (4) *A. macrocarpum*.
 †.†. *Lower pinnae reduced.* (5) *A. acrostichoides*.
b.b. *At least the lower pinnae pinnate to 2-pinnate.* (1)
c. *Fronde small, under 15 c.M. long.* (2)
 §. *Rhizome creeping, often intertwined.* (6) *A. hyalostegium*.
 §.§. *Rhizome short, erect.*
 *. *Fronde ± as broad as long.* (7) *A. Loheri*.
 .. *Fronde ± 2 × as long as broad.* (8) *A. Sarasinorum*.
c.c. *Fronde larger, more than 15 c.M. long when fully developed.*
d. *Teeth of the segments finely aristate, retroflexed.*
 (9) *A. aristulatum*.
d.d. *Teeth whether or not aristate, not retroflexed.*
e. *Lower or costal sori often horseshoe-shaped or hamato-reniform.*
 ×. *Texture herbaceous.* (10) *A. solenopteris*.
 ×.×. *Texture coriaceous.* (11) *A. drepanopterum*.
e.e. *Largest sori hamato-recurvate at the apex only.*
 (12) *A. nitidulum*.
e.e.e. *Sori straight or slightly curved, the lower sometimes double.*
f. *Midribs. of the leaflets beset with distant, subulate, spine-like appendices on the upper side. Fronde ± 20—60 c.M. long.*
 *. *Fronde to 30 c.M. long, to 12 c.M. broad. Pinnae spreading.* (15) *A. gymnogrammoides*.
 .. *Fronde 30—60 c.M. long, 12 c.M. broad or broader. Pinnae erecto-patent.* (14) *A. nigripes*.
f.f. *Appendices wanting or indistinct. Fronde 60—200 c.M. long in the fully developed plants.*
g. *Tertiary segments elliptico-rhomboidal, the base narrowed or ± decurrent.* (15) *A. umbrosum*.
g.g. *Tertiary segments linear-oblong, the base not narrowed, not decurrent.* (16) *A. silvaticum*.

(1) See also *A. macrocarpum*, *Bedd.*

(2) See also *A. aristulatum*, *Copel.*

(1a) **A. nanum**, *Christ*, Philipp. Journ., II, 161.

Rhizome oblique, short or nearly creeping. Stipes few, tufted, slender, fuscous, $\pm \frac{1}{2}$ —1 c.M. long, pubescent like the rachis and clothed with thin, setulose, reddish-brown, spreading scales. Fronds lanceolate, ± 6 c.M. long, 7 m.M. broad, pinnate below the much prolonged, pinnatifid apex, the lobes erecto-patent. Pinnae in ± 5 —6 pairs, erecto-patent, subremote, $2\frac{1}{2}$ —5 m.M. long, the higher equilateral at the base, entire, oblong, the central inequilateral, auricled on the upper side, subrenate, the lowest inequilateral, crenatolobate. Texture flaccido-herbaceous; rachis winged; surfaces puberulous; colour bright-green; veins forked, hardly pinnate. Sori 5—6 to a segment, small, hardly 1 m.M. long, ochraceous; indusium lanceolate, rarely aspidioideo-reniform.

Mindanao.

* (1b) **A. pusillum**, *v. A. v. R.*; *Asplenium pusillum*, *Bl.*, Rac., Flor. Btz., I, 221; C. Chr., Ind. Fil.; *A. confluens*, *Kze*, Bot. Zeit., 1848, 174; Hk. Bk., Syn. Fil., 490.

Rhizome creeping, flexuose, the scales brown, lanceolate, acuminate, small. Stipes seriate, slender, ± 2 —6 c.M. long, deciduously scaly like the rachis, the scales minute, crisped, setiform. Fronds linear-lanceolate, $\pm 7\frac{1}{2}$ — $12\frac{1}{2}$ c.M. long, ± 1 —2 c.M. broad, with ± 5 —6 pairs of pinnae below the terminal one. Terminal pinna long, deeply pinnatifid at the base, narrowed gradually towards the toothed or serrate, acuminate point; lateral pinnae erecto-patent, the highest longest, lanceolate-oblong, bluntish, sessile; lowest pinnae smallest, nearly stalked, remote, oblong, ± 5 —6 m.M. long, ± 5 m.M. broad, blunt, the edge \pm repando-denticulate, the base obliquely cuneate, slightly auricled on the upper side. Texture papyraceous; rachis narrowly winged above; surfaces sparingly covered with few, minute, inconspicuous scales on the costae; main veins oblique, simple, the lower forked. Sori few, minute, the lower anterior of the pinnae often elongate-hippocrepiform, but without sinus between the branches. — The scales of the rhizome being finely cancellate and the cells coloured this plant is decidedly *no* *Asplenium*. (*) — Forbes plant (no. 2894) determined by Raciborski as *Asplenium pusillum*, *Bl.* is a young form of *Diplazium grammitoides*, *Pr.*

Java.

* (1c) **A. Toppingianum**, *Copel.*, Philipp. Journ., II, 127; *As-*

(*) Zollinger's No. 2925 is in the Buitenzorg Herbarium

plenium toppingianum, Copel., Polypod. Philipp., 80; — var. *mixtum* (*mistum*), Copel., l. c.; *Asp. Toppingii*, Copel., MS. in Herb. Bog.

Rhizome creeping, chaffy, the scales lanceolate, pale-brown. Stipes 5—10 c. M. long, scattered, the base brown and chaffy, upward becoming green and naked like the rachis. Fronds 10—15 c.M. long, 2—2½ c.M. broad, the upper half deeply pinnatifid with several erecto-patent, oblong, blunt lobes, the apex toothed to crenate, acuminate, the lower half simply pinnate with ± 6 pinnae on each side. Pinnae oblong, blunt, serrato-crenate, auricled at the base on the upper side. Texture herbaceous, rather firm; surfaces naked; main rachis narrowly winged towards the upper half of the frond. Sori short, straight, curved, or some of them horseshoe-shaped, the lower sometimes double.

Var. mixtum: Larger; teeth of the pinnae more rounded; the lowest pair of pinnae more than 2 × the dimensions, deeply incised.

Luzon.

* (2) **A. gedeanum**, *Christ*, Journ. de Bot., XIX, 68; *Asplenium gedeanum*, *Rac.*, Flor. Btz., I, 223.

Rhizome erect or suberect, densely clothed with brown, lanceolate scales. Stipes firm, erect, 10—50 c.M. long, stramineous, glossy, scaly at the base. Fronds oblong- or lanceolate-deltoid, 15—50 c.M. long, 5—15 c.M. broad, the apex acuminate, toothed to deeply pinnatifid. Lateral pinnae 10—30 on each side, linear to lanceolate, the higher ones sessile, the lower ones distant, subsessile, ± deflexed, often shorter than the next in order, the longest ones spreading, 5—10 m.M. broad, the apex acute or blunt, sometimes acuminate, the upper base truncate or broadly cuneate, often auricled, the lower base cuneate, often slightly auricled, the edge slightly lobed, the lobes sharply toothed. Texture coriaceous; main veins pinnate in the lobes. Sori short, mostly close to the midribs, rarely diplazioid at the apex.

Java; Annam.

(3a) **A. anisopterum** (*anisopteron*), *Christ*, Bull. Herb. Boiss., VI, 962; Philipp. Journ., II, 160; Bull. Ac. Int. Bot., 1907, 135; *Aspidium Fauriei*, *Christ*, var. *elatius*, *Christ*, Bull. Herb. Boiss., VI, 195; *Nephrodium Fauriei*, *Christ*, var. *elatius*, Copel., Polypod. Philipp., 21.

Rhizome oblique. Stipes tufted, few, slender, to ± 25 c.M. long, stramineous, naked, the base black-brown, sparingly clothed with brown, subulate scales. Fronds lanceolate-elongate, to ± 25 c.M. long and 8 c.M. broad, not narrowed at the base. Pinnae alternate, to ± 18 on each side, spreading,

moderately approximate, petiolate, the lower distant, rhomboid-ovate, unequal-sided, lobed, acute, the lower side cuneately cut away, the upper side auriculate; lowest pinnae $\pm 3\frac{1}{2}$ c.M. long, $\pm 2\frac{1}{2}$ c.M. broad at the base. Lobes oblique, oblong, obtuse, ± 7 on each side, denticulate, the lower shallowly lobed again, the lowest deeply incised and sometimes free like the auricle. Texture thin; surfaces naked like the rachis; colour dark-green; veins pinnate in the lobes. Sori scattered, numerous, 3—7 to a lobe, ovate or roundish; indusium reniform.

Luzon; Yunnan.

(5b) **A. benguetense**, *Christ*, Philipp. Journ., II, 161.

Rhizome creeping or oblique, slender, black. Stipes approximate, few, slender, stramineous, 25 c.M. or more long, the base clothed with scattered, lanceolate, brown scales. Fronds lanceolate, ± 30 c.M. long, ± 9 c.M. broad, 2-pinnatifid, not narrowed at the base. Pinnae sessile, spreading, remote, ± 25 below the pinnatifid apex, the lowest subdeflexed, ± 5 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, lanceolate, acuminate, pinnatifid nearly to the costa. Lobes subpectinate, ± 18 on each side, linear-oblong, $\pm 2\frac{1}{2}$ m.M. broad, bluntish, entire or subcrenate. Texture flaccid; rachis and costae stramineous, puberulous; surfaces naked; colour black-green; veins simple, oblique, ± 6 —8 on each side. Sori medial, 6—8 on each side, roundish, hardly 1 m.M. broad; indusium thin, hyaline, roundish-reniform, indistinctly umbonate, caducous. — By its nephrodioid sori a *Dryopteris*, but in all other respects a true *Athyrium*.

Luzon.

(5c) **A. Copelandi**, *Christ*, Philipp. Journ., II, 161.

Rhizome branched, short-creeping, blackish. Stipes ± 9 c.M. long, plumbeo-stramineous, furfuraceo-squamulose like the rachis, the base sparingly clothed with brown, lanceolate-subulate scales. Fronds oblong, ± 14 c.M. long, ± 6 c.M. broad, long-acuminate, not narrowed at the base, the rachis winged towards the apex. Pinnae sessile, ± 15 on each side, falcate, the higher ones adnate, decurrent, the lower remote, coarsely lobed, ± 3 c.M. long, ± 1 c.M. broad. Lobes ± 6 on each side, the lowest anterior largest, triangular-ovate, ± 4 m.M. long, 5 m.M. broad, subacute, subcrenate. Texture coriaceous; surfaces naked; colour black-green, paler beneath, main veins pinnate in the lobes with 3—4 veins on each side, all soriferous.

Sori oblique, 2—3 m.M. long, \pm 1 m.M. broad, ovate-lanceolate, slightly curved; indusium greyish, turgid, persistent.

Luzon.

* (4) **A. macrocarpum**, *Bedd.*, Handb. Ind. Ferns, 165; Ferns S.I., tab. CLII—CLIII; *Asplenium macrocarpum*, *Bl.*, Rac., Flor. Btz., I, 225; *Aspidium macrocarpon*, *Bl.*, Enum., 162.

Rhizome erect or suberect, clothed with lanceolate, brown scales. Stipes \pm tufted, 12—25 c.M. long, firm, erect, stramineous, scaly at the base. Fronds ovate-lanceolate, acuminate, 15—60 c.M. long, 10—25 c.M. broad, pinnate. Pinnae numerous, the lower ones often deflexed, the longest stalked, 1—4 c.M. broad, cut down into several oblong, bluntish, erecto-patent, crenate lobes or even pinnate at the base, the base \pm auricled on the upper side. Secondary segments sometimes pinnatifid. Texture herbaceous; rachis naked or glanduloso-pubescent. Sori copious, roundish or oblong, the costal ones horseshoe-shaped or hamato-reniform.

Trop. Asia, China, Japan.

(5) **A. acrostichoides**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 225, fig. 120, C; *A. thelypteroides*, *Desv.*, *Bedd.*, Handb. Ind. Ferns, 164; *A. allantodioides*, *Bedd.*, Ferns Br.I., II, tab. CCXXI; *Asplenium thelypteroides*, *Mex.*, Hk. Bk., Syn. Fil., 226; *A. acrostichoides*, *Sw.*, (oldest name).

Stipes 15—50 c.M. long, erect, polished, stramineous, furnished at the base with large, lanceolate, thin, light-coloured scales. Fronds \pm 50—90 c.M. long, 15—30 c.M. broad, lanceolate, pinnate. Pinnae numerous, the lower ones gradually reduced, distant, the largest ones spreading or erecto-patent, 2—2½ c.M. broad at or near the base, narrowed gradually towards the apex, the edge deeply cut down into numerous crenate or nearly entire, oblong, blunt, \pm spreading segments. Texture herbaceous; rachis naked or \pm hairy. Sori in close, regular rows, slightly curved, reaching nearly from the costulae to the edge, the lower single or double.

Penang; Br. India, China, North. America.

(6) **A. hyalostegium**, *Copel.*, Philipp. Journ., I, Suppl. 255.

Rhizome creeping, often intertwined, sparingly clothed with minute, ferruginous scales. Stipes caespitose, 4—8 c.M. long, filiform, stramineous or greenish, the base brownish, the upper part clothed like the rachis with short, whitish, silky hairs. Fronds 5—7 c.M. long, 1½—4 c.M. broad, acu-

minate, pinnate. Pinnae ovate, obtuse, the lower ones pinnate again, the higher ones simple, joining. Pinnulae (or segments) narrowly oblong, obtuse, often entire. Texture herbaceous; surfaces sparingly pubescent with short, white hairs; veins simple. Indusium thin, setaceous, rarely asplenioid, often curved and nephrodioid.

Luzon.

(7) **A. Loheri**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1001.

Rhizome short, stout, sparingly clothed like the stipes with setiform scales. Stipes tufted, slender, erect, \pm 18 c.M. long, green, castaneous at the base. Fronds \pm 13 c.M. each way, deltoid, nearly 3-partite at the base, the rachis winged. Pinnae \pm 5 on each side, the higher ones oblong, subacute, \pm deeply lobed, the lowest ones subdeltoid, rather 2-pinnatifid, the posterior side produced. Lobes close, subobtuse, minutely crenulate. Texture thin, pellucid; colour pale-green; surfaces rather naked or hardly puberulous. Sori minute, in 1 or more rows, roundish or broadly reniform; indusium thin, greyish, asplenioid or aspidioid (round), fimbriate.

Luzon.

(8) **A. Sarasinorum**, *Christ*, Fil. Saras., IV, 454; Ann. Bot., XV, 125, tab. XV, fig. 16, a—b; *Asplenium Sarasinorum*, *Copel.*, Polypod. Philipp., 87; — var. *philippinense*, *Christ*, Bull. Herb. Boiss., VI, 154; *Ath. philippinense*, *Christ*, *Copel.*, Philipp. Journ., I, Suppl., 254.

Rhizome short, clothed with long, blackish-brown scales. Stipes tufted, slender, succulent, \pm flaccid, brown, slightly scaly, 10—15 c.M. long. Fronds \pm 10 c.M. long, \pm 4 c.M. broad, oblong-deltoid, acuminate, 3-pinnatifid to 5-pinnate. Pinnae \pm 10 on each side, the lowest the longest. Pinnulae sessile, the lower ones short-stalked, roundish-ovate, divided again, the lobes roundish, broadly crenate or toothed. Texture herbaceous. Sori 3—5 on the ultimate divisions, solitary on the lobes, oblong, slightly recurved; indusium brownish.

Var. philippinense: Lobes deeply incised, the terminal ones linear-lanceolate, moderately acute, \pm $\frac{1}{2}$ m.M. broad.

Celebes, Philippines.

(9) **A. aristulatum**, *Copel.*, Philipp. Journ., I, Suppl., 255; — var. *sphagnicolum*, *Copel.*, l. c.

Rhizome erect, densely clothed with lanceolate, acuminate, cinnamon-

coloured scales like the base of the stipes. Stipes caespitose, \pm 20 c.M. long, slender, the base nigrescent-stramineous, the upper part naked, greenish like the main rachis. Fronds 20—25 c.M. long, 10—15 c.M. broad, acuminate, 5-pinnate. Pinnae lanceolate-ovate, short-stalked, often acute. Pinnulae oblong, obtuse, the rachis narrowly winged. Tertiary segments oblong, adnate, obtuse or truncate, serrate or dentate, the teeth finely aristate, retroflexed. Texture membranaceous; surfaces naked, the upper blackish-green, the lower paler. Sori costal, short, broad, straight or curved; indusium whitish.

Var. sphagnicolum: Stipes filiform, 6—12 c.M. long; fronds 5—10 c.M. long, 2-pinnate.

Luzon.

* (10) **A. solenopteris** (*selenopteris*), *Moore*, *Bedd.*, *Handb. Ind. Ferns*, 166; *A. pectinatum*, *Bedd.*, *Ferns S. I.*, tab. CLV; *A. nigripes*, *Bedd.*, l. c., tab. CLVII; *Asplenium Selenopteris*, *Mett.*, *Farnng.*, VI, *Aspl.*, 196; *Allantodia Solenopteris*, *Kze*, *Linn.*, XXIV, 266; — var. *remotum*.

Rhizome oblique or erect, the scales rather large, ovate-acuminate, dense. Stipes stramineous, slightly scaly, often 25—30 c.M. long. Fronds ovate-acuminate or lanceolate, never deltoid, 25—50 c.M. long. Pinnae distant, erecto-patent, 15—20 c.M. long, stalked, lanceolate-acuminate. Pinnulae to \pm 4 c.M. long, short-stalked, close, bluntish, the edge cut down nearly or quite to the costa; the base more or less obliquely subtruncate. Segments oblong, the lower incised and toothed, the others toothed or serrate. Texture herbaceous, rather flaccid; surfaces naked. Sori oblong, more or less curved, the lower ones not rarely hamate or horseshoe-shaped.

Var. remotum: Pinnulae distant, the lower ones to \pm 2 c.M. apart. Texture somewhat firmer.

South. India, Ceylon; — var. *remotum:* *Java* (*Zippelius*).

(11) **A. drepanopterum**, *A. Br.*, *C. Chr.*, *Ind. Fil.*; *A. drepanopteron*, *Moore*, *Christ*, *Philipp. Journ.*, II, 161; *A. oxyphyllum*, *Moore*, *Bedd.*, *Ferns Br. I.*, I, tab. LXVII; *Asplenium oxyphyllum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 228; *Polypodium drepanopteron*, *Kze*, *Linn.*, XXIII, 278, 318.

Stipes firm, erect, 15—50 c.M. long, stramineous, clothed at the base with large, lanceolate-acuminate, reddish-brown scales. Fronds 30—60 c.M. long, 15—50 c.M. broad, lanceolate, with several distant pinnae on each side, which are 10—20 c.M. long, $2\frac{1}{2}$ —5 c.M. broad. Pinnulae lanceolate, some-

times auricled on the upper side, in the larger forms again pinnatifid, the teeth mucronate. Texture subcoriaceous; rachis firm, naked, stramineous; veins regularly pinnate. Sori in 2 rows on the pinnae or pinnulae, medial; indusium often oblong-reniform.

Luzon; Japan to Yunnan and North. India.

* (12) **A. nitidulum**, *Milde*, Bot. Zeit., 1870, 370; *Asplenium nitidulum*, Moore, Rac., Flor. Btz., I, 229; *Allantodia nitidula*, Kze, Bot. Zeit., 1848, 191.

Rhizome erect or suberect, densely clothed with brown, narrow, lanceolate scales. Stipes to 18 c.M. long, slightly scaly at the base. Fronds deltoid-ovate, 15—50 c.M. long, nearly as broad, acuminate. Pinnae \pm 12 on each side, pinnate, the lower ones stalked, lanceolate, acuminate, to 4 c.M. broad. Pinnulae to 1 c.M. broad, lanceolate, bluntish, the upper base broadly, the lower narrowly cuneate, the edge cut down $\frac{1}{2}$ — $\frac{2}{3}$ -way to the costa into linear-oblong, nearly close lobes. Lobes erecto-patent, rounded and crenate to toothed at the apex. Texture herbaceous to subcoriaceous; rachis pale-coloured like the stipe; main veins pinnate in the lobes; veins few, simple. Sori short, mostly in 2 regular rows, close to the costae, the largest recurved at the apex.

Java.

* (13) **A. gymnogrammoides**, *Bedd.*, Handb. Ind. Ferns, 168; Ferns S.I., tab. CLVI; *Asplenium Filix Femina*, *Bernh.*, var., Rac., Flor. Btz., I, 228; *A. gymnogrammoides*, *Kl.*, *Mett.*, *Farng.*, VI, *Aspl.*, 193, tab. VI, fig. 13—14.

Rhizome erect, slightly scaly, the scales pale-ferruginous or brown, ovate. Stipes 7—15 c.M. long, stramineous-green, sparingly scaly. Fronds 20—30 c.M. long, to 12 c.M. broad, lanceolate-oblong, narrowed gradually towards the apex. Pinnae spreading, oblong, to $4\frac{1}{2}$ c.M. long, the lowest not the largest, subremote, oblong, the apex not or hardly acuminate, the rachis marginate (narrowly winged). Pinnulae trapezoid-oblong, to 1 c.M. long, to 7 m.M. broad, the apex mostly rounded or obtuse, duplicato-inciso-dentate, rarely acuminate, the edge lobed or deeply pinnatifid at the base, the base obliquely cuneate, subtruncate and auricled on the upper side, the lobes inciso-dentate. Texture membranaceo-herbaceous; rachis and surfaces naked but the midribs of the pinnulae beset with subulate, spine-like appendices on the upper side. Sori straight or slightly curved. — The ceylonese specimens are larger and commonly 5-pinnate.

Java; Ceylon.

* (14) **A. nigripes**, *Moore* (not *Bedd.*), *Christ*, *Farnkr. d. Erde*, 224; *Asplenium nigripes*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 230; *Aspidium nigripes & costale*, *Bl.*, *Enum.*, 162 & 170.

Rhizome erect, clothed with dark-brown, lanceolate scales. Stipes tufted, 15—70 c.M. long, stramineous, erect, scaly below. Fronds 30—60 c.M. long, 12—50 c.M. broad, ovate-lanceolate to deltoid, with numerous erecto-patent pinnae on each side. Pinnae lanceolate, pinnate, the apex acuminate, pinnatifid. Lowest pinnulae $1\frac{1}{2}$ —4 c.M. long, oblong, cut $\frac{1}{2}$ -way down or quite to the costa into oblong or ovate, more or less toothed segments. Texture thin, membranaceous or herbaceous; rachis naked, stramineous; upper surface beset with few, distant, subulate, spine-like appendices on the midribs of the segments. Sori oblong or linear-oblong, often slightly curved only, principally in 2 rows near the costae.

Trop. Asia, China, Madagascar.

* (15) **A. umbrosum**, *Pr.*, *Christ*, *Farnkr. d. Erde*, 225; *A. australe*, *Pr.*, *Bedd.*, *Ferns S.I.*, tab. CLVIII; *Diplazium umbrosum*, *Bedd.*, *Handb. Ind. Ferns*, 188; *Asplenium umbrosum*, *J. Sm.*, *Rac.*, *Flor. Btz.*, I, 228; *Polypodium umbrosum*, *Ait.*, (oldest name); — var. *muricatum*; *Asplenium muricatum*, *Mett.*, *Ann. Mus. Bot. L. B.*, II, 239.

Caudex erect, to 25 c.M. high, densely covered with the bases of fallen fronds, roots and brown, lanceolate scales. Stipes 30—80 c.M. long, strong, scaly at the base. Fronds 60—160 c.M. long, 30—120 c.M. broad, ovate-lanceolate to deltoid. Pinnae ovate-lanceolate to deltoid-lanceolate, 7—20 c.M. broad. Pinnulae close, cut down nearly or quite to the costa into unequal-sided, elliptico-rhomboidal segments $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long, 2—5 m.M. broad; the apex of the pinnulae lobed. Segments sessile, the apex rounded, the edge toothed or deeply inciso-crenate, the base narrowed or decurrent. Texture herbaceous; surfaces naked; central ribs of the segments pinnate, veins simple or forked. Sori oblong, copious, the lower often double; indusium inflate, slightly curved, entire. — A variable plant.

Var. muricatum: Stipes muricate.

Trop. Asia & Australia, New Zealand, Africa; — var. *muricatum*: *Java*.

* (16) **A. silvaticum**, *Milde*, *Bot. Zeit.*, 1866, 376; *A. woodwardioides*, *Christ*, *Fil. Saras.*, III, 235; *Asplenium basilare*, *Moore*, *Rac.*, *Flor. Btz.*, I, 229; *A. woodwardioides*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 229; *Diplazium brevisorum*, *J. Sm.*, *Journ. of Bot.*, III, 408; *Allantodia sylvatica*, *Bl.*, (oldest name).

Rhizome erect. Stipes firm, erect, naked, dark-brown, to ± 70 c.M. long or longer. Fronds subdeltoid, 60—200 c.M. long, 30—120 c.M. broad. Lower pinnae to 60 c.M. long, $7\frac{1}{2}$ —20 c.M. broad, pinnate. Pinnulae slightly stalked, 1—3 c.M. broad, narrowed towards the apex, cut down $\frac{2}{3}$ or more on the way to the costa into close or nearly close, linear-oblong, blunt, toothed, subfalcate segments, which are to $\pm 1\frac{1}{2}$ c.M. long, to $\pm \frac{1}{2}$ c.M. broad. Texture herbaceous; rachis and surfaces naked; main veins pinnate in the lobes; veins simple or forked. Sori oblong, short, close to the costulae, the lower ones often double, close to the midribs, not reaching more than $\frac{1}{2}$ -way the edge; indusium ciliate or eroso-fimbriate. — Resembles *Diplazium polypodioides*, *Bl.* in habit (*Bk.*).

Java, Celebes, Philippines.

55. ASPLENIUM, *Linné.*

Sori linear or linear-oblong, straight or slightly recurved, springing from the sides of the veins, oblique to the costa, or occupying the sides of the single central veins of linear ultimate segments, only 1 sorus to a vein. Indusia of the same shape as the sori, springing from the same sides of the veins, opening obliquely towards the costa or occasionally towards the margin.

Rhizome creeping or erect, the scales coarsely cancellate, with the areolae mostly decoloured, i. e. the cells with the side walls conspicuously thickened and mostly brown-coloured, the lumina generally pale or decoloured (*Sadeb.*, in *Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 40, B*). Stipes not articulate to the rhizome, with 1, or 2 free, \pm concave vascular bundles, which mostly unite towards the lamina (or higher) into a single tetrarchous one [$) ($, $) ($, $) ($, $\succ \prec$, \times , \vee , \wedge]. Fronds simple or divided; veins free or rarely anastomosing. *Hk. Bk., Syn, Fil, tab. IV, fig. 38, a—c, h.*

Cosmopolitan.

Arrangement of the sections.

- § 1. **THAMNOPTERIS.** Vein tips united by a \pm marginal strand or an intramarginal line. Fronds simple. *Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 125, E.*
- § 2. **NEOTTOPTERIS.** Veins free. Fronds simple. *Diels, l.c., fig. 125, F—G.*

- § 3. **EU-ASPLENIUM.** Veins free. Fronds divided. Ultimate segments with several to many soriferous veins. Indusium opening mostly towards the costa. Hk. Bk., Syn. Fil., tab. IV, fig. 58, a; Diels, l.c., fig. 126.
- § 4. **DAREA.** Veins free. Fronds divided. Ultimate segments linear or oblong, each with a single, central soriferous vein. Indusium opening towards the margin. Hk. Bk., l.c., tab. IV, fig. 58, b—c; Diels, l.c., fig. 127, F & H.
- § 5. **LOXOSCAPHE.** Like § 4, but sori and indusia much contracted, Davallia-like. Hk. Bk., l.c., tab. II, fig. 18, e; Diels, l.c., fig. 127, J—K.
- § 6. **ASPLENIDICTYUM.** Veins anastomosing towards the margin. Fronds divided. Hk. Bk., l.c., tab. IV, fig. 58, h (*Hemidictyum*).

Key to the sections.

- a.** Veins quite free.
- b.** Sori and indusia contracted, half-cupshaped or semicylindrical, not running along the veins. Fronds divided. § 5. **LOXOSCAPHE.**
- b.b.** Sori and indusia not contracted, oblong or linear, running along the veins.
- c.** Fronds simple. § 2. **NEOTTOPTERIS.**
- c.c.** Fronds divided.
- d.** Ultimate segments linear or oblong, each with a single soriferous vein. Indusium opening towards the margin. § 4. **DAREA.**
- d.d.** Ultimate segments various, each with several to many soriferous veins. Indusium typically and obliquely opening towards the costa or costula. § 3. **EU-ASPLENIUM.**
- a.a.** Veins united at the apex by a transverse, continuous, \pm marginal or intramarginal line. Fronds simple. § 1. **THAMNOPTERIS.**
- a.a.a.** Veins anastomosing more copiously towards the margin. Fronds divided. § 6. **ASPLENIDICTYUM.**

§ 1. **THAMNOPTERIS.**

- a.** Fronds linear, loriform, $1\frac{1}{2}$ —3 c.M. broad. Indusium black. (1) *A. colubrinum.*
- a.a.** Fronds lanceolate or linear-lanceolate, \pm 3—35 c.M. broad. Indusium not black.
- b.** Fronds conspicuously acuminate, rather rostrate. (2) *A. Simonsianum.*

b.b. Fronds obtuse, acute or acuminate, not rostrate.

c. Texture \pm coriaceous throughout; fronds narrowed towards the base.

d. Fronds rather obtuse. (3a) *A. pachyphyllum*.

d.d. Fronds rather acute or short-acuminate.

e. Edges cartilagineous, very prominent. (3b) *A. curtisorum*.

e.e. Edges not or hardly cartilagineous, not conspicuously prominent. (3c) *A. nidus*.

c.c. Texture membranaceous or subcoriaceous; base of fronds broad, scarious or chartaceous. (4) *A. cymbifolium*.

(1) ***A. colubrinum***, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 999; — var. *taeniophyllum*, *Copel.*, Philipp. Journ., II, 131.

Fronds tufted, linear, loriform, to 85 c.M. long, 1½—3 c.M. broad, the apex acuminate, the edge entire, the base narrowed into a short stipe. Texture coriaceous, rigid; costa stout, surrounded on each side by black lines (the costa or the frond?), black-fuscous, prominent beneath, sulcato-concave above; under surface furfuraceous with minute, pale scales; veins prominent, black, numerous, nearly spreading, simple or geminate at the base, \pm 1½ m.M. apart. Sori numerous, occupying the upper half of the frond, slightly immersed, black-brown, each pair separated by a sterile vein; indusium persistent, cartilagineous, black, with a pale margin.

Var. taeniophyllum: Stipes 2—5 c.M. long; fronds narrowed very gradually into the stipe; costa keeled beneath, 2-angular above, concave between the angles at the base; under surface naked; veins inconspicuous; sori often separated by 2 sterile veins.

Luzon; — var. *taeniophyllum*: *Mindoro*.

* (2) ***A. Simonsianum***, *Hk.*, Hk. Bk., Syn. Fil., 191; *Hk.*, Icon. Pl., X, tab CMXXV; *A. nidus* *L.*, var. *Simonsianum*, *Christ*, Bull. Herb. Boiss., VI, 151; *Thamnopteris Simonsiana*, *Moore*, *Bedd.*, Ferns Br.I., II, tab. CCXLVIII.

Fronds 30—45 c.M. long, 2½—4 c.M. broad, tapering above into a distinctly acuminate point and very gradually below into a short stipe, the margin nearly entire, the midrib rounded and prominent below. Texture coria-

ceous; veins erecto-patent. Sori copious, reaching from the midrib to a short distance of the edge.

Philippines; North. India.

(5a) **A. pachyphyllum**, *Hæ.*, Bot. Zeit., 1848, 146; *A. nidus*, L., var. *pachyphyllum*?

Stipes tufted, very short, nigro-paleaceous at the base. Fronds linear-lanceolate, \pm 45—50 c.M. long, 4—5 c.M. broad, rather obtuse, much narrowed towards the base, the margin thickened, reflexed, subrepand, obsoletely-remotely veined. Texture coriaceous, firm, thick; surfaces glossy; veins erecto-patent, often forked. Sori occupying the upper half of the frond distant from each other as well as from the costa and the margin.

Java, New Guinea.

(5b) **A. curtisorum**, *Christ*, Ann. Btz., XIX, 59.

Rhizome stout, rarely clothed with ovate scales. Fronds sessile or subsessile, narrowed gradually towards the base, \pm 50 c.M. long and 9 c.M. broad, shortly acuminate, the edge entire and cartilagineous, very prominent. Texture coriaceous; surfaces glossy, glaucous-green, naked; costa stout, prominent beneath; veins conspicuous, numerous, erecto-patent, forked at the base. Sori at the base of the veins, \pm close to the costa, oblique, \pm 10 m.M. long, occupying the upper half of the frond and reaching to \pm 7 c.M. below the apex, brown; indusium thin, whitish.

Celebes.

* (5c) **A. nidus**, L., Rac., Flor, Btz., I, 209; *A. australasicum*, *Hk.* (not *J. Sm.*), Fil. Exot., tab. LXXXVIII; *A. simplex*, *Mett.* (not *Bl.*), Farng., VI, Aspl., 86; *Thamnopteris Nidus*, *Pr.*, *Bedd.*, Ferns Br.I., II, tab. CXCVII; — var. *musifolium*; *A. musaefolium*, *Mett.*, l.c. 86; — var. *plicatum*; *A. plicatum*, *Zipp.*

Rhizome short. Fronds 40—120 c.M. long, 5—30 c.M. broad, lanceolate, sessile or tapering gradually into a short stipe, the apex acute or short-acuminate, the edge entire. Texture \pm coriaceous; costa rounded or keeled on the back; veins fine, $\frac{3}{4}$ —2 m.M. apart. Sori reaching nearly or quite from the costa to $\frac{1}{3}$ — $\frac{2}{3}$ on the way towards the margin. — *A. phyllitidis*, *Don* is the shortest and narrowest form, under 60 c.M. long, under $7\frac{1}{2}$ c.M. broad.

Var. musifolium: Fronds larger, to $2\frac{1}{2}$ M. long, to 35 c.M. broad, with the sori extending nearly to the edge.

Var. plicatum: Fronds obtuse, rigidly coriaceous, plicate, the edge undulate.

Paleotropical.

(4) **A. cymbifolium**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 999; Copel., Philipp. Journ., II, 150.

Fronds sessile, 55—120 c.M. long, 15—20 c.M. broad in the upper part, slightly narrowed below the middle, the base as broad or broader, 12—15 c.M. broad, (cordate after Christ), subscariose (chartaceous after Copel.), the edge entire, the apex subacute or abruptly short-acuminate. Texture membranaceous (subcoriaceous after Copel.), flaccid; costa very prominent, black or castaneous, woody; veins nearly spreading, slender, 1—2 m.M. apart, simple, connected by a distinct intramarginal line. Sori numerous, reaching from the costa to $\frac{9}{10}$ on the way to the edge, separated by 1—2 sterile veins; indusium yellow (brown after Copel.).

Luzon.

§ 2. NEOTTOPTERIS.

1. *Texture herbaceous to papyraceous.*

‡. *Fronds narrowed below into a rather suddenly truncate or subacute base.*

+. *Rhizome scales few, scattered.* (5) *A. epiphyticum.*

++. *Rhizome scales numerous, dense.* (6) *A. amboinense.*

‡.‡. *Fronds gradually decurrent at the base.*

×. *Fronds lanceolate, not suddenly narrowed into the lower half.*

(7) *A. concolor.*

×.×. *Fronds spathulate, rather suddenly narrowed into the wing-like lower half.*

(8) *A. spathulatum.*

1.1. *Fronds coriaceous or nearly so.*

2. *Stipes naked or nearly so.*

a. *Margin entire or subentire.*

b. *Fronds small, under 30 c.M. long.* (9) *A. microxiphion.*

b.b. *Fully developed fronds more than 30 c.M. long.*

c. *Stipes seriate. Sori erecto-patent.* (10) *A. batuense.*

c.c. *Stipes tufted. Sori patent.* (11) *A. Scortechinii.*

a.a. Margin crenate, serrate or toothed.

+. Rhizome erect, scaly. Fronds finely acuminate.

(12) *A. Mactierii*.

+.+. Rhizome wide-creeping, naked. Fronds acute.

(15) *A. vittaeforme*.

2.2. Stipes more or less scaly.

3. Margin quite entire or obscurely repand.

4. Surfaces naked.

(14) *A. scolopendrioides*.

4.4. Surfaces scaly, at least on the costa beneath.

5. Fronds narrow, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad; surfaces naked, costa scaly.

(15) *A. apoense*.

5.5. Fronds broader, 3—10 c.M. broad; under surface \pm scaly, at least when young.

6. Fronds 3— $3\frac{1}{2}$ c.M. broad.

(16) *A. natunae*.

6.6. Fronds $3\frac{1}{2}$ —10 c.M. broad.

7. Fronds not rounded, not abruptly acuminate at the apex.

(17a) *A. squamulatum*.

7.7. Fronds rounded and abruptly acuminate at the apex.

(17b) *A. Wernerii*.

3.3. Margin crenate, at least towards the apex of the frond.

+. Fronds rather distant, 25—35 c.M. long; costa prominent.

(18) *A. glochidiatum*.

+.+. Fronds densely approximate, 40—50 c.M. long; costa depressed.

(19) *A. comosum*.

(5) ***A. epiphyticum*, Copel.**, Polypod. Philipp., 79.

Rhizome scandent, clothed with few, small scales. Stipes distant, \pm 8 c.M. long, covered with minute, ferruginous scales. Fronds simple, 20—30 c.M. long, \pm 4 c.M. broad, acuminate, crenate or serrate, the base obtuse or subacute. Texture membranaceous; surfaces naked; costa firm, scaly; veins conspicuous, often forked, nearly spreading. Sori linear, not reaching the costa and the margin. — The fronds of the young plants are sometimes 2—3-pinnatisect or entire in the upper and divided in the lower part.

Mindanao.

* (6) ***A. amboinense*, Willd.**, Rac., Flor. Btz., I, 210; *A. fejeense*,

Brack., Hk. Bk., Syn. Fil., 192; *A. simplex*, Bl., Enum., 174; *A. taeniosum*, Kze, Bot. Zeit., 1848, 145; *A. scolopendrioides*, Muell., Pap. Pl., IV, 81.

Rhizome \pm creeping or climbing, the scales dense, brown, lanceolate, acuminate. Stipes 1—15 c.M. long, more or less scaly, especially below. Fronds lanceolate, 40—120 c.M. long, $5\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad, with the apex caudate or acuminate, often proliferous, gradually narrowed below into a rather subtruncate base, the margin entire or nearly so. Texture thin to subcoriaceous; costa slightly scaly below; veins oblique, simple or branched. Sori occupying the lower $\frac{2}{3}$ of the veins or reaching nearly to the edge.

Malaya; Polynesia.

* (7) **A. concolor**, *Hk.*, Hk. Bk., Syn. Fil., 191; Hk., Spec. Fil., III, tab. CLXIV, A; *A. diversifolium*, Bl., forma *concolor*, Rac., Flor. Btz., I, 211; *A. diversifolium*, Bl., forma *indivisa*, C. Chr., Ind. Fil.

Rhizome creeping, clothed with lanceolate scales. Stipes approximate, 2—5 c.M. long, firm, scaly in the lower part. Fronds 15—25 c.M. long, $2\frac{1}{2}$ c.M. or more broad, lanceolate, the point acute, the edge slightly crenato-serrate upwards, the lower part narrowed into the stipe gradually. Texture herbaceous; veins usually simple. Sori numerous, narrow, extending from the midrib to the margin.

Java, Celebes.

* (8) **A. spathulatum**, *Bk.*, Ann. of Bot., VIII, 124.

Stipes short, to 5 c.M. long, approximate, slightly scaly, the scales few, small, lanceolate, \pm adpressed, blackish or brownish. Fronds simple, \pm 25—55 c.M. long, 3—9 c.M. broad in the upper half, which is oblong-lanceolate, cuspidate, crenate or repando-crenate but more conspicuously towards the apex; near the middle the frond is rather suddenly narrowed into the wing-like lower half, which is $1-2\frac{1}{2}$ c.M. broad and decurrent upon the stipe. Texture membranaceous to papyraceous; surfaces naked; costa prominent and clothed beneath with 2 opposite rows of small, distant, \pm spreading scales; veins parallel, nearly spreading, simple or forked, reaching the margin. Sori narrow, occupying the lower $\frac{1}{2}$ — $\frac{2}{3}$ of the veins; indusium narrow, membranaceous, pale-coloured when young, at length recurved.

Sumatra, Malacca.

(9) **A. microxiphion**, *Bk.*, Kew Bull., 1896, 40.

Rhizome erect, the scales dense, lanceolate, membranaceous, dirty-brown.

Stipes tufted, slender, $2\frac{1}{2}$ —5 c.M. long, naked. Fronds simple, lanceolate, 15—20 c.M. long, 1—2 c.M. broad, entire, acuminate, narrowed at the base. Texture subcoriaceous; surfaces naked; veins erecto-patent, distant, simple or forked. Sori linear, not reaching the margin; indusium membranaceous.

Natuna Islands.

* (10) **A. batuense**, *v. A. v. B.*, Bull. Dép. Agr. I.N., 1908, XVIII, 15.

Rhizome stout, woody, creeping, scaleless. Stipes short, distant, to 1 c.M. long. Fronds lanceolate, 40—75 c.M. long, 4—10 c.M. broad, the apex acute or short-acuminate, often proliferous, the edge entire, the lower part narrowed very gradually towards the bluntish base. Texture papyraceous to coriaceous; costa prominent and \pm quadrangular beneath, marginate there with 2 opposite, very narrow wing-like crests in the angles; veins conspicuous, fine, erecto-patent, simple or forked. Sori numerous, occupying nearly the whole frond except the upper $\frac{1}{8}$ and the lower wing-like part, not reaching the midrib and the edge, very irregular in length, separated by 0—6 sterile veins or veinlets; indusium narrow, subcoriaceous, at length distinctly recurved.

Batu Islands (Raap, No. 674).

(11) **A. Scortechinii**, *Bedd.*, Journ. of Bot., XXV, 322.

Stipes tufted, short, erect. Fronds linear-lanceolate, 60—75 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, gradually narrowed below into the stipe, and to the apex into a long, fine point, the margin entire or subentire. Texture coriaceous; surfaces naked or the under surface slightly scaly. Sori patent, occupying the lower $\frac{2}{3}$ of the veins.

Malacca.

(12) **A. Mactierii**, *Bedd.*, Journ. of Bot., XXVI, 5.

Rhizome short, erect, scaly, the scales dark-brown, with a paler margin, lanceolate from a broad base, finely acuminate. Stipes 15—25 c.M. long, pale-yellow, whitish at the base. Fronds 15—25 c.M. long, $2\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, gradually narrowed below, gradually and finely acuminate at the apex, the margin crenate or serrate. Texture subcoriaceous; veins distinct, usually once forked near the base, occasionally again forked towards the apex, not quite reaching the margin. Sori occupying the lower $\frac{2}{3}$ of the veins; indusium broad, prominent.

Penang.

* (13) **A. vittaeforme**, *Cav.*, Hk. Bk., Syn. Fil., 192; *A. sundense*, *Bl.*, *Rac.*, Flor. Btz., I, 209.

Rhizome slender, wide-creeping, naked. Stipes short, erect, naked. Fronds rather distant, lanceolate, 15—45 c.M. long, 2—7½ c.M. broad or narrower, the apex acute, the base narrowed gradually into the stipe, the margin crenate or obscurely toothed. Texture coriaceous, veins simple or forked, close. Sori copious, often reaching from the midrib nearly to the edge. — The plant has sometimes also abnormal fronds, which are finely divided like those of *Teratophyllum*.

Malaya; Polynesia.

(14) **A. scolopendrioides**, *J. Sm.*, Hk. Bk., Syn. Fil., 192; Hk., *Icon. Pl.*, X, tab. CMXXX.

Rhizome clothed with lanceolate, brown, chaffy scales. Stipes ± 2—4 c.M. long, erect. ± scaly. Fronds lanceolate, 50—50 c.M. long, 3—4 c.M. broad, the apex ± suddenly caudate, the base narrowed very gradually, the edge entire. Texture subcoriaceous; veins oblique, commonly simple, rather distant. Sori reaching from the midrib to within 5—7 m.M. of the margin; indusium leaving a distinct, elevated ridge on the frond when it dehisces. — (See also *A. glochidiatum*, *Rac.*).

Borneo, Philippines.

(15) **A. apoense**, *Copel.*, *Polypod. Philipp.*, 79.

Rhizome thick, short, erect. Stipes clustered, 1—5 c.M. long, firm, scaly. Fronds 20—35 c.M. long, 1½—2½ c.M. broad, entire, acute, narrowed gradually into the stipe. Texture coriaceous; surfaces naked; costa firm, scaly; veins inconspicuous, erecto-patent, often forked. Sori linear, occupying the lower ⅔ of the veins.

Mindanao.

(16) **A. natunae**, *Bk.*, *Kew Bull.*, 1896, 41.

Rhizome erect, woody, the scales dense, lanceolate, membranaceous, dirty-brown. Stipes 10—15 c.M. long, brown, slightly scaly above. Fronds simple, 30—45 c.M. long, 3—5½ c.M. broad, lanceolate, narrowed at the base. Texture rigidly coriaceous; under surface minutely scaly; veins close, obscure, immersed, ± 1¼ m.M. apart, reaching to the margin. Sori linear, not reaching the margin; indusium persistent.

Natuna Islands.

* (17a) **A. squamulatum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 210; *Thamnopteris stipitata*, *Pr.*, *Epim.*, 68; — var. sessile.

Rhizome short, creeping or erect. Stipes \pm tufted, 5—20 c.M. long, strong, erect, scaly like the costa, the scales persistent or more or less deciduous. Fronds 50—100 c.M. long, 5—10 c.M. broad, lanceolate, the apex bluntish, acute or acuminate, the edge entire or nearly so, the base narrowed gradually into the stipe. Texture coriaceous; surfaces scaly at least when young, the scales scattered, adpressed, minute; veins not very distinct, simple or 1—2 \times forked, \pm 2 $\frac{1}{2}$ m.M. apart. Sori reaching from near the midrib to 5—15 m.M. of the margin; indusium broad, membranaceous, naked.

Var. sessile: Stipes short or wanting. Fronds 20—25 c.M. long, 3 $\frac{1}{2}$ —5 $\frac{1}{2}$ c.M. broad; veins rather close; indusium \pm coriaceous. — United with the type by intermediate forms (e. g. *forma calvescens*, *Christ*, in *Ann. Btz.*, XX, 111).

Malaya; — var. sessile: *Borneo* (*Teuscher*).

(17b) **A. Wernerii**, *Rst.*, *Fed.*, *Rep.*, V, 39.

Rhizome short-creeping, at length suberect, densely clothed with lanceolate, fusco-ferrugineous, entire or slightly fimbriate scales. Stipes subfasciculate, firm, 10—20 c.M. long, \pm densely brown-scaly. Fronds ovate-elongate, to 45 c.M. long, to 6 c.M. broad, the apex obtuse and abruptly acuminate, the base nearly rounded or short-decurrent upon the stipe, the edge entire or obscurely repand. Texture chartaceous; upper surface olive-green and slightly scaly, lower livid-green and copiously scaly; costa prominent, scaly; veins erecto-patent, simple or forked, \pm 1 $\frac{1}{2}$ m.M. apart, parallel. Sori occupying the lower $\frac{3}{4}$ of the veins; indusium entire.

New Guinea.

* (18) **A. glochidiatum**, *Rac.*, *Bull. Ac. Crac.*, 1902, 62; *A. echinosporum*, *Rac.*, MS. in *Herb. Bog*.

Rhizome creeping, ascending, the scales light-brown, adpressed, lanceolate, acuminate, covered with short hairs near the base. Stipes \pm 1 c.M. apart, 6—10 m.M. long, black, scaly. Fronds simple, 25—35 c.M. long, \pm 5 c.M. broad, narrowed gradually towards both ends, but more or less suddenly at the apex into an acuminate point and towards the base into a narrow wing to the costa, the margin crenate, the lower, narrowed part entire. Texture chartaceous, surfaces glossy, naked in the lower part; costa prominent on both sides, black and scaly at the base, green in the higher part; veins parallel, suboblique,

slightly prominent beneath, simple or forked. Sori linear, occupying the lower $\frac{1}{2}$ — $\frac{2}{3}$ of the veins, often separated by 1—2 sterile veins; indusium thin, entire, pale. — Fronds of the young plants sessile, the lower part gradually narrowed, the base bluntish, the higher part deeply pinnatifid. — The ripe spores are copiously echinate, and the fronds cultivated in the Buitenzorg gardens have the indusia like those of *A. scolopendrioides*, *J. Sm.* — Both species are probably identical.

Borneo.

(19) **A. comosum**, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., 127.

Rhizome creeping, clothed like the stipes with lanceolate, acuminate, black-brown scales. Fronds approximate, numerous, erect, 40—50 c.M. long, linear-lanceolate, long-acuminate, caudate, undulato-crenate in the upper, entire in the lower part, decurrent. Texture coriaceous, lucid; surfaces naked; costa broad, depressed; veins conspicuous, dense, erecto-patent, simple or forked. Sori numerous, \pm 100 on each side, reaching from the costa to the margin, occupying the whole frond; indusium narrow.

New Guinea.

§ 5. EU-ASPLENIUM.

- a.** *Fronds simply pinnate; pinnae entire or incised, but not deeply pinnatifid.*
- b.** *Pinnae short, roundish or oblong, blunt, rarely acute, not more than 2 c.M. long* ⁽¹⁾.
- 1.** *Rachis green, at least in the upper half.*
- 2.** *Rachis naked.*
- 3.** *Pinnae in \pm 6 pairs, $1\frac{1}{2}$ —2 c.M. long.*
- (20) *A. subnormale.*
- 3.3.** *Pinnae in \pm 30 pairs, to $1\frac{1}{4}$ c.M. long.*
- (21) *A. minus.*
- 2.2.** *Rachis hairy or scaly.*
- (22a) *A. Loherianum.*
- 1.1.** *Rachis dark-coloured throughout, not green.*
- §.** *Fronds under 2 c.M. broad* ⁽²⁾.
- ‡.** *Stipe and rachis 3-winged.*
- (22b) *A. Stantonii.*
- ‡.‡.** *Stipe and rachis not 3-winged.*

⁽¹⁾ See also **A. heterocarpum**, *Wall.*, **unilaterale**, *Lam.* & **filipes**, *Copel.*

⁽²⁾ See also **A. normale**, *Don.*

- *. *Sori at length confluent.* (25a) *A. trichomanes.*
- *.*. *Sori not confluent.* (25b) *A. exiguum.*
- §.§. *Fronds broader, 2—4 c.M. broad when fully developed.*
- ×. *Stipe and rachis naked.* (24) *A. normale*
- ×.×. *Stipe and rachis densely fibrillose.* (25) *A. subavenium.*
- b.b.** *Pinnae linear or linear-oblong, acute or acuminate, rarely obtuse, more than 2 c.M. long in the fully developed plants.*
- c.** *Pinnae ± equally cuneate or truncate on both sides at the base.*
- d.** *Rachis naked.*
- e.** *Pinnae few, not more than ± 5 on each side.*
- *. *Pinnae cuneate at the base.* (26) *A. salignum.*
- *.*. *Pinnae cordate at the base.* (27) *A. diversifolium.*
- e.e.** *Pinnae 6 or more on each side.*
- f.** *Texture ± herbaceous or membranaceous.*
- g.** *Pinnae not deeply toothed.*
- h.** *Sori few, distant.* (28) *A. Labillardierii.*
- h.h.** *Sori numerous, close.* (29) *A. remotum.*
- g.g.** *Pinnae deeply toothed.* (30) *A. subaquatile.*
- f.f.** *Texture ± coriaceous.*
- ×. *Rachis narrowly winged in the upper part.*
- (31) *A. sumatranum.*
- ×.×. *Rachis not winged.* (32) *A. vulcanicum.*
- d.d.** *Rachis scaly, chaffy or fibrillose.*
- §. *Pinnae broadly adnate to the rachis. Rachis fibrillose.*
- (33) *A. fuliginosum.*
- §.§. *Pinnae not or narrowly adnate to the rachis. Rachis chaffy or scaly.*
- ‡. *Pinnae in ± 9 pairs, cordate at the base.*
- (34) *A. horizontale.*
- ‡.‡. *Pinnae numerous, subtruncate at the base.*
- (35) *A. longissimum.*
- c.c.** *Pinnae unequal-sided at the base, i. e. the acroscopical side broadly cuneate or truncate and the basiscopical ± narrowly cuneate.*
- 1.** *Texture ± herbaceous or membranaceous.*
- α. *Fronds dimorphous, the fertile ones much longer and broader than the barren ones.* (36) *A. monotis.*
- α.α. *Fronds uniform, the fertile ones ± as large as the barren ones.*
- β. *Pinnae bluntly rounded or subtruncate at the apex.*

*. *Lower pinnae gradually reduced.*

(37) *A. borneense.*

.. *Lower pinnae not reduced.* (38) *A. tenerum.*

β.β. *Pinnae bluntish or acute at the apex* (*).

γ. *Fronde small or middle-sized, under 50 c.M. long.*

δ. *Fronde narrow, under 10 c.M. broad.*

ε. *Pinnae inciso-crenate, bluntish or acute.*

(39) *A. lunulatum.*

ε.ε. *Pinnae coarsely crenate, acute.*

(40a) *A. Steerei.*

ε.ε.ε. *Pinnae coarsely 2-dentate, acuminate.*

(40b) *A. Meyenianum.*

δ.δ. *Fronde broader, more than 10 c.M. broad when fully developed.* (41) *A. prionurus.*

γ.γ. *Fronde large, more than 50 c.M. long when fully developed.*

+. *Pinnae 2—3½ c.M. broad; sori occupying the lower half of the veins.*

(42) *A. griseum.*

+.+. *Pinnae narrower; sori reaching nearly to the edge.* (45) *A. persicifolium.*

1.1. *Texture chartaceous to coriaceous.*

2. *Higher sori subparallel and nearly close to the costae.*

§. *Pinnae more or less deeply lobed, the lobes ± rounded, entire or bluntly crenate, the higher lobes not rarely tooth-like.*

‡. *Fronde small, under 15 c.M. long, under 7½ c.M. broad. Pinnae bluntish or subacute.*

(44) *A. Schoggersii.*

‡.‡. *Fronde larger, more than 15 c.M. long, more than 7½ c.M. broad. Pinnae long-acuminate.*

(45) *A. caudatum.*

§.§. *Pinnae serrate, broadly crenate or ± deeply lobed, the crenations or lobes sharply serrate or toothed.*

×. *Pinnae 4—7½ c.M. long, with the upper base suddenly cuneate.* (46) *A. contiguum.*

(*) See also under **A. tenerum**, Forst.

x.x. *Pinnae 10—18 c.M. long, with the upper base rather truncate or rotundato-cuneate.*

(47a) *A. nigrescens.*

(47b) *A. militare.*

2.2. *Higher sori erecto-patent, not subparallel to the costa.*

3. *Lower pinnae reduced, suborbicular or subcordate.*

(48) *A. pellucidum.*

3.3. *Lower pinnae not or hardly reduced.*

4. *Pinnae suddenly narrowed from a broad base into a narrow point. Stipes ± black.*

(49) *A. dimidiatum.*

4.4. *Pinnae not suddenly narrowed into the point. Stipes mostly green, grey or brown.*

5. *Pinnae narrow, to ± 2½ c.M. broad.*

6. *Pinnae varying from crenulate to inciso-crenate.*

7. *Crenulate.* (50) *A. Cesatianum.*

7.7. *Inciso-crenate.*

8. *Stipe and rachis fibrillose.*

(51) *A. crinicaule.*

8.8. *Stipe and rachis ± scaly or nearly naked.*

(52) *A. acutiusculum.*

6.6. *Pinnae serrate or toothed.*

‡. *Rachis strigoso-paleaceous.*

(53) *A. impressum.*

‡.‡. *Rachis naked or slightly scaly.*

+. *Pinnae ascending. Sori rather regular, ± 5—6-jugate.*

(54) *A. Gardneri.*

+.+. *Pinnae spreading. Sori rather irregular, more numerous.*

(55) *A. adiantoides.*

5.5. *Pinnae broader, 2½—7½ c.M. broad.*

‡. *Pinnae minutely 2-serrate.*

(56) *A. Sancti-Christophori.*

‡.‡. *Pinnae entire, crenate or irregularly and sharply toothed.*

×. Veins distinctly visible beneath.
(57) *A. macrophyllum*.

×.×. Veins sunk, indistinct beneath.
(58) *A. paradoxum*.

c.c.c. Pinnae dimidiate or subdimidiate, i. e. at least the lower $\frac{1}{3}$ of the lower side cut away along the costa.

d. The lower side of the pinnae quite cut away.
(59) *A. heterocarpum*.

d.d. The lower $\frac{1}{3}$ — $\frac{2}{3}$ of the lower side cut away.

e. Stipe and rachis purple or black-brown, sometimes grey.
(60a) *A. unilaterale*.
(60b) *A. filipes*.

e.e. Stipe and rachis green. (61) *A. obscurum*.

a.a. Fronds pinnate; pinnae deeply pinnatifid, or pinnate at the base.

1. Fronds small or middle-sized, under 45 c.M. long.

2. Secondary segments undivided.

3. Pinnae ovate, the lobes cuneate-oblong. (62) *A. denticulatum*.

3.3. Pinnae linear-lanceolate, the lobes linear-cuneate.

(63) *A. Warburgianum*.

2.2. At least the lower secondary segments on the anterior side shallowly or deeply 2—3-fid (¹).

✧. Lowest anterior segment rather broader than long, ± spreading. Sori short, oblong. (64) *A. laciniatum*.

✧.✧. Lowest anterior segment longer than broad, erecto-patent. Sori long, linear. (65) *A. Hagenii*.

1.1. Fronds large, more than 45 c.M. long when fully developed. (³)

(66) *A. horridum*.

a.a.a. Fronds more copiously 2—4-pinnate (²).

b. Ultimate segments small, lanceolate-spathulate, acute or incised with 2—3 sharp, entire teeth at the apex. (67) *A. tenuifolium*.

b.b. Ultimate segments oblong or cuneate-oblong, the outer edge more copiously crenulate to dentate, or incised and crenulate to dentate.

c. Fronds 2—3-pinnate.

d. Rachis pilose, fibrillose or scaly, at least when young.

(¹) See also *A. stereophyllum*, Kze.

(²) With regard to the determination of the species a very difficult group and not always clearly distinguished from group **a.a.**

(³) See also *A. caudatum*, Forst.

- +. *Pinnae rather short, 2 $\frac{1}{2}$ —4 c.M. long.*
 ×. *Pinnae obtuse.* (68a) *A. stereophyllum.*
 ×.×. *Pinnae acuminate.* (68b) *A. Elmeri.*
 +.+. *Pinnae longer, acute or acuminate, 4—8 c.M. long.*
 (69) *A. praemorsum.*
- d.d.** *Rachis naked or slightly scaly.*
- e.** *Lower pinnulae cuneate-oblong, with the outer edge ± rounded, more or less incised.* (70) *A. cuneatum.*
- e'.e'.** *Lower pinnulae lanceolate, rhomboidal or subdeltoid, narrowed towards the apex, the sides more or less incised.*
- f.** *Texture herbaceous. Venation pinnate.*
 (71) *A. bulbiferum.*
- f'.f'.** *Texture subcoriaceous. Venation subflabellate.*
- g.** *Sori short, radiant.* (72) *A. nitidum.*
- g'.g'.** *Sori long, linear.*
- h.** *Pinnae long-acuminate, caudate, the apex nearly filiform.* (75) *A. setisectum.*
- h.h.** *Pinnae less conspicuously acuminate.*
 (74) *A. affine.*
- c'.c'.** *Fronds decoumpound, copiously 3—4-pinnate.* (75) *A. laserpitifolium.*

(20) **A. subnormale**, *Copel.*, Polypod. Philipp., 80; *Perk.*, *Fragm. Philipp.*, tab. IV, fig. B.

Fronds solitary or few on a short, creeping rhizome. Stipes ± 1 $\frac{1}{2}$ —4 c.M. long, naked towards the frond, slender, dark and polished, but shading into a green rachis, with green, lateral lines. Fronds 4—10 c.M. long, to 4 c.M. broad, lanceolate-ovate, acute, pinnate. Distinct pinnae ± 6 pairs, 2 or more pairs indicated by clefts, the lowest whether or not smaller than the next in order, crenate, commonly trapezoid, irregular in shape, the lower half cut away to the midrib at the base, the upper base parallel to the rachis. Texture membranaceous; surfaces naked. Indusium narrow.

Luzon.

* (21) **A. minus**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 211; *A. normale*, *Mett.* (not *Don*), *Ann. Mus. Bot. L.B.*, II, 254.

Rhizome short, erect or creeping, sparingly covered with small, lanceolate scales. Stipes densely crowded, 1—3 c.M. long. Fronds linear-lanceolate, to

20 c.M. long, to $2\frac{1}{2}$ c.M. broad. Pinnae numerous (± 50 on each side), sessile, the lower ones \pm obliquely flabellate, truncate at the base, nearly twice as long as broad, the following subrhomboidal, to $12\frac{1}{2}$ m.M. long, to 5 m.M. broad, the upper base truncate and \pm auricled, the lower narrowly cuneate, the upper and outer edge of all crenate or bluntly toothed. Texture thinly herbaceous; rachis naked, becoming reddish towards the base; both surfaces naked; costae wavy; veins simple, the lower ones on the anterior side forked. Sori 1—3—5 to a pinna, oblong; indusium broad. — Lower pinnae sterile.

Java.

(22a) **A. Loherianum**, *Christ*, Bull. Herb. Boiss., VI, 152.

Rhizome short. Fronds 2—5, short-stalked, ± 10 c.M. long. Pinnae 12—20 on each side, alternate, very distant, ± 5 m.M. long, ovate, deeply crenato-dentate, the teeth obtuse. Texture rather fleshy; rachis slender, flexuose, decumbent, purplish or green at the base, clothed with linear scales, proliferous at the apex; veins subflabellate. Sori oblong, oblique, at length covering the whole under surface of the pinnae.

Philippines.

(22b) **A. Stantonii**, *Copel.*, Philipp. Journ., I, Suppl., 151.

Rhizome erect, short, scaly. Stipes tufted, ± 2 c.M. long, naked, fuscous, triangular, the angles acute, winged. Fronds ± 15 c.M. long, ± 1 c.M. broad, pinnate, the rachis conspicuously 3-winged, not green. Pinnae oblong, obtuse, deflexed, the lower ones flabelliform, somewhat remote. Texture coriaceous; surfaces naked. Sori ± 5 to a pinna, at length filling up the whole under surface of the pinnae.

Luzon.

* (25a) **A. trichomanes**, *L.*, Hk. Bk., Syn. Fil., 196; Bedd., Ferns S.I., tab. CXLVII; — var. *densum*; *A. densum*, *Brack.*, Expl. Exp., XVI, 151, tab. XX, fig. 5.

Stipes densely tufted, 1—10 c.M. long, naked, glossy, chestnut-brown, polished. Fronds 15—50 c.M. long, ± 10 —12 m.M. broad, linear-lanceolate, with 15—50 opposite pairs of sessile, horizontal, \pm orbicular or oblong pinnae, the edges slightly crenate, the 2 sides \pm unequal, the upper side the broadest, the base rounded or broadly cuneate. Texture firm; veins pinnate, inconspicuous; surfaces naked; rachis polished, blackish-brown, slender. Sori linear-oblong, at length covering the whole under surface of the pinnae.

Var. densum: Smaller and narrower.

In the temperate zones and the cool alpine regions of the tropics.

(23b) **A. exiguum**, *Bedd.*, Ferns S.I., 49, tab. CXLVI.

Rhizome short, erect, clothed above with numerous subulate, black scales. Stipes numerous, 7—50 c.M. long, blackish-purple, more or less furnished with long, black, subulate scales. Fronds 4—15 c.M. long, 6—20 m.M. broad, the rachis sulcate above, slightly flexuose, more or less extended at the apex into a naked tail and often bearing a young plant. Pinnae numerous, sessile from a broad base, in the smaller fronds suborbicular and simply crenate, in the larger ones lanceolate-oblong and pinnatifid, cuneato-excised below, the superior and inferior basal auricles 3—4-dentate or crenate, the other lobes 2-dentate or entire; superior pinnae gradually smaller, uppermost ones often very minute but fertile (with 1—2 sori); inferior pinnae gradually smaller, subrotund or flabelliform, always fertile. Texture membranaceous; surfaces minutely hairy when young, at length naked; veins obscure, the apex thickened. Sori of the larger pinnae arranged in a double row, 1 row of 4—5 on each side of the costa and approximate to it, superior auricle bearing 1 ± diverging sorus, rarely 2, sometimes none, lower auricle generally barren, rarely bearing 1—2 sori.

Luzon; Centr. China, South. India.

* (24) **A. normale**, *Don.*, Hk. Bk., Syn. Fil., 197; *A. multijugum*, *Wall.*, Hk., Spec. Fil., III, tab. CLXXXVIII; *Bedd.*, Ferns S.I., tab. CXXXIII.

Stipes tufted, 10—15 c.M. long, blackish, polished. Fronds 20—30 c.M. long, 2½—4 c.M. broad, pinnate. Pinnae in 12—∞ pairs, to 2 c.M. long, the lower ones to ⅔ c.M. broad, deflexed, the apex obtuse, the upper and outer edge inciso-crenate, the upper base truncate and ± auricled, the lower narrowly cuneate, cut away in a ± horizontal line which extends nearly the length of the lower margin of the pinna. Texture rather firm-membranaceous to subcoriaceous; rachis glossy, nearly black; veins simple or forked. Sori oblong or linear-oblong, in 2 unequal, opposite, parallel rows. — A Philippine form has the stipes 2—6 c.M. long and the fronds 10—25 c.M. long, 1—2 c.M. broad.

Malacca, Sumatra, Celebes, Philippines; China, North. India, Hawaii.

(25) **A. subavenium**, *Hk.*, Hk. Bk., Syn. Fil., 198; Hk., Spec. Fil., III, tab. CLXXXI, B; *Bedd.*, Ferns Br.I., II, tab. CCLXXXVIII.

Stipes tufted, 5—8 c.M. long, wiry, blackish, densely fibrillose. Fronds 20—30 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, with very numerous close-placed, horizontal pinnae on each side, which are $\pm \frac{1}{2}$ —1 c.M. broad, blunt at the point, the edge slightly inciso-crenate, the upper side distinctly auricled and narrowed suddenly at the base, the lower narrower below, but nearly equal upwards. Texture coriaceous; veins faint; a distinct midrib only in the lower part of the pinnae; rachis densely fibrillose, especially below. Sori 2—4 on each side, oblong, oblique. — The Malacca plant is perhaps a form of *A. pellucidum*, *Lam.* and different from the Madagascar fern.

Malacca; Madagascar.

* (26) **A. salignum**, *Bl.*, Enum., 175.

Stipes \pm tufted or approximate, greyish-green, 10—20 c.M. long, deciduously scaly in the lower half. Fronds lanceolate, 10—15 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, narrowed towards both ends, simple or with 1—3 pairs of similar, distant but smaller pinnae at the base. Pinnae 6—10 c.M. long, 1—2 c.M. broad, stalked, acuminate, the edge entire, crenate or toothed, but more conspicuously towards the point, the base cuneate. Texture coriaceous or subcoriaceous; under surface covered with few, scattered, adpressed, deciduous scales; veins oblique, simple or forked. Sori whether or not reaching the margin and the costa; indusium persistent.

Java.

* (27) **A. diversifolium**, *Bl.*, Rac., Flor. Btz., I, 211; *A. salignum*, *Hk.* (not *Bl.*), Spec. Fil., III, tab. CLXV; *Hk. Bk.*, Syn. Fil., 199.

Rhizome creeping, clothed with long, lanceolate scales. Stipes approximate, 10—20 c.M. long, slender, slightly scaly below. Fronds 15—50 c.M. long, 15—20 c.M. broad, with a long, linear terminal pinna and 1—5 \pm opposite pairs of lateral ones, which are suberecto-patent, straight or subfalcate, \pm 2 c.M. broad, with the point long-acuminate, the margin faintly crenate or entire, the base cordate. Texture coriaceous; rachis and both surfaces green and naked; veins parallel, \pm distant, obscure. Sori reaching from the midrib nearly to the edge.

Java, New Guinea.

(28) **A. Labillardieri**, *Hæ*, Mett., Farng., VI, Aspl., 95, tab. IV, fig. 11.

Stipes livid, narrowly marginate, afterwards naked. Fronds \pm 30—40 c.M.

long, ovate-oblong, pinnate. Pinnae 12—15-jugate, obliquely patent, 10—13 c.M. long, 2—2½ c.M. broad, linear-lanceolate or elongate-oblong, acuminate, crenulate-serrulate, the base somewhat obliquely cuneate (somewhat broader and slightly rounded on the upper side). Texture membranaceous, thin; surfaces naked; veins erecto-patent, simple or forked at the base, translucent. Sori distant (1—5 sterile veins between 2 sori), narrow, 10—15 m.M. long, reaching nearly the costa and the margin; indusium thin, narrow.

Buru Island.

* (29) **A. remotum**, *Moore*, C. Chr., Ind. Fil.; *A. multilineatum*, *Hk.*, Hk. Bk., Syn. Fil., 199; Hk., Spec. Fil., III, tab. CLXXXIII.

Rhizome creeping. Stipes scattered, firm, erect, greyish-green, ± 15—20 c.M. long. Fronds oblong-lanceolate, 30—50 c.M. long, 15—20 c.M. broad, with a long terminal pinna and 8—18 lateral ones on each side. Pinnae stalked, erecto-patent, 10—15 c.M. long, 1—2 c.M. broad, acuminate, the edge faintly toothed, but more acutely towards the point, the base rather broadly cuneate. Texture slightly fleshy; colour dark-green; rachis and surfaces naked; veins mostly simple, rather spreading, close. Sori numerous, close, reaching from the midrib nearly or quite to the edge.

Java, Celebes, Philippines, New Guinea; Polynesia.

* (50) **A. subaquatile**, *Ces.*, Fel. di Bor., 20, tab. III, fig. 5, a—b.

Rhizome short, tomentose. Stipes numerous, erect, slender, greenish or brownish, naked, 2—10 c.M. long. Fronds 15—30 c.M. long, simply pinnate. Pinnae stalked, alternate or subopposite, erect or erecto-patent, linear, 2½—7½ c.M. long, 2½—5 m.M. broad, the apex narrow, the edge deeply and obliquely toothed with distant, acute teeth, the base narrowed gradually into the stalk. Texture herbaceous; surfaces naked; rachis often slightly winged in the upper part; costae wavy; veins simple, 1 to each tooth, very oblique. Sori short, 1 to each tooth, subparallel and nearly close to the costae.

Borneo.

(31) **A. sumatranum**, *Hk.*, Hk., Bk., Syn. Fil., 199; Hk., Spec. Fil., III, tab. CLXVIII.

Stipes 15—25 c.M. long, firm, erect, grey, naked. Fronds ± 45 c.M. long, ± 15 c.M. broad, with a long, narrow, linear terminal pinna and 6—12 erecto-patent lateral ones on each side, which are ± 15 c.M. long, ± 2 c.M. broad, the point acuminate, the lower part obscurely undulate, the upper

crenate, the base equally cuneate on both sides; upper pinnae connected by a more or less narrow wing. Texture very coriaceous; veins oblique, often once forked. Sori long, reaching nearly from the midrib to the edge.

Sumatra.

* (32) **A. vulcanicum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 215; *A. heterodon*, *Mett.* (not *Bl.*), *Hk.*, *Spec. Fil.*, III, 107; *A. Wightianum*, *Wall.*, *Hk.*, l. c., tab. CLXVII.

Rhizome creeping or erect, the scales dense, brown, lanceolate. Stipes 15—25 c.M. long, erect, naked or deciduously scaly towards the base. Fronds 50—100 c.M. long, 10—20 c.M. broad, oblong, with a linear terminal pinna or a proliferous apex and 6—12 lateral pinnae on each side. Pinnae erecto-patent, to 15 c.M. long, 1—3 c.M. broad, all or only the lower ones on stalks to 1½ c.M. long, the apex acute or acuminate, the edge entire or more or less distinctly crenate to denticulate, the base varying from narrowly to broadly cuneate, equal on both sides, or the upper base sometimes a little broader than the lower and with a tendency to become slightly auricled. Texture coriaceous or subcoriaceous; rachis naked; veins simple or forked. Sori regular, linear, in 2 opposite rows, reaching from the midrib nearly or ½-way to the edge. — A variable plant.

Malaya; Ceylon, South. India.

(33) **A. fuliginosum**, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 205; *A. lugubre*, *Hk.*, 2nd Cent. of Ferns, tab. III.

Stipes tufted, very short, densely clothed with blackish, fibrillose scales. Fronds 15—50 c.M. long, 5—7½ c.M. broad, with 12—24 broadly adnate, horizontal, subfalcate pinnae on each side. Pinnae 1—1½ c.M. broad, the point acute, the edge slightly toothed, the base a little narrowed on both sides in the lower ones, which are gradually reduced in size. Texture thin; rachis densely fibrillose; veins usually once forked. Sori in 2 regular rows near the point of the pinnae.

Borneo.

(34) **A. horizontale**, *Bk.*, *Ann. of Bot.*, VIII, 125.

Stipes 20—25 c.M. long, slender, green, clothed with minute scales, the lower of which are larger, dense, lanceolate, acuminate, pale-brown. Fronds oblong, ± 50 c.M. long, caudate, pinnate. Pinnae ± 9, lanceolate, 7—15 c.M. long, 2½—5 c.M. broad, acuminate, obscurely crenate, horizontal, sessile, the lowest the largest, cordate at the base, auricled on the upper side. Texture

membranaceous: rachis green, slender, slightly scaly; surfaces naked; veins nearly spreading, often forked at the base. Sori medial, linear; indusium broad, membranaceous, naked.

Sumatra.

* (35) **A. longissimum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 212; *Bedd.*, *Ferns Br.I.*, I, tab LXIII; *Hk.*, *Spec. Fil.*, III, tab. CXC; *A. erectum*, *Bl.* (not *Bory*), *Enum.*, 178; *A. Belangeri*, *Bory*, *Bél.*, *Voy. Bot.*, II, 47.

Rhizome creeping, the scales lanceolate, long-acuminate, brown. Stipes approximate, 5—30 c.M. long, erect, dark-coloured, clothed with ciliate scales. Fronds 50—200 c.M. long, 4—15 c.M. broad, lanceolate-elongate, often proliferous at the point, narrowed gradually towards the base. Pinnae numerous, spreading, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, acuminate or subacute, the edge \pm toothed, the base subtruncate, auricled on one or both sides; higher pinnae not rarely unequal-sided at the base. Texture herbaceous to subcoriaceous; rachis dark-coloured, slightly chaffy; veins oblique, the lower however more spreading, forked. Sori very oblique, reaching from the midrib to $\frac{1}{2}$ -way or quite to the margin.

Malaya; North. India, Mauritius, Juan Fernandez.

(36) **A. monotis**, *Christ*, *Schum. & Laut.*, *Flor. deut. Sch.geb.*, *Suppl.*, 43.

Rhizome slender, climbing, yellowish-greenish, naked or sparingly covered with minute, subulate, blackish scales. Stipes distant, 2—8 c.M. long, greenish-stramineous, slightly tinged with red like the rachis. Barren fronds to 12 c.M. long, elongate-lanceolate, long-acuminate, pinnate, narrowed gradually towards the base. Pinnae close, short-stalked, 17—25 on each side, broadly cuneate-oblong, the upper base 2-partite with an ovate-cuneate lobe or auricle, the edge crenate or 2-crenate; lower pinnae growing smaller, distant. Texture thinly herbaceous; surfaces naked; veins simple or forked, blackish. Fertile fronds to 30 c.M. long, the pinnae larger, distant, deeply inciso-dentate, the teeth acute, curved. Sori 1 to each vein, reaching nearly from the costa to the margin; indusium persistent.

New Guinea.

* (37) **A. borneense**, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 203; *Hk.*, *Spec. Fil.*, III, tab. CLXXXVI.

Rhizome short, the scales lanceolate-subulate, brown. Stipes clustered, 2—10 c.M. long, greyish, slightly fibrillose. Fronds 25—90 c.M. long, $2\frac{1}{2}$ —

7½ c.M. broad, with numerous sessile or short-stalked pinnae, the lower ones growing smaller very gradually, the central ones spreading, ± close, 4—7 m.M. broad, the point bluntly rounded, the upper and outer edge incise-lobate to toothed, the upper base broadly cuneate, ± auricled, the lower very narrowly cuneate. Texture herbaceous to subchartaceous; rachis firm, greyish, nearly naked; veins oblique, inconspicuous or more distinctly visible, those of the upper base subflabellate. Sori few, those on the lower side nearly parallel to the costa.

Borneo, Malacca.

* (38) **A. tenerum**, *Forst.*, *Rac.*, *Flor. Btz.*, I, 215; *A. elongatum*, *Sw.*, ? *Bedd.*, *Ferns S.I.*, tab. CCXXIV; — var. *acuminatum*, *Hk.*, *Spec. Fil.*, III, 117; *A. Doreyi*, *Kze.*, *Anal.*, 23.

Rhizome erect, clothed with lanceolate, brown scales at the apex. Stipes tufted, firm, erect, greyish, nearly naked, 5—25 c.M. long. Fronds 15—50 c.M. long, ± 5½—10 c.M. broad, lanceolate-oblong, with 10 to numerous sessile or ± stalked, horizontal pinnae on each side. Pinnae oblong or linear-oblong, ¾—2 c.M. broad, the sides ± parallel in the typical form, the apex bluntly rounded or subtruncate, the edge deeply crenato-dentate throughout; the upper base ± truncate, slightly auricled, the lower cuneate. Texture herbaceous, not rarely firm; rachis naked or slightly scaly; surfaces naked; veins commonly simple. Sori numerous, regular, linear-oblong, parallel, not reaching either the midrib or the edge.

Var. acuminatum: Pinnae acute or nearly acuminate at the point.
Trop. Asia.

* (39) **A. lunulatum**, *Sw.*, *Hk. Bk.*, *Syn. Fil.*, 202; *A. erectum*, *Bory.*, ? *Hk.*, *Fil. Exot.*, tab. LXXII, (var. *proliferum*).

Rhizome short, slightly scaly. Stipes tufted, nearly naked, grey or dark-coloured. Fronds 15—45 c.M. long, ± 3½—4½ c.M. broad, linear-lanceolate, with 10 to numerous horizontal pinnae on each side. Pinnae close, subtrapezoid-oblong, narrowed towards the bluntish or acute apex in the type, ½—1 c.M. broad, the edge incise-crenate throughout; the upper base truncate, slightly auricled, the lower cuneate. Texture herbaceous; rachis naked like the surfaces; veins mostly simple. Sori short, oblong, falling short of both edge and midrib. — Near *A. tenerum*, *Forst.*, but different from it by the less blunt pinnae.

Pantropical.

(40a) **A. Steerei**, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 28.

Stipes dark-grey, 12—15 c.M. long, firm, provided with some few small, brown, scattered scales. Fronds 15—25 c.M. long, 5—6½ c.M. broad, oblong-lanceolate. Pinnae 12—15 on each side, short-stalked, 3½—5 c.M. long, 1—1½ c.M. broad, the apex acute, the margin coarsely crenate, the upper base abrupt with a more or less distinct, large lobe, the lower base cuneate; lower pinnae slightly deflexed. Texture firm-herbaceous; colour dark-green above, brownish beneath; veins distant, simple or once forked. Sori 2—8 in a row on each side of the costa, regular, not reaching midrib and margin.

Philippines.

(40b) **A. Meyenianum**, *Pr.*, Christ, Bull. Herb. Boiss., 2^e Série, VI, 999; *A. pellucidum*, *Lam*, C. Chr., Ind. Fil., p.p.

Rhizome firm, erect, clothed with black, fimbriate, subulate scales. Stipes tufted, black-rufous, 8—12 c.M. long, hispid with short scales like the ebeneous rachis. Fronds to 45 c.M. long, 4—6 c.M. broad, linear, acuminate, the base not or moderately narrowed. Pinnae numerous, subremote, 45—50 on each side, sessile, falcato-rhomboidal, acuminate, 2½—3 c.M. long, the upper base truncate, the lower cuneate, auricled, the edge coarsely 2-dentate; lower pinnae more distant, more blunt. Texture herbaceous, impellucid; upper surface dark-, lower pale-green; veins hidden. Sori very oblique, ± 5 on each side, narrow, reaching the margin and the costae; indusium narrow, brown.

Luzon, Palawan.

(41) **A. prionurus**, *J. Sm.*, Hk. Bk., Syn. Fil., 202.

Stipes tufted, 15—25 c.M. long, firm, erect, slightly scaly below. Fronds 30—45 c.M. long, 15—25 c.M. broad, oblong, with 9—18 pinnae on each side, which are ± 1 c.M. or more broad, very much acuminate at the point, deeply toothed throughout, the base rather unequal, more broadly cuneate on the lower side, with a short petiole on the lower ones. Texture herbaceous; veins usually simple. Sori reaching nearly to the edge.

Philippines.

(42) **A. griseum**, *Copel.*, Polypod. Philipp., 82; *A. insigne*, *Copel.*, Perk., Fragm. Philipp., 185.

Rhizome short, densely clothed with large, brown scales. Stipes firm, slightly chaffy, 40—50 c.M. long, reddish-brown or blackish-green. Fronds 60 c.M. and more long, ± 50 c.M. broad, pinnate. Pinnae 12 or more on

each side, stalked, \pm 20 c.M. long, \pm $3\frac{1}{2}$ c.M. broad, lanceolate, acuminate, subserrate, the upper base broadly and the lower more narrowly cuneate. Texture tough-herbaceous; surfaces naked; veins oblique, often forked, \pm conspicuous. Sori broad, occupying the lower half of the veins, moderately regular.

Mindanao.

(43) **A. persicifolium**, *J. Sm.*, Hk. Bk., Syn. Fil., 202.

Stipe and rachis grey, with few, scattered, minute, grey scales. Fronds oblong-lanceolate, 60—90 c.M. long, often proliferous at the apex. Pinnae distant, ascending, 15—30 on each side, subpetiolate, 10—13 c.M. long, linear-ligulate, acuminate, \pm 1—2 c.M. broad, distinctly inciso-crenate throughout, obliquely cuneate at the base, the lower base narrower; lower pinnae not reduced. Texture thin but firm; surfaces naked; veins fine, forked, not close. Sori regular, 7—10 m.M. long, reaching nearly to midrib and edge.

Celebes, Borneo, Sulu Islands, Philippines.

* (44) **A. Schoggersii**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 14.

Stipes tufted, \pm 4 c.M. long, dark-brown, clothed like the rachis with fine, deciduous scales. Fronds ovate-lanceolate, 8—10 c.M. long. Pinnae \pm 15 on each side, spreading, the lowest shorter, the central ones \pm 2 c.M. long, \pm $\frac{1}{2}$ c.M. broad, the point acute or bluntish, toothed, the edge lobed $\frac{1}{3}$ — $\frac{1}{2}$ -way down, the lobes truncate or rounded and crenate, oblique like the teeth, the base obliquely truncate-cuneate. Texture coriaceous; under surface provided with few, very minute, deciduous, fimbriate scales; veins few, oblique, simple or forked. Sori very small, subparallel and nearly close to the costa, occupying the base of the veins.

Java (Preanger Regencies, Lieut. Col. J. C. F. Schoggers).

* (45) **A. caudatum**, *Forst.*, Rac., Flor. Btz., I, 218; *A. contiguum*, *Bedd.*, Ferns S.I., tab. CXL; *A. truncatilobum*, *Fée*, Christ, Philipp. Journ., II, 164; *A. arayatense*, *Christ*, MS.

Rhizome erect, the scales linear-lanceolate, brown. Stipes tufted, firm, erect, densely clothed with fine, fibrillose, brown scales. Fronds 25—100 c.M. long, 10—25 c.M. broad, with 20—30 pinnae on each side. Pinnae spreading, 1— $2\frac{1}{2}$ c.M. broad, the apex long-acuminate, toothed, the edge cut to \pm $\frac{1}{2}$ -way down or sometimes more at the base into rounded, entire or crenate, oblique

lobes, the base broadly cuneate and auricled on the upper and more narrowly cuneate on the lower side. Texture coriaceous; rachis deciduously scaly; veins very oblique, Sori linear, the higher subparallel and nearly close to the costa, occupying the lowest part of the veins. — Hardly different from *A. horridum*, *Klf.* in the deeply lobed forms.

Pantropical.

* (46) **A. contiguum**, *Klf.*, Hk. Bk., Syn. Fil., 205; Hk., Spec. Fil., III, tab. CXCIV, fig. 1; — var. *lepturus*, Hk., l.c., 156, tab. CXCIV, fig. 3; *A. lepturus*, *J. Sm.*, Mett., Farng., VI, Aspl., 152; — var. *bipinnatifidum*, Christ, Bull. Herb. Boiss., VI, 152; *A. laxivenum*, *Copel.*, Philipp. Journ., II, 152; ? *A. insilitium*, *Brack.*, Christ, Philipp. Journ., II, 166.

Rhizome creeping or ascending. Stipes 15—25 c.M. long, firm, erect, dark-coloured, naked. Fronds lanceolate, acuminate, pinnate, 15—30 c.M. long, the apex pinnatifid. Pinnae 10—50-jugate, subfalcate, short-stalked, 4—7½ c.M. long, ± 1—1½ c.M. broad, acute or acuminate, the edge obliquely and broadly serrate, the lower serratures sharply serrate or toothed, the higher sharp, simple or 2-dentate, the upper base cuneate, sometimes ± auricled, the lower narrowly cuneate in a curve. Texture ± coriaceous, rachis firm, naked or slightly villose, brown; veins oblique. Sori close, subparallel and nearly close to the costa.

Var. lepturus: Pinnae longer, 9—12½ c.M. long, the edge deeply serrate, rather lobed, the lower lobes or serratures sharply serrate, the apex long-acuminate.

Var. bipinnatifidum: Lower pinnae pinnate at the base; pinnulae nearly free, cuneato-rotundate, auricled, deeply toothed.

Java, New Guinea; Hawaii; — var. *lepturus* & *bipinnatifidum*: *Philippines*.

* (47a) **A. nigrescens** *Bl.*, Mett., Farng., VI, Aspl., 151; ? *Rac.*, Flor. Btz., I, 217.

Rhizome and stipes? Fronds 50—90 c.M. long, elongate-oblong, pinnate, the apex pinnatifid. Pinnae numerous, spreading, 10—15 c.M. long, linear, short-stalked, acuminate, the edge broadly and shallowly crenate, but often more deeply towards the apex, the serratures 2—4-dentate except the higher ones, which are often entire and narrow, the upper base broadly and roundly, the lower narrowly cuneate. Texture chartaceous; rachis naked or palea-ceo-villose, dark-coloured; veins repeatedly forked. Sori oblique, subparallel and nearly close to the costa. — Scarcely different from *A. contiguum*, *Klf.* except

in the size of the fronds and the length of the pinnae. The plant found and determined by Raciborski is probably a young form of the latter.

Java, Moluccas.

(47b) **A. militare**, Copel., Philipp. Journ., I, Suppl., 254.

Rhizome creeping, densely clothed with lanceolate, murino-fuligineous, acuminate scales. Stipes 40—50 c.M. long, castaneous, glossy and glabrescent like the rachis. Fronds 60—100 c.M. long, 25—50 c.M. broad. Pinnae short-stalked, remote, 15—18 c.M. long, 2—2½ c.M. broad, the base obliquely cuneate, or truncate on the upper side, the edge serrate, 2-serrate towards the caudate apex; lower pinnae hardly shorter. Texture chartaceous; surfaces naked, the upper rather fuligineous, the lower olive-green; veins forked, spreading, very oblique at the base. Sori curved, costal.

Mindanao.

* (48) **A. pellucidum**, Lam., Rac., Flor. Btz., I, 212; *A. hirtum*, Klf., Christ, Farnkr. d. Erde, 196; Bedd., Ferns Br.I, II, tab. CXCIV; *A. approximatum*, Bl., Enum., 179; — var. *obtusum*, Christ, Ann. Btz., XX, 111, — var. *repressum*, Copel., Polypod. Philipp., 85.

Rhizome erect or suberect, densely clothed with dark-brown, linear, acuminate scales. Stipes varying from short to 10 c.M. long, strong, erect, brownish, deciduously scaly, the scales ± fibrillose. Fronds 20—100 c.M. long, 4—20 c.M. broad, lanceolate. Pinnae numerous, sessile, spreading, acute or acuminate, the edge serrate or toothed, the teeth sometimes toothed again, the upper base truncate, auricled, the lower cuneate; lower pinnae reduced, suborbicular or subcordate. Texture coriaceous; rachis fibrillose or chaffy, the scales dense, dark-coloured, fimbriate, at length deciduous; veins oblique, inconspicuous. Sori short, in regular rows, oblique, occupying the lower half of the veins.

Var. obtusum: Pinnae short, obtuse.

Var. repressum; Fronds under 20 c M. long, almost sessile: pinnae obtuse.

Mataya; Polynesia, Madagascar, Mascarenes, Seychelles.

(49) **A. dimidiatum**, Sw., Hk. Bk., Syn. Fil. 209; ? *A. Zamiae-folium*, Willd., Kze, Farng.; I, tab. XLVIII.

Stipes tufted, 7½—50 c.M. long, blackish, slightly paleaceous, firm, erect. Fronds 15—45 c.M. long, 8—15 c.M. broad, with 6—9 opposite pairs of pinnae. Pinnae ± 2—2½ c.M. broad, the point acuminate, especially the

outer and upper edge sharply inciso-serrate, the upper base rounded or cuneate, the lower very narrowly cuneate. Rachis dark-coloured, firm, slightly scaly; veins close, flabellate; no midribs. Sori radiant, narrow, long. — Resembles *A. adiantoides*, *C. Chr.*, but the pinnae suddenly narrowed from a broad, unequally auricled base into a narrow point.

New Guinea, Timor, Malacca; Tonkin, West. Africa, Trop. America.

(50) **A. Cesatianum**, *Bk.*, Becc., Mal., III, 59; *A. Wightianum*, *Ces.* (not *Wall.*), Fel. Becc. Polin., 4.

Stipes \pm 30 c.M. long, greyish, scaly at the base, the scales few, large, lanceolate-oblong, acuminate, brown. Fronds oblong-lanceolate, simply pinnate, \pm 45 c.M. long. Pinnae multijugate, distant, lanceolate, erecto-patent, crenulate, obscurely stalked, obliquely cuneate at the base, the lower base reduced. Texture firm; surfaces naked; veins conspicuous, erecto-patent, not closely placed, commonly deeply forked. Sori oblong, parallel, medial on the veins; indusium firm, naked, persistent.

New Guinea.

(51) **A. crinicaule**, *Hance*, Hk. Bk., Syn. Fil., 208; *A. Hancei*, *Bk.*, Bot. Zeit., 1869, 130; *A. falcatum*, *Bedd.*, Ferns S.I., tab. CXXI.

Stipes densely tufted, 7—10 c.M. long, erect, dark-brown, fibrillose. Fronds 15—90 c.M. long, 4—12 c.M. broad, with 9—25 horizontal pinnae on each side, which are 4—10 m.M. broad, the point bluntish, the edges irregularly inciso-crenate, the upper base narrowed suddenly, sometimes auricled, the lower obliquely cuneate, sometimes auricled. Texture subcoriaceous; rachis firm, fibrillose; midribs not distinct; veins subflabellate. Sori few, linear, oblique, regular, not reaching to the edge.

Philippines; Br. India, South. China.

* (52) **A. acutiusculum**, *Bl.*, Rac., Flor. Btz., I, 217; *A. anisodontum*, *Pr.*, Hk. Bk., Syn. Fil., 205.

Rhizome erect, the apex clothed with narrow, lanceolate, brown scales. Stipes tufted, firm, erect, \pm scaly, to 30 c.M. long. Fronds 50—70 c.M. long, with numerous sessile or stalked, spreading or subfalcate pinnae on each side. Pinnae acute or acuminate, the lowest hardly smaller, rather larger than the next following ones, 4—10 c.M. long, $\frac{3}{4}$ — $1\frac{1}{2}$ c.M. broad near the base, the edges more or less inciso-crenate, the upper base broadly cuneate or subtruncate, \pm auricled or deeply incised, the lower rather

narrowly cuneate. Texture firm; rachis nearly naked; veins oblique, mostly once forked or the lower acroscopical to 4× branched. Sori in regular rows, reaching nearly the midrib and the edge.

Java, Borneo, Celebes, Philippines, New Guinea; Samoa.

(53) **A. impressum**, *Christ*, C. Chr., Ind. Fil.; *A. canaliculatum*, *Christ*, Warb., Mons., I, 73.

Like *A. contiguum* *Klf.* but smaller. Rachis densely strigoso-paleaceous, at length nearly naked. Pinnae shorter, broader, ovate-cuneate, not 2-aucled, caudato-acuminate, 2-serrate. Veins prominent above, subflabellate. Sori not close to the costae, more spreading, elongate, distant.

Batjan.

(54) **A. Gardneri**, *Bk.*, Hk. Bk., Syn. Fil., 485; *Bedd.*, Ferns S. Br.I., Suppl., tab. CCCLV.

Rhizome short-creeping. Stipes grey, 5—15 c.M. long, with a few small, scattered scales. Fronds lanceolate, 15—30 c.M. long, 7—13 c.M. broad, often gemmiferous; rachis not winged. Pinnae 8—10-jugate, distant, ascending, lanceolate, subpetiolate, acute, central ones 5—8 c.M. long, 1—1½ c.M. broad, toothed, the base obliquely and rather broadly cuneate. Texture thin but firm, supple; surfaces naked; veins erecto-patent, forked, moderately close. Sori regular, parallel, 5—6-jugate, nearer the midribs than the edge, 6—12 m.M. long. — Closely allied to *A. macrophyllum*, *Sw.*

Sumatra; Ceylon, Khasya.

* (55) **A. adiantoides**, *C. Chr.*, Ind. Fil.; *A. falcatum*, *Lam.*, Hk. Bk., Syn. Fil., 208; *Diels*, in *Engl. & Prantl*, Nat. Pfl. Fam., I⁴, fig. 126, G; *A. caudatum*, *Bedd.*, Ferns S.I., tab. CXLIII; *A. zamiaefolium*, *Pr.*, Rel. Haenk., I, 45; *Trichomanes adiantoides*, *L.*, (oldest name).

Stipes tufted, 15—30 c.M. long, erect, grey or brown, naked or slightly scaly. Fronds 15—60 c.M. long, 10—25 c.M. broad. Pinnae horizontal, 6—20 on each side, stalked, alternate or opposite, acute or caudate, 1—2½ c.M. broad, the edge toothed or lobed, the teeth or lobes toothed again, the upper base broadly and the lower narrowly cuneate, the lower sometimes slightly cut away. Texture coriaceous; rachis naked or nearly so; veins very oblique. Sori long, in 1—2 irregular lines, reaching nearly to the edge. — Distinct enough from *A. macrophyllum*, *Sw.* if typical specimens only are shown but gradually passing into it (*Bedd.*).

Trop. Asia, Australia, Polynesia, New Zealand, East. Africa.

(56) **A. Sancti-Christophori**, *Christ*, Schum. & Laut., Flor. deut. Sch.ggeb., 128.

Like *A. macrophyllum*, *Sw.*, but the rachis strigoso-squamulose, the pinnae more numerous, (12—15 on each side), smaller, broadly rhomboidal, inequilateral, auriculate on the upper side, alternate, nearly sessile, growing shorter towards the pinnatifid apex of the frond, the margin not incised but copiously and minutely 2-serrate.

New Guinea.

* (57) **A. macrophyllum**, *Sw.*, Rac., Flor. Btz., I, 216; *Hk.*, Spec. Fil., III, tab. CXCVI—CXCVII; *Bedd.*, Ferns S.I., tab. CXLII; *A. canaliculatum*, *Bl.*, Enum., 180; — var. *minus*.

Rhizome short, the extremity clothed with long, narrow scales. Stipes ± tufted, 15—50 c.M. long, stout, erect, brownish, nearly naked. Fronds 15—80 c.M. long, 12—50 c.M. broad. Pinnae ± 6—12 on each side, 2½—7½ c.M. broad, the terminal one often large, not rarely 5-fid, the point of all acute, the edge mostly sharply and irregularly toothed, not or slightly lobed, the base obliquely and broadly cuneate, the lower base narrower than the upper. Texture ± coriaceous; rachis naked or deciduously chaffy, firm; veins oblique, *distinct* beneath. Sori long, oblique, often reaching from the midrib to the edge. — Near *A. adiantoides*, *C. Chr.*, and scarcely different from the latter in the smaller forms.

Var. minus: Fronds smaller and narrower; lateral pinnae 1—2-jugate.

Malaya; Polynesia, Ceylon, South. India, Comores, Mascarenes; — var. minus: Timor (Forbes, No. 5124^a).

* (58) **A. paradoxum**, *Bl.*, Rac., Flor. Btz., I, 216; *A. zamioides*, *Hk.*, *Bedd.*, Ferns Br.I., II, tab. CXCIII.

Rhizome short-creeping or ascending, the scales short, broad, deltoid, adpressed. Stipes firm, erect, 15—25 c.M. long or longer, greyish to brownish, slightly scaly. Fronds 50—100 c.M. long, 15—25 c.M. broad, with a rather large, oblong-rhomboidal, simple or 5-fid terminal pinna and 8—12 lateral ones on each side, the lowest the largest, subrhomboidal, 2½—5 c.M. broad, mostly entire, crenate or toothed towards the apex, the upper base suddenly narrowed, broadly cuneate, the lower more narrowly cuneate. Texture coriaceous; rachis firm, ± scaly; veins *indistinct* beneath, rather very oblique,

commonly once forked. Sori reaching nearly from the midrib to the edge, sometimes accompanied by smaller ones, which are \pm terminal on the veins, sometimes \pm scolopendrioid. — United with *A. macrophyllum*, Sw. by intermediate forms, but distinguished by the indistinct, hardly visible veins.

Malacca, Sumatra, Java, Borneo.

* (59) **A. heterocarpum**, *Wall.*, Hk. Bk., Syn. Fil., 210; Hk., Spec. Fil., III, tab. CLXXV; Bedd., Ferns S.I., tab. CXXXI; *A. cheilosorum*, Kze, Mett., Farng., VI, Aspl., tab. V, fig. 12—13; C. Chr., Ind. Fil.

Rhizome creeping, scaly at the apex. Stipes firm, erect, brown, polished. Fronds 15—40 c.M. long, $5\frac{1}{2}$ —5 c.M. broad, with numerous closely placed pinnae on each side. Pinnae horizontal, subsessile, $\pm \frac{1}{2}$ — $\frac{2}{3}$ c.M. broad, dimidiate, the lower side being entirely cut away, the upper and outer edge deeply incised, the crenations crenate or toothed again, the upper base nearly truncate; lower pinnae hardly shorter and more or less deflexed. Texture pellucido-herbaceous; rachis like the stipe. Sori 1 or rarely 2 together in the teeth.

Malacca, Borneo, Mindoro; Formosa, North. India.

* (60a) **A. unilaterale**, *Lam.*, Bedd., Handb. Ind. Ferns, 152; *A. resectum*, Sm., Rac., Flor. Btz., I, 214; Hk. & Grev., Ic. Fil., tab. CXIV; Bedd., Ferns, S.I., tab. CXXXII; *A. amoenum*, Pr., Mett., Farng., VI, Aspl., tab. V, fig. 11; — var. *delicatulum*, Bedd., Handb. Ind. Ferns, 155; Ferns S.Br.I., Suppl., tab. CCCLVIII.

Rhizome slender, wide-creeping, the scales small, nearly linear. Stipes scattered, 1—2 c.M. apart, slender, naked, dark-coloured, \pm black-brown and polished like the rachis, some of them sometimes grey and opaque, 10—20 c.M. long. Fronds 15—60 c.M. long, 4—10 c.M. broad, rarely narrower, lanceolate-oblong, acute to long-acuminate. Pinnae stalked or subsessile, 10—30 on each side, trapezoid-lanceolate, horizontal or erecto-patent, the apex bluntish or acute, the upper and outer edge inciso-crenate to dentate, the upper base \pm cuneato-subtruncate, $\frac{1}{2}$ — $\frac{2}{3}$ of the lower side cut away at the base. Texture thinly herbaceous; veins 1—2 \times forked. Sori oblong, occupying the base or the centre of the veinlets.

Var. delicatulum: Fronds and pinnae much smaller and narrower than in the type, the pinnae less cut away at the base on the lower side.

Trop. Asia, Japan, Polynesia, West. Africa and East African Islands; — var. delicatulum: Malacca; Tenasserim.

(60b) **A. filipes**, *Copel.*, Philipp. Journ., III, 54; — var. *minutum*, *Copel.*, l. c., 55.

Rhizome creeping, slender, scaly at the extremity. Stipes seriate, \pm 5 c.M. long, filiform, purple, glossy. Fronds 8—12 c.M. long, acuminate, pinnate. Pinnae \pm 15 on each side, the largest to $1\frac{1}{2}$ c.M. long, 5—4 m.M. broad, oblique or auriculate, the lower edge straight, the upper serrate; lower pinnae hardly reduced, spreading, the higher ones ascending. Texture membranaceous; surfaces naked. Sori few, short. — Near *A. unilaterale*, *Lam.*

Var. minutum: Stipes 1—5 c.M. long; fronds 4—7 c.M. long, diaphanous; sori 1 or sometimes 2 to each pinna.

Negros Island (Philippines).

* (61) **A. obscurum**, *Bl.*, Rac., Flor. Btz., I, 214; *A. serraeforme*, *Mett.*, Farng., VI, Aspl., tab. IV, fig. 15.

Rhizome long, slender, creeping, the apex clothed with short, lanceolate scales. Stipes distant, to 20 c.M. long, erect, greenish like the rachis, scaly at the base. Fronds to 50 c.M. long, to 12 c.M. broad, lanceolate, sometimes 2-pinnate at the base, with \pm 20—50 short-stalked pinnae on each side. Pinnae spreading to erecto-patent, the lowest to 6 c.M. long, to 2 c.M. broad, bluntish or acute, the upper and outer edge crenate to toothed, the upper base cuneate, $\frac{1}{3}$ — $\frac{1}{2}$ of the lower side cut away at the base. Texture thin, \pm hyaline, rachis and both surfaces naked; veins commonly once forked. Sori short, occupying the lower part of the anterior veinlets.

Java.

(62) **A. denticulatum**, *Bl.*, Enum., 186.

Fronds 2-pinnatifid. Pinnae petiolate, small, ovate in general outline, deeply 5-partite or pinnatifid. Lobes cuneate-oblong, obtuse, the apex obscurely denticulate. Texture coriaceous; rachis and costae beneath \pm fibrillose.

Java.

(65) **A. Warburgianum**, *Christ.*, Warb., Mons., I, 72.

Rhizome short, the scales black, subulate. Stipes tufted, \pm 10 c.M. long, clothed like the rachis with lanceolate, brown scales. Fronds \pm 15 c.M. long, \pm 4 c.M. broad, ovate-oblong, pinnate, acuminate, the base not narrowed. Pinnae \pm 5 c.M. long, erecto-patent, linear-lanceolate, deeply pinnatifid with 4—5 linear, cuneate, praemorse segments. Texture coriaceous; surfaces pilose; veins fine, parallel. Sori few, narrow; indusium thin, filiform.

Java.

* (64) **A. laciniatum**, *Don.*, Hk. Bk., Syn. Fil., 211; Hk., Spec. Fil., III, tab. CC, A.; Bedd., Ferns S.I., tab. CXLV.

Rhizome thick, ascending, the scales dense, blackish, subulate, rigid. Stipes tufted or subtufted, 10—20 c.M. long, grey or brown, partially and deciduously scaly. Fronds 15—50 c.M. long, 2½—7½ c.M. broad; main rachis deciduously scaly. Pinnae ± 12—20 on each side, spreading, stalked, ± 1—2 c.M. broad, oblong, narrowed towards the apex, cut down often quite to the costa in the lower part, the upper base truncate, the lower narrowly cuneate. Lobes erecto-patent, rhomboidal-oblong or cuneate, 2—5-fid, the superior basal one (auricle) the largest, broad-flabelliform. Texture ± coriaceous; veins oblique, those of the lower lobes flabellate. Sori few, the higher ones oblique, the lower radiating in the auricle, not reaching either the midrib or the edge.

Celebes; Ceylon, North India, Japan.

* (65) **A. Hagenii**, *v. A. v. R.*, Bull. Dép. Agr. I.N., 1908, XVIII, 14.

Rhizome, stipes and fronds beneath densely scaly when young; the scales linear-lanceolate, acuminate, caducous, the base dilated, the edge finely and shortly toothed, the point very fine, capillary, those of the rhizome and base of the stipes dark-brown, erecto-patent, the others smaller, spreading, crisped, those of the under surface of the fronds the finest, pale-brown. Stipes ± 12 c.M. long, erect, firm, dark-brown. Fronds ± 15 c.M. long or longer, lanceolate-oblong, acuminate. Pinnae subrhomboidal, erecto-patent or nearly horizontal, the lowest mostly the largest, 4—6 c.M. long, 1½—2 c.M. broad, blunt or acute to acuminate, stalked, unequal-sided, deeply pinnatifid or pinnate at the base. Segments 5—4 on each side, oblong or cuneate, erecto-patent, with the outer edge finely toothed, the lower ones ± flabellate-cuneate, with the outer edge flabellato-5-fid and finely toothed. Texture coriaceous; veins indistinct. Higher sori subparallel to the costa, the lower ± parallel to the edges of the segments on which they are placed; indusium membranaceous.

Sumatra (Hagen).

* (66) **A. horridum**, *Kuf.*, Hk. Bk., Syn. Fil., 211; Hk., Spec. Fil., III, tab. CXCIII; *A. caudatum*, *Forst.*, var. *horridum*, *Christ.*, Fil. Saras., I, 15.

Stipes strong, erect, brownish, fibrillose. Fronds 60—90 c.M. long, 12—50 c.M. broad, with numerous (20—50) spreading to erecto-patent pinnae on each side. Pinnae ± 1—2 c.M. broad at the base, narrowed gradually to an acuminate toothed point, cut down throughout more than halfway to the costa into ± uniform, broad lobes, the lobes with the outer edge more or less crenate or

toothed, the upper base broadly cuneate, truncate, cordate or broadly rounded, the lower cuneate in a curve. Texture subcoriaceous; rachis stout, clothed with reddish to blackish, woolly fibres; veins oblique, commonly once forked. Sori nearly parallel and close to the midrib, except those on the lobes which are more spreading. — Near *A. caudatum*, *Forst.*

Malaya.

* (67) **A. tenuifolium**, *Don.*, *Hk. Bk.*, *Syn. Fil.*, 220; *Hk.*, 2nd *Cent. of Ferns*, tab. XXIX; *Bedd. Ferns S.I.*, tab. CXXX.

Stipes tufted, 7—15 c.M. long, slender, naked, slightly ebeneous. Fronds 15—30 c.M. long, 7—13 c.M. broad, oblong-lanceolate, with numerous pinnae on each side, the lower ones 5—7 c.M. long, \pm 2 c.M. broad. Lower pinnulae subdeltoid, cut down to a narrowly winged rachis into spatulate segments \pm 3 m.M. broad, sharply incised on the outer edge. Rachis green, naked, slender; veins solitary. Sori 1—2 to each segment.

Malacca; Br. India.

* (68a) **A. stereophyllum**, *Hae*, *Bot. Zeit.*, 1848, 175; *Mett. Farnng.*, VI, *Aspl.*, 158.

Stipes 7—15 c.M. long, paleaceo-pilose, fusco-paleaceous at the base, afterwards naked. Fronds 12—30 c.M. long, 5—7 c.M. broad, lanceolate-deltoid, acuminate. Pinnae many, short-stalked, trapezio-oblong, mostly obtuse, approximate, subspreading, the lowest \pm the largest, the upper base truncate, the lower subdimidiate-cuneate. Pinnulae rather distant, cuneate or spatulate-linear, the apex incised or toothed, obtuse, the lowest \pm 3-partite, spreading, the higher ones erecto-patent, decurrent, confluent. Texture rigidly coriaceous; surfaces sparingly paleaceo-pilose; rachis subflexuose, inconspicuously marginate, opaque.

Java, Sumatra.

(68b) **A. Elmeri**, *Christ*, *Philipp. Journ.*, II, 164.

Rhizome short, crowned with castaneous, subulate scales. Stipes fasciculate, 7—12 c.M. long, rather deep-black-brown and clothed with spreading, subulate, brown scales like the rachis. Fronds oblong, acuminate, narrowed at the base, \pm 20 c.M. long, 6—8 c.M. broad, 2-pinnate to 3-pinnatifid. Pinnae short-stalked, horizontal, \pm 16—20 on each side, subremote, ovate-elongate, acuminate, the apex coarsely toothed, lanceolate, the rachis short-winged above. Pinnulae few, 2—3 on each side, short-stalked, ovate-

cuneate, obtuse, coarsely toothed, 2—3 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, the teeth ± 8 m.M. long, ± 5 m.M. broad, the lowest anterior lobe acute, approximate to the rachis. Texture herbaceous; colour black-green; veins flabellato-furcate. Sori 2—3 to a segment, lanceolate, ± 4 m.M. broad; indusium narrow, greyish, persistent.

Luzon.

* (69) **A. praemorsum**, *Sw.*, Christ, Farnkr. d. Erde, 204; *A. furcatum*, *Thb.*, Rac., Flor. Btz., I, 218; Bedd., Ferns S.I., tab. CXLIV; — var. *alpestre*, Christ, Ann. Btz., XV, 118; — var. *angustissimum*, Christ.

Rhizome wide-creeping, densely clothed with polished, lanceolate, brown scales. Stipes 8—25 c.M. apart, 10—20 c.M. long, firm, erect, clothed with deciduous, woolly hairs. Fronds 15—45 c.M. long, 7—15 c.M. broad, linear-lanceolate, with 12—20 pinnae on each side. Pinnae lanceolate-deltoid, short-stalked, 4—8 c.M. long, 2—2 $\frac{1}{2}$ c.M. broad, acute, cut down throughout nearly or quite to the costa into sessile segments. Segments linear- or oblong-cuneate, the outer edge rounded and sharply serrate; lower segments often deeply incised. Texture coriaceous; rachis firm, more or less fibrillose like the stipe; veins deep-channelled, \pm flabellate.

Var. alpestre: Stipes and fronds ± 12 c.M. long. Ultimate segments \pm ovate, deeply incised, the outer edge slightly crenate-dentate. Scales of the rhizome very acuminate.

Var. angustissimum: Fronds linear-lanceolate; pinnae erect, the lobes with the apices long-acuminate.

Tropical and subtropical.

* (70) **A. cuneatum**, *Lam.*, Rac., Flor. Btz., I, 219; — var. *subaffine*, Christ, Ann. Btz., XV, 117; — var. *oceanicum*, Christ, l. c.; — var. *tripinnatum*, Christ, Philipp. Journ., II, 165.

Rhizome suberect, clothed with brown, lanceolate scales. Stipes tufted, 12—50 c.M. long, scaly at the base. Fronds 15—100 c.M. long, ± 10 —25 c.M. broad or broader, deltoid-lanceolate, with numerous spreading pinnae on each side. Lower pinnae 2 $\frac{1}{2}$ —4 c.M. or more broad, lanceolate-deltoid, cut down to the rachis into distinct, oblong-cuneate; more or less rounded pinnulae with the outer edge inciso-dentate and the lower part entire or lobed or cut down nearly or quite to the costa. Texture thinly coriaceous; rachis firm. greyish to brownish, nearly naked; veins fine, visible above. Sori subflabellate. linear.

Var. subaffine: Ultimate segments obovate, nearly flabellate, 2 × as large.

Var. oceanicum: Texture firmly coriaceous; rachis often dark-brown or black.

Var. tripinnatum: Fronds 5-pinnatifid; segments short, ovate-cuneate.

Pantropical; — var. *subaffine*: *Celebes*; — var. *oceanicum*: *Malaya*; — var. *tripinnatum*: *Mindanao*.

* (71) **A. bulbiferum**, *Forst.*, Hk. Bk., Syn. Fil., 218; *A. bullatum*, *Wall.*, *Bedd.*, Ferns Br.I., I, tab. LXV; Hk., Icon. Pl., V, tab. CDXXIII.

Stipes tufted, 15—30 c.M. long, firm, erect, greyish, scaly at the base. Fronds 50—60 c.M. long, 20—30 c.M. broad, oblong-deltoid, with many pinnae on each side, erect or pendulous. Pinnae horizontal (erecto-patent after Beddome's drawing), the largest 10—20 c.M. long, 3½—5 c.M. broad, the apex acuminate, toothed or subpinnatifid, the rest cut down to a compressed, winged rachis into many lanceolate-deltoid, unequal-sided pinnulae, which are again cut down into oblong or linear-oblong segments. Segments ½—1½ c.M. long, 2½—5 m.M. broad, slightly toothed. Texture herbaceous, slightly fleshy; surfaces naked, the upper side not rarely copiously proliferous; veins fine, pinnate in the tertiary segments, (the apices thickened and pellucid in the controlled specimens). Sori oblong, at length often filling the whole breadth of the segments.

Penang; *Br. India*, *Australia*, *New Zealand*.

* (72) **A. nitidum**, *Sw.*, *Rac.*, *Flor. Btz.*, I, 219; *Bedd.*, Ferns S.I., tab. CXLVIII, CXLIX; *A. laserpitiiifolium*, *Bedd.*, l. c., tab. CCXXV.

Rhizome suberect, densely clothed with lanceolate, black, glossy scales. Stipes 15—30 c.M. long, firm, erect, greyish to brownish or black, naked. Fronds commonly 60—90 c.M. long, 15—30 c.M. broad, with numerous lanceolate to deltoid, acuminate pinnae on each side. Pinnae ± erecto-patent, the lowest 15—25 c.M. long, 5—8 c.M. broad, cut down to the rachis into numerous stalked, subdeltoid or subrhomboidal pinnulae, which are again cut down into flabellate to cuneate segments, which are inciso-serrate round the outer edge or pinnatifid again. Texture subcoriaceous; rachis firm, naked; veins channelled, conspicuous. Sori short, radiant.

Malaya; *North. India*.

* (75) **A. setisectum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 219; *A. sulcatum*, *Mett.* (not *Lam.*), *Ann. Mus. Bot. L. B.*, II, 236.

Near *A. praemorsum*, *Sw.*, but larger. Fronds to 60 c.M. long or longer, to 22 c.M. broad. Pinnae numerous, erecto-patent, stalked, lanceolate, to 3 c.M. broad or broader, cut down to the rachis into numerous erecto-patent pinnulae, the apex pinnatifid, long-acuminate, caudate, the point nearly filiform. Pinnulae subsessile, lanceolate, 5—4 × as long as broad, the base cuneate, the edge incised and serrate-dentate, the apex acute or acuminate, the teeth fine-pointed. Texture subcoriaceous; rachis polished, naked. Sori linear.

Java.

* (74) **A. affine**, *Sw.*, *Hk. Bk.*, *Syn. Fil.*, 215; *Hk.*, *Spec. Fil.*, III, tab. CCII; *A. spathulinum*, *J. Sm.*, *Bedd.*, *Ferns S.I.*, tab. CCXXVI.

Stipes 15—50 c.M. long, firm, erect, grey, nearly naked. Fronds 50—45 c.M. long, 15—30 c.M. broad, with numerous pinnae on each side, the lower ones lanceolate-triangular, the apex pinnatifid, acute or acuminate. Pinnulae distinct, rhomboidal-triangular, blunt or acute, inciso-serrate or deeply pinnatifid to pinnate again, the base obliquely cuneate; lowest pinnulae acute, ± 5½ c.M. long, ± 1½ c.M. broad, the largest undivided ones 1½—2 c.M. long, under 1 c.M. broad, the higher ones and those of the higher pinnae with the outer edge ± bluntish or ± rounded to truncate, crenate. Texture subcoriaceous; rachis firm, naked; veins subflabellate. Sori copious, linear, the higher ones more oblique than the lower.

Mol'aya; Polynesia, Ceylon, East African Islands.

* (75) **A. laserpitifolium**, *Lam.*, *Rac.*, *Flor. Btz.*, I, 220; *Hk.*, *Spec. Fil.*, III, tab. CCIII; — var. *intermedium*, *Christ*; — var. *subvenustum*, *Christ*, *Philipp. Journ.*, II, 165.

Rhizome creeping, short, densely clothed with spreading, hyaline, yellowish scales. Stipes 15—30 c.M. long, firm, erect, naked, or scaly at the base. Fronds 50—120 c.M. long, to 45 c.M. broad, with numerous pinnae on each side. Lowest pinnae 5—25 c.M. long, 2—15 c.M. broad, deltoid-lanceolate, cut down to the rachis into numerous distinct pinnulae, the lowest of which often 2-pinnate. Ultimate divisions cuneate-oblong, with the outer edge rounded, finely toothed and occasionally incised. Texture herbaceous to chartaceous; rachis slender, naked; veins fine, conspicuous. Sori linear-oblong, irregular.

Var. intermedium: Texture thinly herbaceous; segments broadly ovate, obtuse, coarsely toothed. Sori short, flabellate.

Var. subvenustum: Reduced form; fronds 30—45 c.M. long, much divided, on slender stipes. Leaflets \pm 5 m.M. long or sometimes longer, triangular, flabellate. Sori small, 2—3 to a pinnula (leaflet).

Malaya; — var. *subvenustum*: *Luzon*.

§ 4. DAREA.

a. *Fronds small, to 10 c.M. long.*

b. *Ultimate segments obtuse.*

c. *Fronds 2¹/₂—4 c.M. long, rachis and margins beset with deciduous, black bristles.* (76) *A. Beccarianum*.

c.c. *Fronds 6—10 c.M. long, rachis and margins not bristly.*

(77) *A. Lauterbachii*.

b.b. *Ultimate segments rather acute.*

(78) *A. dichotomum*

a.a. *Fronds larger, 12—60 c.M. long when fully developed.*

1. *Stipes tufted. Lowest secondary segments simple or forked.*

2. *Pinnae 6—12 on each side, subdeltoid, the apex narrowed; secondary segments few.* (79) *A. obtusilobum*.

2.2. *Pinnae many to numerous, linear-oblong, the apex rounded; secondary segments many.* (8¹) *A. Belangeri*.

1.1. *Stipes scattered. Lowest secondary segments rather copiously pinnate.*

(81) *A. scandens*.

(76) **A. Beccarianum**, *Ces.*, Fel. Becc. Polin., 4 & 7.

Rhizome wide-creeping. Stipes approximate, placed in distant groups, erect, \pm 2¹/₂ c.M. long, naked, filiform. Fronds oblong or ovate, \pm 2¹/₂—4 c.M. long, 2-pinnate. Pinnae 8—10 on each side, the segments cuneate, obtuse at the apex, the margin entire or indistinctly crenate with few (1—5) crenations, beset with fugacious, black bristles. Texture weak; rachis bristly like the margin of the pinnae when young. Sori 1 to each ultimate segment.

New Guinea.

* (77) **A. Lauterbachii**, *Christ*, Schum. & Laut., Flor. deut. Sch.geb., 130.

Rhizome wide-creeping, slender, sparingly scaly, bearing distant groups of fronds. Stipes green, sparingly scaly, \pm 3—8 c.M. long. Fronds \pm 6—10 c.M. long, \pm 1¹/₂—2 c.M. broad, lanceolate, acuminate, pinnate. Pinnae

8—12 on each side, subremote, erecto-patent, the higher ones linear-oblong simple, obtuse, entire, the lowest the largest, $\pm 1-1\frac{1}{2}$ c.M. long, 5—6 m.M. broad, the base cuneate, the edge more or less deeply lobed with 2—5 lobes. Lobes not short, oblong to linear-oblong, the largest ± 3 m.M. long, ± 1 m.M. broad, the apex rounded. Texture moderately thick; rachis and surfaces naked; veins flabellato-pinnate, not very distinct, 1 to each ultimate lobe. Sori linear, 1 to each lobe; indusium greyish.

New Guinea.

* (78) **A. dichotomum**, *Hk.*, Hk. Bk., Syn. Fil., 221; Hk., 2nd Cent. of Ferns, tab. XXXIX.

Stipes tufted, $\pm 2\frac{1}{2}$ c.M. long, erect, greenish, deciduously setoso-paleaceous. Fronds 4—10 c.M. long, oblong, acute, sub-3-pinnate. Pinnac 6—12 on each side, 1—2 c.M. long, the base on the lower side cut away, the edge cut down to a narrow, compressed rachis into $\pm 3-7$ linear segments, the lowest of which are mostly 1—2 \times forked. Ultimate segments 3—5 m.M. long, $\frac{1}{2}$ —2 m.M. broad, acute. Texture herbaceous; main rachis compressed, slightly winged, greenish, naked; 1 vein and sorus to each ultimate segment.

Borneo.

(79) **A. obtusilobum**, *Hk.*, Hk. Bk., Syn. Fil., 221; Hk., Icon. Pl., X, tab. M; *A. oceanicum*, *C. Chr.*, Ind. Fil.

Rhizome short, erect, stoloniferous. Stipes tufted, 5—10 c.M. long, naked, greenish or blackish and polished. Fronds 12—15 c.M. long, $1\frac{1}{2}$ —5 c.M. broad, often proliferous at the apex. Pinnac 6—12 on each side, subdeltoid, the lower $\frac{1}{3}$ of the basiscopical side cut away, the largest $1-2\frac{1}{2}$ c.M. long, to 2 c.M. broad, \pm flabellately cut into 3—5 linear, obtuse lobes. Texture thin, herbaceous; rachis green, or black in the lower half; veins 1 to each ultimate segment, covered with scattered, blackish, stellate hairs like the rachis. Sori 1 to each lobe.

Celebes, New Guinea; Polynesia.

* (80) **A. Belangeri**, *Kze.*, Hk. Bk., Syn. Fil., 225; Bedd., Ferns Br.I., II, tab. CCLXXXVII; Hk., Fil. Exot., tab. XLI; *A. tenerum*, *Forst.*, var. *Belangeri*, *Rac.*, Flor. Btz., I, 215; *C. Chr.*, Ind. Fil.; *A. decorum*, *Kze.*, Bot. Zeit., 1848, 176.

Rhizome erect or suberect, the scales brown, lanceolate. Stipes tufted, 5—20 c.M. long, firm, erect, greenish, naked. Fronds 15—50 c.M. long,

4—8 c.M. broad, varying from linear-lanceolate to deltoid-ovate, blunt to caudate, with many to numerous pinnae on each side. Pinnae horizontal, 7—10 m.M. broad, the apex rounded, the edge regularly cut down throughout to a narrowly winged rachis into many linear, erecto-patent segments, the base truncate on the upper side, cuneate below. Segments 1—1½ m.M. broad, simple, the lowest anterior usually forked. Texture coriaceous; veins 1 to each segment, naked or provided with some few fimbriate, minute scales. Sori solitary on the segments. — The young plants have the pinnae less deeply lobed, ± bluntly toothed.

Malaya; Tonkin.

* (81) **A. scandens**, *J. Sm.*, Hk. Bk., Syn. Fil., 224; Hk., 2nd Cent. of Ferns. tab. XXXVII; *A. myriophyllum*, *Muell.* (not *Spr.*), Pap. Pl., IV, 82.

Stipes scattered, varying from very short to 20 c.M. long. Fronds 50—60 c.M. long, 15—50 c.M. broad, with numerous horizontal pinnae on each side. Pinnae ± 3—4 c.M. broad, cut down to a winged rachis into numerous subdistant, ovate-rhomboidal pinnulae, which are cut down to the rachis throughout, with the lowest segments forked or pinnatifid again. Ultimate segments narrowly linear. Texture thin, herbaceous; rachis firm, erect; veins 1 to each ultimate segment. Sori solitary on the segments.

Philippines, Borneo, Ceram, Amboina, New Guinea; Fiji.

§ 5. LOXOSCAPHE.

*. *Stipes pale-coloured; fronds small, 2-pinnate.* (82a) *A. Bakeri.*

*. *Stipes dark-coloured; fronds large, 3—4-pinnate.* (82b) *A. novo-guineense.*

(82a) **A. Bakeri**, *C. Chr.*, Ind. Fil.; *Davallia lanceolata*, *Bk.*, Kew Bull., 1899, 119.

Rhizome slender, short-creeping. Stipes short, slender, stramineous, brownish below. Fronds lanceolate, 2-pinnate, 5—10 c.M. long, 1½—2 c.M. broad, naked; rachis naked, stramineous. Pinnae numerous, contiguous flabellato-pinnate. Segments few, narrowly linear, 1-nerved, 2½—4 m.M. long, the apex dilated. Sori solitary, terminal; indusium persistent, naked.

New Guinea.

(82b) **A. novo-guineense**, *Rst.*, Fed., V, 40.

Rhizome short-creeping, densely clothed with rigid, black, linear-lanceolate, finely acuminate scales. Stipes dense, straight, to 60 c.M. long, terete, naked, black or purple-black, the sides provided with a narrow, pale-coloured stripe. Fronds linear-lanceolate, nearly 1 M. long, \pm 20 c.M. broad, compound. Pinnæ suberecto-patent, short-stalked, subremote, linear-lanceolate, to 12 c.M. long, to 4 c.M. broad, acuminate, truncate at the base. Pinnulae lanceolate, to $2\frac{1}{2}$ c.M. long, to $1\frac{1}{2}$ c.M. broad, the base truncate on the upper, cuneate on the lower side, the apex short-acuminate or obtuse, the anterior side produced. Tertiary divisions simple or 2-fid, the laciniae narrowly linear, nearly 3 m.M. long, $\frac{1}{2}$ m.M. broad, slightly dilated at the apex. Texture thinly herbaceous; surfaces pale-green, nearly naked (clothed here and there with some few black-purple, spinuloso-subulate hairs on the costulae and the margin); veins 1 in each ultimate segment, forked near the apex. Sori subterminal on the longest veinlets; indusium semiovate, opening towards the shortest veinlet.

New Guinea.

 § 6. ASPLENIDICTYUM.

* (83) **A. Finlaysonianum**, *Wall.*, Hk. Bk., Syn. Fil., 245; Hk., Icon. Pl., X, tab. CMXXXVII; *Hemidictyum Finlaysonianum*, *Moore*, Bedd., Ferns Br.I., I, tab. LXXII.

Stipes 15—25 c.M. long, naked or clothed with blackish, subulate-lanceolate scales at the base. Fronds 20—75 c.M. long, acuminate, simply pinnate, with an entire or subrepand terminal pinna, which is often pinnatifid at the base and 1—6 lateral pinnæ on each side, the highest of which are smaller than the terminal one. Lowest pinnæ to 10 c.M. apart, 10—20 c.M. long, 5—8 c.M. broad, erecto-patent, acuminate, the base often unequal, narrowed gradually into a distinct petiole, the edge entire or subrepand, sometimes slightly incised near the base. Texture \pm coriaceous; rachis often slightly winged towards the apex; veins erecto-patent, the highest very oblique. Sori scattered irregularly, $\frac{1}{2}$ —5 c.M. long.

Celebes, Malacca; Br. India.

 56. PHYLLITIS, *Ludwig*.

Characters of *Asplenium* except that the sori and indusia are arranged in

opposite pairs, one originating from the anterior side of a vein or veinlet, the other from the posterior side of the next higher vein or veinlet and so the indusia opening towards each other. — Veins free or anastomosing variously. Hk. Bk., Syn. Fil., tab. IV, fig. 41. (*Scolopendrium*).

A small genus inhabiting different regions and climates.

Arrangement of the sections.

- § 1. **SCOLOPENDRIUM** (*Euscolopendrium*). Midrib present; veins free, not flabellate. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 123, A—B.
- § 2. **CAMPTOSORUS**. Midrib present; veins anastomosing near the base, not flabellate. Diels, l. c., fig. 123, C—D. (Not in Malaya).
- § 5. **ANTIGRAMME**. Midrib present; veins anastomosing towards the apex, not flabellate. Diels, l. c., fig. 124, E. (Not in Malaya).
- § 4. **SCHAFFNERIA**. Midrib wanting; veins free, or anastomosing towards the apex, flabellate. Diels, l. c., fig. 124, F—H. (Not in Malaya).

§ 1. **SCOLOPENDRIUM.**

- §. *Rhizome, stipes and costae quite naked.* (1) Ph. scolopendropsis.
- §.§. *Rhizome, stipes and costae sparingly provided with small or punctiform scales.*
- *. *Rhizome wide-creeping.*
- ×. *Rhizome pale-olivaceous; edges of the fronds entire or undulato-crenate.* (2) Ph. intermedia.
- ×.×. *Rhizome dark-brown; edges of the fronds rather sharply serrate.* (5) Ph. mambare.
- *.*. *Rhizome short-creeping.* (4) Ph. shizocarpa.

(1) **Ph. scolopendropsis**, *v. A. v. R.*, Bull. Dép. Agr. I.N., 1908, XXI, 6; *Asplenium Scolopendropsis*, Muell., Pap. Pl., III, 49; C. Chr., Ind. Fil.

Rhizome creeping, quite naked like the stipes and fronds. Fronds simple, elongate, linear-lanceolate, 20—45 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, narrowed very gradually into an acuminate apex and a long, wingless stipe, the margin more or less sinuato-denticulate, the denticulations rather distant. Texture chartaceous or almost membranaceous; costa stout; veins very spreading, simple or forked, almost parallel. Sori broad, in pairs, traversing the whole width

of the frond from the costa to the margin, those of each pair separated by an ample, empty interstice; the indusia of each pair touching each other with their edge.

New Guinea.

* (2) **Ph. intermedia**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XXI, 6.

Rhizome wide-creeping, angular, pale-olive-brown, rather greenish, covered here and there with very minute, punctiform, dark-brown, scattered scales. Stipes scattered, 5—10 c.M. long, apparently articulate, pale-olive-green, angular, 5—10 c.M. long, naked or sparingly covered with dot-like scales like the stout costa. Fronds linear-lanceolate, simple, rarely forked towards the apex, \pm 25—35 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, narrowed gradually towards both ends, the apex rather caudate, irregularly crenate, the edge irregularly (but more or less broadly and slightly) undulato-crenate, here and there entire. Texture chartaceous, subpellucid; colour olive-green, but paler between the 2 sori of each pair; surfaces naked; veins simple or forked, not close, subspreading or erecto-patent. Sori broad, in pairs, reaching from the costa to some distance from the margin, those of each pair separated by an ample, empty, thin-membranaceous, translucent interstice; the indusia of each pair not touching each other with their edge, at length strongly recurved. — This plant seems to be intermediate between *Ph. mambare* & *scolopendropsis*, *v. A. v. R.*, and specimens recently received from New Guinea (Leg. Dr. Versteeg) quite agree with this.

New Guinea (Treub).

(5) **Ph. mambare**, *v. A. v. R.*, l. c.; *Scolopendrium mambare*, *Bail.*, in *Queensl. Agric. Journ.*, III², 162; *C. Chr.*, *Ind. Fil.*

Rhizome wiry, angular, dark-brown, scaly on the younger portion, the scales few and scattered on the older parts. Fronds erect, linear-lanceolate, sometimes forked at the end, often exceeding 50 c.M. in length, not exceeding $2\frac{1}{4}$ c.M. in width, the edges irregularly sinuate, sometimes the lobes cut down to the costa, all more or less sharply serrate, the apex tapering into an almost filiform cauda and the base into a stipe 5— $7\frac{1}{2}$ c.M. long. Stipe and costa slightly and minutely scaly; veins simple or forked, rather close, parallel. Sori reaching from the costa to near or some distance from the margin.

New Guinea.

(4) **Ph. schizocarpa, v. A. v. R.**, l. c.; *Scolopendrium schizocarpum*, Copel., Philipp. Journ., I, Suppl., 152, tab. IX.

Rhizome short, creeping, clothed towards the apex with fuscous, adpressed scales. Stipes 2—5 c.M. long, black, sparingly scaly. Fronds simple, oblanceolate, 20—50 c.M. long, \pm 5 c.M. broad, caudate-acuminate, entire or undulate, narrowed gradually into the stipe. Texture coriaceous; surfaces naked; veins indistinct, forked. Sori oblique, regular, not reaching either the midrib or edge.

Mindanao.

57. ? TRIPHLEBIA, Baker.

Characters of Phyllitis except that the sori (not the indusia) are not placed *on* but *between* the two contiguous veins, i. e. 1 sorus to each pair of indusia, which open towards each other. Veins free. Diels, in Engl. & Prantl, Nat. Pfl Fam., I⁴, fig. 122, C—D. (*)

East. Malaya, Philippines and New Guinea.

1. *Fronds uniform.*

2. *Fronds simple.*

3. *Fronds deeply crenato-repand.*

3.3. *Fronds entire.*

2.2. *Fronds pinnate.*

1.1. *Fronds dimorphous.*

(1) *T. linza.*

(2) *T. longifolia.*

(3) *T. pinnata.*

(4) *T. dimorphophylla.*

(1) **T. linza, Bk.**, Becc., Mal., III, 42, tab. V; *Asplenium Linza, Ces.*, Fel. Becc. Polin., 4, 7.

Rhizome creeping, naked, slender, grey. Stipes to 5—8 c.M. long, naked, the base dilated, articulate. Fronds lanceolate, naked, \pm 50 c.M. long, the base angustate, the margin deeply and irregularly crenato-repand. Texture

(*) **Triphlebia**, at least in the case of **T. pinnata, Bk.**, is not distinguished by any constant character from *Scolopendrium (Phyllitis)*, to which genus the species should be returned (Copel., Philipp. Journ., I, Suppl., 152).

membranaceous; veins conspicuous, simple or forked. Sori reaching from the costa to the margin; indusium narrow, naked, persistent; receptacle obscure. ⁽¹⁾

New Guinea.

(2) **T. longifolia**, *Bk.*, Copel., Polypod. Philipp., 87; *Scolopendrium longifolium*, *Pr.*, Rel. Haenk., I, 48, tab. IX, fig. 1.

Differs from *T. pinnata*, *Bk.* only in that the fronds are simple and entire, and the limit between the two indusia is marked by a thin, not emphasized line. — Perhaps an undivided form of the young plants of *T. pinnata*, *Bk.*

Borneo, Celebes, Philippines to Papua.

(3) **T. pinnata**, *Bk.*, Copel., Polypod. Philipp., 87; *Hk.*, Icon. Pl., XVII, tab. MDCLII; *Scolopendrium pinnatum*, *J. Sm.*, Kze, Farnkr., I, 124, tab. LVI.

Stipes compressed, greyish. Fronds 50—120 c.M. long, with an entire or finely crenate terminal pinna 10—20 c.M. long, 5—5 c.M. broad, proliferous at the point; below this 1—6 lateral pinnae similar in shape to the terminal one. Texture coriaceous; surfaces naked; veins mostly once forked. Sori numerous, oblique, not reaching either the midrib or edge; indusium ridged where the halves meet.

Borneo, Celebes, Philippines to Papua.

(4) **T. dimorphophylla**, *Bk.*, Becc., mal., III, 42, tab. IV, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 122, C—D; *Asplenium subserratum*, *Ces.*, (not *Bl.*), Fel. Becc. Polin., 4; *Stenochlaena sorbifolia*, *J. Sm.*, var., C. Chr., Ind. Fil.

Rhizome creeping, slender, naked. Stipes short, the base articulate. Fronds dimorphous, both simple and once pinnate. Simple fronds lanceolate, ± 15 c.M. long, the base acute or very gradually narrowed, the edge irregularly crenato-repand. Pinnae of the other fronds small, paucijugate, ± obovate, flabellately cut or inciso-crenate. Texture membranaceous; both surfaces naked; veins erecto-patent, conspicuous, simple or forked. Sori reaching nearly from the costa to the margin; indusium narrow, naked, persistent. ⁽²⁾

New Guinea.

⁽¹⁾ See Schum. & Laut., Flor. deut. Sch.geb., Suppl., 42, 43.

⁽²⁾ See Schum. & Laut., Flor. deut. Sch.geb., Suppl., 43.

TRIBE VII. PHEGPTERIDEAE.

58. NOTHOLAENA, R. Brown.

Sori roundish, oblong or linear-oblong, terminal on the veins, often coalescing afterwards into a continuous, submarginal line, naked or partially covered by the unaltered, often \pm recurved edge of the frond. Spores globoso-tetrahedral.

Rhizome commonly short-creeping. Stipes not articulate to the rhizome. Fronds rather small, divided, glandular, hairy or scaly, often hardly different from *Cheilanthes*; veins free. Hk. Bk., Syn. Fil., tab. VI, fig. 50; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 145, B—E. (*Nothochlaena*).

Tropical and subtropical, but also found in cooler regions, generally inhabiting dry climates.

Arrangement of the sections.

§ 1. **EUNOTHOLAENA.** Fronds not powdery beneath.

§ 2. **CINCINALIS.** Fronds coated beneath with white or yellow powder. (Not in Malaya).

§ 1. EUNOTHOLAENA.

*. Under surface more or less hairy or fibrillose. (1) *N. hirsuta*.

.. Under surface scaly. (2) *N. distans*.

* (1) ***N. hirsuta*, Desv.**, Hk. Bk., Syn. Fil., 372; *Cheilanthes hirsuta*, Mett., Rac., Flor. Btz., I, 147; *Pteris hirsuta*, Poir., (oldest name); — var. *densa*; *N. densa*, J. Sm., Copel., Polypod. Philipp., 97; Kze, Farnkr., I, tab. LXIV, fig. 2, a—c.

Rhizome short-creeping, the scales dense, ferruginous or brown, linear-subulate. Stipes approximate, 3—15 c.M. long, erect, firm, dark-purplish-brown, polished, more or less copiously clothed with whitish or ferruginous, small, spreading or crisped, subulate, fibrillose scales. Fronds 10—20 c.M. long, 4—8 c.M. broad, ovate-elongate, 3-pinnatifid at the base. Pinnae several on each side below the pinnatifid apex, \pm lanceolate, the lower ones rather

distant, \pm deltoid-oblong, equalling or smaller than those next above, pinnate towards the base. Pinnulae sessile, to $1\frac{1}{2}$ c.M. long, to 4 m.M. broad, blunt, crenate, lobed or pinnatifid, the lobes oblong, blunt, small. Texture herbaceous; rachises and lower part of the costae fibrillose; veins sparingly fibrillose, the fibrils whitish or brownish. Sori small, consisting of few or not very numerous capsules, confluent into a rather narrow, more or less interrupted submarginal line, whether or not partially covered by the slightly recurved margin.

Var. densa: Fronds narrower, to 5 c.M. broad, often less divided. Sori broader, extending from the margin nearly or quite to the costules at full maturity.

Trop. Asia, Australia & Polynesia.

* (2) **N. distans**, **R. Br.**, Hk. Bk., Syn. Fil., 572; Hk., Icon. Pl., X, tab. CMLXXX; *Cheilanthes distans*, Mett., Farngr., V, Cheil., 25; — var. *leontopodium*, Christ, Ann. Bot., XV, 169.

Stipes tufted, $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, castaneous, wiry, clothed with ferruginous, linear-lanceolate-subulate scales. Fronds 15—25 c.M. long, 2—3 c.M. broad, 2-pinnate. Lower pinnae distant, deltoid, cut down to the rachis below into oblong, obtuse, entire or slightly lobed pinnulae. Texture subcoriaceous; upper surface pale-green and slightly villose, rachis and under side more or less densely coated with long, linear-lanceolate, ferruginous scales. Pinnae often much curled together and the edge recurved, subindusiform. Sori at length continuous along the margin.

Var. leontopodium: Scales of the under surface very long, white when young.

Philippines; Polynesia, New Zealand, Australia; — var. leontopodium: Celebes.

Species of a dubious systematical place.

(3)? **N. pteridiformis**, **Bk.**, Becc., Mal., III, 49; *Gymnogramme pteridiformis*, Ces., Fel., Becc., Polin., 5 & 8.

Rhizome wide-creeping, slender, clothed with brown, fugacious scales. Stipes 5—15 c.M. long, slender, brown, provided with scattered, ferruginous, furfuraceous, fugacious scales. Fronds pinnate, oblong, 20—25 c.M. long. Pinnae 3—6, distant, glossy, naked, erect or moderately spreading; sterile

pinnae 12—15 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, subfalcate, acute or acuminate, the base obliquely and shortly narrowed, the edge entire; fertile pinnae $\pm \frac{1}{2} \times$ as broad as the sterile ones, long-caudate, the margin eroso-dentate. Veins close, parallel, simple or forked, free, reaching the margin, those of the fertile pinnae dilated at the apex, the fertile ones accompanied by yellow, subpellucid lines on each side, which are connected by a submarginal, moderately conspicuous, yellow strand.

New Guinea.

59. MONACHOSORUM, *Kunze.*

Sori oblong, terminal on the veins, without an indusium, never confluent. Spores 3-lobato-tetrahedral.

Rhizome short-creeping. Stipes approximate, not articulate to the rhizome. Fronds large, finely dissected, naked; veins free. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 117, B & F.

Tropical Asia.

* **M. subdigitatum**, *Kuhn*, Christ, Farnkr. d. Erde, 76, fig. 199; *Polypodium subdigitatum*, *Bl.*, Rac., Flor. Btz., I, 97; *Bl.*, Flor. Jav., II, tab. XCIII; *Bedd.*, Ferns Br.I., II, tab. CCXXIX; *P. davalloides*, *Mett*, Hk., Spec. Fil., IV, 256; *Phegopteris subdigitata*, *Bedd.*, Handb Ind. Ferns, 295; *Aspidium subdigitatum*, *Bl.*, (oldest name).

Rhizome creeping, clothed with reddish, bristle-like hairs. Stipes approximate, firm, stramineous, brown in the lower part, furfuraceous or nearly naked, 30—50 c.M. long. Fronds 50—100 c.M. long, triangular, 3—4-pinnate. Lower pinnae largest, 15—25 c.M. long, deltoid. Pinnulae lanceolate, 3—6 c.M. long, 2—3 c.M. broad. Tertiary divisions stalked, oblong, divided nearly or quite to the costa into oblique, entire or toothed to incised ultimate segments, which are cuneate at the base, $\frac{1}{2}$ —1 c.M. long, 1—3 m.M. broad, the teeth bluntish or acute. Texture thinly herbaceous; rachis furfuraceous; surfaces naked; veins 1 to each ultimate tooth, \pm clavate at the apex. Sori copious, terminal on the veins, \pm oblong.

Tropical Asia.

60a. PHEGOPTERIS, Fée.

Sori roundish or oblong. rarely more than $2 \times$ as long as broad. Indusium wanting.

Rhizome creeping or erect. Stipes not articulate to the rhizome. Fronds various in size and shape, commonly with the habit and appearance of *Dryopteris*; main veins pinnate; veins free as in *Lastraea* or anastomosing as in *Nephrodium*. (*Dryopteris*, p. p., C. Chr., Ind. Fil.).

Tropical and subtropical, also found in cooler climates. (*)

Arrangement of the sections.

- § 1. **EUPHEGOPTERIS**. Veins free. Sori round. Differs from *Lastraea* by the absence of an indusium. Bedd., Handb. Ind. Ferns, fig. 148—152.
- § 2. **LEPTOGRAMMA**. Like § 1, but the sori linear or oblong. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 91, G.
- § 3. **GONIOPTERIS**. Opposite veins of contiguous groups uniting as in *Nephrodium*. Sori round, free. Differs from *Nephrodium* by the absence of an indusium. Hk. Bk., Syn. Fil., tab. V, fig. 4^a, e.
- § 4. **STEGNOGRAMMA**. Like § 3, but the sori linear or oblong, the opposite pairs not confluent. Hk. Bk., l. c., tab. VI, fig. 52, b.
- § 5. **MENISCIUM**. Like §§ 3 & 4, but the sori mostly contiguous in pairs, at least the lower (sometimes the higher) pairs confluent and not rarely produced along the excurrent veinlets. Bedd., l. c., fig. 231—234; Diels, l. c., fig. 92, H.

Key to the sections.

a. Veins free, as in *Lastraea*.

*. Sori round.

.. Sori oblong, sometimes linear.

§ 1. **EUPHEGOPTERIS**.

§ 2. **LEPTOGRAMMA**.

(*) Regarding this genus see the information in "Christ, Farnkr. d. Erde, 266". Its natural place is near **Dryopteris**, from which it differs by the wanting indusium only. A species of **Dryopteris** with fallen or obliterated indusia is not or hardly distinguishable from this genus.

a.a. Veins uniting as in *Nephrodium*.

b. Sori free, i. e. the contiguous pairs never confluent.

c. Sori round.

§ 3. **GONIOPTERIS.**

c.c. Sori oblong or sometimes linear.

§ 4. **STEGNOGRAMMA.**

b.b. At least the lower (sometimes the higher) contiguous pairs of sori confluent and not rarely running along the excurrent veinlets. Sori oblong, rarely round.

§ 5. **MENISCIMUM.**

§ 1. **EUPHEGopteris.**

a. Fronds 1—2 × pinnate.

b. Pinnae cut down $\frac{1}{2}$ -way or more to the costa into close, regular, entire or subentire secondary segments.

c. Stipes not tuberculate at the base.

d. Lower pinnae hardly, if at all, reduced.

e. Texture herbaceous. Lower pinnae with a small gland at the base. (1) *Ph. appendiculata*.

e.e. Texture ± coriaceous. Pinnae without a gland at the base.

×. Lowest pinnae with the lowest segments free, incised, produced, stipule-like. (2a) *Ph. quadriaurita*.

×.×. Lowest pinnae with the lowest segments otherwise. (2b) *Ph. erubescens*.

d.d. Lower pinnae gradually reduced.

*. Sori ± terminal on the veins. Pinnae without a gland at the base. (3) *Ph. Smithii*.

.. Sori nearer the costulae than the edge. Pinnae with a gland at the base. (4) *Ph. auriculata*.

c.c. Stipes tuberculate at the base. (5) *Ph. tuberculata*.

b.b. Pinnae cut down nearly or quite to the costa into toothed or lobed secondary segments.

§. Stipes naked, stramineous; secondary segments sharply toothed.

(6) *Ph. oxyodus*.

§.§. Stipes ± bristly or scaly, dark-brown or blackish; at least the lower secondary segments lobed or incised.

*. Fronds narrow, ± 4 c.M. broad. Pinnae pinnate at the base. (7) *Ph. subobscura*.

.. Fronds broader, 12—25 c.M. broad. Pinnae deeply pinnatifid throughout. (8) *Ph. obscura*.

α.α. Fronds at least deeply 3-pinnatifid.

1. Fronds mostly under 45 c.M. long.

2. Main rachis scaly, or scaly and hairy.

*. Lowest pinnae produced on the lower side. Ultimate segments entire. (9a) *Ph. hirta*.

.. Lowest pinnae not produced on the lower side. Ultimate segments toothed. (9b) *Ph. Copelandi*.

2.2. Main rachis naked or hairy, not scaly.

3. Rachis and under surface slightly pubescent or tomentose.

(10) *Ph. rufescens*.

3.3. Rachis and both surfaces naked.

4. Stipes rather short, castaneous. Lowest pinnae ± equilateral. (11) *Ph. loxoscaphoides*.

4.4. Stipes rather long, green or stramineous. Lowest pinnae with the lower pinnulae on the lower side produced.

5. Sori medial on the veins. (12) *Ph. Hasseltii*.

5.5. Sori terminal on the veins. (15) *Ph. laserpitiifolia*.

1.1. Fronds more than 45 c.M. long when fully developed.

α. Stipes copiously scaly.

*. Texture firm; both surfaces densely setose.

(14) *Ph. cheilanthoides*.

.. Texture thin-herbaceous; costae and veins often hairy.

(15) *Ph. ornata*.

α.α. Stipes naked, hairy or glandular.

β. Texture ± coriaceous (*). (16) *Ph. asperula*.

β.β. Texture ± herbaceous.

γ. Sori submarginal.

*. Fronds large, erect, 45—120 c.M. long.

(17) *Ph. punctata*.

.. Fronds very large, subscandent, 360—450 c.M. long. (18) *Ph. Keraudreniana*.

γ.γ. Sori ± medial on the veins.

δ. Fronds ± 20—50 c.M. broad.

(*) See also under **Ph. punctata, Mett.**

ε. *Stipe and rachis long-hairy.*

(19) *Ph. subconnexa.*

ε.ε. *Stipe and rachis naked or short-pubescent.*

(20) *Ph. distans.*

δ.δ. *Fronds broader, ± 90 c.M. broad.* (21) *Ph. Kingii.*

(1) **Ph. appendiculata, v. A. v. R.**; *Polypodium appendiculatum*, Wall., Hk. Bk., Syn. Fil., 506; (? *Dryopteris cana*, O. Ktze., C. Chr., Ind. Fil.).

Stipes tufted, ± 50 c.M. long, deciduously villose. Fronds 45—60 c.M. long, 20—25 c.M. broad, ovate-lanceolate. Pinnae 7—15 c.M. long, 2—2½ c.M. broad, narrowed from the base to an acuminate point, cut down nearly to the costa into linear-oblong, falcate lobes; lower pinnae with a small gland at the base, slightly reduced. Texture herbaceous; rachis and under surface villose; veins 8—12 on a side. Sori close to the costulae.

Penang; East. Himalaya.

(2a) **Ph. quadriaurita, v. A. v. R.**; *Dryopteris quadriaurita*, Christ, Philipp. Journ., II, 209.

Rhizome probably oblique, the apex crowned with rigid, castaneous, opaque, subulate scales nearly 1 c.M. long. Stipes few or solitary, ± 55 c.M. long, naked, grey-stramineous, terete. Fronds ± 55 c.M. long, ± 24 c.M. broad, deltoid-oblong, hardly narrowed towards the base. Pinnae ± 15 on each side below the pinnatifid apex, sessile, lower opposite, lowest deflexed or decurved, acuminate, ± 12 c.M. long, ± 2½ c.M. broad, cut down to the costa into subdistant, horizontal segments. Secondary segments linear, acute, entire, with acute sinuses between them, ± 1½ c.M. long, ± 3 m.M. broad, lower free, falcate, lowest produced, petiolulate, incised, adpressed to (overlapping?) the rachis, stipule-like. Texture coriaceous, rigid; colour bright-green; costae very short-hairy; veins simple, ± 8—10 on each side, oblique. Sori minute, medial, impressed. — Fronds with the appearance of *Pteris quadriaurita*, Retz.

Mindanao.

* (2b) **Ph. erubescens, J. Sm.**, Christ, Farkr. d. Erde., 272; *Polypodium erubescens*, Wall., Hk. Bk., Syn. Fil., 506; Bedd., Ferns Br.I., II, tab. CCXIII; *Dryopteris erubescens*, C. Chr., Ind. Fil.; *Nephrodium erubescens*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 171; — var. *amboinensis*, Hk. Bk., l. c.

Stipes 50—75 c.M. long, tufted, angular, ± reddish like the rachis. Fronds

50—120 c.M. long, 15—60 c.M. broad, ovate-acuminate, pinnate. Lower pinnae not abbreviated, 1—4 c.M. broad, approximate, sessile, the sides parallel for a long way, the apex gradually acuminate, serrate, the edge deeply pinnatifid nearly to the costa into close, blunt or acute, subfalcate, entire or obscurely serrate segments. Texture rigid, subcoriaceous; rachis, costae (and veins sometimes) beneath sparingly covered with needle-like hairs; veins close, simple, free, soriferous at the very base, the higher ones without sori. Sori in 1 row on each side of the costulae, close to the latter.

Var. amboinensis: Lower pinnae on stalks $2\frac{1}{2}$ c.M. long, conspicuously narrowed at the base. Sori not close to the costulae.

Malaya; North. India, South. China; — var. amboinensis: Amboyna.

* (3) **Ph. Smithii, v. A. v. R.**, Bull. Dép. Agr. I. N., 1908, XVIII, 15; *Dryopteris Smithii, v. A. v. R.*, l. c.

Stipe and rachis naked, stramineous. Fronds \pm 30—35 c.M. long, 6—8 c.M. broad, lanceolate, acuminate, the apex pinnatifid. Pinnae \pm 50 on each side, spreading, the higher ones sessile with a broad base, the central ones largest, subsessile, the base broadly cuneate, the edge cut down nearly to the costa into numerous, subspreading lobes; lower pinnae gradually reduced to mere auricles. Lobes oblong, blunt, the lowest the largest, \pm 5 m.M. long, \pm 1 m.M. broad, the edge entire or slightly crenate, strongly recurved in dried material. Texture subcoriaceous; upper surface hairy on the costae, lower long-hairy on the costae, costulae and veins; veins distinct, simple. Sori round, distant, terminal on the veins, few, submarginal, covered by the recurved edge of the lobes. -- It bears the name of Mr. J. J. Smith, Assistant Curator of the Buitenzorg Herbarium.

Java (Leg.....?, No. 4710).

(4) **Ph. auriculata, J. Sm.** (not *Bedd.*), Christ, Farnkr. d. Erde, 271; *Polypodium auriculatum, Wall.*, Hk. Bk., Syn. Fil., 506; *Bedd.*, Ferns Br.I., II, tab. CCIII; *P. appendiculatum, Wall.*, var. *squamaestipes*, Clarke, Trans. Linn. Soc., Bot., 2nd Series, I, tab. LXXIX, fig. 2; *Dryopteris squamaestipes, C. Chr.*, Ind, Fil.; *Nephrodium auriculatum, Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 171.

Stipes stout, tufted, densely villose with deciduous, white hairs, furnished below with numerous, ovate, black, shining scales. Fronds ample, 90—120 c.M. long, 30—40 c.M. broad, lanceolate-oblong, pinnate, acuminate. Pinnae numerous, \pm $2\frac{1}{2}$ c.M. broad, sessile, furnished at the base beneath with a pro-

minent, hooked gland, the edge pinnatifid nearly to the costa; lower pinnae gradually reduced to mere auricles. Lobes broad, oblong, obtuse, entire, sub-falcate. Texture firm-membranaceous; surfaces sparingly hairy; costae and veins villose. Sori nearer the costulae than the edge; capsules furnished with few, long, weak hairs.

Java; China, North. India.

(5) **Ph. tuberculata**, *v. A. v. R.*; *Dryopteris tuberculata*, *C. Chr.*, *Ind. Fil.*; *Nephrodium tuberculatum*, *Ces.*, *Fel. Becc. Polin.*, 4 & 7; *Polypodium tuberculatum*, *Bk.*, *Ann. of Bot.*, V, 456.

Stipes strong, the base provided with conical, rough, obtuse tubercles, the rest naked, deeply canaliculate, \pm 45 c.M. long. Fronds 90—120 c.M. long, pinnate. Pinnae numerous, 20—25 c.M. long, more than $2\frac{1}{2}$ c.M. broad, cut down to the rachis into numerous linear, falcate, subobtuse lobes, naked on both sides except the costa. Under surface glandular; veins simple. Sori small, in 1 row on each side of the costae.

New Guinea.

(6) **Ph. oxyodus**, *v. A. v. R.*; *Polypodium oxyodon*, *Bk.*, *Journ of Bot.*, XVII, 66; *Dryopteris oxyodus*, *C. Chr.*, *Ind. Fil.*; *Nephrodium oxyodon* (*oxydon*), *Copel.*, *Polypod. Philipp.*, 23.

Rhizome short-creeping, the scales small, erect, lanceolate, membranaceous, dark-brown. Stipes approximate, naked, substramineous, 15—30 c.M. long, slightly scaly near the base. Fronds deltoid-oblong, 15—30 c.M. long, 2-pinnatifid. Pinnae 9—18, lanceolate, 7—10 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, cut down to a broad wing into lanceolate-oblong, sharply toothed lobes 4—5 m.M. broad; higher pinnae sessile, lower stalked. Texture moderately firm; both surfaces naked; main veins pinnate in the lobes; veins 10—12-jugate, many of them 2—4 \times forked. Sori small, round, medial on the anterior fork branches of the veins.

Sulu Islands.

(7) **Ph. subobscura**, *Christ*, *Bull. Herb. Boiss.*, VI, 836; *Dryopteris subobscura*, *C. Chr.*, *Ind. Fil.*

Stipes numerous, tufted, \pm 8 c.M. long, tender, and like the rachis blackish-brown and covered with spreading, narrow, acuminate, hispid bristles. Fronds oblong-deltoid, long-acuminate, 10—15 c.M. long, \pm 4 c.M. broad. Lower pinnae deltoid-ovate, short-stalked, deeply pinnatifid towards the blunt-

ish apex, pinnate at the base, the higher ones ovate, sessile, the terminal one pinnatifid. Lower segments incised, the other ones serrate, falcato-lanceolate, bluntish or short-acuminate. Texture papyraceo-membranaceous; surfaces naked; veins black, conspicuous, the lower pinnate, the higher forked. Sori minute, medial, often submarginal, slightly prominent on the upper surface, sometimes with a depressed indusium.

Sumatra; Yunnan.

(8) **Ph. obscura**, *Fée*, Bull. Herb. Boiss., VI, 196, tab. IV, fig. 5, a—e; *Polypodium obscurum*, *Hk.*, Hk. Bk., Syn. Fil., 508; *Dryopteris sagenioides*, *O. Ktze*, var., C. Chr., Ind. Fil.; *D. obscura*, *O. Ktze*, Rev. Gen. Pl., II, 812; *Nephrodium obscurum*, *Diels*, Copel., Polypod. Philipp., 23.

Stipes tufted, blackish, polished, 18—50 c.M. long, sparingly scaly, the scales dark-coloured, subulate, spreading. Fronds deltoid-oblong, acuminate, \pm 18—45 c.M. long, \pm 12—25 c.M. broad. Pinnae close, \pm opposite, in several pairs below the pinnatifid apex, lowest largest, produced on the lower side, deeply pinnatifid. Lobes 1—2½ c.M. long, regularly, shallowly and roundly lobed, the apex bluntish. Texture thinly papyraceo-herbaceous, flaccid; surfaces (especially the margin and veins) covered with soft, hyaline, purple or reddish hairs, not rarely glandular. Sori small, immersed, medial.

Philippines; Annam,

(9a) **Ph. hirta**, *Christ*, Bull. Herb. Boiss., VI, 195; *Polypodium hirtum*, *Pr.*, Ann. of Bot., V, 458; *Dryopteris setosa*, *C. Chr.*, Ind. Fil.; *Nephrodium hirtum*, *Copel.*, Polypod. Philipp., 25.

Fronds deltoid, decomposed, \pm 50 c.M. long. Lower pinnae the largest, deltoid, produced on the lower side, the central ones lanceolate-oblong. Ultimate segments oblong, entire, 2—2½ m.M. broad. Texture moderately firm; surfaces naked; rachises clothed with copious, adpressed, linear, brown scales; veins paucijugate, erecto-patent. Sori medial.

Philippines.

(9b) **Ph. Copelandi**, *v. A. v. R.*; *Dryopteris Copelandi*, *Christ*, Philipp. Journ., II, 216.

Rhizome probably oblique. Stipes few, rufo-stramineous, \pm 50 c.M. long, clothed like the rachises with minute, setiform, strigillose, reddish-brown paleae besides rather large, ovate, obtuse, diaphanous, pale-brown scales. Fronds deltoid oblong, \pm 40 c.M. long, \pm 50 c.M. broad, 3-pinnatisect. Pinnae ovate-

oblong, ± 15 on each side, acuminate, distant, the lower short-stalked, hardly narrowed at the base, the lowest ± 7 c.M. apart, hardly produced on the lower side, with the lowest anterior pinnulae shorter. Pinnulae ± 15 on each side, approximate, ± 4 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, regularly pinnatifid nearly to the costa. Tertiary segments in ± 10 pairs, rhomboidal-oblong, obtuse, subdecurrent at the base, ± 6 m.M. long, ± 5 m.M. broad, regularly toothed with ± 5 subacute teeth on each side. Texture flaccido-herbaceous; surfaces pubescent with numerous whitish hairs; colour bright-green; costulae pinnate, veins forked (?*). Sori numerous, ± 3 on each side of the costulae.

Luzon.

* (10) **Ph. rufescens**, *Mett.*, Christ, Farnkr. d. Erde, 276; *Polypodium rufescens*, *Bl.*, Hk. Bk., Syn. Fil., 509; *Bl.*, Flor. Jav., II, tab. XCI; *Bedd.*, Ferns S.I., tab. CCXXXVI; *Dryopteris rufescens*, *C. Chr.*, Ind. Fil.; *Nephrodium rufescens*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 175; *Aspidium rufescens*, *Bl.*, (oldest name).

Rhizome creeping, commonly slender, scaly. Stipes 20—45 c.M. long, erect, naked, stramineous (conf. Blume's drawing). Fronds 50—45 c.M. long, 15—50 c.M. broad, subdeltoid. Lowest pinnae largest, erecto-patent, 7—10 c.M. broad, deltoid. Lowest pinnulae lanceolate, stalked, bluntly lobed, deeply pinnatifid or pinnate at the base, those on the lower side the longest, the central ones sessile, decurrent, the highest confluent. Ultimate divisions twice as long as broad or longer, crenate, toothed or lobed. Texture herbaceous to subcoriaceous; rachis and under surface slightly pubescent or tomentose; veins pinnate in the lobes. Sori copious, medial on the veins.

Java; Ceylon, Queensland, New Caledonia.

(11) **Ph. loxoscaphoides**, *v. A. v. R.*; *Polypodium loxoscaphoides*, *Bk.*, Journ. of Bot., XXVIII, 107; *Dryopteris loxoscaphoides*, *C. Chr.*, Ind. Fil.

Stipes short, slender, naked, castaneous. Fronds deltoid-oblong, decomposed, ± 30 c.M. long. Lower pinnae the largest, lanceolate-oblong, subequilateral, 10—15 c.M. long. Ultimate segments ± 2 m.M. long, cuneate with 2 veins, or oblong with a single vein. Texture moderately firm; surfaces naked. Sori solitary and dorsal on the veins, medial in the ultimate segments.

New Guinea.

(*) Christ writes: nervis pinnatis, furcatis.

(12) **Ph. Hasseltii**, *Mett.*, Farngr., IV, Phegop. & Aspid., 13; *Polypodium Hasseltii*, *Bl.*, Flor. Jav., II, 195, tab. XCII; *Dryopteris Hasseltii*, *C. Chr.*, Ind. Fil.

Rhizome oblique. Stipes \pm 20—25 c.M. long, sparingly scaly, stramineous (green after Blume's drawing). Fronds \pm 30 c.M. long and as broad, deltoid, acuminate. Lowest pinnae largest, subopposite, \pm 10 c.M. broad at the base, unequal-sided, the lowest posterior pinnula much the largest. Pinnulae triangular-oblong, with the lowest divisions deltoid-oblong, obtuse, deeply pinnatifid or pinnate at the base, the ultimate segments oblong, blunt, entire or bluntly toothed. Texture coriaceous; rachis and both surfaces naked. Sori medial on the veins.

Java.

(13) **Ph. laserpitiifolia**, *Bedd.*, Handb. Ind. Ferns, Suppl., 84, *Polypodium laserpitiifolium*, *Scort.*, Journ. of Bot., XXV, 524; *Dryopteris laserpitiifolia*, *C. Chr.*, Ind. Fil.

Stipes tufted, fine, stramineous, 50—40 c.M. long, scaly below, naked above. Fronds as long, deltoid-ovate, 5-pinnate; rachis naked. Pinnae numerous, 5—8 c.M. broad. Pinnulae 2—2½ c.M. broad, those on the lower side of the rachis the largest, naked, shining above. Ultimate divisions ovate, oblong, rounded, the base unequal, cuneate on the lower side, the margin broadly lobed; main veins pinnate in the lobes, not reaching the edge. Sori terminal, large, 1—2 on the lowest anterior veins.

Malacca.

(14) **Ph. cheilanthoides**, *v. A. v. B.*; *Polypodium cheilanthoides*, *Bk.*, *Becc.*, Mal., III, 45; *Cheilanthus gigantea*, *Ces.*, Fel. *Becc. Polin.*, 5 & 7; *Dryopteris brunneo-villosa*, *C. Chr.*, Ind. Fil.

Stipes \pm 90 c.M. long, thick, clothed with copious, lanceolate, pale-brown, hairy scales. Fronds ample, deltoid, decomposed. Pinnae lanceolate-oblong. Pinnulae multijugate, sessile, lanceolate. Tertiary segments multijugate, lanceolate, adnate at the base. Ultimate segments small, ovate or semiorbicular, the margin incurved. Texture firm; both surfaces densely setose; veins pinnate. Sori small, occupying the whole under surface of the fertile fronds.

New Guinea.

(15) **Ph. ornata**, *Fée* (not *Bedd.*), *Christ*, Farnkr. d. Erde, 277, fig. 876; *Polypodium ornatum*, *Wall.*, *Bedd.*, Ferns S.I., tab. CLXXI; *Dryopteris*

ornata, C. Chr., Ind. Fil.; *Nephrodium tenericaule*, Hk., Spec. Fil., IV, tab. CCLXIX (not page 142); *N. ornatum*, Christ (not Bk.), Copel., Polypod. Philipp., 26; *Aspidium ornatum*, Christ, Ann. Btz., XV, 130.

Caudex erect, sometimes subarborescent. Stipes, main and lateral rachises clothed with linear-lanceolate scales and muricate or scabrous from their persistent bases, not pilose, viscous or glaucous. Fronds large, to 6 M. long, 3-pinnate, the rachises narrowly winged. Pinnae generally 60 c.M. long or longer, \pm 30 c.M. broad. Pinnulae $2\frac{1}{2}$ —5 c.M. broad, the rachis scabrous and scaly, and furnished with long, white, subulate hairs. Tertiary segments numerous, parallel, 6—7 m.M. broad, linear-oblong from a sessile, square base, connected by a narrow wing, rather distant, the apex blunt or acute, the edges deeply toothed or lobed. Texture thin-herbaceous; costae and often also the veins hairy; veins simple or forked in the ultimate lobes. Sori near the costulae, 1 to each ultimate tooth or lobe, medial on the lowest anterior veinlet.

Malaya; South. China, North. India, North. Australia, Polynesia.

(16) **Ph. asperula**, v. A. v. B.; *Polypodium asperulum*, J. Sm., Hk. Bk., Syn. Fil., 311; *Dryopteris asperula*, C. Chr., Ind. Fil.; *Nephrodium asperulum*, Copel., Polypod. Philipp.; 25.

Stipes firm, terete, brownish, pubescent. Fronds 45—60 c.M. long, 30 c.M. or more broad. Lower pinnae 15—25 c.M. long, 5—8 c.M. broad. Pinnulae lanceolate, \pm 1—1 $\frac{1}{2}$ c.M. broad, unequal-sided, cut down to the rachis below into oblong, pinnatifid, unequal-sided divisions. Texture subcoriaceous; rachis and both surfaces villose; veins immersed. Sori copious, 1 to each ultimate lobe.

Philippines.

* (17) **Ph. punctata**, *Mett.* (not *Bedd.*), Christ, Farnkr. d. Erde, 277; *Polypodium punctatum*, *Thb.*, Hk. Bk., Syn. Fil., 312; *Hypolepis punctata*, *Mett.*, Kuhn, Fil. Afric., 120; *Nephrodium punctatum*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 177; *Dryopteris punctata*, C. Chr., Ind. Fil.; — var. *rugulosa*; *P. rugulosum*, *Lab.*, Hk., Spec. Fil., IV, 272; *Bedd.*, Ferns S.I., tab. CLXX; *H. rugulosa* *J. Sm.*, Bot. Mag., LXXII, Comp. 8; *N. rugulosum*, *Copel.*, Polypod. Philipp., 26.

Rhizome firm, wide-creeping, clothed with reddish hairs or fibrils. Stipes scattered, 30—60 c.M. long, \pm viscous, hairy or fibrillose, at length often naked. Fronds 45—120 c.M. long, 3—4-pinnate. Pinnae lanceolate or deltoid,

erecto-patent, lowest largest, 25—50 c.M. long. Pinnulae close, 10—15 c.M. long. Ultimate divisions numerous, sessile, adnate to the rachis, the apex blunt, the edge crenate, or pinnatifid and crenate. Texture herbaceous; rachis and under surface hairy or fibrillose on the costae and veins. Sori copious, placed near or in the ultimate sinuses, \pm covered by the partially recurved margin as in *Hypolepis*.

Var. rugulosa: Fronds firmer; rachis densely viscid; edges recurved \pm as in *Cheilanthes*.

Nearly pantropical, extending to Japan, New Zealand and the West & East African Islands; not from continental Africa.

* (18) **Ph. Keraudreniana**, *Mann*, Pr. Amer. Ac., VII, 218; *Polypodium Keraudrenianum*, *Gaud.*, Hk. Bk., Syn Fil., 513; *Freye.*, Voy. Ur. Phys., Bot., tab. VII; *Dryopteris Keraudreniana*, *C. Chr.*, Ind. Fil.; *Nephrodium Keraudrenianum*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 177.

Stipes 100—200 c.M. long, stramineous or brownish, polished. Fronds 560—450 c.M. long, subscaudent. Pinnae distant, the lower ones 50—90 c.M. long, 15—40 c.M. broad. Pinnulae lanceolate, \pm spreading, pinnatifid $\frac{1}{2}$ -way down or nearly or quite to the costa into lanceolate-deltoid or oblong, \pm spreading, bluntish, entire, crenate or incised segments. Texture herbaceous, rachises stramineous; both surfaces naked or the under side hairy; veinlets pellucid. Sori submarginal.

Java, Celebes; Hawaii.

* (19) **Ph. subconnexa**, *Christ*, Warb., Mons., I, 85; *Dryopteris subconnexa*, *C. Chr.*, Ind. Fil.

Stipes strong, \pm 50 c.M. long, clothed like the rachis with long, greyish-brown hairs. Fronds \pm 45—50 c.M. long, broad, deltoid. Pinnae very distant, 6—8 on each side, to \pm 25 c.M. long, petiolate. Pinnulae deltoid-ovate, the lowest the longest, the higher ones pinnatifid, the lower pinnate at the base. Tertiary divisions \pm 4 m.M. broad, the higher ones entire, ovate, short-acuminate, the lower coarsely toothed. Texture herbaceous; veins conspicuous, forked. Sori medial, numerous, not close, mixed with white paraphyses. — The whole plant pilose.

Batjan.

* (20) **Ph. distans**, *Mett.*, *Christ*, Farnkr. d. Erde, 274; *Polypodium distans*, *Don*, Rac., Flor. Btz., I, 96; *P. paludosum*, *Bl.*, Flor. Jav., II,

tab. XC; Bedd., Ferns S.I., tab. CLXVIII; *Dryopteris brunnea*, C. Chr., Ind. Fil.; *Nephrodium distans*, Diels, Copel., Polypod. Philipp., 21.

Rhizome creeping, scaly at the extremity. Stipes 45—60 c.M. long, slender, stramineous or brownish, naked or pubescent, the base sparingly clothed with lanceolate scales. Fronds deltoid or ovate, 45—100 c.M. long, 20—50 c.M. broad. Lowest pinnae generally the longest, linear-lanceolate, $3\frac{1}{2}$ —7 c.M. broad, spreading or sometimes deflexed, cut down nearly or quite to the costa into deeply pinnatifid pinnulae. Pinnulae linear-lanceolate, $\frac{1}{2}$ —1 c.M. broad, \pm 1 c.M. apart, the base broadly adnate, slightly decurrent, the lobes blunt or acute, entire or toothed. Texture herbaceous; rachis naked or villose, stramineous or brown; surfaces more or less hairy on the veins; veins pinnate in the lobes. Sori copious, \pm medial, in 1 row on each side of the costules.

Malaya; Formosa, Japan, South. China, North. India.

* (21) **Ph. Kingi**, *Bedd.*, Handb. Ind. Ferns, Suppl., 84; *Dryopteris Kingii*, C. Chr., Ind. Fil.

Fronds large, 3—4-pinnatifid. Pinnae \pm 45 c.M. long. Pinnulae \pm 10 c.M. long, \pm 3—4 c.M. broad, cut down nearly or quite to the rachis into oblong or ovate-oblong, rather deeply lobed tertiary segments \pm 7—8 m.M. broad. Texture membranaceous, rather flaccid; surfaces naked except a few weak setae chiefly on the veins; veins forked. Sori medial, at the forks — A form gathered in Java by Mr. Raap differs from Beddome's plant by the tertiary segments being nearly 5 m.M. broad, with the veins forked or pinnate in the ultimate lobes.

Malacca, Java.

§ 2. LEPTOGRAMMA.

*. *Pinnae pinnatifid* \pm $\frac{1}{2}$ -way down to the costa, the lowest not the longest.

(22) **Ph. totta**.

.. *Pinnae pinnate*, the lowest the longest.

(23) **Ph. opaca**.

* (22) **Ph. totta**, *Mett.*, Christ, Farnkr. d. Erde, 272; *Gymnogramme Totta*, Schl., Rac., Flor. Btz., I, 68; *G. Lowei*, Hk. & Grev., Ic. Fil., tab. LXXXIX; *Leptogramme Totta*, J. Sm., Bedd., Handb. Ind. Ferns, 577; *Grammitis Totta*, Pr., Bedd., Ferns S.I., tab. XLIX; Bau. & Hk., Gen. Fil., tab. LXXII,

B, fig. 7; *Nephrodium Totta*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 91, G.; *Dryopteris africana*, C. Chr., Ind. Fil.; *Polypodium tottum*, Willd., (oldest name).

Rhizome short, creeping or oblique, the scales brown, lanceolate. Stipes subtufted, 10—50 c.M. long, hairy or setulose, scaly below. Fronds 15—50 c.M. long, 5—25 c.M. broad, ovate-oblong, the apex pinnatifid. Pinnae numerous, lanceolate, acute, $\frac{1}{2}$ —5 c.M. broad, the lowest not the longest, short-stalked, cut \pm $\frac{1}{2}$ -way down to the costa into regular, blunt, entire lobes $5\frac{1}{2}$ —5 m.M. broad. Texture herbaceous; rachis and surfaces pilose or setulose; main veins pinnate in the lobes with 5—6 veins on each side. Sori irregular, roundish or more commonly linear-oblong, medial. — Fronds with the aspect of *Dryopteris parasitica*, O. Ktze.

Malaya; Japan, Korea, China, North. India, South. Africa, Madeira, Azores.

* (25) **Ph. opaca**, Mett., Farng., IV, Phegop. & Aspid., 15; *Gymnogramme opaca*, Spr., Rac., Flor. Btz., I, 69; Bedd., Ferns Br.I., II, tab. CCXXXVIII; *G. obtusata*, Bl., Flor. Jav., II, tab. XLIII; *Ph. obtusata*, Christ (not Mett.), Farnkr. d. Erde, 274; *Leptogramme opaca*, Bedd., Handb. Ind. Ferns, 379; *Dryopteris opaca*, C. Chr., Ind. Fil.; *Nephrodium obtusatum*, Diels (not Bedd.), in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 171; *Hemionitis opaca*, Don, (oldest name).

Rhizome suberect or ascending, sparingly scaly. Stipes 30—60 c.M. long, strong, stramineous, glossy, naked or slightly scaly. Fronds 50—100 c.M. long, 50—60 c.M. broad, ovate-deltoid, the apex pinnatifid. Highest pinnae pinnatifid, the lobes broadly crenate; lower pinnae longest, pinnate, stalked, 5—15 c.M. broad. Pinnulae oblong, sessile, decurrent, cut down into rather shallow, slightly toothed or crenated lobes. Texture herbaceous; surfaces naked; rachises and costae commonly slightly hairy; main veins pinnate in the lobes; lower veins soriferous near the base. Sori oblong.

Java; North. India.

§ 5. GONIOPTERIS.

1. Fronds simple. (24) *Ph. holophylla*.

1.1. Fronds pinnate (1).

2. Pinnae entire or crenate (2).

3. Pinnae under 5 c.M. long in the fully developed plants.

4. Pinnae not auricled, 1—6 on each side. (25) *Ph. simplicifolia*.

(1) See also **Ph. canescens**, Mett.

(2) See also **Ph. prolifera**, Kuhn & **diversiloba**, v. A. v. R.

4.4. *Pinnae auricled on the upper side at the base.*

5. *Lateral pinnae \pm 4 on each side.*

(26) *Ph. borneensis.*

5.5. *Lateral pinnae many to numerous.*

6. *Pinnae \pm herbaceous, the lower ones gradually reduced.*

(27) *Ph. debilis.*

6.6. *Pinnae \pm coriaceous, the lower ones not reduced.*

(28) *Ph. aortisora.*

3.3. *Pinnae more than 5 c.M. long in the fully developed plants.*

§. *Pinnae paucijugate.*

†. *Surfaces pubescent.*

(29a) *Ph. xiphioides.*

†.†. *Surfaces naked.*

×. *Rhizome wide-creeping. Pinnae 7—10 c.M. long.*

(29b) *Ph. firmula.*

×.×. *Rhizome short-creeping. Pinnae 12—15 c.M. long.*

(29c) *Ph. Ramosii.*

§.§. *Pinnae multijugate.*

*. *Stipes naked like the rachis. Fronds middle-sized.*

(30) *Ph. rubida.*

.. *Stipes scaly, rachis pubescent or furfuraceous. Fronds large.*

(31) *Ph. arfakiana.*

2.2. *Pinnae toothed or lobed.*

a. *Fronde under 100 c.M. long.*

b. *Pinnae irregularly toothed, or lobed at the very apex.*

(32) *Ph. granulosa.*

b.b. *Pinnae irregularly lobed.*

(35) *Ph. diversiloba.*

b.b.b. *Pinnae regularly toothed or lobed.*

c. *Lower pinnae gradually reduced.*

(34) *Ph. pennigera.*

c.c. *Lower pinnae hardly, if at all, reduced, but the fronds sometimes with a single pair of reduced pinnae at the base.*

d. *Lowest pinnae broadest at the base.*

*. *Texture \pm coriaceous.*

(35a) *Ph. prolifera.*

.. *Texture \pm herbaceous.*

(35b) *Ph. Merrillii.*

d.d. *Lowest pinnae not broadest at the base.*

e. *Pinnae rather short, ovate, obovate or rhomboidal.*

f. *Sori submarginal.*

(36a) *Ph. acromanes.*

f.f. *Sori not submarginal.*

(36b) *Ph. chamaeotaria.*

e.e. Pinnae rather long, linear-acuminate. Sori medial. (36c) Ph. *ceramica*.

a.a. Fronds very large, to 250 c.M. long. (37) Ph. *imponens*.

(24) **Ph. holophylla**, *v. A. v. R.*; *Polypodium holophyllum*, Bk., Journ. of Bot., XXVI, 525 (not XVII, 45); *Dryopteris holophylla*, C. Chr., Ind. Fil.

Rhizome erect. Stipes crowded, naked, erect, \pm 15 c.M. long. Fronds oblong-lanceolate, entire, acuminate, truncate at the base, 15--20 c.M. long, $3\frac{1}{2}$ —5 c.M. broad. Texture moderately firm; under surface slightly hairy on the midrib; main veins distinct from the midrib to the margin, parallel; cross-veins 10—12, arcuate. Sori 2 to each cross-vein. round, minute, superficial.

Borneo.

(25) **Ph. simplicifolia**, *v. A. v. R.*; *Polypodium simplicifolium*, Hk., Hk. Bk., Syn. Fil., 515; *Aspidium simplicifolium*, Hk. (not J. Sm.), Icon. Pl., X, tab. CMXIX; *Dryopteris canescens*, C. Chr., var., C. Chr., Ind. Fil.; *D. simplicifolia*, Christ, Philipp. Journ., II, 206; *Nephrodium simplicifolium*, J. Sm., Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 178.

Stipes approximate, firm, 5—10 c.M. long, villose upwards, scaly below. Fronds 15—50 c.M. long, with a large, lanceolate, acuminate, entire terminal pinna $2\frac{1}{2}$ —4 c.M. broad and below this 1—6 very small, blunt, spreading, distant, entire ones on each side. Texture subcoriaceous; midrib beneath hispid and the prominent veins slightly so; main veins \pm 5 m.M. apart; all the veins joining. Sori minute, \pm medial between the main veins, sometimes Meniscium-like.

Philippines.

(26) **Ph. borneensis**, *v. A. v. R.*; *Polypodium borneense*, Hk., Hk. Bk., Syn. Fil., 514; *Dryopteris labuanensis*, C. Chr., Ind. Fil.

Stipes \pm 15 c.M. long, naked, slender, stramineous. Fronds 15—20 c.M. long, \pm $7\frac{1}{2}$ c.M. broad, with an acuminate terminal pinna \pm $2\frac{1}{2}$ c.M. broad, with shallow, blunt lobes; below this \pm 4 smaller pinnae on each side, $1\frac{1}{2}$ —2 c.M. broad, truncate at the base and auricled on the upper side. Texture subcoriaceous; rachis and both surfaces naked; veins fine, 4—5 on each side. Sori small, in 2 rows.

Borneo.

(27) **Ph. debilis**, *Mett.*, Ann. Mus. Bot. l. B., I, 223, tab. VI, fig. 1; *Polypodium debile*, *Bk.*, Hk. Bk., Syn. Fil., 515; *Aspidium Warburgii*, *Christ*, var. *debile*; *Dryopteris debilis*, *C. Chr.*, Ind. Fil.; *Nephrodium debile*, *Diels*, in Engl. & Prantl., Nat. Pfl.Fam. I⁴, 178.

Stipes slender, slightly hairy. Fronds 15—30 c.M. long, 5—7 c.M. broad, oblong-lanceolate. Pinnae close, \pm 15—20 on each side, the central ones $2\frac{1}{2}$ — $5\frac{1}{2}$ c.M. long, $\frac{1}{2}$ —1 c.M. broad, the point bluntish, the edge slightly crenate, the base truncate, with a distinct, broad, blunt auricle on the upper side; lower pinnae gradually reduced. Texture papyraceo-herbaceous; rachis villose and under surface slightly so; veins 2—3 on a side. Sori small, medial.

Moluccas, New Guinea.

(28) **Ph. aortisora**, *v. A. v. R.*; *Polypodium aortisorum*, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 50; *Dryopteris aortisora*, *C. Chr.*, Ind. Fil.; *Nephrodium aortisorum*, *Copel.*, Polypod. Philipp., 28.

Rhizome short, the scales slender, brown. Stipes clustered, 7—15 c.M. long, slender, grey, slightly pubescent above. Fronds 15—25 c.M. long, 2—4 c.M. broad, linear-lanceolate. Pinnae short-stalked or sessile, 20—40 on each side, 7—10 m.M. broad, elliptical, slightly falcate, obtuse, coarsely crenate, auricled on the upper side, the lower ones deflexed. Texture subcoriaceous; rachis and under side of veins pubescent; main veins pinnate; veins 2—3 on each side, usually the lowest ones only anastomosing. Sporangia scattered, grouping into indefinite sori.

Philippines.

(29a) **Ph. xiphioides**, *v. A. v. R.*; *Dryopteris xiphioides*, *Christ*, Philipp. Journ., II, 201.

Rhizome oblique, creeping. Stipes fragile, approximate, rather close, \pm 30 c.M. long, rufo-stramineous. Fronds broadly ovate, \pm 20 c.M. long, \pm 11 c.M. broad, pinnate, not or hardly narrowed at the base. Terminal pinna \pm 10 c.M. long, \pm $1\frac{1}{4}$ c.M. broad, lanceolate, acuminate, crenate. Lateral pinnae 4—5, similar to the terminal one but shorter, to 7 c.M. long, the lowest not reduced, the highest reduced, approximate to or connected with the terminal one. Texture herbaceous; the whole plant grey-pubescent; colour grey-green; veins 3—4 on each side of the main veins, all anastomosing. Sori small, few.

Mindanao.

(29b) **Ph. firmula**, *v. A. v. R.*; *Polypodium firmulum*, *Bk.*, Kew Bull., 1893, 211; *Dryopteris firmula*, *C. Chr.*, Ind. Fil.

Rhizome woody, naked, wide-creeping. Stipes naked, \pm 45 c.M. long. Fronds oblong-lanceolate, 30—45 c.M. long, simply pinnate. Pinnae paucijugate, sessile, 7—10 c.M. long, 2—5 c.M. broad, lanceolate, acuminate, subentire, the base truncate or broadly cuneate. Texture firm; rachis pubescent; surfaces naked; main veins erecto-patent, parallel; areolae in 3—4 rows between the costae and margin. Sori small, not close, superficial, in 2 rows between the main veins.

Borneo.

(29c) **Ph. Ramosii**, v. A. v. R.; *Dryopteris Ramosii*, Christ, Philipp. Journ., II, 203.

Rhizome short-creeping. Stipes approximate, flexuose, 35—60 c.M. long, rufo-stramineous, clothed with some few, short, scattered scales. Fronds 20—35 c.M. long, oblong, pinnate. Lateral pinnae few, 1—4 on each side below the terminal one, erecto-patent, nearly sessile, ovate-oblong, 12—15 c.M. long, \pm 3 c.M. broad, the base cuneate, the edge subentire or repando-crenate, the apex abruptly caudato-acuminate. Texture herbaceous or papyraceous; rachis naked, provided with roundish axillary buds; surfaces naked, upper dark-green, lower paler; main veins fine but distinct; veins 8—10 on each side of the main ones. Sori small, round, 7—8 on each side near the main veins.

Luzon, Mindoro.

(50) **Ph. rubida**, v. A. v. R.; *Polypodium rubidum*, Hk., Hk., Bk., Syn. Fil., 314; *Dryopteris rubida*, C. Chr., Ind. Fil.; *Nephrodium rubidum*, Copel. (not Hk.), Polypod Philipp., 27; *Goniopteris rubida*, J. Sm., (oldest name).

Stipes firm, erect, glossy, naked, reddish-brown. Fronds 30—45 c.M. long. Pinnae numerous, the lowest stalked, 15—20 c.M. long, 2—2½ c.M. broad, the apex acuminate, the edge nearly entire. Texture subcoriaceous; rachis and both surfaces naked; veins obscure, 8—10 on a side. Sori close to the main veins.

Philippines, ? Borneo, ? Java.

(51) **Ph. arfakiana**, v. A. v. R.; *Polypodium arfakianum*, Bk., Becc., Mal., III, 45; *Dryopteris arfakiana*, C. Chr., Ind. Fil.

Rhizome erect. Stipes short, clothed with more or less spreading, lanceolate, brown scales. Fronds lanceolate-oblong, 90—120 c.M. long, simply pinnate, narrowing gradually towards both ends. Pinnae numerous, alternate, sessile, lanceolate, the medial ones sterile, 7—10 c.M. long, \pm 2½ c.M. broad,

the edge entire or crenate; several lower pinnae gradually becoming smaller, the lowest very small. Rachis brown, rigid, furfureous or pubescent; both surfaces naked, green; main veins with 5—6 veins on each side. Sori semi-globose, occupying the whole under surface of the fertile pinnae.

New Guinea.

(32) **Ph. granulosa**, *v. A. v. R.*; *Polypodium granulatum*, Pr., Rel. Haenk., I, 24, tab. IV, fig. 2; *Nephrodium granulatum*, J. Sm., Bot. Voy. Herald, 428; *Dryopteris granulosa*, C. Chr., Ind. Fil.; — var. *lobata*, Pr., l.c.

Stipes 10—15 c.M. long, clothed like the rachis with a rather rigid, reflexed, dense pubescence, 3-sulcate above, convex below, the base often ± naked. Fronds broadly ovate, 5—8 d.M. long, pinnate. Pinnae alternate, sessile, linear-oblong, narrowed into a long, entire point, the margin obtusely and irregularly dentate, the base cordate, the lower base auriculate, the upper sometimes so; lower pinnae rarely opposite, inclined to become lanceolate, horizontal or deflexed with an upcurved acumen, the medial ones the longest, 7—10 c.M. long, 1½—2 c.M. broad, horizontal, the highest confluent, the terminal one long, broad, unequally and coarsely toothed. Texture membranaceous; colour deep-green; upper surface naked, lower covered with green tubercles or granules and therefore rough; costae pubescent; veins upcurved. Sori nearly round, small, in 2 rows between the main veins.

Var. lobata: Highest lateral pinnae with the acumen abbreviate, with the higher teeth elongate and the apex 4—7-fid, the lobes linear-lanceolate, apiculate, the medial lobe the longest. Terminal pinna not modified.

Timor; Philippines.

(35) **Ph. diversiloba**, *v. A. v. R.*; *Nephrodium diversilobum*, Pr., Copel., Polypod. Philipp., 27; *Dryopteris diversiloba*, Christ, Philipp. Journ., II, 199; *D. canescens*, C. Chr., Ind. Fil., p. p.; — var. *acrostichoides*, Christ, l. c., 200; *Aspidium canescens*, Christ, *forma acrostichoides*, Christ, Ann. Btz., XV, 150—151; *Nephrodium acrostichoides*, J. Sm., Journ. of Bot., III, 411; — subvar. *rhombica* & *lanceola*, Christ, Philipp. Journ., II, 200.

Rhizome elongate, fragile, ± creeping or ascending. Stipes ± approximate, nearly caespitose or subsolitary, flexuose, fragile. Fronds oblong, ± 10 c.M. long, ± 7 c.M. broad, pinnate. Pinnae subpetiolate, 5—7 on each side, ± 4 c.M. long, ± 2½ c.M. broad, rhomboidal or broadly ovate, or prolonged into a lanceolate apex, the base narrowed, subhastate or subcordate, cre-

nate, the edge lobed below the acuminate apex, the lobes very irregular, short or to 2—3 c.M. long, lanceolate, obtuse, $\pm 2\frac{1}{2}$ m.M. broad. Apex of the frond often broad, much elongate, lobate. Texture herbaceous; colour pale-grey-green; the whole plant grey-pubescent. Fertile pinnae hardly contracted, often soriferous towards the apex only. Sori minute, numerous, irregular, rarely seriate, often elongate.

Var. acrostichoides: Barren fronds more or less like those of the type, but the fertile fronds very long-stalked with the pinnae much contracted.

Subvar. rhombea: Fronds ± 7 c.M. long, with the terminal pinna elongate and lobate, acuminate. Lateral pinnae ± 5 on each side, lanceolate-rhomboidal to linear, subpetiolate, truncate at the base, obtuse, crenulate, 4—8 m.M. broad. Sori approximate, seriate or confluent.

Subvar. lanceola: Barren pinnae lanceolate, nearly entire, the apex obtuse or subacute; terminal pinna much elongate, nearly caudate.

Philippines.

(54) **Ph. pennigera, v. A. v. R.;** *Polypodium pennigerum*, Forst., Hk. Bk., Syn. Fil., 317; *Dryopteris pennigera*, C. Chr., Ind. Fil.; *Nephrodium pennigerum*, Pr. (not Hk.), Diels, in Engl. & Prantl, Nat. Pfl.Fam. I⁴, 179.

Stipes tufted, 15—30 c.M. long, slightly scaly. Fronds 45—60 c.M. long, 20—30 c.M. broad. Pinnae 2—2 $\frac{1}{2}$ c.M. broad, cut $\pm \frac{1}{2}$ -way down into blunt, falcate, slightly crenate lobes about 5 m.M. broad; lower pinnae gradually reduced. Texture firm-herbaceous; rachis and both surfaces naked; veins 6—8 on a side. Sori in rows near the costae.

Philippines, New Guinea; Polynesia, New Zealand.

* (35a) **Ph. prolifera, Kuhn,** Christ, Farnkr. d Erde, 270, fig. 851; *Ph. luxurians*, Mett., Farng., IV, Phegop. & Aspid., No. 51; *Polypodium proliferum*, Rb. (not Pr.), Hk. Bk., Syn. Fil., 315; *Meniscium proliferum*, Sw., Hk., 2nd Cent. of Ferns, tab. XV; *Dryopteris prolifera*, C. Chr., Ind. Fil.; *Nephrodium proliferum*, Keys., Copel., Polypod. Philipp., 28; *Hemionitis prolifera*, Retz, (oldest name).

Rhizome stout, creeping. Stipes scattered or approximate, naked, 5—20 c.M. long. Fronds 30—120 c.M. long or longer, ± 15 —30 c.M. broad, erect or decumbent, often elongated and proliferous at the point and from the axils of the pinnae. Pinnae short-stalked, 1—4 c.M. broad, the base broadest, truncate or cordate, sometimes auricled, the edge coarsely crenate or bluntly lobed

to $\pm 2\frac{1}{2}$ m.M. deep, the apex often blunt. Texture subcoriaceous; rachis and under surface naked or slightly pubescent; veins fine, 6—10 on a side. Sori small, medial on the veins, \pm ovate, the lower ones sometimes confluent.

Malaya; Australia, Polynesia, South. China, North. India, Mascarenes, Trop. Africa.

(35b) **Ph. Merrillii**, *v. A. v. R.*; *Dryopteris Merrillii*, Christ, Philipp. Journ., II, 201.

Rhizome erect, stout. Stipes fasciculate, rufo-stramineous, flexuose, 18—20 c.M. long, the base minutely scaly. Fronds ovate-oblong, acuminate, 20—30 c.M. long, \pm 12 c.M. broad, pinnate, the apex elongate, lobed. Barren pinnae rather numerous, close, spreading, petiolulate, falcate-lanceolate, acute, \pm 6 c.M. long, $\pm 1\frac{1}{4}$ c.M. broad, the base truncate not oblique, hastate, with the anterior auricle sometimes elongate, the edge serrato-dentate, the teeth decumbent; lowest pinnae sometimes reduced and deflexed. Texture herbaceous; rachis and surfaces sparingly puberulous; colour bright-green; areolae in 4—5 rows on each side of the costae. Fertile fronds longer stalked, with the pinnae distant, narrowed or contracted and sometimes only 2 m.M. broad. Sori small, close, 4—5-seriate or confluent. — In some respects resembling *Polybotrya appendiculata*, *J. Sm.*, but distinguished by the equal bases of the pinnae.

Palawan, Mindanao.

(36a) **Ph. acromanes**, *v. A. v. R.*; *Dryopteris acromanes*, Christ, Philipp. Journ., II, 200.

Rhizome short, stout. Stipes fasciculate, rufo- or plumbeo-stramineous, slender, 15—30 c.M. long, nearly naked. Fronds oblong, acuminate, \pm 17 c.M. long, \pm 9 c.M. broad, pinnate. Terminal pinna elongate, lobed; lateral pinnae \pm 7 on each side, petiolate, obovate or elongate-rhomboidal, 4—5 c.M. long, \pm 3 c.M. broad, broadest towards the apex, cut $\frac{1}{2}$ -way down or more to the costa, the base broadly truncate; lowest pinnae deflexed, not shorter, narrowed at the base. Lobes \pm 8 on each side, ovate, obtusely rounded, the higher ones increasing in length. Texture herbaceous; surfaces thinly pubescent; colour dark-green; veins in \pm 8 pairs, the lowest anastomosing. Sori large, round, submarginal, bordering the lobes in a single series.

Luzon.

(36b) **Ph. chamaeotaria**, *v. A. v. R.*; *Dryopteris chamaeotaria*, Christ, Philipp. Journ., II, 203.

Rhizome nearly creeping. Stipes approximate, slender, rufo-stramineous, flexuose, 10—15 c.M. long, sparingly puberulous, and clothed with scattered, pale-fuscous, subulate scales. Fronds ovate, \pm 15 c.M. long, \pm 7 c.M. broad, with a large, ovate, acuminate, coarsely lobed terminal pinna \pm 6 c.M. long, \pm 2 c.M. broad. Lateral pinnae 2—3 on each side, petiolulate, similar to the terminal one or reduced, ovate or obtusely rhomboidal, the highest ones adnate. Texture herbaceous; rachis and surfaces minutely puberulous; colour bright-green; veins anastomosing; areolae \pm 4 on each side of the costae. Sori round, in 2 rows on the lobes, \pm 5 to a row.

Luzon.

* (56c) **Ph. ceramica**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 15; *Dryopteris ceramica*, v. A. v. R., l. c.

Stipe and rachis brownish, naked. Fronds deltoid-oblong, \pm 75 c.M. long, \pm 50 c.M. broad near the base. Pinnae \pm 20 on each side below the terminal one. Terminal pinna \pm 15 c.M. long, \pm 1½ c.M. broad at the base, narrowed gradually towards the entire, caudate, acuminate apex, the edge cut \pm ¾-way down to the costa into oblong, blunt, subentire, sub-falcate lobes; higher and central pinnae similar to the terminal one, sessile, broadest and truncate at the base; lower pinnae longer, broadest at the centre, narrowed gradually towards the base, subpetiolate, the lowest pinnae with the lowest lobes subdecurent. Texture membranaceo-herbaceous; surfaces naked or the costae inconspicuously villose; main veins pinnate in the lobes; veins \pm 12 on each side in the longest lobes, the lowest pair anastomosing. Sori medial, close, small, occupying nearly all the veins except the highest ones.

Ceram (Treub).

(57) **Ph. imponens**, v. A. v. R.; *Polypodium imponens*, Ces., Fel. Becc. Polin., 5 & 7; *Dryopteris imponens*, C. Chr., Ind. Fil.

Fronds very large, to 240 c.M. long or longer, to 75 c.M. broad or broader, the main rachis very strong, canaliculate, densely covered with small, brown, rigid, conical, subacute prickles mixed with capillary, deciduous scales and small, yellow hairs. Pinnae to 45 c.M. long, \pm 2½ c.M. broad or broader, the lower ones short-stalked, the others sessile, spreading, rather approximate, broadly linear, acute at both ends, deeply crenate (toothed?) to more than ⅓ on the way to the costa. Texture chartaceous; surfaces opaque; veins close, 15—15 on each side of the main veins. Sori in 1 row on each side, close.

New Guinea.

§ 4. STEGNOGRAMMA.

- §. Terminal pinna very much exceeding the lateral ones, the largest of which at best 5 c.M. long (*). (38) *Ph. canescens*.
- §.§. Terminal pinna not or not very much exceeding the lateral ones, the largest of which more than 5 c.M. long.
- *. Lower pinnae not reduced to mere auricles. (39a) *Ph. stegnogramme*.
- *.*. Lower pinnae dwindling down to mere auricles. (39b) *Ph. Spenceri*.

(38) **Ph. canescens**, *Mett.*, Ann. Mus. Bot. L. B., I, 225 (not Farng. IV, Phegop. & Aspid., 30); *Polypodium canescens*, *Bl.*, Enum., 153; *Gymnogramme canescens*, *Bl.*, Flor. Jav., II, 95., tab. XL; *Nephrodium canescens*, *Christ*, Fil. Saras., II, 208; *Copel.*, Polypod. Philipp., 27, p. p.; *Aspidium canescens*, *Christ*, forma *gymnogrammoides*, *Christ*, Ann. Btz., XV, 150—151; *Dryopteris canescens*, *C. Chr.*, Ind. Fil., p. p: — var. *lobata*, *degener* & *sub-simplicifolia*, *Christ*, Philipp. Journ., II, 198—199.

Rhizome creeping, nearly woody, brownish, bearing the fronds at the extremity. Stipes 10—30 c.M. long, hairy above. Fronds 12½—30 c.M. long, pinnatifid in the upper, pinnate in the lower half. Terminal pinna to 15 c.M. long, to 2½ c.M. broad, narrowed gradually from the base towards the acuminate point, the edge deeply pinnatifid. Lateral pinnae 4—5 or more on each side, sessile, spreading or erecto-patent, ± oblong, the largest ± 4 c.M. long, ± 1—2 c.M. broad, the lowest deflexed, sometimes reduced, the edge not deeply lobed. Texture firm, upper surface deep-green, lower ± greyish-green; both surfaces ± hairy on the veins. Fertile fronds often with the narrowest pinnae. Sori small, oblong or sometimes roundish, ± medial on the veins.

Var. lobata: Pinnae less numerous, broader, the lobes broader, less numerous, deeper; veins flexuose, sometimes irregular; areolae in 1 row on each side. Sori often irregularly elongated. Pubescence grey.

Var. degener: Rhizome nearly creeping. Fronds ± 20—40 c.M. long; texture firmer; colour rather blackish-green. Pinnae ovate, often obovate, narrowed at the base, obtusely crenato-lobate, the terminal one elongate, broad, obtusely lobed; barren pinnae short (± 6 m.M. long), fertile ones often longer (to 2½ c.M. long), stalked, distant. Sori irregularly scattered, ± roundish. Pubescence strigose, brown.

(*) See also **Ph. diversiloba**, *v. A. v. R.*

Var. subsimplicifolia: Smaller, distinguished from the preceding by its distinct, elongated, nearly entire terminal and very slightly developed and auricle-like lateral pinnae.

Java; — the varieties in *Luzon*.

* (39a) **Ph. stegnoگرامme**, *Mett.*, Christ, Farnkr. d. Erde, 268, fig. 846; *Gymnogramme aspidioides*, *Hk.*, Rac., Flor. Btz., I, 70; *Hk.*, Icon. Pl., X tab. CML.; *G. Stegnoگرامme*, *Bl.*, Flor. Jav., II, tab. XLIV; *Stegnoگرامma aspidioides*, *Bl.*, Bedd., Ferns Br.I., I., tab. CXLIX; *Dryopteris stegnoگرامme*, *C. Chr.*, Ind. Fil.; *Nephrodium Stegnoگرامme*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 181.

Rhizome erect, the scales dense, lanceolate. Stipes tufted, long, deciduously hairy. Fronds 30—100 c.M. long, 15—30 c.M. broad, simply pinnate, the apex pinnatifid. Pinnae, lanceolate, often opposite, the lower ones subsessile, often deflexed, acuminate, with the edge pinnatifid $\frac{1}{4}$ — $\frac{1}{3}$ -way down to the costa, the base truncate or broadly rounded; higher pinnae sessile with a broad base. Texture herbaceous; rachis and surfaces hairy, especially on the veins, main veins pinnate; veins 5—10 on each side, the 3—6 lower anastomosing, with an excurrent veinlet reaching to the sinus, all soriferous. Sori linear-oblong, nearly the length of the veins, the lower sometimes produced along the excurrent veinlets as in *Meniscium*. — A form found in Malacca has the fertile fronds contracted.

Java, Malacca; Ceylon, North India.

(39b) **Ph. Spenceri**, *v. A. v. R.*; *Dryopteris Spenceri*, Christ, Philipp. Journ., II, 206; *Nephrodium Spenceri*, Copel., MS.

Rhizome elongate. Stipes 20—30 c.M. long, rufo-stramineous. Fronds 50 c.M. or more long, \pm 12 c.M. broad, oblong, pinnate, with a large, coarsely lobed terminal pinna \pm 12 c.M. long, \pm 3 $\frac{1}{2}$ c.M. broad. Lateral pinnae sessile, oblong, falcate, short-acuminate, \pm crenato-lobate, to \pm 8 c.M. long, \pm 2 c.M. broad, the base unequal-sided, truncate above, semicordate below, overlapping the rachis; lower pinnae dwindling down into mere, triangular auricles (4—10 on each side). Lobes \pm 5 m.M. broad. Texture herbaceous; colour bright-green; the whole plant clothed with scattered, grey hairs; veins distinct, \pm 6 on each side of the main veins, 3—4 lower anastomosing, the terminal pinna with the veins more numerous and anastomosing more copiously. Sori numerous, 4—10 on each side near the main veins (costulae), often oblong.

Mindanao.

§ 5. MENISCIUM. (*)

- a.** *Fronds simple.* (40) Ph. Beccariana.
- a.a.** *Fronds pinnate with 1—2 pinnae on each side.* (41) Ph. triphylla.
- a.a.a.** *Fronds pinnate with 3 or more pinnae on each side.*
- b.** *Fronds 10—25 c.M. long. Pinnae truncate at the base.*
- c.** *Pinnae \pm 5—20-jugate, the barren ones \pm 3—4 c.M. long.*
(42) Ph. Hosei.
- c.c.** *Pinnae \pm 20—30-jugate, the barren ones 1—2 $\frac{1}{2}$ c.M. long.*
(43) Ph. stenophylla.
- b.b.** *Fronds 30—120 c.M. long. Pinnae cuneate or rounded at the base.*
- ‡. *Pinnae narrowed very gradually at the base, with the areolae in 3—5 rows on each side of the costa.* (44) Ph. salicifolia.
- ‡.‡. *Pinnae not narrowed very gradually at the base, with the areolae in 6—20 rows on each side of the costa.* (45) Ph. cuspidata.

(40) **Ph. Beccariana**, v. A. v. R.; *Meniscium Beccarianum*, Ces., Fel. Becc. Polin., 5 & 8; *Dryopteris Cesatiana*, C. Chr., Ind. Fil.

Rhizome creeping, slender, naked, blackish. Stipes angular, dark-brown, strong, slender, 7—15 c.M. long. Fronds simple, lanceolate, \pm 50 c.M. long, \pm 5 c.M. broad, the apex broadly, nearly suddenly acuminate, the margin undulate and obscurely crenate, the base gradually narrowed. Texture subcoriaceous; costa and veins beneath covered with adpressed hairs.

New Guinea; Fiji.

* (41) **Ph. triphylla**, *Mett.*, Christ, Farnkr. d. Erde, 269, fig. 850; *Meniscium triphyllum*, Sw., Rac., Flor. Btz., I, 66; Bedd., Ferns S.I., tab. LVI; *Dryopteris triphylla*, C. Chr., Ind. Fil.; *Nephrodium triphyllum*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 177.

Rhizome creeping, densely clothed with short, spreading hairs and less densely with narrow, lanceolate, ciliate scales. Stipes 1—5 c.M. apart, slender, substramineous, slightly pubescent, those of the barren fronds 6—15 c.M. long, those of the fertile 20—40 c.M. long. Fronds with an oblong-lanceolate terminal pinna 10—15 c.M. long, 2 $\frac{1}{2}$ —4 c.M. broad, the point slightly acuminate, the edge entire or repand, the base cuneate or rounded; below this

(*) See also **Ph. simplicifolia**, v. A. v. R., **prolifera**, Christ, **stegogramme**, *Mett.* & **Merrillii**, v. A. v. R.

1—2 similar but smaller, short-stalked pinnae on each side, the point acute, the base rounded, the fertile ones smaller than the barren. Texture herbaceous or subcoriaceous; veins beneath slightly pubescent; areolae 6—9 between the midrib and edge.

Malaya; Queensland, South. China, North. India.

* (42) **Ph. Hosei**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XXI, 7; *Meniscium Hosei*, Bk., Journ. of Linn. Soc., Bot., XXII, 230; *Dryopteris Hosei*, C. Chr., Ind. Fil.; — var. *sumbensi* v. A. v. R., l. c.

Rhizome short-creeping. Stipes approximate, slender, stramineous, pubescent, 5—15 c.M. long, those of the fertile fronds much longer. Fronds oblong-lanceolate, simply pinnate, to \pm 15 c.M. long. Pinnae 5—10 on each side, lanceolate, straight, erecto-patent, sessile, entire or obscurely crenate, those of the barren fronds 3—4 c.M. long, 5—9 m.M. broad, those of the fertile ones to $2\frac{1}{2}$ c.M. long, 4— $4\frac{1}{2}$ m.M. broad, the base truncate and auricled on 1 or both sides. Texture firm-herbaceous; surfaces naked, \pm verruculose; rachis pubescent; veins 2—5-jugate. Sori afterwards confluent. — Quite agreeing with Hooker's coloured drawing of *Meniscium proliferum*, Sw. (*Ph. prolifera*, Kuhn).

Var. sumbensis: Pinnae more numerous, 12—16-jugate, rather falcate, serrato-crenate. Fertile fronds narrowed, \pm $2\frac{1}{2}$ —5 c.M. broad, with the pinnae rather contracted, 2—3 m.M. broad, strongly falcate, the edges entire, \pm recurved when dry. Sori in 1 row on each side of the costae. — Rhizome not seen. Barren fronds sometimes provided with soriferous, but not contracted pinnae.

Borneo; — var. *sumbensis*: *Sumba Island (Karita, Teysmann No. 10693).*

(43) **Ph. stenophylla**, v. A. v. R.; *Meniscium stenophyllum*, Bk., Journ. of Bot., XXIX, 108; *Dryopteris brevipinna*, C. Chr., Ind. Fil.

Rhizome short-creeping. Stipes slender, naked, 7—10 c.M. long, those of the fertile fronds much longer. Fronds lanceolate, simply pinnate, 20—25 c.M. long. Pinnae sessile, 20—30-jugate, oblong, obscurely crenate, those of the barren fronds to $2\frac{1}{2}$ c.M. long, 5—9 m.M. broad, with the base truncate and auricled on the upper side, those of the fertile ones smaller, with a considerable space between them. Texture firm; surfaces naked, green; main veins pinnate with 2—3 distinct, raised veins on each side. Sori semiglobose, soon confluent.

Borneo.

* (44) **Ph. salicifolia**, *Mett.*, Fil. Lechl., II, 22; *Meniscium salicifolium*, *Wall.*, Hk. Bk., Syn. Fil., 591; *Bedd.*, Ferns Br.I., II, tab. CCVII; Hk., Icon. Pl., X, tab. XM; *Dryopteris salicifolia*, *C. Chr.*, Ind. Fil.; *Nephrodium salicifolium*, *Diels*, in Engl. & Prantl., Nat. Pfl.Fam., I⁴, 178.

Stipes 30 c.M. or more long, stramineous or brownish, glossy. Fronds 50—45 c.M. long, to 50 c.M. broad. Pinnae numerous, 2—4 c.M. apart; 10—20 c.M. long, $\frac{1}{2}$ —2 c.M. broad, narrowed very gradually from the middle to both ends, the edge nearly entire, the lowest the largest. Texture subcoriaceous; rachis and both surfaces naked; main veins oblique; areolae 5—5 between the midrib and edge; excurrent veinlets mostly free. Sori oblong, formed of 2 confluent, rounded ones.

Sumatra, Malacca.

* (45) **Ph. cuspidata**, *Mett.*, Farnng., IV, Phegop. & Aspid., 25; *Meniscium cuspidatum*, *Bl.*, Rac., Flor. Btz., I, 65; *Bl.*, Flor. Jav., II, tab. XLV; *Bedd.*, Ferns Br.I., II, tab. CCCIX; *Nephrodium urophyllum*, *Keys.*, *Diels*, in Engl. & Prantl., Nat. Pfl.Fam., I⁴, 178, p. p.: *Dryopteris urophylla*, *C. Chr.*, var., *C. Chr.*, Ind. Fil.; *D. cuspidata*, *Christ*, Philipp. Journ., II, 205.

Rhizome creeping, the scales lanceolate. Stipes 20—60 c.M. long, stout, scaly at the very base, blackish in the lower part, yellowish above. Fronds 50—120 c.M. long, 15—50 c.M. broad, simply pinnate. Pinnae often short-stalked, 2—7 $\frac{1}{2}$ c.M. broad, the point acuminate, the edge entire or toothed, the base cuneate or rounded. Texture subcoriaceous; rachis and both surfaces naked or nearly so, but the upper surface sometimes covered with white, cretaceous dots; areolae 6—20 between the midrib and edge. Sori round or oblong, in 2 rows between the main veins, the pairs mostly contiguous, at least the lower or higher pairs confluent, the others not rarely free; sporangia setose. — This species is very near *Dryopteris urophylla*, *C. Chr.*, but must be specifically distinct, even if the genera were united (Copel.).

Malaya; North. India.

60b. DICTYOPTERIS, *Presl.*

Sori round or suboblong, very rarely more than 2 × as long as broad. Indusium wanting.

Rhizome creeping or erect. Stipes not articulate to the rhizome. Fronds various, commonly with the habit and appearance of *Aspidium*; veins anasto-

mosing more or less copiously, commonly irregularly, but never as in *Nepbrodium*. (*Aspidium*, p. p., C. Chr., Ind. Fil.). (*)

Tropical.

Arrangement of the sections.

- § 1. **ARCYPTERIS**. Sori round or suboblong; veins anastomosing generally as in *Aspidium*, rarely as in *Campteria* or *Pleocnemia*. Hk. Bk., Syn. Fil., tab. V, fig. 48, o (as *Dictyopteris*).
- § 2. **DIGRAMMARIA**. Sori oblong, 2—3 × as long as broad, simple, or joining at the base when the veins fork; veins anastomosing as in *Pleocnemia*. Bau. & Hk., Gen. Fil., tab. XCIV, fig. 1—4 (as *Stenosemia*).

§ 1. **ARCYPTERIS**.

- a.* Sori marginal or nearly so, or placed in arches within the margin,
 *. Sori confined to the margins of the lobes. Venation as in *Aspidium*.
 (1) *D. pteroides*.
 .. Sori placed in horseshoe-shaped arches. Venation as in *Campteria*.
 (2) *D. andaiensis*.
- a.a.* Sori placed in rows parallel to the main veins or costulae.
b. Fronds 3—5-fid or pinnate, but then the lowest pinnae forked at the base on the lower side.
c. Stipes winged. (3) *D. Bryanti*.
c.c. Stipes not winged.
d. Included free veinlets few or wanting.
 (4) *D. subdecurrens*.
d.d. Included free veinlets copious. (5) *D. Barberi*.
b.b. Fronds deeply pinnatifid or pinnate, but then the lower pinnae pinnatifid or more compoundly divided.
§. Fronds pinnatifid. (6) *D. Beccariana*.
§.§. Fronds 2-pinnatifid.
 *. Lateral pinnae 5—12 on each side. (7) *D. saxicola*.

(*) See the note under **Phegopteris**. The natural place of this genus is near **Pleocnemia** and **Aspidium**, from which it differs by the exindusiate sori only.

.. *Lateral pinnae more numerous.* (8) *D. macrodonta.*

§.§.§. *Fronds at least 2-pinnate. Venation ± as in Pleocnemia.*

(9) *D. ferruginea.*

a.a.a. *Sori scattered irregularly. (*)*

1. *Fronds simple, lobed, 3-partite or pinnatifid.*

2. *Fronds under 30 c.M. long.*

3. *Under surface densely hairy; fronds simple.* (10) *D. vitis.*

3.3. *Under surface naked; fronds at best ternate.*

4. *Stipes naked.* (11) *D. labrusca.*

4.4. *Stipes minutely fibrillose.* (12) *D. lamoensis.*

2.2. *Fronds more than 30 c.M. long.*

§. *Fronds ternate.* (13) *D. Hancockii.*

§.§. *Fronds deeply pinnatifid.*

*. *Lobes ± 4-5 c.M. broad.* (14a) *D. heterosora.*

.. *Lobes ± 10 c.M. broad.* (14b) *D. Bolsteri.*

1.1. *Fronds pinnate; pinnae entire, crenate, lobed or pinnatifid, the lowest pinnatifid or forked at the base.*

α. *Pinnae paucijugate.*

β. *Lateral pinnae ± 2-jugate.* (15) *D. pentaphylla.*

β.β. *Lateral pinnae ± 6-jugate.*

γ. *Lowest pinnae 2-partite.* (16) *D. Whitfordi.*

γ.γ. *Lowest pinnae 3-partite.* (17) *D. polycarpa.*

α.α. *Pinnae multijugate.* (18) *D. irregularis.*

(1) **D. pteroides**, Pr., Tent. Pterid., 194, tab. VIII, fig. 6 & 15; *Phegopteris Brongniartii*, Mett., Farng., IV, Phegop. & Aspid., 51; *Aspidium Brongniartii*, Diels, Copel., Polypod. Philipp., 55; C. Chr., Ind. Fil.; *Polypodium Brongniartii*, Bory, Hk. Bk., Syn Fil., 518; *Tectaria irregularis*, Copel., var. *Brongniartii*, Copel., Philipp. Journ., II, 417; *P. pteroides*, Pr., (oldest name).

Fronds 60—90 c.M. long, 30—45 c.M. broad, the apex pinnatifid with deep, lanceolate, repand lobes; below this numerous distinct pinnae, the lowest 15—25 c.M. long, lanceolate, cut $\frac{3}{4}$ of the way down throughout into acute, subentire, lanceolate lobes 1— $5\frac{1}{2}$ c.M. long, $\frac{1}{2}$ —1 c.M. broad. Texture subcoriaceous; areolae copious, with few free veinlets. Sori rather small and confined to the margin of the lobes.

Malaya.

(*) See also **D. Bryanti**, v. A. v. R.

FILICES.

(2) **D. andaiensis**, v. A. v. R.; *Polypodium andaiense*, Bk., Becc. Mal., III, 45; *Aspidium andaiense*, C. Chr., Ind. Fil.

Stipes strong, brown, \pm 90 c.M. long, the base densely clothed with linear, elongate, crisped, brown scales. Fronds ample, deltoid, 2-pinnate. Pinnae oblong, lanceolate, the lowest the longest, \pm 50 c.M. long. Pinnulae $2\frac{1}{2}$ —4 c.M. long, lanceolate, adnate at the base, the margin conspicuously crenate. Surfaces naked; venation rather as in *Campteria*; main veins erecto-patent; higher veins curved, free, paucijugate. Sori small, semiglobose, placed in horseshoe-shaped arches.

New Guinea.

(3) **D. Bryanti**, v. A. v. R.; *Aspidium Bryanti*, Copel., Polypod. Philipp., 54; C. Chr., Ind. Fil.; *Tectaria Bryanti*, Copel., Philipp. Journ., II, 412.

Stipes 20—40 c.M. high, strong, reddish-brown, winged, the scales scattered, spreading, the wing entire, not narrowed, with a truncate base \pm 1 c.M. broad on each side. Fronds 30—50 c.M. long, ovate, 3-fid with the lateral segments approximate, or often 5-fid with the lowest segments distant. Terminal segment 20—30 c.M. long, \pm 10 c.M. broad, acuminate, entire or subrepand; lateral segments similar, 10—18 c.M. long, 5—7 c.M. broad. Texture papyraceous; main veins curved, reaching nearly the margin; transversal veins arching, conspicuous; areolae large, regular, divided into irregular smaller ones, with free veinlets. Sori mostly in rows along the main veins.

Philippines.

(4) **D. subdecurrens**, v. A. v. R.; *Phegopteris subdecurrens*, Luerss., Bot. Centralbl., XI, 50; *Aspidium subdecurrens*, C. Chr., Ind. Fil.

Stipes dark-red-brown, opaque, naked like the rachis and costae. Fronds \pm 40 c.M. or more long, \pm 20 c.M. or more broad, elliptic, pinnate. Pinnae \pm 5 on each side below the deeply pinnatifid apex, \pm $4\frac{1}{2}$ c.M. broad, entire, the lowest pair stalked, deltoid, acuminate, obliquely lanceolate, forked at the base on the lower side. Terminal pinna large, rhomboidal, narrowed gradually at the base, 5-lobate, the lowest lobes large, the central lobe 5-lobate again with a large, ovate central part and 2 small lateral lobes. Costae prominent; main veins slightly curved, distinct nearly to the edge; areolae large, with included smaller areolae, without or rarely with some few free veinlets. Sori in 2—4 regular rows between the main veins, forming 2 rows on each side of the costae.

Singapore.

(5) **D. Barberi**, *Bedd.*, Handb. Ind. Ferns, 298; Ferns Br.I., II, tab. CCCXXIII; *Phegopteris Barberi*, *Mett.*, Linn., XXXVI, 125; *Aspidium Barberi*, *C. Chr.*, Ind. Fil.; *Copel.*, Polypod. Philipp., 34; *Polypodium Barberi*, *Hk.*, Hk. Bk., Syn. Fil., 517; *Tectaria Barberi*, *Copel.*, Philipp. Journ., II, 414.

Stipes tufted, 15—30 c.M. long, slightly scaly at the base. Fronds palmately 5-lobed or simply pinnate with a large, simple or 3-fid terminal segment and 1—4 pairs of lateral pinnae, the upper ones oblong-lanceolate, 10—15 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, nearly entire, the lowest pair with a large, lanceolate lobe at the base on the lower side. Texture subcoriaceous; both surfaces naked; areolae rather large and regular, with copious free veinlets. Sori copious, principally in 2 rows near the main veins.

Malaya.

(6) **D. Beccariana**, *v. A. v. R.*; *Polypodium Beccarianum*, *Ces.*, Fel. Becc. Polin., 5 & 8; *Aspidium Beccarianum*, *Diels*, C. Chr., Ind. Fil.

Stipes paleaceous, strong, the unwinged lower part $2\frac{1}{2}$ —5 c.M. long, the higher part broadly winged, ± 30 c.M. and more long. Fronds 60—90 c.M. long, ± 60 c.M. broad, deeply pinnatifid. Lobes 7—8 c.M. broad, linear oblong, acuminate, subrepand, the terminal one ovate-lanceolate, 3-fid at the base, with narrowed lobes. Texture membranaceous; lateral veins of the lobes numerous, distinct to the edge; transversal veinlets arcuate; areolae hexagonal; free veinlets rather numerous. Sori small, in 1 row on each side of the costulae.

New Guinea.

(7) **D. saxicola**, *v. A. v. R.*; *D. petrophyta*, *Moore*, Ind., 318; *Phegopteris petrophyta*, *Mett.*, Ann. Mus. Bot. L. B., I, 225; *Polypodium petrophytum*, *Bl.*, Flor. Jav., II, 163, tab. LXXI; *Aspidium saxicola*, *Bl.*, Enum., 160; *C. Chr.*, Ind. Fil.

Stipes ± 50 c.M. long, fusco-purpureous, paleaceo-furfuraceous like the rachis, sulcate above, minutely tomentose in the groove. Fronds 30—75 c.M. long, to ± 30 c.M. or more broad, ovate, 2-pinnatifid. Terminal pinna ovate, deeply lobed; lobes erecto-patent, falcate, blunt, the lowest again lobed. Pinnae 5—12 on each side, sessile, ± 4 c.M. broad, \pm falcate, oblong-lanceolate, acuminate or acute, cut $\frac{1}{3}$ — $\frac{2}{3}$ -way down into oblong, erecto-patent, entire, blunt lobes; lowest pinnae 6—12 c.M. broad, with the lobes on the lower side (except the basal ones) produced, the base sessile, subtruncate or broadly cuneate; higher pinnae slightly decurrent. Texture membranaceous;

surfaces nearly naked or slightly furfuraceous or scaly. Sori confined to the lobes, in 1 row on each side of the main veins or costulae.

Java.

(8) **D. macrodonta**, *Pr.*, Tent. Pterid., 194; *Phegopteris macrodonta*, *Mett.*, Christ, Farnkr. d. Erde, 266; *Polypodium macrodon*, *Reinw.*, Hk. Bk., Syn. Fil., 318; *P. Cumingianum*, *Hk.*, Spec. Fil., V, 105; *Aspidium macrodon*, *Keys.* (not *Diels*), *Copel.*, Polypod. Philipp., 34; *C. Chr.*, Ind. Fil.; *Tectaria irregularis*, *Copel.*, var. *macrodon*, *Copel.*, Philipp. Journ., II, 417.

Rhizome decumbent. Stipes 30 c.M. or more long, slightly scaly below. Fronds 60—100 c.M. long, 30—30 c.M. broad, the apex deeply pinnatifid; below this numerous lanceolate pinnae, the lowest the largest, 15—20 c.M. broad, cut down to a narrow wing on the rachis into pinnatifid, close, lanceolate segments. Texture thin; both surfaces naked; areolae copious, without free veinlets. Sori small, in rows near the main veins.

Malaya; Fiji.

* (9) **D. ferruginea**, *v. A. v. R.*, Hort. Bog.; *Phegopteris ferruginea*, *Mett.*, Ann. Mus. Bot. L. B., 224, tab. VI, fig. 2; *Polypodium ferrugineum*, *Bk.*, Hk. Bk., Syn. Fil., 318; *Aspidium ferrugineum*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I^A, 186; *A. Zippelianum*, *C. Chr.*, Ind. Fil.; *Sagenia ferruginea*, *Christ*, MS. in Herb. Bog.

Stipes \pm 25—45 c.M. long, clothed densely below, and less so above, with ferrugineous, lanceolate, spreading or deflexed scales. Fronds \pm 30—40 c.M. long, \pm 20—30 c.M. broad. Pinnae erecto-patent, the higher ones pinnatifid, the central and lower pinnate. Lowest pinnae largest, deltoid, with the pinnulae on the lower side elongate, again pinnate, the segments pinnatifid. Texture herbaceous; rachises and surfaces fibrillose with sprinkled, ferrugineous or reddish, hair-like scales; lower veins anastomosing, forming costal and costular areolae \pm as in *Pleocnemia*, otherwise free; areolae without free veinlets. Sori in 1 row on each side of the costulae or main veins of ultimate segments.

New Guinea.

* (10) **D. vitis**, *v. A. v. R.*; *Nephrodium Vitis*, *Rac.*, Bull. Ac. Crac., 1902, 60; *Sagenia Vitis*, *Christ*, Ann. Bot., XX, 108, tab. III, fig. 2; *S. rufipubis*, *Christ*, MS. in Herb. Bog.; *Aspidium vitis*, *C. Chr.*, Ind. Fil.

Rhizome thick, oblique, the scales subulate, brown. Stipes tufted, 20—

25 c.M. long, slender, flexuose, canaliculate, dark-brown or blackish, hairy or naked, but clothed towards the base with linear, dark-purple scales, which make the stipes rough when they fall. Fronds \pm 5—10 c.M. long, \pm 3—9 c.M. broad, simple, broadly cordate with a deep sinus, the margin entire or undulate, the basal lobes sometimes meeting or overlapping. Texture subcoriaceous, fragile; upper surface naked or hairy, lower densely clothed with spreading, reddish hairs; main veins 10—12 on each side, conspicuous and prominent like the costa, straight, reaching the margin; veins hidden, forming moderately regular, tetra- or pentagonal areolae with free veinlets. Sori punctiform, scattered irregularly.

Borneo.

* (11) **D. labrusca**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 16; *Polypodium Labrusca*, Hk., Rac., Flor. Btz., I, 109; Hk., Spec. Fil., V, 73, tab. CCLXXXV, B; *Nephrodium Labrusca*, Rac., Bull. Ac. Crac., 1902, 61; *Sagenia Labrusca*, Christ, Ann. Btz., XX, 108; *Aspidium labrusca*, Christ, Ann. Btz., XV, 137; C. Chr., Ind. Fil.; *Tectaria Labrusca*, Copel., Philipp. Journ., II, 410; — var. *Boschae*, Rac., l. c., 61; — var. *ternata*, v. A. v. R., l. c.

Rhizome woody, erect, the scales linear- or lanceolate-subulate. Stipes crowded 10—25 c.M. long, firm, naked, erect, glossy, the base dark-coloured and slightly scaly. Fronds 7—18 c.M. long, 6—15 c.M. broad. Texture subcoriaceous; surfaces naked; costae and main veins distinct; areolae fine, visible, copious, with free veinlets. Sori minute, scattered irregularly. —

Forma typica: Fronds simple, cordate with a deep sinus, the edge entire to sinuate, the point acute.

Var. Boschae: Stipes 20—40 c.M. long. Fronds 5-lobed, 10—35 c.M. each way, the base slightly cordate.

Var. ternata: Fronds 3-foliate. Terminal segment hastato-5-fid with a broad, sinuate central lobe and 2 narrow, \pm entire lateral ones. Lateral segments free or connected with the terminal one, upcurved, unequal-sided, acuminate, the upper base cuneate, slightly rounded, the lower cordate and broadly rounded, the edges \pm entire.

I think Raciborski's Nusa-Kembangan plant would better be treated as a distinct species.

Borneo, Celebes; — var. *Boschae*: *Borneo*; — var. *ternata*: *Nusa-Kembangan (Java)*.

* (12) **D. lamoensis**, v. A. v. R.; *Aspidium lamoense*, Copel., Polypod.

Philipp., 55; C. Chr., Ind. Fil., *Tectaria irrigua*, Copel., Philipp. Journ., II, 413, p. p.

Rhizome ascending. Stipes \pm 10—25 c.M. long, erect, minutely fibrillose like the rachis, the base covered with some few small scales. Fronds 15—25 c.M. long, subdeltoid, commonly 3-foliate. Terminal pinna 10—15 c.M. long, simple and lanceolate or more commonly deltoid and deeply 5-furcate, the central lobe 2—5 c.M. broad, narrowed gradually towards the acute or acuminate apex, the edge entire or inconspicuously undulate, the lateral lobes suberect, 6—10 c.M. long. Lateral pinnae erecto-patent, short-stalked, 6—10 c.M. long, forked at the base on the lower side, the terminal lobe \pm acuminate, the lateral one shorter, more or less bluntish. Texture herbaceous; surfaces naked; main veins hardly reaching the margin; areolae with free veinlets. Sori numerous, scattered irregularly, roundish or oblong.

Luzon.

(13) **D. Hancockii**, v. **A. v. R.**; *Polypodium Hancockii*, Bk., Ann. of Bot., VIII, 127 (not Journ. of Bot., XXIII, 106); *Aspidium sumatranum*, C. Chr., Ind. Fil.

Stipes firm, slender, glossy, naked, brown. Fronds deltoid, ternate, \pm 45 c.M. long. Terminal pinna cordate-ovate, 20—25 c.M. long, 15—20 c.M. broad, pinnatifid $\frac{1}{2}$ -way down. Lateral pinnae petiolate, ovate, 12—15 c.M. long, the lower side produced, distinctly auricled. Texture membranaceous; surfaces naked; rachis naked, glossy, castaneous; main veins curved; areolae hexagonal, copious, with numerous free veinlets. Sori semiglobose, superficial.

Sumatra.

(14a) **D. heterosora**, **Bedd.**, Handb. Ind. Ferns, 302; *Polypodium heterosorum*, Bk., Hk. Bk., Syn. Fil., 506; *Aspidium vastum*, Bl., var., C. Chr., Ind. Fil.

Stipes short, brown, with spreading, linear scales. Fronds oblong-spathulate, 60—90 c.M. long, deeply pinnatifid, the lowest foot only a narrow wing to the costa; above this 3—4 pairs of distant, oblong-lanceolate, acuminate lobes 4—5 c.M. broad at the middle, confluent in a narrow wing, the lowest simple or 2-partite. Texture thin but firm; surfaces naked; ribs naked, brownish; main veins distinct to the edge, with distinct arches between them, and copious small areolae with free, included veinlets. Sory very abundant, small, irregular, often confluent. — ? An exindusiate form of *Aspidium vastum*, Bl.

Malacca.

(14b) **D. Bolsteri**, v. A. v. R.; *Aspidium Bolsteri*, Copel. Philipp. Journ., I, Suppl., 252; ? *Tectaria Bryanti*, Copel., l. c. II, 412, p. p.

Rhizome rather ascending, woody, the extremity clothed with rather long, apiculate, castaneous scales. Stipes close, \pm 50 c.M. long, long-winged, the unwinged base \pm 5 c.M. long, the wing \pm $1\frac{1}{2}$ c.M. broad. Fronds 60—80 c.M. long, ovate in outline, pinnatifid. Lobes \pm 5-jugate, ovate, spreading, \pm 30 c.M. long, \pm 10 c.M. broad, acuminate, entire, distinctly narrowed at the base, the costal wing $1\frac{1}{2}$ —3 c.M. broad. Texture coriaceo-papyraceous; surfaces naked; main veins reaching nearly the margin, arcuate; transversing veins inconspicuous in the fertile region. Sori superficial, in 2 rows between the transversing veins, 4—10 to a row, occupying the whole under surface of the fertile segments.

Mindanao.

* (15) **D. pentaphylla**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 16; *Aspidium pentaphyllum*, v. A. v. R., l. c.

Stipes \pm 50 c.M. long, like the rachis brown or blackish-purpureous, minutely fibrillose and deciduously scaly, the scales few, scattered, lanceolate-subulate, brown or purpureous. Fronds 30—45 c.M. long, 15—25 c.M. broad, with a terminal pinna and 2 lateral ones on each side. Terminal pinna 15—20 c.M. long, simple, broadly ovate, acuminate, the edge undulate with rounded or sharp lobes, the base broadly cuneate or rounded, not rarely purveyed with 1—2 lanceolate, bluntly rounded or acuminate, erecto-patent segments. Lateral pinnae erecto-patent, distant, broadly lanceolate, unequal-sided, the lower side broader than the upper, the apex acuminate, the edge entire or undulate, the base broadly cuneate to rounded; lower pinnae with an oblong or sublanceolate, blunt or acuminate lobe at the base on the lower side. Texture subcoriaceous, subpellucid; costae and main veins prominent; veins visible; areolae numerous, with free veinlets; surfaces naked. Sori numerous, scattered irregularly, often confluent. — According to Copeland's conception of the generic name this plant should be called *Tectaria pentaphylla*, v. A. v. R.

New Guinea (Treub).

(16) **D. Whitfordi**, v. A. v. R.; *Aspidium Whitfordi*, Copel., Polypod. Philipp., 55; C. Chr., Ind. Fil.; *Tectaria irregularis*, Copel., Philipp. Journ., II, 416, p. p.

Rhizome short, suberect, densely scaly at the apex. Stipes 20—30 c.M.

long, dark-brown at the base, which is clothed with linear scales, the rest naked, stramineous. Fronds 30—45 c.M. long, deltoid, acuminate, pinnate, the apex pinnatifid with broadly crenate and obscurely serrate lobes. Pinnae \pm 6 on each side, erecto-patent, lanceolate, acuminate, cut \pm $\frac{1}{2}$ -way down to the costa into oblong, obtuse, entire or serrate lobes; lowest pinnae stalked, deltoid, with a free, basal pinnula on the lower side resembling one of the upper pinnae. Texture papyraceous; surfaces naked; veins laxly anastomosing, without free veinlets. Sori round or oblong, irregularly scattered but wanting near the costae.

Luzon.

(17) **D. polycarpa**, *Bedd.*, Handb. Ind. Ferns, 300; *Phegopteris polycarpa*, *Mett.*, Linn., XXXVI, 124; *Polypodium polycarpum*, *Bk.*, Hk. Bk., Syn. Fil., 506; *Aspidium Kuhnii*, *C. Chr.*, Ind. Fil.

Stipes \pm 15 c.M. long, rubro-stramineous, slightly pilose upwards. Fronds deltoid, 2-pinnatifid, 15—45 c.M. long. Pinnae 6-jugate, lowest largest, deltoid, unequal-sided, stalked, 3-partite, central ligulate, acuminate, crenate. Texture membranaceous; surfaces glabrous; main veins distinct to the edge; areolae with free veinlets. Sori copiously scattered, terminal or compital.

Malacca.

* (18) **D. irregularis**, *Pr.*, Tent. Pterid., 194, tab. VIII, fig. 7; *Bedd.*, Ferns Br.I., I, tab. LXXVII; *D. difformis*, *Moore*, *Bedd.*, Handb. Ind. Ferns, 300; *Phegopteris difformis*, *Mett.*, *Christ*, *Farnkr. d. Erde*, 267, fig. 843; *Ph. irregularis*, *Mett.*, *Ann. Mus. Bot. L. B.*, I, 225; *Aspidium difforme*, *Bl.*, *Enum.*, 160; *A. irregulare*, *C. Chr.*, Ind. Fil.; *Polypodium difforme*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 118; *Tectaria irregularis*, *Copel.*, *Philipp. Journ.*, II, 416, p. p.; *P. irregulare*, *Pr.* (oldest name), *Bl.*, *Flor. Jav.*, II, tab. LXXII; — var. eu-irregularis, *Copel.*, l. c.

Rhizome erect or ascending, densely clothed with lanceolate, very long-acuminate, brown scales. Stipes tufted, \pm 50—50 c.M. long, erect, scaly below. Fronds 50—200 c.M. long, the apex pinnatifid with the lobes crenate or subrepand. Upper pinnae lanceolate, broadly crenate or lobed, the lobes blunt, falcate, reaching \pm $\frac{1}{4}$ — $\frac{1}{2}$ -way down to the costa; lower pinnae stalked, often deltoid, to \pm 30 c.M. long, pinnatifid often more than $\frac{1}{2}$ -way down, the lowest forked at the base on the lower side with the fork branch deflexed and sometimes free, to 12—15 c.M. long. Texture subcoriaceous; surfaces naked; areolae copious, with few free veinlets. Sori very abundant.

Var. eu-irregularis: Fronds simply pinnate; sori small and very numerous.

Malaya; — var. eu-irregularis: *Philippines*.

§ 2. DIGRAMMARIA.

(19) **D. ambigua, v. A. v. R.;** *Digrammaria ambigua*, Pr., Tent. Pterid., 117, tab. IV, fig. 12; *Phegopteris philippinensis*, Mett., var. 1, Mett., Farng., IV, Phegop. & Aspid., 27; *Stenosemia aurita*, J. Sm. (not Pr.), Bau. & Hk., Gen. Fil., tab. XCIV, fig. 1—4; *Aspidium ambiguum*, Diels, Copel., Polypod. Philipp, 39; C. Chr., Ind. Fil.; *Tectaria ambigua*, Copel., Philipp. Journ., II, 415, p. p.; *Gymnogramme ambigua*, Hk., Hk., Bk., Syn. Fil., 378.

Stipes slender, nearly naked. Fronds 45—60 c.M. long, \pm 30 c.M. broad, oblong-deltoid, 2-pinnatifid. Pinnae lanceolate, $2\frac{1}{2}$ —4 c.M. broad, the lower ones slightly stalked, cut $\frac{1}{2}$ -way down to the costa or more into linear-oblong lobes, the fertile lobes contracted. Texture thinly herbaceous; colour dark-green; rachis and both surfaces thinly clothed with long, jointed hairs; veins of the lobes often forked, the lower ones forming costal areolae. Sori oblong, simple, or confluent when the veins fork.

Philippines.

61. DIPTERIS, *Reinwardt*.

Sori punctiform or round, on the back of united veinlets. Indusium wanting.

Rhizome creeping. Stipes scattered, not articulate to the rhizome. Fronds flabellato-dichotomous, i.e. divided nearly or quite to the base in 2 halves, each half once or more times dichotomously divided again with radiating lobes. Primary ribs radiating, dichotomous; secondary ribs (if present) \pm parallel to the primary ones; veins anastomosing copiously. Diels, in Engl. & Prant. Nat. Pfl.Fam., I⁴, fig. 108.

Tropical Asia and Polynesia.

α . Sori in 1 row on each side of the costae.

(1) *D. Lobbiana*.

$\alpha\alpha$. Sori scattered irregularly.

b. Ultimate segments entire.

c. Fronds small, 2—3 × furcate with 6—8 ultimate segments.

(2) *D. Nieuwenhuisii*.

c.c. Fronds middle-sized, 4—5 × furcate with 16 or more ultimate segments.

(3) *D. quinquefurcata*.

b.b. Ultimate segments toothed, fronds rather large.

(4) *D. conjugata*.

* (1) ***D. Lobbiana*, Moore** (not *Bedd.*), *Christ*, *Farnkr. d. Erde*, 123; *Bedd.*, *Ferns Br I.*, II, tab. CCXXXIII; *D. bifurcata*, *Christ* (not *Bk.*), MS. in *Herb. Bog.*; *Polypodium bifurcatum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 562; *P. Lobbianum*, *Hk.*, *Spec. Fil.*, V, 100 (not IV, 226); *Journ. of Bot.*, V, tab. XI; — var. *Ridleyi*, *Christ*, *Ann. Btz.*, XX, 126.

Rhizome creeping, scaly, the scales bristle-like, subulate, blackish, small. Stipes to 50 c.M. long or longer. Fronds 15—50 c.M. long, ± as broad, divided in 2 halves, which are 5—4 × flabellately dichotomous. Ultimate lobes linear, acute, ± 5—8 m.M. broad, the edges entire. Texture coriaceous, central ribs prominent; areolae fine, in 1 or more rows. Sori small, 1—5 to each main areola, placed in 1 ± irregular row on each side near the costae of the lobes.

Var. *Ridleyi*: Fronds 10—15 c.M. long; ultimate lobes 3—4 m.M. broad. Sori punctiform.

Malaya; — var. *Ridleyi*: *Borneo*.

* (2) ***D. Nieuwenhuisii*, Christ**, *Ann. Btz.*, XX, 124, tab. III, fig. 3.

Rhizome creeping, blackish-brown, the scales firm, bristle-like, dark-brown, adpressed. Stipes distant, 50 c.M. long or longer, firm, brown, naked. Fronds ± 14 c.M. long, ± 22 c.M. broad, flabellate, 2-partite. Each half cuneate at the base, 2-fid, with the inner segment forked and the outer simple or forked again. Ultimate lobes ± triangular-oblong, acuminate, ± 2½ c.M. broad at the base, entire, 2-costate. Texture rigidly coriaceous; surfaces naked; costae prominent; veins horizontal, connecting the costae as to form quadrangular primary areolae with smaller, included secondary ones. Sori dot-like, scattered irregularly.

Borneo.

* (3) **D. quinquefurcata**, *Christ*, Farnkr. d. Erde, 123, fig. 350; *Polypodium quinquefurcatum*, *Bk.*, Journ. of Linn. Soc., Bot., XXIV, 260.

Rhizome creeping, densely scaly, the scales blackish, subulate, bristle-like. Stipes straight, firm, naked, very long. Fronds cuneate-flabellate, 4—5 × dichotomous with 16 or more linear-lanceolate ultimate segments to 50 c.M. long, 12—20 m.M. broad. Texture rigidly coriaceous; primary veins stout, raised, forming 1 row of large, subquadrangular primary areolae on each side of the costae of the segments; secondary areolae numerous, fine. Sori near de costae, scattered irregularly, confined to the primary areolae, forming groups of 4—12 to an areola. — Very near *D. Lobbiana*, *Moore*.

Borneo, Lingga Islands.

* (4) **D. conjugata**, *Reinw.*, *Christ*, Farnkr. d. Erde, 123, fig. 548; *D. Horsfieldii*, *Bedd.*, Ferns Br.I., II, tab. CCCXXI; *D. Dipteris*, *Christ*, MS. in Herb. Bog.; *Polypodium Dipteris*, *Bl.*, *Rac.*, Flor. Btz., I, 108; *Bl.*, Flor. Jav., II, tab. LXXXI; *P. Horsfieldii*, *R.Br.*, *Hk.*, Spec. Fil., V, 99; — var. *alpina*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 991.

Rhizome creeping, densely clothed with blackish, adpressed, subulate scales. Stipes 1—100 c.M. apart, 70—200 c.M. long, firm, naked, glossy, erect. Fronds 30—90 c.M. long, more broad than long, divided quite to the base in 2 halves, the main lobes of each reaching $\pm \frac{3}{4}$ of the way down, once or twice forked again, with the edges broadly and obliquely toothed. Texture firm, coriaceous; surfaces naked, glossy, the lower often glaucous; primary ribs prominent, dichotomously forked; areolae copious. Sori minute, abundant, scattered irregularly, especially plentiful near the main veins.

Var. alpina: Fronds 3 × smaller; lobes few, obtuse, obscurely toothed.

Trop. Asia & Polynesia; — var. *alpina*; *Laguna (Philippines)*.

TRIBE VIII. GYMNOGRÁMMEAE.



62. CONIOGRAMME, *Fée*.

Sori linear, running along the veins, without paraphyses. No indusium.

Rhizome creeping. Stipes not articulate to the rhizome. Fronds pinnately divided; under surface not powdery; veins free in malayan material, simple or more commonly forked, pinnately arranged. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 158, A—B. — See the information in Philipp. Journ., II, 171, (Christ).

Paleotropical, and extra-tropical in Asia.

* **C. fraxinea**, *Diels* (not *Fée*), Copel., Polypod. Philipp., 66; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 158, A—B; *Gymnogramme javanica*, *Bl.*, Rac., Flor. Btz., I, 68; *G. fraxinea*, *Bedd.* (not *Don*), Ferns S.Br.I., Suppl., 24; *Syngramme fraxinea*, *Bedd.*, Handb. Ind. Ferns, 586; *Neurogramme fraxinea*, *Christ*, Farnkr. d. Erde, 65; *Diplazium fraxineum*, *Don*, (oldest name); — var. *serrulata*; *G. serrulata*, *Bl.*, Enum., 115; *Bedd.*, Ferns Br.I., I, tab. LVII & Ferns S.I., tab. CCXXXII (as *G. javanica*); (? var. *Copelandi*, *Christ*, Philipp. Journ., II, 171).

Rhizome creeping, sparingly clothed with very small, deltoid scales. Stipes firm, naked, glossy, 30—120 c.M. long. Fronds 50—150 c.M. long, varying from simply pinnate to 5-pinnate. Pinnae erecto-patent, 10—50 c.M. long, stalked, the simple ones 2—5 c.M. broad, lanceolate, acuminate, narrowed at the base, the edge entire. Pinnulae of the same shape as the simple pinnae, but smaller. Texture herbaceous; rachis and both surfaces naked or nearly so; veins distinct, close, simple or once forked. Sori yellow, commonly forked, beginning at the costa, but falling short of the edge.

Var. serrulata: Edges finely toothed.

Trop. Asia, Australia, Polynesia & Africa, Japan, China, Madagascar.

63. CEROPTERIS, *Link.*

Sori linear or linear-oblong, running along the veins, not quite reaching their tips, without paraphyses. No indusium.

Rhizome commonly short-creeping. Stipes not articulate to the rhizome. Fronds divided, at least the fertile ones coated beneath with white or yellow,

waxy powder; veins free, generally forked or pinnate. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 159, C. (*Neurogramme*).

Chiefly neotropical.

✧. Both barren and fertile fronds powdery beneath.

+ . Fronds 2-pinnatifid.

(1) *C. tartarea*.

+ . + . Fronds 3-pinnatifid.

(2) *C. calomelanos*.

✧ . ✧ . Fertile fronds only powdery beneath.

(5) *C. chrysosora*.

* (1) **C. tartarea**, *Link*, C. Chr., Ind. Fil.; *Gymnogramme tartarea*, *Desv.*, Hk. Bk., Syn. Fil., 584; *Acrostichum tartareum*, *Cav.*, (oldest name); — var. *ochracea*; *G. ochracea*, *Pr.*, Rel. Haenk., I, 17; *G. tartarea*, *Desv.*, var. *ochracea*, Hk. Bk., l. c.

Stipes tufted, firm, 15—50 c.M. long, dark-chestnut-brown, scaly towards the base. Fronds 50—60 c.M. long, 15—50 c.M. broad, deltoid-oblong, 2-pinnatifid. Pinnae lanceolate, the lowest the largest, sometimes sharply deflexed. Secondary segments oblong, obtuse, entire, united, or the lower free, whether or not toothed. Texture subcoriaceous; under surface powdery; powder dense, white.

Var. ochracea: Segments very regular; powder bright-yellow.

In cultivation. — *Orig. loc.*: Trop. America.

* (2) **C. calomelanos**, *Und.*, C. Chr., Ind. Fil.; *Gymnogramme calomelanos*, *Klf.*, Rac., Flor. Btz., I, 69; Hk., Gard. Ferns, tab. L; Bau. & Hk. Gen. Fil., tab. XXXVII; *G. Boerlageana*, *Christ*, MS. in Herb. Bog.; *Neurogramme colomelanos*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 264; *Acrostichum calomelanos*, *L.*, (oldest name); — var. *chrysophylla*; *G. chrysophylla*, *Klf.* Enum. Fil., 74; *G. calomelanos*, *Klf.*, var. *chrysophylla*, Hk. Bk., Syn. Fil., 585.

Rhizome short, oblique or erect, the scales brown, lanceolate. Stipes tufted, firm, angular, 5—50 c.M. long, black or brown and polished like the rachis. Fronds 10—90 c.M. long, 5—50 c.M. broad, deltoid-oblong or lanceolate-triangular, 3-pinnatifid. Pinnae close, linear-lanceolate, the lowest mostly the largest, to 2—5 c.M. broad, stalked, the apex pinnatifid. Lower pinnulae lanceolate or deltoid-oblong, the lowest the largest, short-stalked, or sessile with a narrowed base, often deeply pinnatifid into oblong or deltoid lobes; lobes entire or broadly toothed to serrate, the lowest anterior one often the largest.

Texture coriaceous; rachis naked; upper surface naked, lower densely coated with white, waxy powder. Sori linear, forked, at length filling up nearly the whole under surface of the lobes, the ripe sporangia visible as blackish points amongst the powder.

Var. chrysophylla: Powder yellow.

Much cultivated, and escaped from gardens; very common now in the neighbourhood of European settlements. — Orig. loc.: Trop. America, West. Africa, Natal.

(3) **C. chryosora, v. A. v. R.;** *Gymnogramme chryosora, Bk.*, Journ. of Linn. Soc., Bot. XXIV, 260; C. Chr., Ind. Fil.

Rhizome short-creeping. Stipes approximate, castaneous, naked, $\pm 10-15$ c.M. long, the fertile fronds on the longest stipes. Fronds moderately firm, rather rigid, 7—10 c.M. long. Lowest pinnae much the largest and most compound, ultimate segments obovate-cuneate, 2—5½ m.M. broad. Both surfaces naked. Fertile fronds similar, but with the under surface densely coated with golden, waxy powder. Sori at length confluent, covering nearly the whole breadth of the fertile segments, but do not reach the tips.

Borneo.

64. ANTROPHYUM, *Haukfuss.*

Sori long, linear, carried along the veins, irregularly longitudinal or sub-flabellate, superficial or more commonly immersed, free or reticulate but often imperfectly. Paraphyses present. No indusium.

Rhizome short-creeping. Fronds simple, stalked or sessile, not articulate to the rhizome; veins copiously anastomosing, forming longitudinally elongated, \pm hexagonal areolae; costa wanting, at least in the upper half of the frond. Christ, Farnkr. d. Erde, fig. 141 & 142. (*)

Pantropical.

a. *Sori marginal or very near the margin, not scattered over the frond.*

§. *Sori quite marginal, in 1 line on each side.* (1) *A. marginale.*

§.§. *Sori near the margin, in ± 3 approximate lines on each side.*

(2) *A. Clementis.*

a.a. *Sori not marginal, nor very near the margin, more or less scattered over the frond.*

(*) The species are variable and not always clearly distinguished.

- b.** *Paraphyses linear, the apex abruptly thickened, clavate, obpyriform or capitate.*
- c.** *Fully developed fronds mostly under 2 c.M. broad.*
- ‡. *Fronds blunt or rounded at the apex.* (3) *A. immersum.*
- ‡.‡. *Fronds acute or acuminate at the apex.* (4) *A. parvulum.*
- c.c.** *Fully developed fronds 2 c.M. or more broad.*
- d.** *Fronds sessile or short-stalked, i.e. the unwinged part of the stipe very short with regard to the length of the whole frond.*
- e.** *Costa wanting towards the apex of the frond only.*
(5) *A. costatum.*
- e.e.** *Costa wanting throughout or at best present in the lower half of the frond.*
- f.** *Fronds thinly chartaceous. Paraphyses globose at the apex.* (6) *A. strictum.*
- f.f.** *Fronds coriaceous. Paraphyses clavate at the apex.*
(7) *A. semicostatum.*
- d.d.** *Fronds rather long-stalked, i.e. the unwinged part of the stipe not very short with regard to the length of the whole frond.*
- *. *Sori superficial.* (8) *A. latifolium.*
- *.*. *Sori immersed.*
- ×. *Texture subcoriaceous, rather thick.*
(9) *A. plantagineum.*
- ×.×. *Texture membranaceous, pellucid.* (10) *A. elongatum.*
- b.b.** *Paraphyses linear, the apex not or hardly thickened.*
- 1.** *Fully developed fronds at best ± 2 c.M. broad.*
- *. *Sori few, in 2—6 lines.* (11) *A. subfalcatum.*
- *.*. *Sori rather copious.* (12) *A. coriaceum.*
- 1.1.** *Fully developed fronds ± 2½—10 c.M. broad.*
- 2.** *Fronds broadly spathulate, rather small, to ± 10 c.M. long.*
(15) *A. ovatum.*
- 2.2.** *Fronds ligulate, lanceolate or spathulate, more than 10 c.M. long in the fully developed plants.*
- 3.** *Sori ± straight, continuous or interrupted, not reticulate.*
- ×. *Fronds ligulate-lanceolate. Costa wanting, or only distinct towards the base of the frond.*
(14) *A. callifolium.*
- ×.×. *Fronds spathulate-lanceolate. Costa more distinct in the lower half of the frond.* (15) *A. alatum.*

3.3. *Sori mostly flexuose, short or long, reticulate.*

4. *Costa wanting or present, but then not black.*

(16) *A. reticulatum.*

4.4. *Costa present, black.*

(17) *A. Cumingii.*

1.1.1. *Fully developed fronds \pm 15 c.M. broad.*

(18) *A. spathulatum.*

(1) **A. marginale**, *Bl.*, Flor. Jav., II, 80, tab. XXXIV, fig. 1 & 2; (? *A. callifolium*, *Bl.*, C. Chr. Ind. Fil.).

Rhizome short-creeping, reddish-brown, the scales setiform. Fronds sessile, 4—25 c.M. long, 8—20 m.M. broad, simple, linear-lanceolate, sometimes falcate, narrowed towards both ends, the apex sometimes unequally 2-fid, the edge entire, or at best undulate in the fertile fronds. Texture coriaceous; no midrib; surfaces naked. Sori marginal, continuous or interrupted, slightly immersed; paraphyses linear.

Java.

(2) **A. Clementis**, *Christ*, Philipp. Journ., II, 175.

Fronds elongate-lanceolate, caudate, \pm 30 c.M. long (incl. the stipe), \pm 2½ c.M. broad at the middle, narrowed gradually towards both ends, the margin bounded by a narrow, hyaline line. Texture diaphanous; costa present in the lower part of the frond, yellow, at best 8 c.M. long; areolae long, vertical, \pm 2½ c.M. long, \pm 2 m.M. broad. Sori narrow, filiform, in \pm 3 rows on each side near the margin, irregular, varying from ½ to 7 c.M. long, not joining, slightly prominent on the upper surface of the frond. — Colour of the fronds emerald-green.

Mindanao.

* (3) **A. immersum**, *Mett.*, Rac., Flor. Btz., I, 63; *A. pumilum*, *Klf.*, Hk. & Grev., Ic. Fil., tab. XLVI; *A. obtusum*, *Bl.* (not *Klf.*), Flor. Jav., II, 80, tab. XXXIV, fig. 4; *Hemionitis immersa*, *Bory*, (oldest name).

Rhizome short, slender, scaly. Fronds dense, 2—6 or rarely to 10 c.M. long, \pm 8—15 m.M. broad, elongate-oblong to cuneate or spathulate, narrowed gradually from a blunt or rounded apex to the base or a stipe to 2½ c.M. long. Texture coriaceous; no distinct midrib. Sori immersed, often forked or joining; paraphyses clavate or capitate at the apex.

Malaya; Mascarenes.

* (4) **A. parvulum**, *Bl.*, Flor. Jav., II, 78, tab. XXXIV, fig. 3; Bedd., Ferns Br.I., II, tab. CCLXVII.

Rhizome short-creeping, the scales weak, blackish, subulate or subcapillary, crowded more densely at the base of the younger fronds. Fronds close, erect, sessile or stalked, $2\frac{1}{2}$ —10 or rarely to 20 c.M. long, 4—10 or rarely to 20 m.M. broad, ligulate, the apex acute or acuminate, the edge entire, the base narrowed gradually into the stipe. Texture submembranaceous, subpellucid; surfaces naked; costa none, or slightly visible in the lower $\frac{1}{3}$ of the frond; areolae longitudinal, oblong to linear. Sori linear, long and subcontinuous, or short and interrupted, immersed; paraphyses clavate or obpyriform at the apex.

Java, Penang.

* (5) **A. costatum**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 17, tab. II, fig. 2, a—d.

Rhizome short, the scales brown, linear-lanceolate, acuminate, subdenticulate. Fronds dense, sessile, spathulate-cuneate, 10—25 c.M. long, 3—5 c.M. broad, the fully developed ones broadest near the blunt apex, narrowed gradually to the base, the sides nearly straight, entire. Texture coriaceous; costa distinct, but lost in the upper $\frac{1}{8}$ — $\frac{1}{4}$ of the frond. Sori short or long, mostly free, slightly sunk, the edges of the grooves slightly raised; paraphyses linear, the apex capitate-clavate.

Borneo (Hallier, No. 5540).

(6) **A. strictum**, *Mett.*, Ann. Mus. Bot. L. B., IV, 171, tab. VII, fig. 1—3.

Rhizome scaly like the base of the stipes, the scales black, denticulate. Fronds spathulate-lanceolate, 15—50 c.M. long, 2 — $4\frac{1}{2}$ c.M. broad, the base narrowed gradually into the winged stipe, the apex short-acuminate. Texture thinly chartaceous but firm, rigid; colour pale-green; costa visible in the lower half; areolae in 6—8 rows on each side, the central ones moderately elongate, the lateral \pm as broad as long. Sori immersed in the numerous grooves, continuous or interrupted, hardly reticulate; paraphyses branched, the apices globoso-capitate.

New Guinea.

* (7) **A. semicostatum**, *Bl.*, Rac., Flor. Btz., I, 65; *Bl.*, Flor. Jav., II, tab. XXXIII; Christ, Farnkr. d. Erde, fig. 142.

Rhizome creeping, the scales long, linear, narrowed from the base towards

the apex, thin, spreading, 1—2 c.M. long. Fronds 15—50 c.M. long, 2—10 c.M. broad, ligulate, broadest $\pm \frac{1}{3}$ on the way down, acute or acuminate, narrowed gradually to the base or a short stipe, the edge entire. Texture coriaceous; a distinct, often blackish costa in the lower $\frac{1}{4}$ — $\frac{1}{2}$ of the frond; areolae large, long. Sori copious, often joining; paraphyses minute, the apex clavate.

Malaya; Polynesia, Ceylon.

* (8) **A. latifolium**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 64; *Bedd.*, *Ferns Br.I.*, II, tab. CLXXVI; *A. Boryanum*, *Bl.*, *Flor. Jav.*, II, tab. XXXI.

Rhizome short-creeping, the scales fine, linear-lanceolate, denticulate, blackish-brown, at length deciduous. Stipes approximate, 7—20 c.M. long, compressed. Fronds 8—25 c.M. long, 4—12 c.M. broad, roundish, ovate or obovate, the apex more or less suddenly acuminate with a moderate-long point, the edge entire, slightly repand or acutely but not deeply sinuato-lobate towards the apex, bordered by a narrow, pellucid line, the base cuneate or suddenly narrowed and slightly decurrent. Texture coriaceous; no midrib; areolae longitudinal. Sori mostly copious, not often joining; paraphyses linear, the apex clavate-capitate.

Malaya; North. India, South. China.

* (9) **A. plantagineum**, *Klf.*, *Hk. Bk.*, *Syn. Fil.*, 395; *A. reticulatum*, *Bedd.*, *Ferns S.I.*, tab. LII; *Hemionitis plantaginea*, *Cav.*, (oldest name).

Rhizome short; scales linear-lanceolate, acuminate, brown, with the margin beset with sharp, short, spreading, distant teeth. Stipes distinct, approximate, $2\frac{1}{2}$ —15 c.M. long. Fronds spatulate- or ovate-lanceolate, 6—30 c.M. long, $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad, broadest $\pm \frac{1}{3}$ — $\frac{1}{2}$ of the way down, the point acute or acuminate, the margin entire, often slightly thickened, the base narrowed gradually, or sometimes more suddenly in the smaller forms. Texture more or less coriaceous; no midrib or an indistinct one at the base. Sori short or long, deeply sunk, often forked and anastomosing, the edges of the grooves more or less raised or not raised; paraphyses capitate at the apex.

Malaya; Polynesia, North. India, Ceylon.

(10) **A. elongatum**, *Fée*, *Hk.*, *Spec. Fil.*, V, 168; *A. parvulum*, *Bl.*, *var. elongatum*, *Bl.*, *Flor. Jav.*, II, 79.

Stipes 10—15 c.M. long. Fronds elongate-lanceolate, much acuminate, 20—30 c.M. long, $2\frac{1}{2}$ —4 c.M. broad. Texture submembranaceous, pellucid;

no midrib; veins very slender; areolae long, narrow. Sori linear, simple, subcontinuous or short and interrupted; paraphyses clavate or obpyriform at the apex.

Java.

* (11) **A. subfalcatum**, *Brack.*, Hk. Bk., Syn. Fil. 592; *A. Brookkei*, *Hk.*, 2nd Cent. of Ferns, tab. LXXIX.

Fronde 15—50 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, linear or linear-lanceolate, the point acute, the edge entire, the lower part narrowed very gradually to the base or a short stipe, an indistinct midrib in the lower part. Texture \pm coriaceous; surfaces naked; areolae longitudinal, about 2—3 on each side, several times longer than broad. Sori moderately immersed, in 2—6 interrupted lines, hardly uniting; paraphyses linear.

Malaya; Polinesia.

* (12) **A. coriaceum**, *Wall.*, Hk. Bk., Syn. Fil., 393; *A. reticulatum*, *Bedd.*, Handb. Ind. Ferns, 401; Ferns S.I., tab. CCXXXI; *Hemionitis coriacea*, *Don*, (oldest name).

Fronde \pm 15—45 c.M. long, 1—2 c.M. broad, narrowed gradually from the centre to both ends, the apex acute. Texture very thick, surfaces naked; no midrib; areolae long and narrow; veins raised on the upper surface. Sori rather copious, immersed, sometimes confluent; paraphyses linear.

Malaya; North. India.

* (13) **A. ovatum**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 17, tab. II, fig. 1, a—e.

Rhizome short, clothed with linear-subulate or -lanceolate, shortly toothed or entire, brown or black scales. Stipes close, 1—4 c.M. long, firm, erect. Fronds broadly ovate-spathulate, \pm 8—10 c.M. long, the broadest part \pm ovate, 4—6 c.M. broad (at or above the middle), more or less suddenly narrowed into an acuminate point and the lowest \pm cuneate part of the frond, the edge entire, subpellucid. Texture coriaceous; a slender midrib in the lowest part; areolae 1— $1\frac{1}{2}$ c.M. long, to 5 m.M. broad. Sori not copious, slightly sunk, rarely joining, not long, nearer the edge than the central part of the frond, subflabellate; edges of the furrows not raised; paraphyses linear, spirally twisted.

Sumatra (Forbes, No. 2923a).

* (14) **A. callifolium**, *Bl.*, Rac., Flor. Btz., 62; *Bl.*, Flor. Jav., II, tab. XXXV; *A. falcatum*, *Bl.*, l. c., tab. XXXII.

Rhizome creeping; scales linear-lanceolate, acuminate, the margin sharply serrate or subserrate. Fronds approximate, ligulate-lanceolate, 10—50 c.M. long, $2\frac{1}{2}$ —8 c.M. broad, broadest $\pm \frac{1}{3}$ — $\frac{1}{2}$ of the way down, the apex bluntish, acute or acuminate, the margin bordered by a narrow, yellowish, diaphanous line, the lower part narrowed gradually into the short, flat stipe. Texture fleshy or coriaceous; costa wanting, or only present towards the base. Sori linear, continuous or interrupted, free; paraphyses linear, (tortuose in *A. falcatum*, *Bl.*).

Malaya; Polynesia.

* (15) **A. alatum**, *Brack.*, Wilk., Expl. Exp., XVI, 64; *A. callae-folium*, *Bl.*, var. *alatum*, Rac., Flor. Btz., I, 65; *A. semicostatum*, *Bl.*, Hk., Spec. Fil., V, 168, p. p.

Near *A. callifolium*, *Bl.*, but the fronds spatulate-lanceolate, relatively shorter and broader, narrowed gradually into the short, winged stipe, with a more distinct costa in the lower part; paraphyses linear. — Distinct enough from *callifolium* if typical specimens only are viewed, but passing into it by intermediate forms, and it is impossible to draw the line between them.

Malaya, Polynesia.

* (16) **A. reticulatum**, *Hf.*, Hk. Bk., Syn. Fil., 593; Christ, Farnkr. d. Erde, fig. 141; *Hemionitis reticulata*, *Forst.*, (oldest name).

Fronds 15—50 c.M. long, $2\frac{1}{2}$ —8 c.M. broad, broadest about $\frac{1}{3}$ — $\frac{1}{2}$ on the way down, the apex blunt, acute or short-acuminate, the margin \pm pellucid, the lower part narrowed gradually to the base or a short stipe. Texture thin-coriaceous; surfaces naked; costa wanting or present, green, usually visible in the lowest $\frac{1}{4}$ — $\frac{1}{3}$ of the frond; veins distinct. Sori copious, short or long, joining; paraphyses linear. — Very variable, with forms often approaching the other species.

Malaya; Polynesia, Trop. Australia, North India, Madagascar.

(17) **A. Cumingii**, *Fée*, Hk. Bk., Syn. Fil., 593; *A. Grevillei*, *Balf.*, C. Chr., Ind. Fil.

Fronds to \pm 45 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, ligulate, \pm equal in width in the central half, narrowed very gradually to the base and an acute point, with a black midrib in the lower quarter, the central areolae long,

and not more than $1\frac{1}{2}$ m.M. broad. Sori slender, deep-sunk, often joining, the edges of the furrows rather raised; paraphyses linear.

Philippines; Polynesia.

* (18) **A. spathulatum**, v. **A. v. R.**, Bull. Dép. Agr. I. N., 1908 XVIII, 17.

Rhizome creeping; scales dense, brown, linear-lanceolate, the margin beset with sharp, short, spreading, subremote teeth. Fronds 45—55 c.M. long, \pm 15 c.M. broad, broadest \pm $\frac{1}{3}$ of the way down to the base; upper half obovate, short-acuminate at the point, narrowing gradually below into the linear-cuneate lower half and the short stipe. Texture coriaceous; surfaces naked; the lowest part with a distinct, black, but not prominent midrib; central areolae 1—5 c.M. long, \pm 4 m.M. broad. Sori long, very numerous, often joining, deeply sunk; edges of the furrows not raised; paraphyses linear. *Lingga Islands* (Teysmann, No. 16656).

65. LOXOGRAMME, *Presl.*

Sori linear-oblong or linear, forming an acute angle with the costa, rarely subparallel to it; receptacle crossing the areolae. No indusium.

Rhizome more or less creeping. Fronds not articulate to the rhizome, simple and entire; veins immersed, invisible, (so the main ones when present), anastomosing irregularly to form areolae with or without included free veinlets. *Bedd., Handb. Ind. Ferns, fig. 228—229. (*)*

Paleotropical.

a. *Fronds uniform or nearly so.*

b. *Sori subparallel to the costa.*

(1) *L. parallela.*

b.b. *Sori erecto-patent.*

c. *Fronds rather long-stalked.*

(2) *L. grandis.*

c.c. *Fronds sessile or short-stalked.*

d. *Costa flattened above, prominent beneath.*

(3) *L. involuta.*

d.d. *Costa prominent above, flattened beneath.*

(*) Diels referred **Loxogramme** to **Polypodium**, § **Selliguea**, from which it is distinguished by the fronds adherent to the rhizome and the hidden or wanting main veins.

- e.* Sori short (4) *L. lanceolata*.
e.e. Sori long. (5) *L. Blumeana*.
a.a. Fronds distinctly dimorphous, the fertile ones much the narrowest.
 †. Fronds small, under 3 c.M. long. (6) *L. conferta*.
 †.†. Fronds larger, 10—30 c.M. long. (7) *L. iridifolia*.

(1) ***L. parallela*, Copel.**, Polypod. Philipp., 68; *Polypodium parallelum*, C. Chr., Ind. Fil.

Rhizome short, creeping, the scales greyish-reddish-brown, lanceolate, acuminate. Fronds linear, \pm 15 c.M. long, \pm 6 m.M. broad, sessile, erect, acuminate, entire, narrowed gradually to the base. Texture coriaceous; surfaces naked; costa distinct beneath; veins immersed. Sori oblong, in 1 row on each side near the costa, subparallel to it.

Luzon.

* (2) ***L. grandis*, Copel.**, Philipp. Journ., III, 35; *Gymnogramme grandis*, Rac., Flor. Btz., I, 72; *Polypodium Raciborskii*, C. Chr., Ind. Fil.

Rhizome short-creeping, thick, the scales long, lanceolate. Stipes approximate, 10—12 c.M. long. Fronds 70—90 c.M. long, to 10 c.M. broad, lanceolate, acute, the edge entire, the lower part narrowed gradually into the stipe. Texture coriaceous; costa prominent; main veins hidden, parallel; areolae with free veinlets. Sori linear, oblique, reaching nearly to the edge.

Nusa Kembangan (Java).

* (3) ***L. involuta*, Pr.**, Copel., Polypod. Philipp., 68; Bedd., Ferns S.I., tab. L; *Gymnogramme involuta*, Hk., Hk. Bk., Syn. Fil., 387; *Polypodium involutum*, Mett., Christ, Farnkr. d. Erde, 106; *P. scolopendrinum*, C. Chr., Ind. Fil.; *Antrophyum involutam*, Bl., Flor. Jav., II, 87; — var. *gigas*, Copel., Philipp. Journ., III, 35.

Rhizome creeping, the scales large, brown, lanceolate. Fronds scattered, 15—50 c.M. long, 2—5 c.M. broad, ligulate-lanceolate, the point acuminate, the edge entire or somewhat repand, \pm involute, the lower part narrowed very gradually to the base or a short stipe. Texture thick, coriaceous, flaccid; surfaces naked; costa broad, compressed, flattened above, keeled beneath; veins fine, obscure; areolae with few free veinlets. Sori oblique, reaching from the midrib nearly to the margin.

Var. gigas: Fronds to 65 c.M. long, to 8 c.M. broad, caudate, the margin hardly involute, slightly crisped.

Trop. Asia, China, Melanesia; — var. gigas: Laguna (Philippines).

* (4) **L. lanceolata, Pr.**, Copel., Polypod. Philipp., 68; Bedd. Ferns S.I., tab. LI; *Gymnogramme lanceolata, Hk.*, Rac., Flor. Btz., I, 71; *Antrophyum lanceolatum, Bl.*, Flor. Jav., II, tab. XXXVI; *Polypodium loxogramme, Mett.*, Farng., I, Polypod., 112; C. Chr., Ind. Fil.

Rhizome slender, wide-creeping, the scales small, dense, brown, linear-lanceolate. Fronds 15—30 c.M. long, 5—20 m.M. broad, the point acute or acuminate, the edge entire, the lower $\frac{1}{3}$ — $\frac{1}{2}$ narrowed gradually to the base or a short stipe. Texture coriaceous, flaccid; surfaces naked; costa slightly prominent above, flattened beneath; areolae with few free veinlets. Sori oblique, short, near the costa, falling well short of the margin.

Trop. Asia, Polynesia, Japan, Abessinia, Guinea, Mascarenes, Natal.

* (5) **L. Blumeana, Pr.**, Tent. Pterid., 215; *L. avenia & coriacea, Pr.*, l. c.; Bedd., Handb. Ind. Ferns, 593; *Gymnogramme avenia, Bk.*, Rac., Flor. Btz., I, 72; *Antrophyum avenium & coriaceum, Bl.*, Flor. Jav., II, tab. XXXVII; *Polypodium avenium & Blumei, Mett.*, Farng., I, Polypod., 113 & 114; *P. Blumeanum, C. Chr.*, Ind. Fil.; — var. *Forbesii, Rac.*, l. c., 73.

Rhizome creeping, the scales linear-lanceolate, acuminate, brown. Fronds \pm 10—60 c.M. long, \pm 1—7 c.M. broad, linear- or spatulate-lanceolate, acute or acuminate, the edge entire, the lower part narrowed gradually to the acutely cuneate base or a short stipe to 2 c.M. long. Texture coriaceous; surfaces naked; colour mostly darker in the smaller forms (*L. avenia, Pr.*: \pm 10—30 c.M. long, \pm 1—2 c.M. broad), commonly paler in the larger ones (*L. coriacea, Pr.*, = *Blumeana typica*: \pm 25—60 c.M. long, \pm 2 $\frac{1}{2}$ —7 c.M. broad); costa prominent above, flattened beneath; areolae fine, small, with several free veinlets. Sori erecto-patent, reaching nearly from the midrib to the edge.

Var. Forbesii: Like *L. coriacea, Pr.*, but the fronds with the lower half broad, abruptly narrowed at the broadly cuneate or rounded base.

Malaya; Japan; — var. Forbesii: Sumatra (Forbes, Nos. 2234 & 2262a).

(6) **L. conferta, Copel.**, Polypod. Philipp., 68; *Drymoglossum confertum, Copel.*, Perk., Fragm. Philipp., 187; *Polypodium Copelandii, C. Chr.*, Ind. Fil.

Rhizome slender, creeping, clothed with small, thin, brown, lanceolate, acuminate scales. Fronds crowded, on stalks 1—7 m.M. long. Sterile fronds

elliptical, to ± 2 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, naked, entire, cordate, the apex rounded. Texture thick; veins anastomosing; areolae few, without free veinlets. Fertile fronds linear-oblongate, $\pm 2\frac{1}{2}$ c.M. long, $\pm 3\frac{1}{2}$ m.M. broad, the apex rounded, the base narrowed into the stipe. Sori superficial, medial, afterwards filling up the whole under surface.

Mindanao.

* (7) **L. iridifolia**, *Copel.*, Philipp. Journ., I, Suppl., 149; *Gymnogramme iridifolia*, *Christ*, Ann. Btz., XV, 170, tab. XVI, fig. 26, a—b; *Rac.*, Flor. Btz., I, 75; *Polypodium iridifolium*, *Diels*, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 316; C. Chr., Ind. Fil.

Rhizome slender, creeping, the scales brown, lanceolate. Stipes very short, 0—2 m.M. long, 5—15 m.M. apart. Barren fronds 10—50 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, lanceolate, entire, the point acuminate, the lower part narrowed gradually to a rather broad, rounded or truncate base. Texture firm; costa prominent; areolae fine, with free veinlets. Fertile fronds as long as the barren ones or longer, 4—15 m.M. broad, narrowly lanceolate or nearly linear. Sori linear, close, nearly parallel to the costa.

Java, Celebes, Mindanao.

66. SYNGRAMMA, *J. Smith.*

Sori linear, occupying the veins, (some of them sometimes oblong or round), free or joining, with paraphyses. No indusium.

Rhizome creeping or erect, setaceous, hairy or fibrillose. Stipes not articulate to the rhizome. Fronds naked, simple, rarely pinnate; veins pinnately arranged, free or anastomosing more or less copiously.

Malaya and Polynesia.

Arrangement of the sections.

- § 1. **AUSTROGRAMME.** Veins and sori free. (Not in Malaya).
 § 2. **TOXOPTERIS.** Veins uniting $1 \times$ at or near the margin. Sori free. *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 135, D.
 § 3. **CALLOGRAMME.** Veins uniting $2-5 \times$ near the margin. Sori free, or joining like the veins. *Diels*, l.c., fig. 135, E.

- § 4. **HEMIGRAMMA.** Veins anastomosing more or less copiously. Sori free or reticulate. Fronds simple or irregularly incised, dimorphous, the barren ones often forming a rather dense rosette. Main veins present.
- § 5. **DICTYOGRAMME.** Veins and sori as in § 4. Fronds simple or divided, uniform, the barren ones not in a dense rosette. No main veins.

§ 2. **TOXOPTERIS.**

- a.** *Fronds simple, entire, crenate or slightly toothed.*
- b.** *Fronds oblong, not more than 4 × as long as broad.* (1) *S. valleculata.*
- b.b.** *Fronds lanceolate, ligulate or linear, more than 4 × long as broad.*
- c.** *Fronds rounded at the apex.* (2) *S. obtusifolia.*
- c.c.** *Fronds not decidedly rounded at the apex.*
- d.** *Fronds narrow-linear, under 5 m.M. broad, the apex suddenly acute.* (3) *S. Dayi.*
- d.d.** *Fronds broader, at least the barren ones more than 5 m.M. broad, the apex ± acuminate.*
- e.** *Fronds entire; barren fronds not much broader than the fertile ones.* (4) *S. Lobbiana.*
- e.e.** *Fronds crenate or slightly toothed; barren fronds distinctly broader than the fertile ones.* (5) *S. borneensis.*
- a.a.** *Fronds simple, conspicuously toothed with ± spreading, sharp or blunt-oblong, cartilaginous teeth or spines.* (6) *S. cartilagens.*

(1) **S. valleculata, C. Chr.,** Ind. Fil.; *Gymnogramme valleculata*, *Bk.*, Journ. of Bot., XXVI, 325.

Rhizome woody, short-creeping, clothed with short, stiff, subulate, brown scales. Stipes glossy, naked, castaneous, 15—25 c.M. long. Fronds oblong, entire, 10—13 c.M. long, 5½—5 c.M. broad at the middle, deltoid or rather rounded at the base, the apex deltoid or obtuse, the margin revolute. Texture rigid, thick; costa castaneous; surfaces naked; veins immersed, obscure, anastomosing near the margin. Sori immersed, erecto-patent, reaching nearly to the edge.

Borneo.

* (2) **S. obtusifolia**, *J. Sm.*, Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 257; *Gymnogramme obtusifolia*, *Hk.*, Rac., Flor, Btz., I, 70; *Hk.*, Spec. Fil., V, tab. CCIC.

Rhizome creeping, furnished with short, black hairs. Stipes 5—8 c.M. long, naked, flexuose. Fronds 15—25 c.M. long, $\frac{1}{2}$ —1 $\frac{1}{2}$ c.M. broad, ligulate, the apex rounded, the edge entire, the lower part gradually narrowed. Texture \pm coriaceous; surfaces naked; costa prominent; veins distinct or hidden, simple or forked, connected near the edge. Sori medial, often occupying the whole veins.

Java; Sumatra.

(3) **S. Dayi**, *Bedd.*, Handb. Ind. Ferns, Suppl., 100; *Gymnogramme Dayi*, *Bedd.*, Journ. of Bot., XXVI, 5, tab. CCLXXIX, B, fig. 1—4.

Rhizome creeping, slightly fibrillose. Stipes 2 $\frac{1}{2}$ —4 c.M. long. Fronds narrow-linear, subentire or obscurely crenate, 10—18 c.M. long, \pm 4 m.M. broad, narrowed gradually at the base. Texture coriaceous; veins obscure, simple or forked, the apices running into a transverse, continuous or interrupted submarginal vein. Sori linear, narrow, occupying the veins. — The young fronds are less coriaceous, with the veins also less hidden.

Malacca.

* (4) **S. Lobbiana**, *J. Sm.*, Diels, in Engl. & Prantl, Nat. Pfl. Fam. I⁴, 257; *Gymnogramme Lobbiana*, *Hk.*, Hk. Bk., Syn. Fil., 586; *Hk.*, Spec. Fil., V, tab. CCC; — var. *latior*.

Rhizome creeping, woody, the scales small, brown, subulate. Stipes whether or not approximate, 5—50 c.M. long, glossy, dark-coloured. Fronds 12—25 c.M. long, 1—2 c.M. broad, linear-lanceolate, entire, the apex acuminate, the base narrowed or slightly rounded. Texture coriaceous; surfaces naked; veins simple or forked, united by transverse veinlets near the edge. Sori copious.

Var. latior: Fronds 2—4 c.M. broad.

Borneo, Malacca.

* (5) **S. borneensis**, *J. Sm.*, Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 257, fig. 155, C & D; *Gymnogramme borneensis*, *Hk.*, Hk. Bk., Syn. Fil., 586; *Hk.*, Spec. Fil., V, tab. CCCI.

Rhizome creeping, woody, densely clothed with nearly black, fibrillose scales. Stipes nearly wanting or to 5 c.M. long. Fronds 10—45 c.M. long,

$\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, ligulate, narrowed to both ends, but more gradually to the base, the edge slightly toothed or crenate; barren fronds the broadest. Texture coriaceous; surfaces naked; veins immersed, $1-2 \times$ forked; connecting veinlets indistinct, submarginal.

Borneo, Celebes, Amboyna; Fiji.

* (6) **S. cartilagens**, *Diels*, Bull. Dép. Agr. I. N., 1908, XVIII, 18; *Gymnogramme cartilagens*, *Bk.*, *Hk. Bk.*, Syn. Fil., 586; *Ces.*, Fel. di Bor., tab. II, fig. 5—3a; *Neurogramme cartilagens*, *Christ*, Farnkr. d. Erde, 63, fig. 156; *Acrostichum Borneense*, *Burck*, Ann. Btz., IV, 99; *Elaphoglossum borneense*, *C. Chr.* Ind. Fil.

Rhizome short-creeping, densely clothed with small, black or dark-brown hairs or fibres towards the apex. Stipes approximate, $\frac{1}{2}$ —4 c.M. long, (the fertile fronds on the longest stipes in the fully developed plants), fibrillose at the base, the fibrils deciduous. Fronds decidedly dimorphous in the adult plants, but less so in the younger ones; barren fronds $\pm 10-50$ c.M. long, $\pm 2-5$ c.M. broad, lanceolate or linear-lanceolate, narrowed gradually from the middle towards both ends, the apex acuminate, the edge cartilagineous, decoloured, densely fringed with small, cartilagineous, short-oblong, bluntish, spreading or erecto-patent teeth or serratures in the younger or smaller forms, sharply serrato-dentate throughout or towards the base only in the adult or largest forms, which are not rarely subentire towards the apex. Texture coriaceous; surfaces naked; costa prominent beneath, more or less distinctly canaliculate towards the base on the upper side in the larger and sometimes also in the smaller forms; veins close, fine, simple or forked, connected by the cartilagineous edge or by fine, transversal veinlets near the margin. Fertile fronds longer and narrower than the barren ones, not rarely subsimilar to the latter in the smaller forms, $\pm 15-60$ c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, linear-lanceolate or linear, narrowed very gradually towards both ends in the larger forms. Sori occupying the upper $\frac{1}{2}$ — $\frac{9}{10}$ of the fertile fronds, reaching from the costa to the cartilagineous margin, at length often confluent. — The description is taken from the specimens gathered by Mr. Teuscher (described by Dr. Burck as *Acrostichum Borneense*) and Dr. Nieuwenhuis (determined by Dr. Christ).

Borneo.

§ 3. CALLOGRAMME.

✧. *Fronde simple.*

+ . *Fronde entire, oblong, acuminate.* (7) *S. alismifolia.*

+ . + . *Fronde irregularly repand, spathulate-oblong, acute.* (8) *S. vittaeformis.*

✧ . ✧ . *Fronde ternate or quinate.* (9) *S. quinata.*

* (7) ***S. alismifolia***, **J. Sm.**, Copel., Polypod. Philipp., 69; Lond. Journ. of Bot., IV, 168, tab. VII—VIII, fig. B; Bedd., Ferns Br.I., II, tab. CCXL; *Gymnogramme alismaefolia*, *Hk.*, Hk. Bk., Syn. Fil., 386; *Neurogramme alismaefolia*, *Christ*, Farnkr. d Erde, 64; *Diplazium alismaefolium*, *Pr.*, (oldest name); — var. *Wallichii*; *G. Wallichii*, *Hk.*, Hk. Bk., l. c.; *Hk.*, Spec. Fil., V, tab. CCCII; *S. Wallichii*, *Bedd.*, Ferns Br.I., II, tab. CLIII.

Rhizome creeping. Stipes 10—4½ c.M. long, firm, glossy, chestnut-brown, naked, or scaly towards the base. Fronds 12—30 c.M. long, 5—8 c.M. broad, oblong, acuminate, entire, cuneate or rounded at the base. Texture thinly coriaceous; surfaces naked; costa distinct; veins fine, numerous, nearly spreading, uniting 2—3 × near the margin. Sori extending from the midrib to where they join.

Var. *Wallichii*: Fronds narrower, 2—5 c.M. broad, the base narrowed, the veins less spreading.

Malaya; — var. *Wallichii*: *Singapore, Borneo.*

(8) ***S. vittaeformis***, **J. Sm.**, Copel., Polypod. Philipp., 69; Lond. Journ. of Bot., IV, 168, tab. VII—VIII, fig. A; *Gymnogramme vittaeformis*, *Mett.* (not *Hk.*), Hk. Bk., Syn. Fil., 386.

Stipes subtufted, 30—45 c.M. long, naked, flexuose, dark-brown. Fronds 20—30 c.M. long, 5½—7 c.M. broad, spathulate-oblong, the edge irregularly repand, the point acute. Texture subcoriaceous; surfaces naked; veins fine, parallel, forming 1—2 rows of distinct areolae near the edge. Sori extending from the midrib to the edge.

Philippines.

* (9) ***S. quinata***, **Carr.**, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 257; *Gymnogramme quinata*, *Hk.*, Hk. Bk., Syn. Fil., 387; *Hk.*, Spec. Fil., V, tab. CCXCVII; *Neurogramme quinata*, *Christ*, Farnkr. d Erde, 64; — var. *ternata*, *Christ*, Fil. Saras., III, 250.

Rhizome creeping. Stipes 15—40 c.M. long, pale or brown, glossy, naked. Fronds 15—50 c.M. long, 10—20 c.M. broad, quinate. Central pinna the largest, 15—50 c.M. long, 3½—5 c.M. broad, oblong-lanceolate, slightly stalked, the point acuminate, the edges entire, the base narrowed gradually; lateral pinnae subsessile, smaller, the lowest deflexed, distinct or originating from the base of the next higher ones. Texture coriaceous; surfaces naked; veins fine, parallel, uniting near the edge. Sori on all the veins, extending from the midrib to where they join.

Var. ternata: Fronds ternate.

Malaya; Salomon Islands; — var. ternata: Celebes, Sumatra.

§ 4. HEMIGRAMMA.

* (10) **S. Zollingeri**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 257; *Hemionitis Zollingeri*, *Kurz*, Rac., Flor. Btz., I, 64; *Christ*, Farnkr. d. Erde, 60, fig. 147; *Hemigramma Zollingeri*, *Christ*, Philipp. Journ., II, 170; *H. latifolia*, *Copel.*, Philipp. Journ., II, 406, p.p.; — var. *Teysmannii*.

Rhizome erect, the scales dense, brown, with the edges fringed or ciliate. Fronds dimorphous, simple or irregularly incised. Sterile fronds often in a dense rosette, 8—25 c.M. long, 5—10 c.M. broad, sessile or short-stalked, the broadest part nearer the blunt or short-acuminate apex than the base, the lower part narrowed gradually, the very base rounded or truncate. Texture membranaceo-herbaceous; main veins distinct nearly to the edge; venation ± as in *Aspidium*; areolae numerous, fine, with free veinlets. Fertile fronds linear-lanceolate or ligulate, 6—10 c.M. long, 5—7 m.M. broad, the edge subentire or slightly repand, the stipe 4—15 c.M. long. Sori in 2—5 rows, parallel to the costa, united by short, transversal bars.

Var. Teysmannii: Fronds coriaceous, more or less cordate at the very base.

Malaya; — var. Teysmannii: Amboyna (Teymann).

§ 5. DICTYOGRAMME.

§. *Fronds simple. Sori joining.*

*. *Fronds narrowed at the base.*

(11) *S. lanceolata.*

.. *Fronds more or less rounded at the base.*

(12) *S. Hosei.*

S.S. Fronds pinnate, sometimes 2-pinnate at the base.

+. Pinnae short, herbaceous, unequal-sided at the base. Sori free.

(13) *S. Boerlageana*.

+.+. Pinnae long, rather coriaceous, narrowed gradually at the base. Sori often confluent.

(14) *S. pinnata*.

(11) **S. lanceolata**, *Diels* (not *J. Sm.*), in Engl. & Prantl, Nat. Pfl.Fam. I⁴, 257; *Hemionitis lanceolata*, *Hk.*, *Hk. Bk.*, Syn. Fil., 598; *Hk.*, 2nd Cent. of Ferns, tab. LV; *S. Hookeri*, *C. Chr.*, Ind. Fil.

Rhizome short, ascending. Stipes 15—25 c.M. long, firm, erect, brown, naked, glossy. Fronds 15—25 c.M. long, 5½—6½ c.M. broad, ovate or oblong-lanceolate, the point acute, the edge entire or repand, the base narrowed. Texture coriaceous: surfaces naked; midrib distinct; areolae copious, elongate, oblique. Sori copiously reticulate.

New Guinea; Fiji.

(12) **S. Hosei**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 257; *Hemionitis Hosei*, *Bk.*, Journ. of Bot., XXIX, 108.

Rhizome stout, short-creeping. Stipes naked, glossy, castaneous, 15—30 c.M. long. Fronds simple, lanceolate, ± 30 c.M. long, 3½—5 c.M. broad at the middle, more or less rounded at the base. Texture subcoriaceous; surfaces naked; veins fine, distinct, erecto-patent, anastomosing in the barren fronds mainly in the outer ⅓, but in the rather narrower fertile fronds, in which the slender sori are confined to a band ± ½ c.M. broad remote from both midrib and edge, they anastomose copiously.

Borneo.

* (13) **S. Boerlageana**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 19, tab. III (the paraphyses not quite correct).

Stipes tufted, naked, ± 15—25 c.M. long, firm, erect. Fronds pinnate, ± 10—15 c.M. long, triangular, 6—12 c.M. broad. Terminal pinna ± 3—6 c.M. long, simple, lanceolate or more commonly hastate, 1½—4 c.M. broad, narrowed gradually towards the bluntish apex, the base broadly cuneate, the lateral lobes spreading, deltoid, blunt; lateral pinnae in 5—5 pairs, the lowest the largest, 1½—2½ c.M. broad, spreading, unequal-sided, the upper base broadly cuneate, truncate or cordate, the lower more narrowly cuneate, the apex gradually narrowed, blunt, the edge slightly irregularly wavy; higher pinnae erecto-

patent, sublanceolate, but distinctly unequal-sided. Texture herbaceous; surfaces naked; no main veins; veins distinct, anastomosing irregularly; areolae oblong, oblique, resembling those of *Acrostichum*. Sori linear, straight, curved or flexuose, irregularly oblique, intermixed with shorter ones, which are roundish or oblong; capsules intermixed with numerous, densely crowded, jointed, linear paraphyses, which have the upper part narrowly ellipsoidal, deciduous.

Amboyna (Boerlage).

* (14) **S. pinnata**, **J. Sm.**, Lond. Journ. of Bot., IV, 168, tab. VII—VIII, fig. C; *Gymnogramme pinnata*, *Hk.*, Hk. Bk., Syn. Fil., 390; *Hemionitis elongata*, *Brack.*, Wilk., Expl. Exp., XVI, tab. VIII.

Stipes subtufted, 15—50 c.M. long, firm, erect, glossy. Fronds 30 c.M. or more long, nearly as broad, pinnate. Pinnae erecto-patent, to 15—25 c.M. long, 1—2½ c.M. broad, narrowed gradually to both ends, cuneate at the base, the lower ones stalked, simple or pinnate, the edges entire or nearly so. Texture coriaceous; rachis and surfaces naked; areolae narrow, irregular, usually in 2 rows between the costa and edge. Sori linear or oblong, irregular, often confluent.

East-Malayan Isles; Polynesia, Trop. Australia.

67. HEMIONITIS, *Linné*.

Sori linear, continuous along the veins, reticulate, without paraphyses. No indusium.

Rhizome short. Stipes not articulate to the rhizome. Fronds simple or lobed in malayan material, the fertile ones on the longest stipes; veins distinct, anastomosing copiously, bearing the sori throughout. *Hk. Bk.*, Syn. Fil., tab. VI, fig. 59, a; *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam., I⁴, fig. 158, C—D.

A small genus, almost confined to the tropics.

✧. *Barren fronds lanceolate or cordate.*

+ . *Barren fronds lanceolate, repand.*

+ . + . *Barren fronds cordate, entire or crenate.*

✧ . ✧ . *Barren fronds palmately 5-lobed.*

(1) *H. gymnopteroidea.*

(2) *H. arifolia.*

(3) *H. palmata.*

(1) **H. gymnopteroidea**, *Copel.*, Polypod. Philipp., 67; *Hemigramma latifolia*, *Copel.*, Philipp. Journ., II, 406, p. p.; (? *Syngramma Zollingeri*, *Diels*); — var. *major*; *Hemigramma Zollingeri*, *Christ*, var. *major*, *Christ*, Philipp. Journ., II, 170.

Rhizome short, erect, scaly at the extremity. Fronds clustered, the sterile sessile or subsessile, 7—17 c.M. long, $2\frac{1}{4}$ —5 c.M. broad, oblanceolate, obtuse or acute, repand and crisped at the edge, broad or contracted at the base. Texture papyraceo-herbaceous; colour dark-green with light patches; surfaces naked except on the usually scaly costa; areolae rather large, with copious free veinlets. Fertile fronds on stipes 5—15 c.M. long, the fronds 5—10 c.M. long, 2—4 c.M. broad, the lamina often interrupted. Soriferous veins 1—3 on each side, \pm parallel to the costa, occasionally anastomosing; entire lower surface soon covered by the sporangia.

Var. major: Stipes of barren fronds to 16 c.M. long, scaly. Barren fronds simple or pinnatifid, or pinnate at the base. Segments to 3 on each side. Fertile fronds much contracted, 8 c.M. long; segments to 2 on each side. Texture herbaceo-membranaceous.

Philippines.

* (2) **H. arifolia**, *Moore* (not *Bedd.*), *Christ*, *Farnkr. d. Erde*, 60; *H. cordata*, *Hk. & Grev.*, *l.c. Fil.*, tab. LXIV; (not *Rb.*), *Rac.*, *Flor. Btz.*, I, 65; *Bedd.*, *Ferns S.I.*, tab. LIII; *H. cordifolia*, *Rb.*, *Hk.*, *Fil. Exot.*, tab. XXXV; *Asplenium arifolium*, *Burm.*, (oldest name).

Rhizome erect, the scales linear-lanceolate. Stipes erect, polished, reddish-brown or black, furnished with spreading hairs. Fronds dimorphous. Barren fronds cordate, $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, often nearly as broad, with a deep sinus at the base, the edge entire or crenate, the point blunt, the stipe 4—10 c.M. long. Fertile fronds sagittate, the lobes acute; lateral lobes \pm deflexed, the edges entire, straight or concave; the stipe 20—25 c.M. long. Texture herbaceous or coriaceous, \pm flaccid; each lobe with a central rib; veins anastomosing copiously. Sori reticulate.

Trop. Asia.

* (3) **H. palmata**, *L.*, *Hk. Bk.*, *Syn. Fil.*, 398; *Diels*, in *Engl. & Prantl*, *Nat. Pfl. Fam.*, I⁴, fig. 138, C—D.

Rhizome erect. Stipes of barren fronds 5—10 c.M. long, of the fertile fronds 15—30 c.M. long, glossy, dark-chestnut-brown, clothed with soft, yellowish hairs. Fronds 5—15 c.M. each way, palmatifid with 5 nearly

equal, deltoid-lanceolate, crenato-lobate segments, which are sometimes sub-entire or lobed $\frac{1}{3}$ of the way down, those of the barren fronds blunt, of the fertile more acute. Texture herbaceous; surfaces hairy; central ribs radiate, 1 to each segment.

In cultivation. — *Orig. loc.: Trop. America.*

TRIBE IX. GRAMMITIDEAE.

1950

68. MONOGRAMMA, *Schkuhr.*

Sori linear, placed on or close to the costa, often protected by 1—2 lateral outgrowths of leaf-tissue.

Fronds narrow, linear, grass- or rush-like, simple or rarely subdichotomous, not articulate to the creeping rhizome, mostly with no other veins than the costa. Hk. Bk., Syn. Fil., tab. VI, fig. 51; Diels in Engl. & Prantl., Nat. Pfl.Fam., I⁴, fig. 157, B—G.

Tropical.

Arrangement of the sections.

- § 1. **EUMONOGRAMMA.** Fronds more or less approximate; lateral veins wanting.
§ 2. **PLEUROGRAMME.** Fronds tufted; lateral veins present.

§ 1. **EUMONOGRAMMA.**

- ✧. *Fronds gradually dilated towards the apex. Sori with 1 raised outgrowth at the side.* (1) *M. dareicarpa.*
✧✧. *Fronds not gradually dilated towards the apex. Sori flanked by 2 raised outgrowths.*
+. *Sori once or twice interrupted.* (2) *M. trichoidea.*
+ +. *Sori not interrupted.* (3) *M. paradoxa.*

(1) **M. dareicarpa**, *Hk.*, Hk. Bk., Syn. Fil., 375; Hk., Spec. Fil., V, tab. CCLXXXVIII, A.

Rhizome slender, creeping, the scales minute, narrow, brown. Fronds $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, 1 — $1\frac{1}{2}$ m.M. broad near the top, subacute, narrowed gradually downwards but without a distinct stipe, the fertile taller and relatively more slender than the barren ones. Sorus along 1 side of the costa, covered by an outgrowth from it reaching nearly to the margin, occupying the higher part of the frond, not more than 6 m.M. long.

Borneo, Philippines, New Guinea.

* (2) **M. trichoidea**, *J. Sm.*, Hk. Bk., Syn. Fil., 375.

Rhizome slender. Fronds approximate, thread-like. 7—10 c.M. long, the fertile ones dilated in 2—5 places for a space of \pm 5—6 m.M., and there bearing the sori with a raised edge on each side.

Philippines, Borneo.

* (3) **M. paradoxa**, *Bedd.*, Handb. Ind. Ferns, 373; *M. Jung-huhnii*, *Hk.*, Rac., Flor. Btz., I, 57; *Bedd.*, Ferns S.I., tab. CCX; *Hk.*, Spec. Fil., V, tab. CCLXXXIX, B; *Pleurogramme paradoxa*, *Fée*, (oldest name).

Rhizome slender, the scales narrow, lanceolate, finely acuminate. Fronds densely approximate, grass-like, flaccid, 5—30 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ m.M. broad. Sori sunk in a furrow with a raised edge on each side, occupying a large portion of the costa or confined to its upper half.

Ceylon to Hawaii and Australia.

§ 2. PLEUROGRAMME.

- | | |
|--------------------------------------|---------------------------|
| ‡. <i>Sori interrupted.</i> | (4) <i>M. interrupta.</i> |
| ‡.‡. <i>Sori continuous.</i> | |
| + . <i>Sori medial.</i> | (5) <i>M. intermedia.</i> |
| + . + . <i>Sori rather marginal.</i> | (6) <i>M. Loheriana.</i> |
-

(4) **M. interrupta**, *Blk.*, Ann. of Bot., V, 482; *Pleurogramme interrupta*, *Christ*, Farnkr. d. Erde, 55.

Stipes densely tufted, wiry, naked, 5—8 c.M. long. Fronds linear, $2\frac{1}{2}$ —4 c.M. long, \pm 2 m.M. broad. Texture rigidly coriaceous, thick; surfaces naked; veins quite hidden. Sori filling the whole space between the midrib and margin, broken up into a series of pits of unequal length, in which the sporangia are immersed.

New Guinea.

(5) **M. intermedia**, *Copel.*, Philipp. Journ., I, Suppl., 255.

Rhizome erect, slender. Fronds tufted, 4—6 c.M. long, 3—5 m.M. broad, obtuse, the lower part narrowed gradually towards the base. Texture coriaceous; surfaces naked; costa narrow, sulcate above; veins hidden. Sori continuous, medial, immersed, the groove opening towards the margin.

Negros (Philippines).

(6) **M. Loheriana**, *v. A. v. B.*; *Pleurogramme Loheriana*, Christ, Bull. Herb. Boiss., 2e Série, VI, 1006; (? *Vittaria*).

Rhizome short, erect, caespitose. Fronds fasciculate, to 8 c.M. long, to $\pm 2\frac{1}{2}$ m.M. or more broad, ligulate, often obtuse, narrowed at the base into a short stipe. Texture rigidly coriaceous; surfaces naked; costa slender but prominent; veins hidden. Sori narrow, deeply immersed, protected by a very broad outgrowth of leaf-tissue and therefore (apparently) marginal. — Near *P. pusilla*, Christ (*Vittaria pusilla*, Bl).

Luzon, Mindoro.

69. VITTARIA, J. Smith.

Sori long, linear, continuous, commonly sunk in marginal, submarginal or intramarginal, mostly 2-lipped furrows.

Rhizome creeping, rarely erect. Fronds linear, tape- or grass-like, simple or rarely forked, often pendulous, whether or not articulate to the rhizome: veins simple, rarely forked, erecto-patent, often very oblique, free or united by the receptacle, mostly invisible. (*)

Tropical and subtropical.

Arrangement of the sections.

§ 1. **EUVITTARIA.** Sori immersed, \pm marginal, the upper (outer) lip of the furrows not much broader than the lower (inner) one, *not rolled* over the sorus when young. Hk. Bk., Syn. Fil., tab. VI, fig. 56, a; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 157, J—K.

§ 2. **TAENIOPSIS.** Sori rarely superficial, mostly immersed, submarginal or intramarginal, the lips of the furrows \pm wanting or the outer one much broader than the inner and *rolled* over the sorus when young, or the sorus protected by the recurved margin of the frond. Hk. Bk., l. c. fig. 56, b—c; Diels, l. c., fig. 157, L, M & Q.

§ 1. EUVITTARIA.

a. *Fronds simple, or some of them occasionally forked.*

b. *Fronds sessile or subsessile, under 1 c.M. broad in most of the specimens.*

(*) There is no genus whose species are more difficult of determination (Copel.).

- c.** *Fronds generally* $1\frac{1}{2}$ —3 m.M. broad.
- ‡. *Costa distinct, fronds under 1 m.M. broad.* (1) *V. caudiformis*.
- ‡.‡. *Costa indistinct, fronds more than 1 m.M. broad.*
- *. *Fronds small, to 12 c.M. long, 1—1 $\frac{1}{2}$ m.M. broad.*
(2) *V. lloydiifolia*.
- *.*. *Fronds longer, 15—25 c.M. long, 1 $\frac{1}{2}$ —3 m.M. broad.*
(3) *V. angustifolia*.
- c.c.** *Fronds mostly 3—10 m.M. broad.*
- d.** *Sori confined to the upper part of the frond.*
(4) *V. crassifolia*.
- d.d.** *Sori long, continuous, not confined to the upper part of the frond.*
- e.** *Rhizome scales grey.* (5) *V. rigida*.
- e.e.** *Rhizome scales blackish.*
- f.** *Fronds erect.* (6) *V. ensiformis*.
- f.f.** *Fronds pendulous.* (7) *V. elongata*.
- b.b.** *Fronds distinctly stalked, more than 1 c.M. broad in most of the specimens.*
- *. *Barren fronds entire.*
- ×. *Rhizome scaly.* (8a) *V. zosterifolia*.
- ×.×. *Rhizome naked, ± setose at the apex.* (8b) *V. pachystemma*.
- *.*. *Barren fronds crenulato-undulate. Rhizome sparingly setose.*
(8c) *V. crispomarginata*.
- a.a.** *Fronds mostly 2—4-fid, the segments simple or forked again.*
- ‡. *Fronds rather small, sessile or subsessile.* (9a) *V. Bensei*.
- ‡.‡. *Fronds long, stalked.* (9b) *V. Merrillii*.

* (1) **V. caudiformis**, *Bl.*, Enum., 200.

Rhizome creeping, scaly, the scales hair-pointed. Fronds linear-filiform, under 1 m.M. broad, lax, pendulous, elongate, acuminate, revolute. Texture coriaceous; costa visible. Sori marginal.

Java.

* (2) **V. lloydiifolia**, *Rac.*, Flor., Btz., I, 59.

Rhizome short-creeping, the scales dense, short, thin, subentire, brown, narrowed gradually from the base to the fine, acute point. Fronds linear, 1—10 m.M. apart, 3—12 c.M. long, 1—1 $\frac{1}{2}$ m.M. broad, acute, narrowed

gradually towards the base, which is $\pm \frac{1}{2}$ m.M. broad. Texture rigid; costa nearly invisible. Lips of the grooves equal.

Sumatra, Java, Borneo, Celebes.

* (3) **V. angustifolia**, *Bl.*, Rac., Flor. Btz., I, 60.

Rhizome short-creeping, the scales dense, thin, brown, nearly entire, narrowed gradually from the base to the fine, subulate point. Fronds approximate, linear, acute, 15—25 c.M. long, $1\frac{1}{2}$ —3 m.M. broad, narrowed gradually towards the base, which is $\pm \frac{1}{2}$ — $1\frac{1}{2}$ m.M. broad. Texture rigid; costa nearly invisible.

Malacca, Sumatra, Java, Borneo.

* (4) **V. crassifolia**, *Bk.*, Kew Bull., 1893, 212.

Rhizome short-creeping, woody. Fronds densely approximate, sessile, linear, to 10—12 c.M. long, ± 5 m.M. broad, narrowed gradually at the base. Texture coriaceous, very thick; surfaces naked. Sori in the upper part of the frond, the upper lip of the groove projecting a little beyond the lower.

Borneo, Banca.

(5) **V. rigida**, *Kuf.*, Enum. Fil., 193; Bau. & Hk., Gen. Fil., tab. LXVIII; *V. plantaginea*, *Hk. & Grev.*, Ic. Fil., tab. CLXXXVII.

Rhizome creeping. Fronds close, linear, subfalcate, 25—40 c.M. long, 5—8 m.M. broad, the base narrowed, clothed with linear-lanceolate, grey, glossy scales. Texture rigid; costa hardly visible. Sori marginal, continuous, reaching nearly to the apex.

Malaya; Polynesia.

* (6) **V. ensiformis**, *Sw.*, Ges. Nat. Fr., II, 154, tab. VII, fig. 1; *V. ensifolia*, *Bl.*, Enum., 198; *V. elongata*, *Sw.*, var. *ensiformis*, Rac., Flor. Btz., I, 59; *V. intermedia*, *Bl.*, l.c., 199; — var. *dilatata*.

Rhizome creeping, the scales black, thin, narrowed gradually from the base to the acuminate, hair-pointed apex, the margin serrulate. Fronds sessile, approximate, erect, straight or falcate, 10—30 c.M. long, ± 4 —7 m.M. broad, narrowed gradually towards both ends. Texture rigid; costa whether or not visible; veins immersed. Sori marginal; both lips of the grooves equal or the upper projecting a little beyond the lower.

Var. dilatata: Fronds ± 1 — $1\frac{1}{2}$ c.M. broad.

Java, Sumatra; Mascarenes.

* (7) **V. elongata**, Sw., Christ, Farnkr. d. Erde, 55, fig. 125; Bedd., Ferns S.I., tab. XXI; *V. elongata*, Sw., var. *genuina*, Rac., Flor., Btz., I, 59.

Rhizome creeping; scales dense, black, narrowed gradually from the base to a fine, acuminate point, the edge serrulate. Fronds linear, 20—180 c.M. long, 3—10 m.M. broad, pendulous, the apex bluntish or more commonly acute, the lower part narrowed gradually towards the base. Texture subcoriaceous; costa distinct; veins hidden or indistinct. Sori long; lips of the grooves equal or the upper projecting a little beyond the lower.

Trop. Asia & Polynesia, New South Wales.

* (8a) **V. zosterifolia**, Willd., Spec., V, 406; *V. elongata*, Sw., var. *zosterifolia*, Rac., Flor., Btz., I, 59, — var. *pellucida*.

Rhizome woody, wide-creeping; scales blackish, narrowed gradually from the base to the acuminate, hair-pointed apex, the edge serrulate. Stipes scattered, 1—5 c.M. apart, slender, 5—25 c.M. long. Fronds lanceolate or linear-lanceolate, 1—1½ c.M. broad, 20—120 c.M. long, narrowed gradually towards both ends, the apex acute, the base decurrent upon the stipe. Texture coriaceous; costa distinct; veins visible but not very distinct. Sori long, reaching nearly the base and apex of the fronds; lips of the grooves equal or the upper projecting beyond the lower.

Var. pellucida: Texture thinner, pellucid; veins conspicuous.

Malaya; Polynesia, Comores, Mascarenes.

(8b) **V. pachystemma**, Christ, Philipp. Journ., II, 174.

Rhizome creeping, the extremity crowned with black, crisped setae, the rest naked. Stipes approximate, rufo-stramineous like the costa, ± 6 c.M. long. Fronds 25—50 c.M. long, falcate, linear-lanceolate, acute (not long-acuminate) ± 1¼ c.M. broad. Texture succulent; colour lurid-glaucous; costa flat but distinct. Sori submarginal, continuous from the base to the apex; indusium (? lips of the groove) narrow, pale-coloured.

Mindanao.

(8c) **V. crispomarginata**, Christ, Bull. Herb. Boiss., 2e Série, VI, 1007.

Rhizome slender, creeping, sparingly clothed with thin, short, flaccid, brown setae. Stipes approximate, slender, often filiform, green, 8—10 c.M. long, widening gradually into the base of the frond. Fronds to ± 25 c.M. long, the fertile ones the longest. Barren fronds linear-lanceolate, to 1½ c.M.

broad, long-acuminate, with the margin distinctly crenulato-undulate. Texture flaccid; colour livid-glaucous; costa distinct, white; veins few, very oblique, nearly parallel to the costa. Fertile fronds flat, entire. Sori marginal, the lips of the grooves \pm 1 m.M. broad, whitish.

Luzon.

* (9a) **V. Bensei**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 19; XXI. tab. II, fig. 1, a—e.

Rhizome short-creeping, rather climbing, the scales dense, thin, dark-brown, narrowed gradually from the base to a fine, acuminate point, the margin serrulate. Fronds sessile, approximate, few, \pm horizontal, \pm 10 c.M. long, forked near the narrowed base. Segments linear, falcate, \pm decurved, 2—3 m.M. broad, bluntish or acute, simple or forked again. Texture coriaceous; costae indistinct; veins hidden. Sori long, continuous; lips of the grooves equal.

Java (Preanger Regencies, Capt. H. L. Bense).

(9b) **V. Merrillii**, *Christ*, Philipp. Journ., II, 174.

Rhizome horizontally creeping, hardly branched, not caespitose, clothed with erect, black-brown setae. Stipes distant, pendulous, setulose at the base, slender, flexuose, fusco-stramineous, \pm 20 c.M. long. Fronds \pm 60 c.M. or more long, 1—2 c.M. broad, linear, narrowed gradually to the base, simple and acuminate, but the apex often 2—4-fid, the segments falcate, acute, \pm 1½ c.M. long. Texture flaccid; colour pale-green, opaque; costa slender, often inconspicuous. Sori narrow, strictly marginal, with the lips of the grooves pale-coloured.

Mindanao, Mindoro.

§ 2. TAENIOPSIS.

1. *Fronds clothed with long, whitish, stiff hairs.*

(10a) *V. hirta*.

1.1. *Fronds minutely puberulous.*

(10b) *V. subcoriacea*.

1.1.1. *Fronds naked.*

2. *Fronds to 4 m.M. broad.*

3. *Fronds under 2 m.M. broad, the point acute or acuminate.*

+. *Fronds erect, 10—15 c.M. long, at best \pm 1 m.M. broad.*

(11a) *V. parvula*.

(11b) *V. alternans*.

+.+. Fronds pendulous, to 70 c.M. long, $\pm 1-1\frac{1}{2}$ m.M. broad.
(12) *V. longicoma*.

3.3. Fronds 2—4 m.M. broad, the apex blunt or rounded.

4. Rhizome scales dark-brown; midrib keeled above.

(13) *V. pumila*.

4.4. Rhizome scales pale-brown; midrib slightly sulcate above.

5. Veins simple.

(14) *V. debilis*.

5.5. Veins forked.

6. The outer lip of the groove rather thin at the edge.

(15) *V. pusilla*.

6.6. The outer lip of the groove thick at the edge.

(16) *V. sulcata*.

2.2. Fronds mostly 4—10 m.M. broad.

✧. Fronds 10—13 c.M. long, submembranaceous.

(17) *V. amboinensis*.

✧✧. Fronds 15—120 c.M. long.

×. Coriaceous.

(18a) *V. lineata*.

××. Carnoso-herbaceous.

(18b) *V. taeniophylla*.

2.2.2. Fronds 1—3 c.M. broad in most of the specimens.

✧. Costa stout, especially towards the base. Fronds under 50 c.M. long.

(19) *V. scolopendrina*.

✧✧. Costa slender, inconspicuous. Fronds 50—60 c.M. long when fully developed.

(20) *V. philippinensis*.

(10a) **V. hirta**, *Fée*, Mém., X, 12, tab. XXXI, fig. 4.

Fronds linear, nearly filiform, bluntish, clothed with long, stiff, whitish, spreading hairs. Sori extra-marginal; sporangia roundish.

Borneo.

(10b) **V. subcoriacea**, *Christ*, Philipp. Journ., II, 175.

Rhizome (probably) short-creeping, clothed like the base of the fronds with slender, black-brown, setiform scales. Fronds tufted, linear, ± 55 c.M. long, ± 8 m.M. broad, narrowed gradually into a stipe 5—5 c.M. long, the apex broadly subacute, the edge broadly reflexed, adpressed. Texture succulent, brittle; colour black-castaneous in the lower part of the frond, ochraceous-green above; surfaces minutely rugulose and puberulous; costa strong below,

rather inconspicuous and flattened upwards. Sori continuous from below the middle to the apex, quite covered by the reflexed margin.

Palawan.

(11a) **V. parvula**, *Bory*, *Bél.*, *Voy. Bot.*, II, 35.

Fronds erect, linear, subulate, 10—12½ c.M. long, ± 1 m.M. broad. Texture rather rigid.

Java.

(11b) **V. alternans**, *Copel.*, *Philipp. Journ.*, I, *Suppl.*, 157.

Rhizome slender, creeping, densely clothed with minute, subulate, purple-black scales. Fronds seriate, whether or not close, sessile, 10—15 c.M. long, at the utmost 1 m.M. broad, ± ½ m.M. thick, apparently ecostate. Sori intramarginal, in deep grooves, interrupted, not protected by the margin.

Mindanao.

* (12) **V. longicoma**, *Christ*, *Ann. Btz.*, XX, 129.

Rhizome creeping, the scales black, linear, hair-like, finely acuminate, ± 5 m.M. long. Fronds numerous, to 70 c.M. long and longer, 1—1½ m.M. broad, linear, acuminate, the base narrowed. Texture firm; costa prominent, ± keel-shaped; veins hidden, very oblique. Sori narrow, reaching nearly the apex and base of the fronds, continuous, placed in submarginal grooves; lips of the grooves elevated, the outer produced, rolled over the sorus when young.

Borneo.

(13) **V. pumila**, *Mett.*, *Hk. Bk.*, *Syn. Fil.*, 517,

Rhizome short, the scales hardly 5 m.M. long, rather rigid, blackish-brown, lanceolate-subulate, serrulate at the base. Fronds linear-spathulate, 5½—5 c.M. long, ± 3 m.M. broad, obtuse, falcate or straight, narrowed towards the base, the edge much recurved. Texture coriaceous; costa very prominent beneath, keeled above; veins immersed, oblique, short. Sori hidden by the revolute edge, medial between the midrib and margin.

Borneo.

(14) **V. debilis**, *Kuhn*, *Hk. Bk.*, *Syn. Fil.*, 518.

Rhizome creeping, the scales linear, pale-brown. Fronds sessile, tufted, ligulate-linear, obtuse, 2½—5 c.M. long, 2½—4 m.M. broad, narrowed gradually at the base. Texture firm, subcoriaceous; costa distinct beneath from

the base to the apex, slightly sulcate above; veins subimmersed, oblique, simple, free, the apices not or hardly thickened, hardly produced beyond the receptacle in the fertile region. Sori in rather deep grooves between the thickened costal parenchyma and the margin; the margin \pm thin, much produced beyond the costal parenchyma, which is \pm half as broad as the lamina, i. e. the leaf-tissue beyond the soral groove in transversal section \pm as in *V. pusilla*, *Bl.*, but rather spreading.

Borneo.

* (15) **V. pusilla**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 60; *Bull. Dép. Agr. I. N.*, 1908, XXI, 7, tab. II, fig. 2, a—c; *V. falcata*, *Kze*, *Hk. Bk.*, *Syn. Fil.*, 595; *Taenitis pusilla*, *Mett.*, *Ann. Mus. Bot. L. B.*, IV, 172, tab. VII, fig. 11—14; *Pleurogramme pusilla*, *Christ*, *Philipp. Journ.*, II, 175; — var. *minor*; *V. minor*, *Fée*, *Hk.*, *Spec. Fil.*, V, 185.

Rhizome ascending, the scales membranaceous, yellow-brown, lanceolate, not acuminate. Fronds multiseriate, 2—12 c.M. long, \pm 3—4 m.M. broad and nearly as thick, simple or occasionally forked, straight or falcate, linear-ligulate, the apex blunt or rounded, the lower part narrowed gradually to the base. Texture coriaceous, firm; midrib \pm distinct beneath, but lost in the not abruptly thickened fertile region, slightly sulcate above; veins free, oblique, forked, transversed by the receptacle in the fertile part. Sori confined to the upper $\frac{1}{3}$ — $\frac{1}{2}$ of the frond, deeply sunk in submarginal grooves, the thickened costal parenchyma nearly as broad as the lamina or produced a little beyond the margin, which is rather thin, i. e. the leaf-tissue beyond the soral groove in transversal section rather narrowly linear with the base \pm as broad as the extremity or the latter narrowed.

Var. minor: Fronds shorter, less than 3 m.M. broad.

Borneo, Java, Malacca; Ceylon, Queensland.

* (16) **V. sulcata**, *Kuhn*, *Bedd.*, *Handb. Ind. Ferns*, 408; *Bull. Dép. Agr. I. N.*, 1908, XXI, 7, tab. II, fig. 5, a—c; *Taeniopsis falcata*, *Bedd.*, *Ferns Br. I.*, II, tab. CLXXV; *Taenitis simplicivenia*, *Ces.*, *Fel. Becc. Polin.*, 5 & 8.

Rhizome short-creeping, the scales dense, pale-brown, lanceolate. Fronds crowded, ligulate, obtuse, sessile, narrowed towards the base, 5—10 c.M. long, 3—4 m.M. broad. Texture coriaceous; midrib visible in the lower, barren part of the frond, quite lost in the thickened, fertile region; veins forked or rarely with 2 branches, not reaching the margin. Sori in deep, intramarginal furrows, occupying the upper half of the frond. The margin very thick, much

produced beyond the costal parenchyma, i.e. the leaf-tissue beyond the soral groove in transversal section broadly rhomboidal with the basal part \pm as broad as the extremity or the latter broader, more or less obliquely truncate.

Malaya; Ceylon, Polynesia.

(17) **V. amboinensis**, *Fée*, Bedd., Handb. Ind. Ferns, 407; *Taeniopsis amboinensis*, *Bedd.*, Ferns Br.I., I, tab. CXVII.

Rhizome flexuose, contorted, scaly, the scales rigidly toothed at the margin. Fronds stalked, 10—15 c.M. long, 7—10 m.M. broad, falcate, acuminate, tapering gradually downwards. Texture submembranaceous; surfaces naked; costa slender, disappearing below the apex. Sori submarginal, cuticle of the margin resembling a spurious indusium.

Amboyna, Mindoro; Martaban.

* (18a) **V. lineata**, *Sm.* (not *Sw.*), *Rac.*, *Flor. Btz.*, I, 60; *Taeniopsis lineata*, *J. Sm.*, *Bedd.*, Ferns S.I., tab. LIV; *Pteris lineata*, *L.*, (oldest name); — var. *trichoides*, *Christ*, in *Warb.*, *Mons*, I, 57.

Rhizome creeping; scales dense, linear-lanceolate, the base broad, the apex suddenly acuminate into a fine point, the colour blackish-brown. Fronds linear, 15—120 c.M. long, 4—9 m.M. broad, stipitate, narrowed gradually to both ends. Texture coriaceous, costa visible, running from the base to the apex; veins immersed, parallel, oblique. Sori intramarginal, in shallow grooves, protected by the margin of the frond when young.

Var. trichoides: Differs from the type in having the fronds \pm $\frac{1}{2}$ m.M. broad, and the margin so folded over the sorus, that it may scarcely appear.

Trop. Asia, Japan; Trop. & South. Africa; America, from Florida to South. Brasilia; — var. *trichoides: Philippines.*

(18b) **V. taeniophylla**, *Copel.*, *Philipp. Journ.*, I, Suppl., 157.

Rhizome creeping, the scales dense, lanceolate, subentire, brown, at length whitish. Fronds close, 30—55 c.M. long, 5—7 m.M. broad, acuminate, narrowed gradually towards the base. Texture carnosu-herbaceous; costa distinct to the apex, prominent beneath; veins few, hidden. Sori dorsal, superficial.

Luzon.

* (19) **V. scolopendrina**, *Thw.*, *Rac.*, *Flor. Btz.* I, 59; *V. scolopendroides*, *Mett.* (not *Thw.*), *Christ*, *Ann. Btz.*, XIX, 59; *Taeniopsis scolopendrina*, *J. Sm.*, *Bedd.* Ferns S.I., tab. CCXII; *Pteris scolopendrina*, *Bory*, (oldest name).

Fronds 50—45 c.M. long, 1—5 c.M. broad, the point acute, the edge entire, the lower part narrowed very gradually to the base. Texture thick: midrib thick, blackish, grooved, sometimes nearly lost upwards; veins oblique, fine, parallel. Sori in broad, continuous, submarginal grooves, with the firm, unaltered edge of the frond at first rolled over it.

Samoa to Ceylon, Br. India, South. Africa.

(20) **V. philippinensis**, *Christ*, Bull. Herb. Boiss., 2e Série, VI, 1007.

Rhizome creeping, caespitose, crowned with numerous, long, slender, flexuose, setaceous scales. Stipes subfasciculate, \pm 5 c.M. long, widening gradually into the frond. Fronds \pm 50—60 c.M. long, \pm 1 $\frac{1}{4}$ c.M. broad, long-acuminate. Texture flaccido-coriaceous; colour light-green, opaque; costa slender, inconspicuous: veins very oblique, inconspicuous. Sori narrow, intra-marginal, impressed, reaching from the base to the apex of the frond, more or less covered by the \pm recurved, indusium-like margin.

Luzon.

70. TAENITIS, *Willdenow.*

Sori long, linear, continuous or interrupted, superficial or more or less immersed, in 1 line on each side, parallel to the costa or the margin; paraphyses linear, the terminal cell large and thickened; spores globose to tetrahedral. No indusium.

Rhizome hairy. Stipes not articulate to the rhizome. Fronds simple or more commonly pinnate; veins anastomosing copiously. *Hk. Bk.*, Syn. Fil., tab. VI, fig. 57; *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam., I⁴, fig. 161, C—D. Tropical Asia & Fiji.

-
- ✧. *Fronds simple, obtuse.* (1) **T. obtusa.**
 - ✧✧. *Fronds pinnate; pinnae acute or acuminate.*
 - +. *Pinnae \pm 1—4 $\frac{1}{2}$ c.M. broad.* (2) **T. blechnoides.**
 - +..+. *Pinnae \pm $\frac{1}{2}$ —1 c.M. broad.* (3) **T. stenophylla.**
-

(1) **T. obtusa**, *Hk.*, *Hk. Bk.*, Syn. Fil., 396; *Hk.*, Icon. Pl., X, tab. CMXCIV; *T. blechnoides*, *Sw.*, var., C. Chr., Ind. Fil.

Rhizome creeping. Stipes 5—8 c.M. long, slender, glossy, naked. Fronds $5\frac{1}{2}$ —5 c.M. long, 1—2 c.M. broad, undivided, ovate-oblong, the apex blunt, mucronate, the edge entire (sometimes irregularly incised from the effect of injury), the base \pm rounded. Texture subcoriaceous; areolae large, oblique. Sori in a continuous or slightly interrupted line \pm 3 m.M. from the edge, often united at the apex.

Borneo.

* (2) **T. blechnoides**, Sw., Rac., Flor. Btz., I, 61; Bedd., Ferns Br.I., I, tab. LIV; *Pteris blechnoides*, Willd., (oldest name); — var. *interrupta*, Hk., Spec. Fil., V, 188; *T. interrupta*, Hk. & Grev., Ic. Fil., tab. LXIII; — var. *latior*, Christ Ann. Btz., XX, 128.

Rhizome creeping, the scales capillary, short. Stipes 20—70 c.M. long, firm, naked, glossy. Fronds simply pinnate, 15—60 c.M. long, 15—50 c.M. broad. Pinnae \pm 5—7 or more on each side below the terminal one, 15—50 c.M. long, 1—5 c.M. broad, the fertile ones not rarely the narrowest, lanceolate-elongate, the point acuminate, the edge entire or wavy, \pm thickened, the base cuneate; lower pinnae stalked. Texture coriaceous; areolae copious, oblique, without free veinlets. Sori commonly in a continuous line \pm midway between the midrib and edge. — Fronds of the young plants not rarely simple, similar to the pinnae of the adult fronds, but mostly broader.

Var. interrupta: Pinnae smaller; sori commonly in an interrupted line nearer the edge than the midrib, sometimes submarginal.

Var. latior: Pinnae few (\pm 5—5), to \pm $4\frac{1}{2}$ c.M. broad.

Trop. Asia, Fiji; — var. *interrupta*: *Singapore, Lingga, Borneo*; — var. *latior*: *Borneo, Banca*.

* (5) **T. stenophylla**, Christ, Ann. Btz., XX, 129.

Near *T. blechnoides*, Sw. Pinnae \pm 8 on each side, \pm 12 c.M. long, \pm $\frac{1}{2}$ —1 c.M. broad, linear-lanceolate, the fertile ones the narrowest. Costa narrow; veins very fine. Sori submarginal, hardly immersed, continuous or slightly interrupted.

Borneo.

TRIBE X. POLYPODIEAE.

71. POLYPODIUM, *Linné*.

Sori round, oblong or linear, many on each side, superficial or sunk in cavities; the oblong ones often, the linear ones always dorsal, mostly oblique to the costa and margin; the round and oblong ones dorsal, submarginal or on marginal outgrowths, sometimes immersed in saccate, indusium-like depressions of the margin, rarely covered by the raised edges of the cavities or by long-stalked, peltate scales. No indusium.

Rhizome creeping or erect. Stipes articulate to the rhizome, the articulation not always visible in the smaller species. Fronds simple, pinnatifid or pinnate, rarely more compoundly divided; surfaces naked, scaly or hairy, the hairs not stellate; veins free or anastomosing variously.

A very large genus, inhabiting principally the tropics.

Arrangement of the sections and groups.

§ 1. **EUPOLYPODIUM** (incl. *Grammitis*). Veins free. Sori superficial, or immersed in dorsal or submarginal pits, but not as in § 2. Hk. Bk., Syn. Fil., tab. V, fig. 48, a—b; Diels, in Engl. & Prantl. Nat. Pfl.Fam., I⁴, fig. 162.

a. Integrifoliae. Fronds simple, the margin entire, crenate, repand or toothed.

b. Furcatae. Fronds typically forked, the branches simple or forked again.

c. Pinnatifidae. Fronds lobed or pinnatifid. ⁽¹⁾

d. Pinnatae. Fronds \pm pinnate. ⁽¹⁾

e. Compositae. Fronds at least 2-pinnatifid.

§ 2. **PROSAPTIA** (incl. *Acrosorus*) Veins free. Sori terminal on the veins, immersed in \pm marginal, saccate, indusium-like depressions or in adnate, retroflexed portions of the margin. Fronds simple, crenate, lobed to pinnatifid, at best once pinnate. Diels, l. c., fig. 115, C—H (under *Davallia*). ⁽²⁾

⁽¹⁾ The limit between *c. Pinnatifidae* and *d. Pinnatae* being not always distinctly marked, it will often be advisable to consult both groups.

⁽²⁾ Regarding this section see the information in Christ, Farnkr. d. Erde, 305. — Copeland thinks the section would better be treated as 2 distinct genera, **Prosaptia** & **Acrosorus**. Anyhow, the species of this group are *no Davalliae*.

- § 3. **GONIOPHLEBIUM**. Veins forming 1 or more rows of regular areolae \pm as in *Nephrodium*, each with a simple or rarely forked, included, excurrent free veinlet. Sori terminal on the included veinlets, commonly in the costal areolae only. Hk. Bk., l. c., fig. 48, h; Diels l. c. fig. 163, B & D.
- a.* **Crypsinus**. Fronds simple. (Not in Malaya).
- b.* **Synammia**. Fronds pinnatifid. (Not in Malaya).
- c.* **Marginaria**. Fronds pinnate; pinnae broadly adnate to the rachis.
- d.* **Schellolepis** (*Schellopsis*). Fronds pinnate; at least the lower pinnae \pm stalked, or sessile with a cuneate to truncate, not adnate base.
- § 4. **PHLEBODIUM**. Veins forming more or less regular areolae, each with 1—5 included, excurrent free veinlets, which bear the sori on their united tips; costal areolae always destitute of free veinlets and sori. Hk. Bk., l. c., fig. 48, g; Diels, l. c., fig. 163, F & G. (Not in Malaya).
- § 5. **CAMPYLONEURON**. Main veins connected by parallel, straight or curved, transverse veinlets forming regular, subquadrangular areolae, each with commonly 2 or more included, excurrent free veinlets. Sori subterminal on the free veinlets, 1 to each. Hk. Bk., l. c., fig. 48, j; Diels, l. c., fig. 163, H. (Not in Malaya).
- § 6. **PLEOPELTIS**. Veins anastomosing commonly more or less copiously, forming several to numerous, mostly more or less irregular areolae; included free veinlets (if present) directed towards the midrib or more generally spreading in various directions, rarely excurrent but then not as in § 5. Sori round or oblong, very rarely more than $3 \times$ as long as broad. Hk. Bk., l. c., fig. 48, m.; Diels, l. c., fig. 164, B & E. (*)
- a.* **Myrmecophila**. Rhizome inflated, \pm cylindrical or tuberous, soon cavernous and inhabited by ants, the scales peltato-orbicular or wanting. Fronds rarely simple, commonly pinnatifid or pinnate.
- *. **Aspidopodium**. Sori not on marginal tooth- or lobe-like outgrowths. Rhizome-scales present. Diels, l. c., fig. 166, D—E.
- ****Lecanopteris**. Sori on marginal tooth- or lobe-like outgrowths. Rhizome scales wanting. Diels, l. c., fig. 169.
- b.* **Atactosia**. Rhizome not as in *a.* Rhizome scales very rarely roundish or wanting, generally oval, oblong, ovate, lanceolate, linear or subulate. Fronds simple.

(*) This section would perhaps better be treated as a distinct genus.

- *. **Lepisorus**. Main veins indistinct, not rarely wanting.
- **.. **Pleuridium** (incl. *Microsorium*). Main veins distinct nearly or quite to the edge.

c. Chrysopteris. Rhizome and scales as in **b**. Fronds lobed, pinnatifid or pinnate.

- *. **Phymatodes**. Fronds stipitate, lobed or pinnatifid.
- **.. **Phymatopsis**. Fronds stipitate, pinnate.
- ***. **Drynariopsis**. Fronds sessile, lobed, pinnatifid or pinnate. Fertile segments not contracted.
- ****. **Aglaomorpha**. Like *** but the sori restricted to the much contracted higher segments.

§ 7. **SELLIGUEA**. Venation as in § 6. Sori linear or linear-oblong, more than 3 \times as long as broad, continuous or interrupted, rarely abbreviated or round, more or less oblique to the costa. Fronds simple or pinnate. Hk. Bk., l. c., tab. VI, fig. 52, e—f. (*)

Key to the malayan sections and groups.

- a.** Veins quite free.
 - b.** Sori superficial, or immersed in dorsal or submarginal pits, but not as in **b.b.** § 1. **EUPOLYPODIUM**.
 - c.** Fronds simple, lobed, pinnatifid or forked.
 - d.** Fronds simple. § 1, a. **Integrifoliae**.
 - d.d.** Fronds lobed or pinnatifid. § 1, c. **Pinnatifidae**.
 - d.d.d.** Fronds 1—2 \times forked. § 1, b. **Furcatae**.
 - c.c.** Fronds \pm pinnate or more compoundly divided.
 - *. Fronds pinnate, the pinnae distinct or slightly confluent at the base. § 1, d. **Pinnatae**.
 - *.*. Fronds at least 2-pinnatifid. § 1, e. **Compositae**.
 - b.b.** Sori immersed in saccate or indusium-like, marginal or submarginal depressions. Fronds simple to deeply pinnatifid. § 2. **PROSAPTIA**.
- a.a.** Veins free, forked, the anterior branch of the veins not rarely reaching to the posterior branch of the next higher vein so as to form a row of regular costular areolae; the posterior veinlet always free towards the margin of the frond. Fronds deeply pinnatifid. **P. Cesatianum**, Bk.
- a.a.a.** Veins anastomosing more or less copiously.

(*) This section would perhaps better be treated as a distinct genus.

- 1.** Veins forming 1 or more rows of regular areolae \pm as in *Nephrodium*, each with a single, simple or rarely forked, included, excurrent free veinlet. Sori terminal on the free veinlets, often confined to the costal areolae only. Fronds pinnate in malayan material.

§ 3. **GONIOPHLEBIUM.**

§. Pinnae broadly adnate to the rachis. § 3, c. **Marginaria.**

§.§. At least the lower pinnae stalked, or sessile with a cuneate to truncate, not adnate base. § 3, d. **Schellolepis.**

- 1.1.** Veins forming commonly more or less irregular areolae. Included free veinlets wanting or present but then spreading in various directions, rarely directed towards the costa or the margin only, but not \pm as in 1.

2. Sori linear or linear-oblong, more than $3 \times$ as long as broad, more or less oblique to the costa and margin, continuous or interrupted. Fronds simple or pinnate. § 7. **SELLIGUEA.**

2.2. Sori round or oblong, very rarely more than $3 \times$ as long as broad, if much elongated *then always* parallel to the costa or the margin. § 6. **PLEOPELTIS.**

3. Rhizome scales peltato-orbicular or wanting. Rhizome inflated, \pm cylindrical or tuberos, soon cavernous and inhabited by ants. § 6, a. **Myrmecophila.**

†. Fronds simple, pinnatifid or pinnate. Sori not on marginal outgrowths. Rhizome scales peltato-orbicular.

§ 6, a*. **Aspidopodium.**

†.†. Fronds pinnatifid or pinnate. Sori on specialized, marginal, tooth- or lobe-like, mostly inflexed outgrowths of the fertile segments. Rhizome scales wanting.

§ 6, a**. **Lecanopteris.**

3.3. Rhizome scales oval, oblong, ovate, lanceolate, linear or subulate, very rarely wanting or orbicular. Rhizome not as in **3**, solid.

4. Fronds simple. § 6, b. **Atactosia.**

*. Main veins indistinct, often wanting.

§ 6, b*. **Lepisorus.**

.. Main veins distinct nearly or quite to the edge.

§ 6, b**. **Pleuridium.**

4.4. Fronds lobed, pinnatifid or pinnate.

§ 6, c. **Chrysopteris.**

- ‡. Fronds stipitate.
- ×. Fronds lobed or pinnatifid.
 - § 6, c*. **Phymatodes.**
- ×.×. Fronds pinnate. § 6, c**. **Phymatopsis.**
- ‡.‡. Fronds sessile.
 - *. Sori not on much contracted fertile segments.
 - § 6, c***. **Drynariopsis.**
 - *.*. Sori restricted to the much contracted higher segments.
 - § 6, c****. **Aglaomorpha.**

§ 1. EUPOLYPODIUM. a. **Integrifoliae** ⁽¹⁾.

- a. Sori round, not oblong when young; receptacle round.
- b. Fronds small, under $7\frac{1}{2}$ c.M. long.
 - c. Fronds roundish or oblong, nearly as broad as long, naked.
 - (1) P. flabellivenium.
 - c.c. Fronds linear or linear-lanceolate, much longer than broad.
 - d. Fronds naked or sparingly scaly. (2) P. oblanceolatum.
 - d.d. Fronds hairy. (3) P. trichopodum.
- b.b. Fronds larger, more than $7\frac{1}{2}$ c.M. long.
 - §. Sori in 1 regular row on each side of the costa.
 - *. Fronds under 20 c.M. long, naked. (4a) P. oleandroides.
 - *.*. Fronds longer, hairy. (4b) P. ornatissimum.
 - §.§. Sori in 1 or more irregular rows, or scattered irregularly.
 - *. Fronds membranaceous, villose. (5) P. Friderici et Pauli.
 - *.*. Fronds coriaceous, rather naked. (6) P. pleiosoroides.
- a.a. Sori oblong, at least when young; receptacle oblong.
 - 1. Stipes and fronds naked ⁽²⁾.
 - 2. Fronds narrow, under 5 m.M. broad.
 - ‡. Fronds subterete, rush-like. (7) P. bisulcatum.
 - ‡.‡. Fronds flattened, not rush-like.
 - *. Fronds short, under 10 c.M. long. Sori subimmersed.
 - (8) P. Maxwellii.
 - *.*. Fronds longer, more than 10 c.M. long when fully developed. Sori deeply immersed.

⁽¹⁾ See also **P. ludens**, Bk.

⁽²⁾ See also **P. Koordersti**, Christ.

- ×. *Rhizome short-creeping.* (9a) *P. locellatum.*
 ×.×. *Rhizome erect.* (9b) *P. paucisorum.*
- 2.2.** *Fronds 5—15 m.M. broad.*
- 3.** *Sori medial between the midrib and edge.* (10) *P. Stanleyanum.*
- 3.3.** *Sori nearer the midrib than the edge.*
- 4.** *Fronds sessile or subsessile* ⁽¹⁾.
- 5.** *Veins simple.* (11) *P. diminutum.*
- 5.5.** *Veins forked.*
- 6.** *Fronds tufted.* (12) *P. malaicum.*
- 6.6.** *Fronds scattered or approximate.*
- 7.** *Sori rather close.* (13) *P. australe.*
- 7.7.** *Sori rather distant.* (14) *P. adspersum.*
- 4.4.** *Fronds decidedly stalked.*
- §.** *Fronds short, to ± 15 c.M. long.* (15) *P. subselligueum.*
- §.§.** *Fronds longer, more than 15 c.M. long.*
- ×. *Rhizome scales broad.* (16a) *P. fasciatum.*
 ×.×. *Rhizome scales narrow.* (16b) *P. subfasciatum.*
- 1.1.** *Stipes or fronds more or less distinctly hairy, tomentose, furfuraceous or scaly, at least on the midrib or the margin* ⁽²⁾.
- 2'.** *Fronds narrow, to ± 5 m.M. broad.*
- 3'.** *Fronds short, to 5 c.M. long.*
- 4'.** *Stipes short or wanting. Fronds slightly hairy or ciliate.*
- 5'.** *Fronds minute, under 3 c.M. long.*
- 6'.** *Under surface slightly pubescent.* (17) *P. vittariifolium.*
- 6'.6'.** *Both surfaces covered with brown, scattered hairs.* (18) *P. Merrillii.*
- 5'.5'.** *Fronds somewhat larger, ± 3½—5 c.M. long when fully developed.*
- *. *Fronds sessile.* (19) *P. universe.*
 .. *Fronds short-stalked, the stipe scaly.* (20) *P. sucklingianum.*
- 4'.4'.** *Stipes distinct, 1 c.M. long or longer. Fronds densely hairy.* (21) *P. lasiosorum.*

⁽¹⁾ See also *P. subfasciatum*, *Rst.*

⁽²⁾ See also *P. australe*, *Mett.* & *ornatissimum*, *Rst.*

3.3. Fully developed fronds \pm 5—15 c.M. long.

†. Sori scattered. (22) *P. pilosiusculum*.

†.†. Sori seriate.

*. Sori ultimately confluent.

+. Fronds very narrow, \pm 2—3 m.M. broad.

×. Fronds entire, the hairs scattered.

(23) *P. Christii*.

×.×. Fronds distantly crenato-dentate, the hairs often placed in stellate clusters.

(24) *P. Koordersii*.

+.+. Fronds somewhat broader, 4—5 m.M. broad.

(25) *P. parasiticum*.

.. Sori rather distant.

+. Sori superficial,

×. Stipes tufted.

△. Fronds membranaceous, hairy.

(26) *P. hirtellum*.

△.△. Fronds coriaceous, slightly furfuraeous on the midrib.

(27) *P. subvenosum*.

×.×. Stipes seriate, approximate.

(28) *P. congenerum*.

+.+. Sori immersed.

(29) *P. caespitosum*.

2.2. Fully developed fronds 5—10 m.M. broad.

α. Fronds under $12\frac{1}{2}$ c.M. long.

β. Hairs of the fronds in stellate clusters.

×. Upper surface hairy. (30a) *P. Jagorianum*.

×.×. Upper surface naked. (30b) *P. pleurogrammoides*.

β.β. Hairs of the fronds not in clusters, scattered.

γ. Veins simple or once forked.

δ. Sori round.

ε. Texture thin, submembranaceous.

(31) *P. Reinwardtii*.

ε.ε. Texture firmer, subcoriaceous.

*. Surfaces thinly hairy. (32) *P. Hookeri*.

.. Surfaces densely hairy.

(33) *P. mollipilum*.

δ.δ. Sori oblong.

(34) *P. Havilandii*.

$\gamma.\gamma.$ Veins pinnate or $2 \times$ forked.

+. Stipes very short. (35) *P. knutsfordianum*.

+.+. Stipes distinct, to 8 c.M. long or longer.

(56) *P. intromissum*.

$\alpha.\alpha.$ Fully developed fronds more than $12\frac{1}{2}$ c.M. long.

$\beta.$ Stipes scabrous; fronds nearly naked. (57a) *P. scabristipes*.

$\beta'.\beta'.$ Stipes hairy; fronds naked. (57b) *P. dolichosorum*.

$\beta'.\beta'.\beta'.$ Stipes hairy; fronds hairy, at least on the costa.

$\gamma'.$ Sori in 1 regular row on each side of the costa.

$\delta'.$ Margin ciliate. (58) *P. setosum*.

$\delta'.\delta'.$ Margin not ciliate. (59) *P. pubinerve*.

$\gamma'.\gamma'.$ Sori in 1—2 irregular rows on each side of the costa (*). (40) *P. diplosorum*.

2'.2'.2'. Fully developed fronds more than 10 m.M. broad.

p. Sori in 1 row on each side of the costa.

q. Veins simple or once forked.

r. Fronds entire or slightly repand.

s. Fronds \pm sessile or subsessile.

(41) *P. padangense*.

s.s. Fronds distinctly stalked. (42) *P. obscurum*.

r.r. Fronds sinuate. (43) *P. Warburgii*.

q.q. Veins pinnate or $2-3 \times$ forked. (44) *P. setigerum*.

p.p. Sori in 2—3 rows on each side of the costa.

(45) *P. sumatranum*.

(1) **P. flabellivenium**, *Bk.*, Hk. Bk., Syn. Fil., 322; Christ, Farnkr. d. Erde, fig. 207; Ces., Fel. di Bor., tab. III, fig 7—7a.

Rhizome wiry, slender, creeping, the scales thin, brown, lanceolate. Stipes $2\frac{1}{2}$ —5 c.M. long, filiform, naked, black, glossy. Fronds roundish or broadly oblong, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long and nearly as broad, the point subacute, the edge thick, cartilagineous, deeply and sharply toothed, or with shallow, rounded lobes or sometimes subentire, the base slightly cordate, broadly rounded or \pm cuneate. Texture coriaceous; surfaces naked; veins subflabellate, once or more times forked, the ultimate forks sometimes uniting before they reach the border. Sori 1—6 to a frond, all in the upper half, not far from the margin.

Borneo.

(*) See also **P. setosum**, *Pr.*

(2) **P. oblanceolatum**, *Blk.*, Ann. of Bot., VIII, 128.

Stipes short, filiform, naked. Fronds simple, $5\frac{1}{2}$ —5 c.M. long, \pm 5 m.M. broad, oblanceolate, obtuse, crenulate, the base narrowed gradually. Texture membranaceous; surfaces naked or covered with some few brown, subulate scales; veins visible, not close, forked. Sori superficial, subglobose, in 1 row on each side, medial between the midrib and margin.

New Guinea.

(5) **P. trichopodium**, *Muell.*, Journ. of Bot., XXVIII, 107.

Rhizome short-creeping, the scales dense, small, ovate, brown, imbricate. Stipes slender, approximate, brown, wiry, clothed with inconspicuous, fine, spreading hairs. Fronds simple, linear, obtuse, entire, $2\frac{1}{2}$ —5 c.M. long, 4—5 m.M. broad, narrowed gradually to the base. Texture thick, moderately firm; surfaces clothed with fine, spreading, brown hairs; veins erecto-patent, forked. Sori round, superficial, costal.

New Guinea.

(4a) **P. oleandroides**, *Blk.*, Ann. of Bot., VIII, 128.

Stipes short, nearly $2\frac{1}{2}$ c.M. long, the scales dense, subulate, brown, recurved. Fronds lanceolate, 12—15 c.M. long, 10—15 m.M. broad, entire, obtuse, narrowed at the base, the margin ciliate with few, subulate, brown scales. Surfaces naked; veins conspicuous, erecto-patent, forked near the base. Sori semiglobose, superficial, close, in 1 row on each side near the costa.

New Guinea.

(4b) **P. ornatissimum**, *Rst.*, Fed., Rep., V, 41.

Rhizome slender, caespitose. Stipes densely tufted, slender, 4—6 c.M. long, clothed with soft, spreading, rather long, purple-black hairs. Fronds pendulous, linear-lanceolate, to 60 c.M. long, 1 c.M. broad, acute, entire or slightly undulate, narrowed very gradually towards the base. Texture membranaceous; both surfaces clothed with moderate-long, single or tufted, purple-black hairs, the upper side covered with many, round, white, peltate, cretaceous scales; costa black, prominent beneath; veins 1—2 \times forked, thickened at the apex. Sori terminal, in 1 row on each side near the costa; (receptacle oblong?); sporangia naked.

New Guinea.

(5) **P. Friderici et Pauli**, *Christ*, Ann. Bot., 148, tab. XVI, fig. 21, a & b.

Rhizome short, the scales brown. Stipes tufted, rigid, castaneous, \pm 3 c.M. long, densely clothed with dark-coloured, spreading hairs. Fronds 15—20 c.M. long, \pm 1½—2 c.M. broad, lanceolate, the apex and base narrowed, the point bluntish. Texture membranaceous, thin; colour green, paler beneath; midrib prominent; main veins zigzag, pinnate; vein tips thickened; both surfaces villose. Sori terminal on the 5—4 lower veins so as to form several irregular rows on each side of the midrib.

Celebes.

(6) **P. pleiosoroides**, *Copel.*, Polypod. Philipp., 120.

Rhizome short, creeping, clothed with broad, lanceolate, acuminate, stramineous scales. Stipes slender, 2—4 c.M. long, densely provided with stramineous, spreading hairs. Fronds lanceolate, 10—15 c.M. long, 1—1½ c.M. broad, narrowed towards both ends, the margin sinuate especially in the lower part. Texture coriaceous; surfaces naked, but the edges ciliate with a few short hairs; veins free, immersed, hidden, pinnate or to \pm 4 \times forked. Sori round, large, superficial, in 1—3 irregular rows, or altogether scattered.

Philippines.

(7) **P. bisulcatum**, *Hk.*, Hk. Bk., Syn. Fil., 521; *Grammitis bisulcata*, *Hk.*, Icon. Pl., X, tab. CMXCVIII.

Rhizome creeping, slender, the scales narrow, brown. Fronds scattered, sessile or short-stalked (stalks under 2½ c.M. long), resembling the stipe of a rush, subterete, 15—25 c.M. long, less than 2½ m.M. broad and thick, with 5 furrows, of which the 2 lowest are the deepest. Texture coriaceous; surfaces naked; no veins. Sori oblong, vertical, alternate, placed in the deepest furrows.

Borneo.

* (8) **P. Maxwellii**, *Bk.*, Kew Bull., 1893, 211.

Rhizome short-creeping, the scales dense, erect, lanceolate, membranaceous, pale-brown. Stipes short, naked, approximate. Fronds simple, 2½—6½ c.M. long, \pm 3½—4 m.M. broad, \pm entire, obtuse, narrowed at the base. Texture firm; surfaces naked; veins distant, erecto-patent, forked, not reaching the margin. Sori in 1 row on each side, subdistant, rather close to the costa, subimmersed, oblong.

Borneo.

(9a) **P. locellatum**, *Bk.*, Journ of Bot., XXVIII, 108.

Rhizome short-creeping, the scales dense, small, ovate-lanceolate, pale-brown. Stipes slender, naked, wiry, erect, 5—8 c.M. long. Fronds simple, linear, entire, 12—15 c.M. long, 4—4½ m.M. broad at the middle, narrowed gradually to both ends. Texture thick, rigid; surfaces naked; veins hidden. Sori oblong, oblique, immersed in deep pits, forming a long row on each side nearer the midrib than the margin.

New Guinea.

(9b) **P. paucisorum**, *Copel.*, Philipp. Journ., II, 157, tab. III, B.

Rhizome erect, short, scaly. Fronds tufted, sessile, 8—12 c.M. long, 5—4 m.M. broad, sublinear, subacute, entire, narrowed gradually towards the base. Texture subcoriaceous; surfaces naked; veins forked, hidden. Sori in 1 row on each side, medial between the costa and margin, few, distant, deeply immersed, tending slightly towards oblong.

Mindoro.

(10) **P. Stanleyanum**, *Bk.*, Journ. of Bot., XXVIII, 107.

Stipes naked, 1—1½ c.M. long. Fronds linear, obtuse, entire, 7—13 c.M. long, 6—9 m.M. broad, narrowed gradually from the middle to the base. Texture very thick and rigid; surfaces naked; veins quite hidden. Sori globose, superficial, in 1 row on each side medial between the midrib and margin, confined to the upper part of the frond.

New Guinea.

(11) ? **P. diminutum**, *Bk.*, Hk. Bk., Syn. Fil., 507.

Rhizome short-creeping, the scales dense, firm, lanceolate, dark-brown. Fronds subsessile, narrow, ligulate, blunt, 10—15 c.M. long, 5—10 m.M. broad at the middle, narrowed gradually to the base. Texture rigidly coriaceous, thick; surfaces naked, pale-green; midrib very prominent beneath; veins close, immersed, obscure, simple, reaching the edge, erecto-patent (very oblique). Sori oblong, oblique, placed close to the midrib in the upper 1/3 of the frond.

A plant found in New Guinea differs a little from the Lord-Howe's-Island typical plant in scales, stipes and sori.

(12) **P. malaicum**, *v. A. v. B.*; *P. sessilifolium*, *Hk.*, Hk. Bk., Syn. Fil., 522; *Hk.*, Spec. Fil., IV, 168 (not 251), tab. CCLXXII, A; *Bedd.*, Ferns Br.I., I, tab. CXIX; *P. subvenosum*, *Bk.*, var., C. Chr., Ind. Fil.; *Grammitis sessilifolia*, *J. Sm.*, Hist. Fil., 181.

Fronds sessile, tufted, 7—25 c.M. long, 5—7 m.M. broad, narrowed gradually towards the base, the edge entire or slightly undulate, the point narrowed, bluntish. Texture subcoriaceous; surfaces naked; veins oblique, forked, not reaching the margin. Sori oblong, distant, in 1 row on each side near the midrib, occupying the anterior veinlets.

Malaya.

(13) **P. australe**, *Mett.*, Hk. Bk., Syn. Fil., 322; Christ, Farnkr. d. Erde, fig. 204; *Grammitis australis*, *R. Br.*, Prod., 146; *G. Billardieri*, *Willd.*, Kze., Anal., tab. IX, fig. 2; *P. Billardieri*, *C. Chr.*, Ind. Fil.

Rhizome creeping, copiously short-branched, forming a dense mass. Fronds short-stalked or subsessile, 2½—15 c.M. long, 5—15 m.M. broad, the smaller ones spatulate, the larger ligulate, obtuse at the apex, tapering gradually towards the base. Texture coriaceous; stipes and surfaces naked or slightly ciliate, the edge not bordered; veins oblique, often forked. Sori copious, nearly close, in 1 row on each side, nearer the midrib than the edge, ± parallel to the costa, terminal on the anterior veinlets.

Philippines, New Guinea; Australia, Antarct. America, South. Africa.

(14) **P. adpersum**, *Bl.*, Hk., Spec. Fil., IV, 168; *Grammitis adpersa*, *Bl.*, Flor. Jav., II, 115, tab. XLVIII, fig. 2; *P. hirtellum*, *Bl.*, var., *C. Chr.*, Ind. Fil.

Rhizome creeping or ascending, the scales ovate-oblong, acuminate, entire, ochraceous, membranaceous, deciduous. Fronds sessile or very short-stalked, scattered, linear-lanceolate, 7—15 c.M. long, 6—8 m.M. broad, bluntish, the edge subrepand, the base narrowed. Texture coriaceous; surfaces naked, but the upper side sparingly covered with blackish dots; costa distinct in the lower, obsolete in the upper half; veins hidden, forked, not reaching the margin. Sori rather distant, oblong or subrotund, near the midrib, terminal on the short, anterior veinlets.

Java, Singapore.

(15) **P. subselliguum**, *Bk.*, Journ. of Bot., XXVIII, 107.

Stipes naked, 3½—5 c.M. long. Fronds simple, linear, entire, 12—15 c.M. long, 6—7 m.M. broad at the middle, narrowed gradually to both ends. Texture thick; surfaces naked; veins quite hidden. Sori oblong, superficial, 4—7 m.M. long, confined to the middle of the frond, in 1 row on each side, close to the midrib and parallel to it.

New Guinea.

* (16a) **P. fasciatum**, *Pr.* (not *Mett.*), *Rac.*, *Flor. Btz.*, I, 88, *Grammitis fasciata*, *Bl.*, *Flor. Jav.*, II, 117, tab. XLIX, fig. 1.

Rhizome strong, horizontally creeping, clothed with broad, grey or ferruginous scales. Stipes $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. apart, $2\frac{1}{2}$ —10 c.M. long, rigid, naked, slightly scaly at the base, dark-colored. Fronds to 30—45 c.M. long, 5—10 m.M. broad, linear, narrowed gradually towards both ends, the edge entire. Texture coriaceous; surfaces naked; midrib prominent, conspicuous; veins immersed, once or rarely twice forked, with the branches reaching nearly to the edge. Sori oblong (elliptical when young), close to the midrib.

Malaya.

(16b) **P. subfasciatum**, *Rst.*, *Fed.*, *Rep.*, V, 41.

Rhizome short-creeping, slender, clothed with pale-rufescent, narrow, lanceolate, entire scales. Stipes 0—2 c.M. long, stramineous, naked. Fronds linear-lanceolate, to 25 c.M. long, 12 m.M. broad, narrowed gradually towards the base, the margin entire, reflexed (towards the base?). Texture coriaceous, thick, the margin thinner; surfaces naked, upper bright-green, lower paler, costa prominent beneath; veins immersed, subdistinct, 1—2 \times forked or subpinate. Sori at the base of the anterior veinlets, in 1 row on each side near the costa, suboblong, confined to the upper part of the frond, subimmersed; capsules setose.

New Guinea.

(17) **P. vittariifolium**, *C. Chr.*, *Ind. Fil.*; *P. minimum*, *Bk.*, *Journ. of Bot.*, XVII, 41.

Fronds densely tufted, simple, sessile, ligulate, entire, obtuse, 1— $2\frac{1}{2}$ c.M. long, to $2\frac{1}{2}$ m.M. broad, narrowed gradually from the middle to the base. Texture subcoriaceous; upper surface naked, lower slightly pubescent; veins simple, erecto-patent, reaching nearly to the margin. Sori round, superficial, close, forming a single row close to the midrib, confined to the upper $\frac{1}{2}$ — $\frac{2}{3}$ of the frond, in the finest fronds 8—12 in a row.

Borneo.

(18) **P. Merrillii**, *Copel.*, *Polypod. Philipp.*, 119.

Rhizome short, erect, the scales small, brown. Fronds minute, sessile or narrowed into a short stipe, linear-oblongate, obtuse, entire, to $2\frac{1}{2}$ c.M. long, under 2 m.M. broad. Texture coriaceous; surfaces clothed with a pubescence of short, straight, red-brown hairs mostly confined to the upper

side and margin; costa conspicuous; veins simple, immersed. Sori superficial, large, ± 6 on a frond, near the costa, forming a single wavy line.

Philippines.

(19) **P. universe**, *Bk.*, Ann. of Bot., V, 464.

Fronds sessile, densely tufted, simple, linear, $3\frac{1}{2}$ —5 c.M. long, 2—4 m.M. broad above the middle, narrowed gradually towards the base. Texture rigidly coriaceous; surfaces furnished with deciduous, brown hairs; costa distinct; veins hidden. Sori oblong, crowded, in 1 row on each side, confined to the upper part of the frond, filling up the whole space between the midrib and margin.

Penang.

(20) **P. sucklingianum**, *Bk.*, Ann. of Bot., VIII, 128.

Stipes short, clothed with subulate, brown, spreading scales. Fronds simple, linear, $3\frac{1}{2}$ —5 c.M. long, ± 5 m.M. broad, obtuse, crenate, the base narrowed. Surfaces naked; the margin minutely ciliate; veins distant, erecto-patent, simple. Sori oblong, superficial, oblique, close, in 1 row on each side near the costa.

New Guinea.

* (21) **P. lasiosorum**, *Hk.*, Spec. Fil., IV, 166; *Bedd.*, Ferns Br. I., II, tab. CLXXII; *Grammitis pusilla*, *Bl.*, var. *lasiosora*, *Bl.*, Flor. Jav., II, 110, tab. XLVI, fig. 6; *P. hirtellum*, *Bl.*, var., C. Chr., Ind. Fil.

Rhizome short. Stipes tufted, ± 1 —2 c.M. or more long, slender, clothed with spreading, ferruginous hairs. Fronds spatulate- or ligulate-lanceolate, 2—4 c.M. long, $2\frac{1}{2}$ —5 m.M. broad, the apex rounded, the edge entire, the base gradually narrowed. Texture membranaceous to subcoriaceous; surfaces densely villose and ciliate with patent, soft, ferruginous hairs more or less deciduous in age; veins not discernible. Sori round, commonly occupying the upper half of the frond, crowded in 2 lines close to the costa; capsules copiously intermixed with hairs.

Java.

(22) **P. pilosiusculum**, *Pr.* (not *Hk.*), *Hk.*, Spec. Fil., IV, 168; *Grammitis pilosiuscula*, *Bl.*, Flor. Jav., II, 107, tab. XLVI, fig. 2.

Rhizome creeping, the scales membranaceous, ovate, obtuse, entire, yellowish-fuscous. Stipes 1— $2\frac{1}{2}$ c.M. long, slender, dirty-brown, approximate,

clothed with setiform hairs. Fronds linear or narrowly lanceolate, \pm 8 c.M. long or shorter, \pm 7½ m.M. broad, the apex obtuse, the edge entire or subrepand, the base narrowed. Texture coriaceous; surfaces moderately covered with \pm ferrugineous, setiform hairs. Sori small, scattered, oblong, hardly immersed, \pm 5—6 to each frond.

Java.

(23) **P. Christii**, *Copel.*, Polypod. Philipp., 118.

Rhizome short, the scales deciduous, stramineous. Stipes clustered, 0—15 m.M. long, clothed with brown or reddish, short hairs. Fronds linear, \pm 5—8 or rarely to 15 c.M. long, 2—3 m.M. broad, obtuse or subacute, narrowed very gradually downward, entire, or subentire. Texture subcoriaceous; surfaces bearing scattered, stiff, nearly black hairs; veins immersed, forked. Sori oblong, oblique, superficial, ultimately confluent.

Mindanao.

(24) **P. Koordersii**, *Christ*, Ann. Btz., XV, 144, tab. XV, fig. 19, a—c.

Fronds tufted, stalked or sessile, linear, 6—8 c.M. long, 2—3 m.M. broad, slightly and remotely crenato-dentate, rounded at the apex, narrowed at the base. The whole plant furnished and ciliate with black, strigose hairs, which often are placed in stellate groups or clusters; costa conspicuous; veins simple except a short, basal, soriferous branch. Sori oblong, oblique, approximate, close to the costa, ultimately confluent, confined to the upper part of the frond.

Celebes. — A similar, but quite naked plant was found in Penang.

* (25) **P. parasiticum**, *Mett.*, Hk. Bk., Syn. Fil., 519; Bedd., Ferns S.I., tab. CLXV; Ann. Btz., XV, tab. XV, fig. 19, d; *P. mediale*, *Bk.*, var., C. Chr., Ind. Fil.

Fronds tufted, 5—8 c.M. long, 4—5 m.M. broad, sessile or on stalks to 1 c.M. long, narrowed very gradually below, the edge entire or slightly undulate, the point bluntish. Texture coriaceous; both surfaces when young rather thickly clothed with long, soft hairs; veins immersed, forked. Sori roundish, oblong when young, placed on the anterior branches of the veins, medial between the costa and margin, forming 1 row on each side of the costa, ultimately confluent.

Java, Celebes, Philippines.

* (26) **P. hirtellum**, *Bl.*, Rac., Flor. Btz., I, 86; Bedd., Ferns Br. I., II, tab. CCXII; *Grammitis pusilla*, *Bl.* (*typica* & *var. alpestris*), Flor. Jav., II, 109—110, tab., XLVI, fig. 4—5; *P. pusillum*, *Christ*, Ann. Btz., XV, 145.

Rhizome short, ascending, the scales broadly lanceolate. Fronds linear-lanceolate, densely tufted, 5—15 c.M. long, \pm 2 $\frac{1}{2}$ —5 m.M. broad, narrowed gradually towards both ends, the apex bluntish, the edge entire or slightly repand, the stipe 1—4 c.M. long, slender, brown, provided with soft, spreading, reddish hairs. Texture firm-membranaceous; surfaces hairy like the stipe; veins distinct or more or less hidden, forked near the base, sometimes simple, not reaching the margin, with the branches \pm of the same length. Sori rather distant, in 1 row on each side of the slightly wavy midrib, close to it or nearer the latter than the margin.

Malaya; New Caledonia, South. China.

(27) **P. subevenosum**, *Bk.*, Hk. Bk., Syn. Fil., 520; Bedd., Ferns Br.I., II, tab. CCCXXIII.

Stipes tufted, very short, black, wiry. Fronds ligulate, 7—10 c.M. long, \pm 2—4 m.M. broad, the apex bluntish, the edge slightly repand, the lower part narrowed gradually. Texture subcoriaceous; midrib black, slightly furfureous below; veins simple, short, indistinct. Sori roundish, tending slightly towards oblong, in a long row on each side close to the costa.

Malacca.

* (28) **P. congenerum** (*congener*), *Pr.* (not *Hk.*), Rac., Flor. Btz., I, 88; *P. alatellum*, *Christ*, Fil. Saras., IV, 441; *Grammitis congener*, *Bl.*, Flor. Jav., II, 108, tab. XLVI, fig. 3, a—b.

Rhizome creeping, the scales pale-rufous. Stipes approximate, wiry, more or less densely clothed with rigid, short, spreading, red-brown hairs. Fronds linear-lanceolate, 7 $\frac{1}{2}$ —12 $\frac{1}{2}$ c.M. long, 2 $\frac{1}{2}$ —5 m.M. broad, narrowed very gradually towards both ends, the apex bluntish, the edge entire. Texture coriaceous or thinner; upper surface naked or covered with black dots corresponding with the apices of the veinlets, under surface moderately provided with short, blackish-fuscous hairs; veins branched with 2—4 branches. Sori roundish, elliptical when young, superficial, in 1 row on each side, close to the midrib, placed on the lowest anterior branch of the veins.

Malaya.

* (29) **P. caespitosum**, *Mett.*, Rac., Flor. Btz., I, 87; *Grammitis caespitosa*, *Bl.*, Flor. Jav., II, 106, tab. XLVI, fig. 1.

Rhizome short, scaly. Stipes approximate, 1—2 c.M. long, clothed with numerous, very short hairs. Fronds 5—10 c.M. long, to 4 m.M. broad, linear, narrowed towards both ends. Texture coriaceous; surfaces naked; veins once forked. Sori roundish, elliptical when young, deeply immersed, rather distant, \pm medial between the midrib and margin.

Java, Philippines.

(50a) **P. Jagorianum**, *Mett.*, Hk. Bk., Syn. Fil., 507.

Rhizome 5—10 c.M. long, clothed with pale-brown chaff. Stipes tufted, short or wanting, at best 2 c.M. long, densely clothed with reddish hairs. Fronds ligulate, $2\frac{1}{2}$ —10 c.M. long, 5—8 m.M. broad, entire, obtuse, narrowed gradually towards the base. Texture coriaceous; surfaces clothed (especially near the margin and apex) with firm, usually fascicled, reddish hairs; veins forked, almost invisible. Sori near the midrib, close, oblong, on the anterior fork branch of the veins; capsules setaceous.

Philippines.

(50b) **P. pleurogrammoides**, *Rst.*, Fed., Rep., V, 42.

Rhizome ascending, branched, clothed with ovate-lanceolate, obtuse, slightly adpressed, \pm fuscous, peltate scales, bearing the fronds towards the apex. Stipes scattered or subtufted, nearly $\frac{1}{2}$ c.M. long, purple-black, adpressed to the rhizome. Fronds linear-ligulate, \pm 5 c.M. long, 6—7 m.M. broad, the edge slightly repando-crenate, the base narrowly cuneate. Texture coriaceous; upper surface naked, lower clothed principally near the margin with purple-black, deciduous, forked or stellate hairs, the incisions provided with single, simple, thicker, incurved hairs; costa greenish, prominent on both sides, evanishing in the upper $\frac{1}{3}$; veins immersed, forked. Sori at the base of the anterior veinlets, immersed, oblique, in 1 row near the costa; capsules setose, the setae subulate, geminate, black-purple. — Near *P. Jagorianum*, *Mett.*

New Guinea.

* (31) **P. Reinwardtii**, *Pr.* (not *Mett.*), Rac., Flor. Btz., I, 85; *Grammitis Reinwardtii*, *Bl.*, Flor. Jav., II, 114, tab. XLVIII, fig. 1.

Rhizome slender, creeping, the scales dense, lanceolate, obtuse. Stipes 2—5 c.M. long, blackish-brown, densely clothed with reddish, spreading hairs. Fronds lanceolate or linear-lanceolate, 5— $12\frac{1}{2}$ c.M. long, 5—10 m.M. broad, narrowed towards both ends, the apex bluntish, the edge entire or crenato-repand. Texture thin, pellucid; surfaces (especially the edge and midrib)

hairy like the stipe; costa conspicuous, straight; veins erecto-patent, forked at the base, the soriferous branches very short, the sterile ones not reaching the edge. Sori round, elliptical when young, in a long row on each side, close to the midrib.

Java, Philippines.

(32) **P. Hookeri**, *Brack.*, Hk. Bk., Syn. Fil., 319; *P. setigerum*, *Hk. & Arn.*, Beech. Voy., tab. XXI, A.

Stipes tufted, $1\frac{1}{4}$ —4 c.M. long, clothed with soft, spreading hairs. Fronds narrow-lanceolate, 7—10 c.M. long, 5—10 m.M. broad, tapering at both ends, the point acute or bluntish, the edge entire. Texture rather thinly subcoriaceous; surfaces thinly clothed with short, soft hairs; veins simple or forked. Sori round, in a long row on each side, close to the midrib.

Philippines; Polynesia, North. Australia.

(33) **P. mollipilum**, *Bk.*, Journ. of Bot., XXVIII, 107.

Stipes slender, brown, wiry, $2\frac{1}{2}$ —4 c.M. long, clothed with fine, spreading hairs. Fronds simple, linear, entire, 7—10 c.M. long, 6—7 m.M. broad at the middle, narrowed gradually to both ends. Texture moderately firm and thick; surfaces densely clothed with fine, spreading, brown hairs; veins forked. Sori round, superficial, costal.

New Guinea.

(34) **P. Havilandii**, *Bk.*, Trans. Linn. Soc., Bot., 2nd Series, IV, 255.

Stipes caespitose, slender, nearly $2\frac{1}{2}$ c.M. long, clothed with fine, soft, spreading hairs. Fronds simple, lanceolate, 5—8 c.M. long, $\pm 7\frac{1}{2}$ m.M. broad at the middle, obscurely crenate, narrowed gradually towards both ends. Texture subcoriaceous; surfaces pubescent; veins obscure, forked. Sori oblong, oblique, subremote, in 1 row on each side close to the costa.

Borneo.

(35) **P. knutsfordianum**, *Bk.*, Journ. of Bot., XXVIII, 107.

Rhizome short-creeping, the scales dense, small, pale-brown, ovate, imbricate. Stipes very short, densely clothed with spreading, brown hairs. Fronds simple, linear, entire, 7—10 c.M. long, 6—7 m.M. broad, obtuse, narrowed gradually to the base. Texture firm; surfaces clothed with few, spreading, brown hairs; veins distinct, with 2 short anterior branches. Sori round, superficial, in 2 rows near the midrib.

New Guinea.

(36) **P. intromissum**, *Christ*, Fil. Saras., IV, 440; Ann. Btz., XV, tab. XV, fig. 20, a—b.

Stipes filiform, 8 c.M. and more long. Fronds 5—8 c.M. long, 7—10 m.M. broad, ovate-lanceolate, apex and base rounded. Texture papyraceous, thin; surfaces clothed like the stipe with reddish-brown, spreading hairs; costa distinct; veins forked and the branches branched again. Sori in 1 row on each side near the midrib, on the lowest veinlets, at length confluent.

Celebes.

(37a) **P. scabristipes**, *Blk.*, Journ. of Bot., XXVIII, 108.

Stipes brown, wiry, scabrous, 5—8 c.M. long. Fronds simple, linear, entire, 12½—15 c.M. long, 8—9 m.M. broad at the middle, narrowed towards the base. Texture thick, rigid; surfaces nearly naked; veins forked, erecto-patent. Sori oblong, oblique, rather immersed, in 1 row on each side nearer the midrib than the margin.

New Guinea.

(37b) **P. dolichosorum**, *Copel.*, Philipp. Journ., I, Suppl., 159, tab. XVI.

Rhizome wide-creeping, clothed with rather large, ovate, fulvous scales. Stipes firm, black-brown, ± 4 c.M. long, clothed with short, reddish hairs, or naked towards the base. Fronds linear-lanceolate, ± 15 c.M. long, ± 1 c.M. broad, narrowed towards both ends, the apex blunt, the edge entire, the base cuneate. Texture coriaceous; surfaces naked, opaque; veins often forked. Sori oblong, oblique, nearer the midrib than the margin, slightly immersed.

Mindanao.

* (38) **P. setosum**, *Pr.* (not *Bl.*), Rac., Flor. Btz., I, 87 p.p.; *Grammitis setosa*, *Bl.*, Flor. Jav., II, 116, tab. XLVIII, fig. 3; *P. diplosorum*, *Christ*, var., C. Chr., Ind. Fil.; — var. *calvum*, *Copel.*, Philipp. Journ., II, 136.

Rhizome short-creeping, scaly, the scales lanceolate, yellow-brown, Stipes wiry, 2—5 c.M. long, dirty-brown, clothed with spreading, soft, red-brown hairs. Fronds approximate, erect, 12½—40 c.M. long, ± 7—10 m.M. broad, linear-lanceolate, narrowed gradually towards both ends, the margin and costa on both sides more or less clothed with dark-reddish-brown hairs. Texture ± coriaceous; veins slightly conspicuous, 2—3 × forked, the soriferous branch short; costa prominent below. Sori round (elliptical when young), approximate, placed in 1 row on each side ± close to the midrib, someti-

mes accompanied by a few others placed in an outer row close to the former.

var. calvum: Upper surface sparingly pilose.

Java, Celebes, Philippines; — var. *calvum:* *Mindoro.*

(39) **P. pubinerve**, *Christ*, Ann. Btz., XV, 146; *Grammitis pubinervis*, *Bl.*, Flor. Jav., II, 117, tab. XLVIII, fig. 4.

Rhizome suberect, scaly. Stipes approximate, clothed with short, spreading hairs. Fronds linear-lanceolate, 20—25 c.M. long, \pm 10 m.M. broad, narrowed towards both ends. Texture coriaceous; costa prominent below, hairy; surfaces naked; veins pinnate, conspicuous. Sori oblong, close to the costa, rather distant, occupying the upper $\frac{1}{2}$ — $\frac{2}{3}$ of the frond.

Java, Celebes.

(40) **P. diplosorum**, *Christ*, Ann. Btz., XV, 146; *Grammitis hirta*, *Bl.*, Flor. Jav., II, 111, tab. XLVII, fig. 1.

Rhizome short, clothed with blackish scales. Stipes tufted, rigid, light-brown, 3—4 c.M. long, furnished like the costa with short hairs, sometimes only asperous. Fronds ligulate, \pm 12 $\frac{1}{2}$ —20 c.M. long, \pm 7 m.M. broad, moderately narrowed towards both ends, the edge entire or slightly wavy. Texture coriaceous; surfaces naked or short-hairy; costa blackish and prominent beneath; veins hidden, pinnate on the anterior side; veinlets 3—4, the lowest forked. Sori superficial, round, elliptical when young, in 1 or 2 irregular rows between the midrib and edge, rather distant, the inner row close to the costa.

Java, Borneo, Celebes.

(41) **P. padangense**, *Bl.*, Journ. of Bot., XVIII, 215.

Rhizome short-creeping, the scales dense, imbricate, large, pale-brown, lanceolate, membranaceous. Stipes almost obsolete. Fronds ligulate, entire, \pm 15 c.M. long, 1—1 $\frac{1}{2}$ c.M. broad at the middle, narrowed gradually to the base. Texture very thick and coriaceous; surfaces almost naked, but the margin ciliate with few, minute, black hairs; veins forked. Sori round, slightly immersed, placed in 1 row on each side near the thick midrib, a short space from each other, confined to the upper half of the frond.

Sumatra.

(42) **P. obscurum**, *Mett.*, Farng., I, Polypod., 38; *Grammitis obscura*, *Bl.*, Flor. Jav., II, 113, tab. L, fig. 1; *P. Reinwardtii*, *Pr.*, var., C. Chr., Ind. Fil.

Rhizome creeping, the scales small, scariose, pale-fuscous, lacerate. Stipes approximate, slender, 1—4 c.M. long, clothed with short, fine hairs. Fronds 8—15 c.M. or more long, 1—1½ c.M. broad, linear-lanceolate, the apex obtuse, the edge entire or slightly repand, the lower part narrowed. Texture coriaceous; colour deep-green; the margin and the distinct costa on both sides provided with short, dark-reddish-brown hairs; veins simple or forked, not reaching the margin. Sori suboblong or oval, rather distant, in 1 row on each side near the costa, slightly immersed.

Java.

(43) **P. Warburgii**, *Christ*, Warb., Mons., I, 59.

Rhizome short, the scales adpressed, ovate, inconspicuous, scattered. Stipes short, tufted, to ± 2 c.M. long, clothed with long, black, spreading hairs. Fronds narrow, lanceolate-ligulate, ± 12 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, the apex acuminate, the margin sinuate, the base narrowed gradually. Texture chartaceous; costa and margin clothed or ciliate with black hairs; veins conspicuous, prominent, forked. Sori on the lower veinlets, large, oblong, 30—40 on each side, occupying the upper $\frac{1}{2}$ of the frond.

Batjan.

* (44) **P. setigerum**, *Bl.*, Rac., Flor. Btz., I, 86; *Hk.*, Icon. Pl., X, tab. CMXLI; *Grammitis fasciculata*, *Bl.*, Flor. Jav., II, 112, tab. XLVII, fig. 2.

Rhizome short, densely clothed with brownish scales. Stipes tufted or approximate, 5—10 c.M. long, reddish-black, densely clothed with long, soft, spreading, ferruginous or reddish hairs. Fronds 20—45 c.M. long, 1—2 c.M. broad, linear, the point acute, the base narrowed, the edge entire. Texture thick, coriaceous, succous, thin and translucent when dry; both surfaces hairy like the stipe; veins pinnate or 2—3 \times forked, the soriferous branches the shortest. Sori round, in 1 row on each side near the midrib, distant.

Singapore, Java, Moluccas, Philippines.

* (45) **P. sumatranum**, *Bk.*, Bull. Dép. Agr. I. N., 1908, XVIII, 19; *P. subpleiosorum*, *Rac.*, Flor. Btz., I, 89.

Rhizome short, creeping, the scales dense, lanceolate, pale-brown. Stipes approximate, 2—6 c.M. long, wiry, erect, densely clothed with minute, rigid, spreading, brown hairs. Fronds linear-lanceolate, 15—40 c.M. long, 1½—4 c.M. broad, narrowed gradually towards both ends, the edge repando-crenate to irregularly undulate with the crenations to 4 m.M. broad, to 2—3 m.M.

deep. Texture coriaceous; the edges and midrib beneath very sparingly covered with distant, minute, inconspicuous, blackish-brown hairs; veins hidden, pinnate with 1—3 veinlets on each side, which are very rarely joining at the apex. Sori round, \pm elliptical when young, nearly superficial (very slightly and indistinctly immersed), in 2—3 \pm irregular rows on each side of the midrib. — Beccari's and Raciborski's plants in the Buitenzorg herbarium are quite the same plants.

Java, Sumatra.

§ 1. EUPOLYPODIUM. b. **Furcatae.**

* (46) **P. subdichotomum**, *Rac.*, Bull. Dép. Agr. I. N., 1908, XVIII, 20; XXI, tab. III.

Rhizome short, the scales membranaceous, dense, ferruginous, narrowed gradually from the base to the apex, the edge entire. Stipes tufted or approximate, short, clothed like the whole frond with long, spreading, red-brown hairs. Fronds 10—15 c.M. long, once or more times forked, tapering very gradually into the slender stipe. Segments of the fronds linear, 2—5 m.M. broad, with a gradually narrowed apex. Texture coriaceous; veins hidden, short, forked. Sori in 1 row on each side near the costa, large, alternate, nearly confluent, filling up the whole under surface of the ultimate segments.

Sumatra (Forbes Nos. 2408 & 2497).

§ 1. EUPOLYPODIUM. c. **Pinnatifidae.**

a. *Sori solitary on the lobes. The lower edge of the lobes folded over the sori.*

+ . *Fronds approximate.* (47) *P. gracillimum.*

+ . + . *Fronds densely tufted.* (48) *P. cucullatum.*

a.a. *Sori whether or not solitary. The lower edge of the lobes not folded over the sori* (*).

b. *Fronds narrow, to 5 m.M. broad.*

† . *Fronds rather small, to 2½ c.M. long or somewhat longer.*

(49) *P. alternidens.*

† . † . *Fronds longer, 8—30 c.M. long when fully developed.*

(50) *P. subpinnatifidum.*

b.b. *Fully developed fronds 5—10 m.M. broad.*

c. *Lobes irregular, entire.*

(51) *P. Loherianum.*

(*) See also under **P. cucullatum**, *Nees & Bl.*

c.c. Lobes rather regular, toothed with 2—5 sharp teeth.

(52) *P. denticulatum*.

b.b.b. Fully developed fronds more than 15 m.M. broad.

×. Fronds sessile, the margin deeply pinnatifid. (53) *P. Burbidgei*.

×.×. Fronds stalked, the margin cut at best $\frac{1}{2}$ -way down to the costa.

(54) *P. ludens*.

(47) **P. gracillimum**, *Copel.*, Polypod. Philipp., 121.

Rhizome creeping or ascending, the scales small, brown, lanceolate. Fronds approximate, to \pm 12 c.M. long and 4 m.M. broad, nearly sessile, pinnatifid nearly to the rachis into small, acute, distant, erecto-patent, decurrent, alternate segments. Surfaces naked. Sori 1 to each segment, medial, the lower side of the lobe folded over the sorus.

Philippines.

* (48) **P. cucullatum**, *Nees & Bl.*, Rac., Flor. Btz., I, 89; Christ, Farnkr. d. Erde, fig. 211; *Grammitis cucullata*, *Bl.*, Flor. Jav., II, 119, tab. L, fig. 3; *Callymodon cucullatus*, *Pr.*, Bedd., Ferns S.I., tab. CCXXXIII; — var. *planum*, *Copel.*, Philipp. Journ., II, 157.

Rhizome erect, the scales linear-lanceolate. Stipes densely tufted, very short or nearly wanting. Fronds 5—13 c.M. long, \pm 3—10 m.M. broad, narrowed towards both ends, the edge deeply pinnatifid. Lobes linear-oblong, blunt, entire, erecto-patent, numerous, rather distant, 1—2 m.M. broad, the higher ones fertile, rather shorter and broader, with the lower edge reflexed so as to clasp the sorus. Texture slightly coriaceous; surfaces and midrib often hairy; veins simple, 1 to each lobe. Sori 1 to each lobe, medial.

Var. planum. Segments flattened, large, approximate, orbicular-oblong. Sori large, subsuperficial.

Malaya; Ceylon, Fiji, New Caledonia; — var. planum: Mindoro.

(49) **P. alternidens**, *Ces.*, Fel. di Bor., 25, tab. II, fig. 4—4a.

Rhizome short, sparingly fibrillose. Fronds 5—8, erect, $2\frac{1}{2}$ c.M. long or longer, under $2\frac{1}{2}$ m.M. broad, linear, sessile or subsessile. Lobes 20—24, rather distant, alternate, oblique, dentiform, obtuse. Texture rigid; surfaces naked. Sori oblong, 1 to each lobe.

Borneo, Celebes.

* (50) **P. subpinnatifidum**, *Bl.*, Rac., Flor. Btz., I, 91; *Grammitis subpinnatifida*, *Bl.*, Flor. Jav., II, 118, tab. XLIX, fig. 2.

Rhizome erect, short, clothed with ovate-lanceolate scales. Stipes tufted, slender, 2—4 c.M. long, covered with soft, spreading, red-brown hairs. Fronds linear, 8—30 c.M. long, 3—5 m.M. broad, deeply lobed, the apex and base narrowed. Lobes with the upper edge nearly straight and horizontal, the lower edge oblique and the apex blunt, entire or slightly emarginate, the sinuses \pm roundish. Texture subcoriaceous; both surfaces naked or the midrib hairy beneath; veins forked in the lobes. Sori 1 or rarely 2 to a lobe.

Java, Malacca.

(51) **P. Loherianum**, *Christ*, Bull. Herb. Boiss., VI, 197, tab. III, fig. 2, a—b.

Rhizome short-creeping, scaly. Stipes 1—5 c.M. long, brown, naked. Fronds ligulate-lanceolate, \pm 15 c.M. long, 7½—10 m.M. broad, narrowed very gradually into the stipe, the edge shallowly lobed (rather conspicuously and deeply toothed), the apex bluntish. Lobes deltoid, irregularly shaped, obtuse, to 3 m.M. long. Texture coriaceous; surfaces naked; costa and veins inconspicuous; main veins pinnate in the lobes. Sori round or oblong, in 1 row on each side nearer the midrib than the edge, medial on the lower anterior veins, rarely accompanied by a second, submargina. row.

Philippines.

* (52) **P. denticulatum**, *Pr.*, Rac. Flor. Btz., I, 89; *Grammitis denticulata*, *Bl.*, Flor. Jav., II, 121, tab. L, fig. 4.

Rhizome short, erect, the scales short, brown, lanceolate. Fronds sessile or short-stalked, 4—10 c.M. long, 6—8 m.M. broad, deeply pinnatifid. Lobes deltoid-oblong, erecto-patent, \pm falcate, the point acute, the edge with 2—5 often sharp teeth. Texture coriaceous; surfaces naked; veins with 1—2 branches. Sori 1—3 to a lobe.

Java.

* (53) **P. Burbidgei**, *Bk.*, Journ. of Bot., XVII, 42.

Rhizome suberect, the scales lanceolate or linear, membranaceous, brown, ciliate. Fronds tufted, sessile, lanceolate, deeply pinnatifid, to 30 c.M. long, 2½—5½ c.M. broad at the middle, cut down to a broad wing to the costa into numerous close, entire, obtuse, spreading or ascending segments 2½—5½ m.M. broad at the base, the lower segments growing gradually smaller and

the basal part of the costa bordered only by a moderate-long, entire wing \pm 4—5 m.M. broad at the top and narrowed gradually to the base. Texture subcoriaceous; under surface \pm pubescent on the costa, the rest glabrous; veins distant, simple, erecto-patent, reaching nearly to the edge. Sori semi-globose, slightly immersed, commonly in a single row in the wing of the costa, not or hardly prominent on the upper surface.

Borneo, New Guinea.

(54) **P. ludens**, *Bk.*, Ann. of Bot., VIII, 129.

Rhizome short-creeping, the scales dense, broad-lanceolate, pale-brown. Stipes \pm 5—7 $\frac{1}{2}$ c.M. long, clothed with subulate, recurved, brown scales. Fronds lanceolate, 15—20 c.M. long, 2—3 c.M. broad, the base narrowed, the margin entire or cut nearly $\frac{1}{2}$ -way down to the costa into irregular lobes. Surfaces naked; veins forked, erecto-patent, immersed, obscure. Sori oblong, superficial, oblique, in 1 row on each side near the costa.

Java.

§ 1. EUPOLYPODIUM. d. **Pinnatae.**

a. *Pinnae contiguous, dilated and often confluent at the base; lower pinnae sometimes distant.*

b. *Fronds narrow, to \pm 12 $\frac{1}{2}$ m.M. broad.*

c. *Pinnae entire, the sori solitary on the pinnae.*

§. *Sori terminal on the pinnae.*

*. *Sori superficial.* (55) *P. streptophyllum.*

.. *Sori rather deeply immersed.*

×. *Sori quite terminal on the pinnae.*

(56) *P. triangulare.*

×.×. *Sori occupying the upper half of the pinnae.*

(57) *P. subtriangulare.*

§.§. *Sori at the base of the pinnae.*

+. *Fronds small, under 5 c.M. long, under 5 m.M. broad.*

(58) *P. musgravianum.*

+.+. *Fronds larger, \pm 10—13 c.M. long, \pm 7 $\frac{1}{2}$ m.M. broad.*

(59) *P. conjunctisorum.*

c.c. *At least the largest pinnae \pm crenate or repand, or the sori not solitary on the pinnae.*

- d.** *Stipes naked.*
- e.** *Stipes under 1½ c.M. long; fronds erect.*
- f.** *Costae forked.* (60) *P. cornigerum.*
- f.f.** *Costae pinnate.* (61) *P. inconspicuum.*
- e.e.** *Stipes longer; fronds pendulous.* (62) *P. undosum.*
- d.d.** *Stipes hairy.* (63) *P. minutum.*
- c.c.c.** *Pinnae sharply toothed.* (64a) *P. serrato-dentatum.*
- c.c.c.c.** *Pinnae bluntly toothed.* (64b) *P. Merritti.*
- b.b.** *Fully developed fronds more than 12½ m.M. broad (*).*
- 1.** *Texture herbaceous or papyraceous; fronds often pendulous.*
- ‡.** *Surfaces hairy.*
- ***. *Pinnae entire or slightly crenate to repand.*
- ×**. *Pinnae very narrow, ± 1 m.M. broad.*
(65) *P. subrepandulum.*
- ×.×**. *Pinnae broader, ± 2 m.M. broad.*
- +**. *Fronde short, to ± 6 c.M. long. Sori distant.*
(66) *P. subminutum*
- +.+**. *Fronde longer, 6—10 c.M. long, sori copious, close.*
(67) *P. lancifolium.*
- *.*.** *Pinnae toothed or lobed on 1 or both sides.*
- Δ.** *Pinnae incised on the upper side only, the apex entire or slightly toothed.* (68) *P. subsecundo-dissectum.*
- Δ.Δ.** *Pinnae incised on both sides.* (69) *P. subfalcatum.*
- ‡.‡.** *Surfaces ± naked or glabrescent, but the rachis often hairy.*
- ◇.** *Sori superficial or slightly immersed.*
- +**. *Rachis naked.* (70a) *P. repandulum.*
- +.+**. *Rachis hairy.* (70b) *P. erythrotrichum.*
- ◇.◇.** *Sori deeply immersed.*
- +**. *Sori round.*
- ×**. *Pinnae narrow, ± 1—2 m.M. broad.*
(71) *P. brevifrons.*
- ×.×**. *Pinnae broader, ± 4—8 m.M. broad.*
(72) *P. papillosum.*
- +.+**. *Sori oblong.* (73) *P. obliquatum.*
(74) *P. Schenkii.*

(*) See also **P. inconspicuum**, *Bl.*

1.1. *Texture chartaceous to coriaceous; fronds often erect.* (*)

2. *Pinnae ± hairy or scaly beneath.*

α. *Fronds to $3\frac{1}{2}$ c.M. broad.*

β. *Pinnae entire.*

γ. *Sori immersed.* (75) *P. decipiens.*

γ.γ. *Sori superficial or hardly immersed.*

δ. *Stipes hairy.* (76a) *P. mollicomum.*

(76b) *P. pseudoarticulatum.*

δ.δ. *Stipes scaly.*

ε. *Surfaces scaly.* (77) *P. malaccanum.*

ε.ε. *Surfaces pubescent.*

(78) *P. brachyphlebium.*

β.β. *Pinnae crenate or toothed.* (79) *P. lividum.*

α.α. *Fully developed fronds more than $3\frac{1}{2}$ c.M. broad.*

+. *Fronds nearly sessile.* (80) *P. khasyanum.*

+.+. *Fronds distinctly stalked.* (81) *P. celebicum.*

2.2. *Pinnae naked, the rachis often so.*

3. *Rachis naked.*

4. *Pinnae entire or slightly crenate.*

5. *Sori superficial.*

6. *Sori submarginal.* (82) *decrescens.*

6.6. *Sori not submarginal.*

7. *Pinnae ± $1\frac{1}{4}$ — $1\frac{3}{4}$ c.M. long, slightly and irregularly crenate.*

(83) *P. Cesatianum.*

7.7. *Pinnae ± $3\frac{1}{2}$ — $4\frac{1}{2}$ c.M. long, entire.*

(84) *Curtisii.*

5.5. *Sori ± immersed.*

6. *Sori solitary and terminal on the pinnae.*

(85) *P. davalliaceum.*

6.6. *Sori otherwise, 3 or more to a pinna.*

7. *Pinnae sublanceolate, narrowed near the base.* (86) *P. duriusculum.*

7.7. *Pinnae not sublanceolate, not narrowed near the base.*

(*) See also *P. obliquatum*, Bl.

s. *Stipes very short; fronds to 2 c.M. broad.*

(87) *P. barathrophyllum.*

s'.s. *Stipes 5—8 c.M. long; fronds 2½—4 c.M. broad.*

(88a) *P. blechnoides.*

s'.s'.s. *Stipes to 10 c.M. long; fronds more than 5 c.M. broad.*

(88b) *P. multicaudatum.*

4.4. *Pinnae toothed.*

‡. *Sori round, confluent.* (89) *P. sibirongae.*

‡.‡. *Sori oblong, distant.* (90) *P. solidum.*

3.3. *Rachis hairy on 1 or both sides.*

p. *Sori superficial or hardly immersed.*

‡. *Fronde ± 10—15 c.M. long. Sori slightly immersed.* (91) *P. subobliquatum.*

‡.‡. *Fronde ± 20—40 c.M. long. Sori quite superficial.* (92) *P. nutans.*

p.p. *Sori distinctly immersed.*

q. *Sori submarginal.*

×. *Stipes winged.* (93) *P. Leysii.*

×.×. *Stipes not winged.* (94) *P. stenobasis.*

q.q. *Sori not submarginal.*

r. *Rims of the cavities elevated.*

(95) *P. craterisorum.*

r.r. *Rims of the cavities not elevated.*

s. *Pinnae subentire.* (96) *P. decorum.*

s.s. *Pinnae crenate or toothed.*

(97) *P. Lobbianum.*

a.a. *Most of the pinnae distinctly separate at the base.*

b. *Sori solitary on the pinnae.* (98) *P. clavifer.*

b'.b'. *Sori not solitary on the pinnae.*

c. *Fronde rather small, under 15 c.M. long, under 1½ c.M. broad.*

(99) *P. Schefferi.*

c'.c'. *Fronde larger, more than 15 c.M. long, more than 1½ c.M. broad.*

d. *Pinnae entire, crenate or undulate.*

e. *Pinnae entire, blunt. Stipes clothed with reddish hairs.*

(100a) *P. venulosum.*

e.e. *Pinnae undulate, subacute. Stipes clothed with fuscous hairs.* (100b) *P. macrum.*

d.d. *Pinnae serrate or toothed.*

*. *Pinnae adnate at the base.* (101a) *P. halconense.*

.. *Pinnae not adnate at the base.* (101b) *P. harpophyllum.*

* (55) ***P. streptophyllum***, *Bk.*, Journ. of Bot., XVII, 42.

Fronds sessile, densely tufted, linear, 7—10 c.M. long, \pm 5 m.M. broad, cut down to the rachis into very numerous, ligulate, entire, slightly twisted, adnate, contiguous pinnae \pm $\frac{1}{2}$ m.M. broad, with a single medial vein; lower pinnae gradually reduced. Texture firm; colour green; surfaces naked. Sori solitary and terminal on the pinnae.

Borneo, Singapore.

(56) ***P. triangulare***, *Scort.*, Bedd., Journ. of Bot., XXV, 324, tab. CCLXXVIII, fig. 1.

Rhizome erect, short, scaly. Fronds tufted, sessile, linear, narrowed at both ends, 15—20 c.M. long, \pm 5 m.M. broad, cut down to the rachis into distichous, obliquely triangular pinnae; barren pinnae obtuse, the fertile ones acute; upper surface convex, lower flat. Texture coriaceous; surfaces naked, the lower whitish; veins hidden. Sori solitary, terminal, sunk in a deep pit near the apex of the higher pinnae, the edge on both sides of the sorus folded inwards.

Malacca.

* (57) ***P. subtriangulare***, *v. A. v. R.*, Bull. Dép. Agr. I.N., 1908, XVIII, 20.

Fronds tufted, sessile, linear, narrowed at both ends, $7\frac{1}{2}$ —15 c.M. long, \pm 5 m.M. broad, cut down to the rachis into distichous, oblique, oblong, blunt pinnae $1-1\frac{1}{2}$ m.M. broad, dilated at the base; upper surface convex, lower flat; edge of the barren pinnae slightly recurved. Texture coriaceous; surfaces naked, but the costa slightly clothed beneath with scattered, reddish, deciduous hairs; veins hidden. Sori confined to the upper half of the frond, solitary and terminal on the segments, deeply sunk in longitudinally oblong cavities, occupying the upper half of the segments, the inner edge of each cavity forming with the much recurved, meeting lateral edges a subcylindrical hollow quite enclosing the sorus when young.

Amboyna (Boerlage).

(58) **P. musgravianum**, *Bk.*, Journ. of Bot., XXVIII, 108.

Stipes tufted, very slender, nearly naked, 1—2½ c.M. long. Fronds linear, 2½—4 c.M. long, 3—5½ m.M. broad, cut down to the rachis into ovate, adnate, erecto-patent pinnae. Texture firm; surfaces naked. Sori large, round, superficial, solitary at the base of the pinnae.

New Guinea.

(59) **P. conjunctisorum**, *Bk.*, Ann. of Bot., VIII, 129.

Fronds linear, 10—13 c.M. long, ± 7½ m.M. broad, simply pinnate, narrowed gradually at the base. Pinnae ovate-linear, entire, obtuse, contiguous, dilated at the base. Texture rigidly coriaceous; veins hidden. Sori large, subglobose, superficial, solitary at the base of the pinnae.

New Guinea.

(60) **P. cornigerum**, *Bk.*, Hk. Bk., Syn. Fil., 508.

Fronds tufted, subsessile, erect, ligulate, 7—10 c.M. long, 5—7 m.M. broad, pinnate. Pinnae 20—30-jugate, diminishing in size towards both ends of the frond, almost triangular, subacute, with the upper edge repand. Texture subcoriaceous; surfaces naked; veins 1 to each pinna, forked, falling far short of the apex. Sori large, round, 1 to each pinna, close to the rachis.

Malacca; Ceylon.

* (61) **P. inconspicuum**, *Bl.*, Rac., Flor. Btz., I, 90; *P. parvulum*, *Bl.* (not *Bory*), Flor. Jav., II, 187, tab. LXXXVII, C.

Rhizome erect or ascending, the scales narrow, lanceolate. Stipes under 1 c.M. long. Fronds commonly subsessile, 5—20 c.M. long, 6—12 (rarely to 30) m.M. broad, linear-lanceolate, narrowed towards both ends, cut down to the rachis into oblong to linear, erecto-patent or subspreading pinnae. Pinnae to 2½ m.M. broad, dilated at the base, the largest or longest ones repand or crenate, the point bluntish. Texture herbaceous to subcoriaceous; rachis and both surfaces naked or deciduously hairy; veins simple. Sori 1—6 (rarely to 12) to a pinna, medial or costal.

Sumatra, Java, New Guinea, Philippines.

(62) **P. undosum**, *Bk.*, Journ. of Bot., XXVIII, 108.

Stipes naked, wiry, 2½—4 c.M. long. Fronds pendulous, lanceolate, simply pinnate, 12—15 c.M. long, 10—12½ m.M. broad. Pinnae numerous, adnate, not approximate, subtrapezoidal, 2—4 m.M. broad, obtuse, deeply crenate

especially on the upper edge. Texture coriaceous; surfaces naked; veins hidden. Sori round, superficial, filling up nearly the whole under surface of the pinnae.

New Guinea.

* (65) **P. minutum**, *Bl.*, Rac. Flor. Btz., 94; *Bl.*, Flor. Jav., II, 188, tab. LXXXVII, D.

Rhizome erect, the scales very small. Stipes tufted, 4—8 m.M. long, clothed with fine, long, soft, whitish hairs. Fronds 3—8 c.M. long, 5—10 m.M. broad, linear-lanceolate, narrowed towards both ends. Pinnae oblong, $\frac{1}{2}$ — $1\frac{1}{2}$ m.M. broad, to \pm 6 m.M. long, blunt, erecto-patent, slightly decurrent, hardly crenate, the lower ones reduced. Texture thin-membranaceous, hyaline; surfaces and margin hairy; veins few, simple, short. Sori 1—5 to a pinna, alternate, \pm costal.

Malaya.

* (64a) **P. serrato-dentatum**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 20.

Rhizome short-creeping, the scales \pm linear-subulate, pale-brown. Stipes approximate, 1—2 c.M. long, clothed with short, spreading hairs. Fronds linear-lanceolate, \pm 10—15 c.M. long, 8—12 m.M. broad. Pinnae triangular, erecto-patent, slightly confluent, 3—4 m.M. broad at the base, the apex acute, the edge strongly recurved and conspicuously serrate-dentate with \pm 4 oblique teeth on each side; higher and lower pinnae diminishing gradually in size. Texture rigidly coriaceous; upper surface dark-coloured, lower paler, naked; rachis black, hairy; veins immersed, 1 to each tooth, forked at the base with a short, anterior branch, the apex of the principal branch black like the costa (when seen towards the light). Sori small, oblong when young, 1 to each tooth, terminal on the short anterior veinlets, subcostal.

? *Java (True habitat unknown, Forbes, No. 244).*

(64b) **P. Merritti**, *Copel.*, Philipp. Journ., I Suppl., 255.

Rhizome short-creeping, stout, clothed with lanceolate, glanduloso-ciliate, fulvo-ferrugineous scales. Stipes 1—2 c.M. long, clothed like the whole plant with short, scattered, hyaline, subglandulous hairs. Fronds 10—17 c.M. long, \pm 1 c.M. broad, hardly narrowed at the apex. Segments oblong, often obtuse, decurrent-confluent at the base, the edges obtusely toothed with 1—3 teeth on each side. Texture firm-herbaceous; veins simple, inconspicuous. Sori 2—5 to a segment, round, superficial, costal, sometimes confluent.

Mindoro.

* (65) **P. subrepandum**, *Christ*, Ann. Btz., XX, 119.

Rhizome short, erect, scaly. Stipes tufted, short, to 2 c.M. long, clothed with long, spreading, brownish hairs. Fronds 6—10 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, acuminate, narrowed at the base, pinnate. Pinnae subpatent, decurrent, slightly confluent at the base, \pm 2 m.M. apart, \pm 12—25 on each side, \pm 1 m.M. broad, crenate, rather slightly wavy, the apex subacute. Texture herbaceous, firm; rachis and surfaces short-hairy; costulae pinnate; veins simple. Sori few, 1—2 on each side of the costules, slightly immersed, terminal on the veins.

Borneo.

* (66) **P. subminutum**, *v. A. v. R.*; ? *P. minutum*, *Bk.* (not *Bl.*), *Hk. Bk.*, Syn. Fil., 528; ? *P. parvulum*, *Bedd.*, Ferns S.I., tab. CLXVI.

Rhizome erect, clothed with lanceolate, brown scales. Stipes tufted, 4—8 m.M. long, clothed with spreading, long, soft, pale-ochraceous hairs. Fronds 5—6 c.M. long, lanceolate, acuminate, $1\frac{1}{2}$ —2 c.M. broad. Pinnae \pm 2 m.M. broad, linear, \pm erecto-patent, bluntish, slightly repando-crenate, subcontiguous, \pm decurrent, the lower ones gradually reduced. Texture papyraceo-herbaceous; surfaces \pm hairy like the stipe; veins simple, short. Sori 2—5 on each side, distant, alternate, superficial, medial between the midrib and edge.

Java (Forbes).

* (67) **P. lancifolium**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 21.

Like *P. subminutum*, *v. A. v. R.* but the hairs shorter, white or pale-reddish, the fronds 6—10 c.M. long, the pinnae sometimes nearly horizontal. Sori copious, close, occupying the whole under surface of the pinnae.

Sumatra (Hagen).

(68) **P. subsecundo-dissectum**, *Zoll.*, *Hk. Bk.*, Syn. Fil., 528; (? *P. subfalcatum*, *Bl.*, var. *semiintegrum*, *Copel.*, Philipp, Journ., II, 138).

Stipes tufted, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long. Fronds 10—20 c.M. long, 2— $2\frac{1}{2}$ c.M. broad, cut down nearly to the rachis into numerous, close, subdimidiate, oblong pinnae, which are quite entire or slightly toothed towards the point on the posterior side, but cut down nearly to the midrib on the anterior side into 4—6 oblong lobes. Texture papyraceo-herbaceous; both surfaces clothed with soft, spreading, yellowish hairs; veins abbreviate. Sori close to the midrib. — Copeland's plant in very small, under 10 c.M. long.

Java, ? *Mindoro*; *New Caledonia*.

* (69) **P. subfalcatum**, *Bl.*, *Rac.*, *Flor.*, 94; *Bl.*, *Flor. Jav.*, II, 186, tab. LXXXVII, A—B; *Bedd.*, *Ferns Br.I.*, II, tab. CLXXXIX, A.

Rhizome erect, clothed with short, inconspicuous scales. Stipes densely tufted, 1—2 c.M. long, more or less clothed with soft, spreading, reddish hairs. Fronds linear-lanceolate, 8—25 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, narrowed towards both ends. Pinnae \pm erecto-patent, contiguous, straight or falcate, slightly decurrent at the base, the edge distinctly, bluntly or sharply toothed, the apex bluntish or acuminate; lower pinnae gradually reduced. Texture papyraceo-herbaceous; surfaces slightly and the rachis more densely hairy like the stipe; veins simple. Sori superficial, \pm 3—6 on each side, 1 to each tooth, rather distant.

Malaya; North. India, South. China.

* (70a) **P. repandulum**, *Mett.*, *Hk. Bk.*, *Syn. Fil.*, 528; *P. obliquatum*, *Hk.*, *Spec. Fil.*, IV, 190, p. p.; *P. subfalcatum*, *Bl.*, *var. glabrum.*, *Bedd.*, *Ferns Br I.*, I, tab. LXXVI; II, tab. CLXXXIX, B; *Ctenopteris repandula*, *Kze.*, (oldest name).

Stipes tufted, very short, naked. Fronds 10—15 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, cut down to the rachis throughout into erecto-patent, blunt, repandocrenate pinnae 1—2 m.m. broad. Texture firm-herbaceous; rachis and both surfaces \pm naked; veins simple. Sori distant, slightly immersed, terminal on the veins, filling nearly the whole space between the midrib and edge.

Amboyna, Borneo; Ceylon.

(70b) **P. erythrotrichum**, *Copel.*, *Philipp. Journ.*, I, *Suppl.*, 160, tab. XX.

Rhizome short-creeping, clothed with short, subciliate, deciduous, fuscous scales. Stipes approximate, 2—5 c.M. long, densely clothed like the rachis with red-purple hairs. Fronds pendulous, 30—50 c.M. long, $2\frac{1}{2}$ —4 c.M. broad. Pinnae very numerous, spreading, \pm 2 m.M. broad, hardly dilated at the base, entire, obtuse, the lower reduced gradually to mere teeth. Texture papyraceo-herbaceous; surfaces glabrescent; costae sinuate (flexuose?) towards the apex; veins simple, erecto-patent. Sori oblong, nearly parallel to the costae, to 10 on each side, subimmersed, the edges of the cavities slightly raised.

Mindanao, Mindoro.

(71) **P. brevifrons**, *Scort.*, MS.; *P. repandulum*, *Mett.*, var. *malayanum*, *Bedd.*, in *Journ. of Bot.*, XXXI, 226.

Like *P. repandulum*, *Mett.*, but the sori deeply sunk in cavities with raised margins.

Malacca.

* (72) **P. papillosum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 90; *Bl.*, *Flor. Jav.*, II, 191, tab. LXXXIX, A—B; *Hk.*, *Icon. Pl.*, X, tab. CMXLVI; *Christ*, *Farnkr. d. Erde*, fig. 254.

Rhizome horizontally creeping, the scales lanceolate, acute. Stipes scattered, \pm 5—20 c.M. long, erect, slender, rigid, naked. Fronds linear-lanceolate, 20—60 c.M. long, 4—8 c.M. broad, cut down to the rachis into numerous spreading pinnae. Pinnae linear, blunt, entire or crenate to toothed, 4—8 m.M. broad, dilated at the base, the lowest not reduced. Texture papyraceo-herbaceous, thin; rachis and both surfaces naked; veins conspicuous, forked. Sori in 1 row on each side near the edge, deeply immersed, much prominent on the upper surface so as to form conspicuously elevated, conical or saccate papillae.

Malaya.

* (73) **P. obliquatum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 93; *Bl.*, *Flor. Jav.*, II, 181, tab. LXXXV, B; *Bedd.*, *Ferns S.I.*, tab. CLXVII.

Rhizome short, the scales lanceolate, ciliate. Stipes tufted, $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, rigid, naked or pilose. Fronds 20—70 c.M. long, \pm $2\frac{1}{2}$ —6 c.M. broad, linear-lanceolate, cut down to the rachis into entire or slightly wavy, close, subspreading, bluntish pinnae $2\frac{1}{2}$ —4 m.M. broad, dilated at the base, the lower ones growing shorter, reduced, deltoid, very blunt. Texture herbaceous to subcoriaceous; rachis naked or hispid; surfaces naked; veins simple. Sori elliptical, oblique, 4—6 on each side of the costae, immersed, somewhat prominent on the upper surface, the edges of the cavities distinctly raised and covering the sori when young.

Malaya; Ceylon, South India.

(74) **P. Schenkii**, *Harr.*, *Journ. of Linn. Soc.*, Bot. XVI, 31; (? *P. obliquatum*, *Bl.*).

Rhizome thick, horizontal, densely clothed with slender, brown scales. Stipes clustered, 5—8 c.M. long, brown, pubescent, firm, erect. Fronds 15—30 c.M. long, $5\frac{1}{2}$ —5 c.M. broad, oblong, pinnate, narrowing gradually to both

ends. Pinnae numerous, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, $3\frac{1}{2}$ —5 m.M. broad, linear, obtuse, entire, broadened and nearly confluent at the base, the lowest ones small, triangular. Texture membranaceous; rachis pubescent; surfaces naked; veins simple, free, falling short of the margin. Sori deeply immersed, oblong, 4—8 in a row on each side of the midribs, oblique, nearly covered by the recurved margins of the depressions.

Philippines.

* (75) **P. decipiens**, *Mett.*, *Rac.*, *Flor. Btz.*, I, 93; *P. cryptosorum*, *C. Chr.*, *Ind. Fil.*

Rhizome short-creeping, densely clothed with lanceolate-subulate, ciliate, black-brown scales. Stipes tufted, short or 1—2 c.M. long, densely furnished with long, spreading, reddish hairs. Fronds linear-lanceolate, 10—40 c.M. long, $1\frac{1}{4}$ — $2\frac{1}{2}$ c.M. broad, narrowed towards both ends, cut down nearly to the costa into linear or lanceolate, nearly close, entire, blunt lobes. Texture coriaceous, rather fleshy; surfaces thinly beset with red-brown hairs, the midrib more densely; veins zigzag throughout or only at the base; veinlets translucent, 5—14-jugate, simple or the lower forked. Sori submarginal, deeply immersed in elliptical cavities.

Malaya.

* (76a) **P. mollicomum**, *Nees & Bl.*, *Rac.*, *Flor. Btz.*, I, 91; *Bl.*, *Flor. Jav.*, II, 184, tab. LXXXVI, B; — var. *fuscatum*; *P. furcatum* *Bl.*, l.c., tab., LXXXVI, C; *P. fuscatum*, *Bl.*, l.c., 185; *C. Chr.*, *Ind. Fil.*; *Bedd.*, *Ferns Br. I.*, II, tab. CCCXXIV.

Rhizome erect or suberect, densely clothed with shining, adpressed, lanceolate scales. Stipes tufted, 2—8 c.M. long, densely purveyed with soft, spreading, reddish hairs. Fronds 7—50 c.M. long, $1\frac{1}{4}$ —3 c.M. broad, cut down nearly or quite to the rachis throughout into close, entire, blunt pinnae 1—4 m.M. broad, the base often dilated, the lower ones shorter and broader, close. Texture more or less coriaceous; rachis and both surfaces hairy; veins simple, obscure. Sori superficial, forming a close row on each side of the midribs, at length often filling up nearly the whole under surface.

Var. fuscatum: Lower pinnae distant, narrowed towards the base, the very base mostly much dilated again.

Malaya.

(76b) **P. pseudoarticulatum**, *Copel.*, *Philipp. Journ.*, I, Suppl., 160, tab. XVIII.

Rhizome short-creeping, clothed with narrow, black, ciliate scales. Stipes 2—4 c.M. long, puberulous. Fronds 10—20 c.M. long, $1\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, acuminate or caudate, the rachis black, short-pubescent. Pinnae linear, spreading, often separated by more than their own breadth, the longest ± 2 m.M. broad, blunt, entire, dilated at the base, the lower reduced, more distant, hardly decurrent. Texture coriaceous; under surface clothed with more or less deciduous, reddish hairs; veins hidden, simple. Sori nearly terminal on the veins, nearly parallel to the costae, hardly immersed, ± 22 on the longest pinnae.

Luzon.

(77) **P. malaccanum**, *Bk.*, Ann. of Bot., VIII, 129.

Stipes densely tufted, slender, scaly, $2\frac{1}{2}$ —5 c.M. long, the scales brown, subulate, spreading, deciduous. Fronds lanceolate, 12—15 c.M. long, $1\frac{1}{4}$ — $1\frac{1}{2}$ c.M. broad. Pinnae multijugate, linear, entire, obtuse, the base dilated, the lower ones smaller. Texture moderately firm; surfaces covered with subulate, brown scales; veins simple, erecto-patent, 6—8-jugate in the central pinnae. Sori round, superficial, in 1 row on each side of the costae.

Malacca.

(78) **P. brachyphlebium**, *Bk.*, Ann. of Bot., VIII, 129.

Stipes short, slender, tufted, clothed with short, spreading, subulate scales. Fronds lanceolate, ± 15 c.M. long, $\pm 1\frac{1}{2}$ — $1\frac{3}{4}$ c.M. broad, narrowed at the base. Pinnae contiguous, multijugate, linear, obtuse, entire, $\pm 2\frac{1}{2}$ m.M. broad, dilated at the base. Texture moderately firm; surfaces pubescent; veins short, erecto-patent, simple. Sori semiglobose, superficial, in 1 row on each side of the costae.

Sumatra.

(79) **P. lividum**, *Mett.*, Ann. Mus. Bot. L. B., II, 221.

Rhizome short, subascending, the scales membranaceous, ferruginous, oblong-lanceolate. Stipes short, clothed with short, rigid hairs. Fronds 5—15 c.M. long, $\pm 1\frac{1}{4}$ — $1\frac{1}{2}$ c.M. broad, lanceolate, narrowed gradually towards both ends, the edge pinnatifid nearly to the costa. Segments numerous, spreading, decurrent, confluent, linear-oblong, the apex narrowed, obtuse, crenate; lowest segments decurrent upon the stipe. Texture thinly coriaceous; colour bluish-olivaceous; under surface hairy like the stipe, the hairs glanduliform;

veins subimmersed. Sori immersed, medial between the costulae and the margin.

Java.

(80) **P. khasyanum**, *Hk.*, Hk. Bk., Syn. Fil., 325; Hk., Icon. Pl., X, tab. CMXLIX; Bedd., Ferns Br.I., II, tab. CLXXIII.

Stipes tufted, very short, fringed with soft hairs. Fronds 30 c.M. or more long, \pm 4 c.M. broad, flaccid, cut down nearly to the costa into linear-oblong, entire or slightly undulate, blunt lobes 5—8 m.M. broad. Texture thick, coriaceous; rachis and both surfaces hairy; veins simple. Sori sunk, in rows of 4—6 on each side of the costulae, elliptico-rotundate.

Malacca; Khasya, Assam.

* (81) **P. celebicum**, *Bl.*, Flor. Jav., II, 179, tab. LXXXIV, B.

Rhizome short, woody, reddish-brown, scaly, the scales densely imbricate, membranaceous, scariose, hairy, pale-reddish-brown. Stipes 7—13 c.M. long, or shorter, approximate (tufted, Hk. Bk., Syn. Fil., 330), reddish-brown, clothed with brownish, spreading hairs like the rachis. Fronds commonly nutant, 20—40 c.M. long, 5—8 c.M. broad, lanceolate, narrowed towards both ends. Pinnae close, nearly horizontal, $3\frac{1}{2}$ —5 m.M. broad, numerous, entire, bluntish, subconfluent at the base, the lower ones reduced. Texture coriaceous; costae indistinct, slightly hairy beneath like the margin; veins simple. Sori immersed, distant, 8—12 on each side, nearer the edge than the midrib, \pm oblong, the cavities ciliate at the margin, the ciliae covering the sori when young.

Malaya.

(82) **P. decrescens**, *Christ*, Ann. Btz., XIX, 55; Philipp. Journ., I, Suppl., 159, tab. XVII.

Rhizome oblique, short-creeping, clothed with brown, ovate, acute scales. Stipes short, brown like the rachis. Fronds approximate, nearly tufted, 20 c.M. long (incl. the stipe). Pinnae 15—20 on each side, often unequal, 2 c.M. long, 2 m.M. broad, spreading, blunt, not close, connected at the base by a narrow wing; lower pinnae suddenly reduced so as to form an undulate wing to the rachis. Texture papyraceo-coriaceous; surfaces naked, under side \pm glaucous; veins hidden. Sori \pm 9 on a side, round, submarginal.

Celebes, Mindoro, Luzon.

* (83) **P. Cesatianum**, *Bk.*, Journ. of Bot., XVII, 42; *P. papillosum*, *Ces.* (not *Bl.*), Fel. di Bor., 25.

Rhizome wide-creeping, woody, the scales dense, brown, small, narrowed gradually from a rounded, peltate base towards the acuminate point. Stipes scattered, brownish, naked, glossy, wiry, slender, ± 10 c.M. long. Fronds \pm lanceolate, 15—20 c.M. long, $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, narrowed towards both ends, or the base nearly as broad as the central part of the frond, the edge cut down nearly to the costa. Pinnae spreading, 3—5 m.M. broad, the apex blunt, the edge slightly, not regularly crenate, the base dilated, the sinuses as broad as the lobes. Texture coriaceous; surfaces naked; costa prominent below; costulae distinct, \pm wavy; veins oblique, forked, free or united near the base by the arched anterior veinlets, which are nearly parallel to the costulae and bear a very short soriferous branch directed towards the edge. Sori round, small, superficial, ± 6 —8 on each side, nearer the costulae than the edge, 1 between each pair of veins. — The description is taken from a specimen gathered by Hallier (No. 3312) and determined by Christ (Ann. Btz., XX, 120).

Borneo.

(84) **P. Curtisii**, *Bk.*, Journ. of Bot., XIX, 367.

Rhizome wide-creeping, the scales dense, large, lanceolate, brown, membranaceous. Stipes brown, naked, 5—8 c.M. long. Fronds oblanceolate, 20—25 c.M. long, 7—9 c.M. broad at the middle, cut down nearly or quite to the rachis. Pinnae close, entire, linear, $2\frac{1}{2}$ —4 c.M. long, ± 4 —5 m.M. broad above the dilated deltoid base; lower pinnae reduced so as to form a repand wing to the rachis. Texture subcoriaceous; surfaces naked; veins hidden. Sori round, medial, superficial, 6—9 on each side of the costa in the fully developed central pinnae.

Sumatra.

(85) **P. davalliaceum**, *Muell. & Bk.*, Journ. of Bot., XXVIII, 108.

Rhizome short-creeping, the scales small, brown, ovate-lanceolate. Stipes wiry, naked, $2\frac{1}{2}$ —5 c.M. long. Fronds lanceolate, simply pinnate, 15—20 c.M. long, to $2\frac{1}{2}$ c.M. broad. Pinnae adnate, lanceolate or oblique-cuneate, $1\frac{1}{2}$ — $2\frac{1}{2}$ m.M. broad, entire or obscurely crenate on the upper edge. Texture coriaceous; surfaces naked; veins hidden. Sori solitary, immersed, sub-terminal on the pinnae, the edges of the latter folded round them.

New Guinea.

(86) **P. duriusculum**, *Christ*, Ann. Btz., XV, 151, tab. XVI, fig. 25, a—c.

Rhizome short. Fronds densely tufted, \pm 10 c.M. long, lanceolate, acuminate. Pinnae erecto-patent, \pm 1 c.M. long, 5—6 m.M. apart, lanceolate, slightly crenate, the base narrowed and more or less decurrent, the apex blunt; upper and lower pinnae reduced. Texture very firm; surfaces naked, dark-coloured; veins hidden. Sori on the broadest part of the pinnae, 5—6 to a pinna, immersed, confluent.

Celebes.

* (87) **P. barathrophyllum**, *Bk.*, Journ. of Bot., XXIX, 107.

Rhizome erect. Stipes tufted, very short. Fronds lanceolate, to \pm 30 c.M. long, $1\frac{1}{4}$ —2 c.M. broad, narrowed gradually towards both ends, cut down to a narrow wing into entire, linear-oblong segments \pm 2 m.M. broad above the dilated base. Texture thick, flexible; surfaces naked or the margin ciliate; veins hidden. Sori medial, semiglobose, 6—8 in a row on each side of the costae, sunk in distinct, round cavities, which do not look as papillae on the upper surface of the frond.

Borneo.

(88a) **P. blechnoides**, *Hk.*, Hk. Bk., Syn. Fil., 531; *P. contiguum*, *Brack.*, Expl. Exp., XVI, tab. II, fig. 1; *Grammitis blechnoides*, *Grev.*, (oldest name).

Rhizome stout, the scales dense, linear. Stipes 5—8 c.M. long, rigid, erect, naked. Fronds 20—30 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, cut down very nearly to the rachis throughout. Fertile pinnae confined to the upper half of the frond, longer and narrower than the barren ones, which are bluntish, \pm 3—3 $\frac{1}{2}$ m.M. broad, growing gradually shorter and broader downwards. Texture coriaceous; rachis and both surfaces naked. Sori oblong, immersed, in rows near the midribs.

Java, Borneo, New Guinea; Queensland, Polynesia.

(88b) **P. multicaudatum**, *Copel.*, Philipp. Journ., I, Suppl., 160, tab. XIX; *P. obliquatum*, *Copel.* (not *Bl.*), Polypod. Philipp., 122.

Rhizome short-creeping, clothed with rather long, ciliate, fuscous scales. Stipes approximate, to 10 c.M. long, sparingly clothed with reddish-black hairs. Fronds 50—60 c.M. long, often more than 6 c.M. broad, the rachis naked. Pinnae erecto-patent, narrowed gradually from the rather suddenly dilated base towards the caudiform apex, the margin entire; lower pinnae reduced. Texture subcoriaceous; surfaces naked; veins simple, oblique. Sori often restricted to

the central part of the pinnae, 4—10 on each side of the costae, reaching nearly from the midrib to the margin, immersed, the margins of the cavities conspicuously crateriform.

Mindanao.

(89) **P. sibarongae**, *Christ*, Ann. Btz. XIX, 36.

Rhizome short, oblique, clothed with thin, subulate, brown scales. Stipes short, tufted. Fronds linear-lanceolate, \pm 15 c.M. long, \pm $1\frac{1}{2}$ c.M. broad, acuminate. Pinnae numerous, 30—40 on each side, often opposite, 8 m.M. long, 5 m.M. broad, oblong, lower deltoid, higher toothed, the teeth deep, large, \pm 3 on a side, deltoid, acute. Texture coriaceous; rachis ebeneous, rigid; colour dark; surfaces naked, under side glaucous; veins hidden. Sori medial, large, round, confluent, 1 to each tooth.

Celebes.

(90) **P. solidum**, *Mett.*, Hk. Bk., Syn. Fil., 324; *Mett.*, Farng., I, Polypod., tab. I, fig. 1—3; *Ctenopteris solida*, *Kze.* (oldest name).

Rhizome stout, creeping, clothed with lanceolate, ferruginous scales. Stipes $2\frac{1}{2}$ c.M. long, firm, erect, naked. Fronds 10—20 c.M. long, 2 c.M. broad, cut down nearly to the midrib into close, distinctly toothed, bluntish lobes \pm 5 m.M. broad. Texture coriaceous; both surfaces naked; veins once or more times forked. Sori oblong, subimmersed, 1—4 on each side, medial between the costulae and the margin.

Java, Philippines.

(91) **P. subobliquatum**, *Christ*, Bull. Herb. Boiss., VI, 197; *P. obtusissimum*, *C. Chr.*, Ind. Fil.

Rhizome short-creeping, slightly scaly. Stipes approximate, 1—2 c.M. long, brown, clothed like the rachis with short, cinnamon-coloured hairs. Fronds 10—15 c.M. long, \pm 2 c.M. broad, lanceolate-oblong. Pinnae obtuse, alternate, \pm 2 m.M. broad, dilated at the base, remote, reduced towards the ends of the frond. Texture coriaceous; colour rufescent; surfaces naked; veins oblique, simple, inconspicuous. Sori 4—6 on each side, medial, oval, hardly immersed, the margins of the cavities not crateriform.

Luzon.

* (92) **P. nutans**, *Bl.*, Rac., Flor. Btz., I, 92; *Bl.*, Flor. Jav., II 182, tab. LXXXVI, A.

Rhizome creeping, clothed with dark-brown, narrow, lanceolate, ciliate scales. Stipes 2—8 c.M. long, provided with very short reddish hairs. Fronds lanceolate, 20—40 c.M. long, $1\frac{1}{2}$ —3 c.M. broad, narrowed towards both ends, the edge cut down to the rachis into entire, blunt pinnae 1—3 m.M. broad. Texture coriaceous; surfaces \pm naked; rachis hairy; veins hidden, simple or forked. Sori round (elliptical when young), in 1 row on each side of the midribs, at length filling up the whole under surface.

Malacca, Sumatra, Java; New Caledonia.

(93) **P. Leysii**, *Bk.*, Journ. of Bot., XVII, 66.

Rhizome short-creeping, the scales small, dense, dark-brown, lanceolate. Stipes winged nearly or quite down to the base, $2\frac{1}{2}$ —5 c.M. long to where the pinnae begin. Fronds lanceolate, 15—25 c.M. long, $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad above the middle, narrowed gradually from the middle to the base, cut down to the rachis into very numerous, crowded, adnate, linear, subentire or slightly repand pinnae 2— $2\frac{1}{2}$ m.M. broad. Texture firm; surfaces naked; rachis minutely pubescent beneath; veins simple, erecto-patent, 10—12-jugate in the central pinnae, not reaching the margin. Sori terminal on the veins, globose, submarginal, distinctly immersed.

Sulu Islands.

(94) **P. stenobasis**, *Bk.*, Ann. of Bot., VIII, 130.

Rhizome hardly creeping, the scales lanceolate, brown, membranaceous. Stipes nearly naked. Fronds lanceolate, 15—25 c.M. long, 3—4 c.M. broad, narrowed at the base. Pinnae linear, multijugate, $\pm 2\frac{1}{2}$ m.M. broad, obscurely crenate, dilated and adnate at the base. Texture coriaceous; surfaces naked; rachis greenish, pubescent; veins simple, obscure. Sori in 1 row on each side of the costae, submarginal, deeply immersed.

Sumatra.

(95) **P. craterisorum**, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 31; (? *P. celebicum*, *Bl.*).

Rhizome densely clothed with brown, lanceolate scales. Stipes clustered, 1—4 c.M. long, hirsute with brown hairs, firm, erect. Fronds 20—30 c.M. long, $3\frac{1}{2}$ —5 c.M. broad, oblong or oblong-oblancoelate, tapering gradually to both ends, pinnate. Pinnae numerous, linear, ± 5 m.M. broad, obtuse, entire, dilated and \pm confluent at the base, passing below into the narrow, lobed wing of the stipe. Texture membranaceous; rachis pubescent; surfaces naked;

veins simple, not quite reaching the margin. Sori confined to the upper pinnae, immersed, oval, 5—10 in a row on each side of the midrib, \pm medial between this and the margin, the edge of each cavity forms a projecting ring, but does not cover the sorus.

Philippines.

(96) **P. decorum**, *Brack.*, Hk. Bk., Syn. Fil., 351; Bedd., Ferns S.I., tab. CCXXXVIII, A; *P. nutans*, *J. Sm.* (not *Bl.*), Journ. of Bot., III, 394.

Rhizome short, creeping, densely ferrugineo-squamose. Stipes approximate, $2\frac{1}{2}$ c.M. or less long, naked. Fronds 15—30 c.M. long, $1\frac{1}{4}$ — $2\frac{1}{2}$ c.M. broad, linear-lanceolate, narrowed towards both ends, cut down nearly or quite to the rachis into numerous pinnae. Central pinnae longest, rather spreading, linear-oblong, $\pm 2\frac{1}{2}$ —3 m.M. broad, blunt, the lower ones gradually becoming blunter, shorter and broader. Texture coriaceous; surfaces naked, or the rachis hispid; costulae and veins indistinct. Sori oblong, 2—8 on each side, medial between the costulae and the edge, moderately filling up the space between both, apical on the short veins, sunk in hairy cavities.

Malaya; Ceylon, Polynesia.

(97) **P. Lobbianum**, *Hk.*, Spec. Fil., IV, 226 (not V, 100), tab. CCLXXVIII, B; *P. pediculatum*, *Bk.*, C. Chr., Ind. Fil.; *P. inarticulatum*, *Copel.*, Philipp. Journ., I, Suppl., 160.

Rhizome short-creeping, stout, densely scaly. Fronds sessile, 10—15 c.M. long, 2—3 c.M. broad, lanceolate-acuminate. Pinnae numerous, close, 1 — $1\frac{1}{2}$ m.M. broad, the edge with distinct, rounded crenations or teeth; lower pinnae gradually reduced. Texture coriaceous; surfaces naked; rachis wiry, ebeneous, sparingly and deciduously hairy; veins simple, indistinct, 1 to each crenation, short. Sori immersed, \pm medial, 1 to each vein.

Borneo, Mindanao.

(98) **P. clavifer**, *Hk.*, Hk. Bk., Syn. Fil., 354; *P. claviferum*, *Hk.*, Becc., Mal., III, 47; *Grammitis clavifer*, *Hk.*, 2nd Cent. of Ferns, tab. V.

Stipes densely tufted, very short. Fronds 10—30 c.M. long, $\pm 12\frac{1}{2}$ m.M. broad, pinnate throughout. Pinnae dimorphous, distant; barren pinnae linear-subulate or ligulate, fertile ones spatulate, with the lower half narrowly linear, the upper half oblong-elliptical, with recurved sides. Texture coriaceous; rachis ciliate with deciduous hairs; pinnae tipped with 1—3 erect

bristles; veins 1 to each pinna. Sori large, oblong, solitary on the pinnae, placed in the centre of the dilated fertile part.

Borneo, New Guinea.

* (99) **P. Schefferi**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 21.

Rhizome short, clothed with dark-brown, entire, lanceolate scales. Stipes \pm tufted, $1\frac{1}{2}$ —5 c.M. long, clothed with very short, spreading, erecto-patent or adpressed gland-like hairs. Fronds linear, narrowed towards both ends, 5—10 c.M. long, 5—10 m.M. broad, pinnate. Pinnae erecto-patent, irregularly rhomboidal-oblong in the narrowest, ovate-oblong in the broadest forms, the higher ones rather confluent, subentire or crenate, the central decurrent, distant, crenate, crenato-dentate or crenato-pinnatifid with few crenations, the lower much reduced, narrow, entire, distant. Texture firm-herbaceous, fragile when dry, pellucid; rachis and surfaces papillose, the papillae clavate, pellucid, yellowish-brown, subadpressed; costae pinnate with \pm 1—3 simple veins on each side. Sori round, 1 to each vein, close to the costae, at length often contiguous. — Differs from *P. glandulosum*, *Hk.*, Spec. Fil., IV, 194, tab. CCLXXVI, A, in having all the pinnae distant, not confluent, and the opposite sori contiguous, costal, though terminal on the short veins.

Java (Scheffer).

* (100a) **P. venulosum**, *Bl.*, Rac., Flor. Btz., I, 92; *Bl.*, Flor. Jav., II, 180, tab. LXXXV, A.

Rhizome short-creeping, clothed with dark, lanceolate, ciliate scales. Stipes subtufted or to 1 c.M. apart, $2\frac{1}{2}$ —12 c.M. long, densely covered with deciduous or persistent, spreading, reddish hairs. Fronds linear-lanceolate, 30—70 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, narrowed gradually towards both ends, cut down to the rachis throughout into entire, linear, blunt, horizontal, sessile pinnae 2—3 m.M. broad, the base often slightly narrowed. Texture herbaceous to subcoriaceous; rachis villose; surfaces naked or villose; veins parallel, simple. Sori copious, superficial or immersed, round, elliptical when young.

Malaya.

(100b) **P. macrum**, *Copel.*, Polypod. Philipp., 121.

Rhizome short, erect, densely clothed with brown scales. Stipes \pm tufted, 1— $2\frac{1}{2}$ c.M. long, blackish-brown, covered with hairs of the same colour. Fronds linear-lanceolate, acute, pinnate, \pm 25 c.M. long, \pm $2\frac{1}{2}$ c.M. broad.

FRICES.

Pinnae narrow, subacute, undulate, subfalcate, decurrent, the higher ones confluent, the lower distant, reduced. Texture subcoriaceous; rachis \pm hairy; surfaces naked or nearly so; veins simple. Sori \pm elliptical, terminal on the veins, medial between the costae and the margin, (oblique?), superficial.

Philippines.

(101a) **P. halconense**, *Copel.*, Philipp. Journ., II, 138, tab. II, B.

Rhizome creeping, clothed with grey-fuliginous, lanceolate scales. Stipes approximate, \pm 5 c.M. long, clothed like the rachis with straight, castaneous hairs. Fronds \pm 20 c.M. long, \pm 4 $\frac{1}{2}$ c.M. broad, pinnate, abruptly narrowed at both ends. Pinnae suberecto-patent, straight or curved, \pm 2 $\frac{1}{2}$ c.M. long, \pm 3 $\frac{1}{2}$ m.M. broad, obtuse, decurrent, toothed, only the superior ones confluent, the lower rather remote, abruptly reduced to mere auricles, the teeth reaching $\frac{1}{2}$ — $\frac{2}{3}$ -way down to the costa, distant, spreading. Texture coriaceo-papyraceous; costae and apices of the teeth setiferous; veins simple, hidden. Sori superficial, occupying the bases of the veins, often opposite, and the pairs not rarely confluent.

Mindoro.

(101b) **P. harpophyllum**, *Zenk.*, Hk. Bk., Syn. Fil., 511; Christ, Ann. Bot., XV, 150, tab. XVI, fig. 22, a—b.

Rhizome wide-creeping, slender. Stipes slender, fragile, black-brown, rough with short, dark scales, like the rachis, \pm 10 c.M. long, \pm 5 c.M. apart. Fronds \pm 45 c.M. long, oblong-lanceolate, pinnate. Pinnae ligulate-acuminate, falcate, serrate, 3 $\frac{1}{2}$ —5 c.M. long, \pm 1—1 $\frac{1}{2}$ c.M. broad, subpetiolate; upper base parallel to the rachis, auricled, lower cuneate-truncate. Texture subcoriaceous; surfaces glabrous; veins 2—3-furcate. Sori medial between the edge and midribs, terminal on the lowest anterior veinlets.

Celebes; South. India.

§ 1. EUPOLYPODIUM. c. **Compositae.**

a. Texture thin, hyaline.

(102) *P. gedeanum.*

a.a. Texture thick, not hyaline.

b. Stipes naked.

(103) *P. tamariscinum.*

b.b. Stipes hairy.

c. Rachis naked.

(104) *P. bipinnatifidum.*

c.c. *Rachis hairy.***d.** *Pinnae irregularly divided.* (105) *P. millefolium.***d.d.** *Pinnae regularly pinnatisect.***e.** *Sori occupying the base of the secondary segments.*
(106) *P. tenuisectum.***e.e.** *Sori not occupying the base of the secondary segments.***f.** *Fertile pinnulae linear.* (107a) *P. taxodioides.***f.f.** *Fertile pinnulae spathulate.* (107b) *P. Yoderi.** (102) **P. gedeanum**, *Rac.*, Flor. Btz., I, 96.

Rhizome erect or suberect, slender. Stipes tufted, 1—2 c.M. long, densely clothed with spreading, yellow-brown hairs. Fronds linear-lanceolate, narrowed towards both ends, to 12—18 c.M. long, 2—3 c.M. broad. Pinnae 5 m.M. apart, decurrent, unequal-sided, subdimidiate, deeply pinnatifid, the upper side with 5—6, the lower with 1—5 segments, the lower base cut away and without segments. Segments linear, rounded, entire, separated by narrow spaces between them, 3—4 m.M. long, 1—1½ m.M. broad. Texture very thin, hyaline; rachis, veins and edges furnished with very soft, white, spreading hairs; veins free, 1 to each segment. Sori solitary at the base of the ultimate segments.

Java.

(103) **P. tamariscinum**, *Klf.*, Hk. Bk., Syn. Fil., 538; Diels, in Engl. & Prantl., Nat. Pfl.Fam., I⁴, fig. 162, H—K; *Adenophorus Tamarisci*, *Hk. & Grev.*, Ic. Fil., tab. CLXXV; *A. bipinnatus*, *Gaud.*, Hk. & Grev., l. c., tab. CLXXIV.

Rhizome stout, creeping, scaly. Stipes 5—8 c.M. long, naked, wiry, flexuose. Fronds 10—20 c.M. long, 5½—8 c.M. broad, ovate-lanceolate. Pinnae close, erecto-patent, ½—1 c.M. broad, cut down to the rachis into simple or forked, linear or oblong lobes. Texture subcoriaceous; both surfaces naked; veins 1 to each lobe. Sori copious, 1 to each ultimate segment, near the apex.

Malaya; Polynesia.(104) **P. bipinnatifidum**, *Bk.*, Journ. of Bot., XXVIII, 109.

Stipes very slender, 5½—5 c.M. long, clothed with fine, spreading, brown hairs. Fronds lanceolate-oblong, 12—15 c.M. long, ± 2½ c.M. broad, pendulous, 2-pinnatifid. Pinnae distant, linear, cut down to a narrowly winged

rachis into small, oblong or ovate lobes. Texture coriaceous; surfaces naked; veins 1 to each lobe. Sori round, superficial, 1 to each lobe.

New Guinea.

* (105) **P. millefolium**, *Bl.*, Rac., Flor. Btz., I, 95; *Bl.*, Flor. Jav., II, 190, tab. LXXXVIII, B.

Rhizome short, slender, erect, hidden by the stipes and roots. Stipes tufted, $2\frac{1}{2}$ —8 c.M. long, densely clothed with spreading, ferruginous or pale-coloured hairs, like the rachis. Fronds lanceolate, \pm 15—20 c.M. long, \pm $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, narrowed towards both ends. Pinnae erecto-patent, \pm stalked, 7—10 m.M. broad, lanceolate, cut down nearly or quite to the rachis into oblong or linear-oblong to spatulate, erecto-patent segments \pm 5—10 m.M. long, 1—3 m.M. broad. Segments blunt, the edge entire, toothed or pinnatifid, the base narrowed, slightly decurrent. Ultimate lobes 1—3 on each side. Texture subcoriaceous; under surface slightly hairy; veins simple. Sori roundish, terminal on very short veinlets, \pm 1—3 to each pinnula.

Sumatra, Java, Philippines.

* (106) **P. tenuisectum**, *Bl.*, Rac., Flor. Btz., I, 95; *Bl.*, Flor. Jav., II, 189, tab. LXXXVIII, A; Christ, Farnkr. d. Erde, fig. 243; *Hk.*, 2nd Cent. of Ferns, tab. XXI; — var. *paucisetosum*, *Rst.*, in *Fed.*, Rep., V, 42.

Rhizome short-creeping, densely clothed with lanceolate scales. Stipes approximate, 5—5 c.M. long, clothed with soft, spreading, ferruginous hairs. Fronds linear-lanceolate, narrowed towards both ends, 15—25 c.M. long, $5\frac{1}{2}$ —6 c.M. broad. Pinnae close, to 8 m.M. broad, cut down quite to the rachis or a narrow wing into linear, regular, entire, erecto-patent segments 2—3 m.M. apart, 3—4 m.M. long, $\frac{1}{2}$ —1 m.M. broad. Texture subcoriaceous; rachis erect, straight, pilose; surfaces naked or the veins sparingly hairy beneath; veins simple, hidden, 1 to each lobe. Sori broader than the lobes, 1 to each lobe, at the base.

Var. paucisetosum: Ultimate segments narrower, provided at the apex with 1 or rarely 2 setae.

Malacca, Java to Samoa and the Philippines; — var. *paucisetosum:* *New Guinea.*

(107a) **P. taxodioides**, *Bk.*, Journ. of Bot., XVII, 42.

Rhizome filiform, wide-creeping. Stipes wiry, brown, $2\frac{1}{2}$ —4 c.M. long, clothed like the rachis with fine, spreading, brown hairs. Fronds lanceolate,

regularly 2-pinnate, 15—20 c.M. long, $4\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad. Pinnae numerous, close, lanceolate, \pm 6—7 m.M. broad, regularly pectinato-pinnate with narrow, linear, parallel, erecto-patent, 1-nerved pinnulae not more than $\frac{1}{2}$ m.M. broad. Texture rather rigid; surfaces naked. Sori confined to a few of the uppermost pinnae, solitary in the centre of the pinnulae.

Borneo.

(107b) **P. Yoderi**, *Copel.*, Philipp. Journ., I, Suppl., 161.

Rhizome filiform, sparingly clothed with brown, ovate-lanceolate scales. Stipes woody, clothed with reddish-black hairs, or glabrescent towards the base. Fronds 15—25 c.M. long, lanceolate, the rachis pilose. Pinnae erecto-patent, \pm 5 m.M. broad, contiguous, the rachis winged. Barren pinnulae narrow, hardly $\frac{1}{4}$ m.M. broad at the base, somewhat broader towards the apex, often bearing a single terminal hair; fertile pinnulae spatulate. Sori solitary on the pinnulae, occupying the dilated portion.

Mindoro, Panay.

§ 2. PROSAPTIA.

a. *Fronds under $1\frac{1}{2}$ c.M. broad. Sori 1 to each tooth or segment. (Acrosorus).*

b. *Fronds toothed, lobed, or pinnatifid nearly or quite to the costa, the teeth or segments deltoid, acute or bluntish, not narrowed at the base.*

c. *Fronds sharply toothed in the fertile region.*

(108a) *P.* Schlechteri.

c.c. *Fronds bluntly lobed in the fertile region.*

(108b) *P.* exaltatum.

(108c) *P.* tortile.

b.b. *Fronds pinnatifid nearly or quite to the costa, the fertile segments sub-clavate, narrowed at the base.*

(109) *P.* Sarasinorum.

a.a. *Fronds more than $1\frac{1}{2}$ c.M. broad when fully developed. Sori rarely 1, commonly many to each segment. (Euprosaptia).*

1. *Fronds lobed to \pm $\frac{2}{3}$ -way down to the costa. (110a) *P.* serraeforme.*

(110b) *P.* polymorphum.

1.1. *Fronds pinnatifid nearly or quite to the costa.*

2. *Lobes entire, broadly toothed or sinuate. (111) *P.* contiguum.*

2.2. *Lobes crenate or serrate towards the apex.*

- 3.** *Base of frond crenate or repand. Sori 18—25 on the largest lobes.* (112a) *P. cryptocarpum.*
- 3.3.** *Base of frond sinuate. Sori ± 9 on the largest lobes.* (112b) *P. Toppingii.*

(108a) **P. Schlechteri**, *v. A. v. R.*; *Prosaptia Schlechteri*, Christ, Schum. & Laut., Flor. deut. Sch.ggeb., Suppl., 41, tab. I, fig. B, c; *Davallia Schlechteri*, C. Chr., Ind. Fil.

Rhizome short. Fronds sessile, tufted, numerous, ± 25 c.M. long, ± 1/2 c.M. broad, linear-ligulate, acuminate, narrowed at the base, the edge rotundato-crenate in the lower, sharply toothed towards the fertile higher part. Texture coriaceous, rigid; surfaces naked, upper bright-green, lower ochraceous; costa immersed; veins oblique, hidden. Sori solitary on the teeth, oblong, the depressions saccate, ovate, not scariose, consisting of the partially reflexed margin of the lobes.

New Guinea.

* (108b) **P. exaltatum**, *Copel.*, MS. in Herb. Bog.; *Davallia exaltata*, *Copel.*, Polypod. Philipp., 52; C. Chr., Ind. Fil.; *Acrosorus exaltatus*, *Copel.*, Philipp. Journ., I, Suppl., 159, tab. XV.

Rhizome short, ascending, the scales dense, linear, brown. Fronds sessile, crowded, pendulous, 30—50 c.M. long, 5—9 m.M. broad, narrowed towards the base, the lower 1/3 part sterile, pinnatifid almost to the costa, the lower segments reduced. Segments unequally triangular, oblique, the apex bluntish. Texture coriaceous; surfaces naked, the lower densely covered with irregularly scattered, black or dark-brown dots; veins hidden. Sori solitary, deeply immersed in terminal, saccate, indusium-like depressions, which form much prominent, laterally compressed projections on the upper surface.

Mindanao (Copeland, No. 1478).

(108c) **P. tortile**, *v. A. v. R.*; *Acrosorus Merrilli*, *Copel.*, Philipp. Journ., II, 136.

Differs from *P. exaltatum*, *Copel.*, in the fronds being smaller, with narrower sterile segments and more reduced and twisted fertile segments giving the fertile apical part of the frond an appearance of distinctness wanting in *P. exaltatum*, *Copel.*

Mindoro.

(109) **P. Sarasinorum**, v. A. v. R.; *P. Friderici et Pauli*, Christ, Ann. Bot., XIX, 37 (not XV, 148); *Davallia Friderici et Pauli*, Christ, l. c., XV, 94, tab. XIV, fig. 10, a—d; C. Chr., Ind. Fil.; *Acrosorus friderici et pauli*, Copel., Philipp. Journ., I, Suppl., 159. ✓

Rhizome woody, ascending, the scales few, ovate, reddish-brown. Fronds numerous, 15—20 c.M. long, ligulate, \pm 6 m.M. broad, narrowed towards the apex, pinnate. Pinnae deltoid, bluntish, \pm 2 m.M. each way, close, 60—90 on each side, the lower ones gradually becoming shorter. Texture thick, spongy, but firm-coriaceous when dry; veins inconspicuous, hidden, 1 to each pinna. Sori occupying the upper half of the fronds, 1 to each pinna. Soriferous pinnae slightly narrowed near the base, the upper part subclavate, thickened, with the sori solitary, immersed in \pm deep, marginal, subterminal depressions.

Celebes.

* (110a) **P. serraeforme**, J. Sm., Journ. of Bot., III, 394; *Davallia alata*, Bl., Rac., Flor. Bot., I, 132; C. Chr., Ind. Fil.; *D. Emersoni*, Hk. & Grev., Ic. Fil., tab. CV; *D. serraeformis*, Wall. (oldest name); *Prosaptia Emersoni*, Pr., Bedd., Ferns S.I., tab. XX; *Pr. serraeformis*, Christ, Bull. Herb. Boiss., VI, 143; *Pr. alata*, Christ, Copel., Philipp. Journ., I, Suppl., 158, tab. XIV, B.

Rhizome short, the scales dense, ciliated. Fronds sessile, tufted or subtufted, linear-lanceolate, 15—30 c.M. long, 2—5 c.M. broad, narrowed towards both ends, cut \pm $\frac{1}{2}$ — $\frac{2}{3}$ -way down to the costa into numerous, linear-oblong, spreading or suboblique, blunt lobes, except in the lower part, which is often entire, repand or sinuate with short, deltoid lobes. Texture coriaceous; under surface covered with scattered, short, acute glands or hairs. Sori 1—12 to a lobe, closely placed round the edge, especially near the apex, deeply sunk in submarginal, indusium-like depressions.

Trop. Asia, Samoa.

(110b) **P. polymorphum**, v. A. v. R.; *Prosaptia polymorpha*, Copel., Philipp. Journ., II, 136.

Differs from *P. serraeforme*, J. Sm. by the broader and less truncate segments and the sori immersed in obtuse teeth. Fronds to \pm 40 c.M. long, 3 c.M. broad, acuminate. Segments oblong, obtuse or subacute, the lower gradually passing into a wavy wing 5—8 c.M. long, entire at the base and narrowed into a short, slender stipe. Sori \pm 5 to a segment, not restricted to the apical part.

Mindoro.

* (111) **P. contiguum**, **J. Sm.**, Journ. of Bot., III, 394; *Davallia contigua*, Spr., Rac., Flor. Btz., I, 152; C. Chr., Ind. Fil.; *Prosaptia contigua*, Bedd., Ferns S.I., tab. XIX; *Trichomanes contiguum*, Forst., Prod., 84; — var. monosorum, Copel., in Perk., Fragm. Philipp., 179; — var. pectinatum; *D. pectinata*, Bl. (not Sm.), Enum., 229.

Rhizome short, the scales dense, brown, ciliate. Fronds sessile or on stipes to 7 c.M. long, tufted or subtufted, 15—50 c.M. long, 2—5 c.M. broad, linear-lanceolate, narrowed towards both ends, incised nearly or quite to the rachis into linear-oblong, blunt, contiguous or slightly joined segments \pm 2—4 m.M. broad, slightly dilated at the base, the edge entire but remotely, shallowly and broadly toothed towards the apex, the lower segments shorter, deltoid. Texture coriaceous; under surface sparingly hairy, especially on the rachis. Sori not numerous, 1 to each tooth, more or less deeply sunk in \pm marginal, indusium-like depressions.

Var. monosorum: Segments narrower, quite entire, 1—1½ m.M. broad. Sori solitary and terminal on the segments.

Var pectinatum: Segments undulate. Sori marginal, all round the edge.

Malaya; Polynesia; — var. monosorum: *Mindanao, New Guinea* — var. pectinatum: *Java*.

* (112a) **P. cryptocarpum**, **v. A. v. R.**; *Prosaptia cryptocarpa*, Copel., Philipp. Journ., I, Suppl., 158, tab. XIV, a & d.

Rhizome short, rather creeping, clothed with black-brown, lanceolate, acuminate scales. Stipes 1—1½ c.M. long, minutely pubescent, the base black. Fronds 20—55 c.M. long, 3½—5 c.M. broad, the apex acute to caudate, the edge deeply pinnatifid, the costal wing \pm ¼—1½ c.M. broad, the base gradually narrowed, crenate to repand. Lobes numerous, \pm spreading, 2—3 m.M. broad, obtuse or acute, slightly repand or serrate towards the apex, dilated at the base. Texture coriaceous; upper surface naked, lower provided with scattered, short, red, subulate hairs, especially on the costa; costulae inconspicuous; veins hidden, simple. Sori 18—25 on the largest lobes, marginal or submarginal.

Mindanao.

(112b) **P. Toppingii**, **v. A. v. R.**; *Prosaptia toppingii*, Copel., Philipp. Journ., I, Suppl., 158, tab. XIV, c.

Near *P. cryptocarpum*, **v. A. v. R.** Fronds 20—50 c.M. long, 3—4

c.M. broad, acute, the costal wing 2—3 m.M. broad, the base sinuate. Lobes triangular-lanceolate, subacute, \pm 5 m.M. broad, obtusely serrate. Sori \pm 9 on the largest lobes, marginal.

Luzon.

§ 5. GONIOPHLEBIUM. c. **Marginaria.**

(115) **P. Prainii**, *C. Chr.*, Ind. Fil.; *Goniophlebium Prainii*, *Bedd.*, Journ. of Bot., XXXI, 226.

Rhizome stout, densely clothed with long, finely acuminate, brown scales, which are very iridescent in the broader part near the peltate base. Stipes 45—60 c.M. long, firm, erect, naked. Fronds deltoid-lanceolate, \pm 30 c.M. long, \pm 25 c.M. broad, pinnate. Pinnae numerous, \pm 2 c.M. broad, narrow-lanceolate from a broad base, separate, the margin slightly crenate especially towards the apex. Texture somewhat papyraceous; surfaces naked except some minute scales and a few hairs on the rachis below; veins prominent on both sides; areolae in 2 series. Sori in 1—2 rows on each side of the costae.

Malacca.

§ 3. GONIOPLEBIUM. d. **Schellolepis.**

- 1.** *Pinnae acutely cuneate or gradually narrowed at the base.*
- §. *Sori in 2—3 rows on each side of the costae.* (114) *P. Korthalsii.*
- §.§. *Sori in 1 row on each side of the costae.*
- ×. *Pinnae subentire.* (115a) *P. integriore.*
- ×.×. *Pinnae crenate, serrate or toothed.* (115b) *P. persicifolium.*
- 1.1.** *Pinnae broadly and abruptly cuneate at the base.*
- *. *Pinnae paucijugate.* (116a) *P. Koningsbergeri.*
- *.*. *Pinnae multijugate.* (116b) *P. verrucosum.*
- 1.1.1.** *Pinnae cordate, broadly rounded, truncate or the higher adnate at the base (*).*
- 2.** *Surfaces more or less densely hirsute, hairy or tomentose.*
- a.** *Veins forming 1 row of areolae on each side of the costae.*
- b.** *Sori scarcely immersed.* (117a) *P. molliculum.*
- b.b.** *Sori distinctly immersed.* (117b) *P. pallens.*
- a.a.** *Veins forming 2—3 rows of areolae on each side of the costae.* (118) *P. Beddomei.*
- 2.2.** *Surfaces naked.*

(*) See also **P. verrucosum**, *Wall.*

3. *Sori superficial.***4.** *Sori in 1—2 rows on each side of the costae.***5.** *Pinnae narrowly triangular.* (119a) *P. argutum.***5.5.** *Pinnae not uniformly narrowed from the base to the apex.*
(119b) *P. pseudoconnatum.***4.4.** *Sori in 3 rows on each side of the costae.*(120) *P. lampongense.***3.3.** *Sori more or less immersed.*+. *Terrestrial. Sori subimmersed.* (121a) *P. benguetense.*+.. *Epiphytical. Sori deeply immersed.*(121b) *P. subauriculatum.*

* (114) **P. Korthalsii**, *Mett.*, Hk. Bk., Syn. Fil., 345; *Goniophlebium Korthalsii*, *Bedd.*, Handb. Ind. Ferns, Suppl., 91.

Rhizome creeping, clothed with small, linear-lanceolate-subulate, ciliate, reddish-brown scales. Stipes 12—15 c.M. long, smooth, reddish-brown. Fronds 50—40 c.M. long, to \pm 50 c.M. broad. Pinnae lax, spreading or erecto-patent, 12—15 c.M. long, $1\frac{1}{2}$ —4 c.M. broad, lanceolate-acuminate, the base cuneate, articulate, the edge slightly crenate. Texture subcoriaceous; rachis and both surfaces naked; areolae in 3—4 rows. Sori \pm 2—5-serial, superficial or hardly immersed.

Java, Sumatra, Malacca.

* (115a) **P. integriore**, *Copel.*, Philipp. Journ., II, 139.

Rhizome creeping, woody, clothed with lanceolate-acuminate, castaneofulvous scales or glabrescent, and not calcareous. Stipes naked, 15—25 c.M. long, brown-stramineous. Fronds \pm 50 c.M. long. Pinnae linear, to \pm 20 c.M. long, to \pm $1\frac{1}{2}$ c.M. broad, stalked, the base narrowly acuminate, the edge entire or crenulate, the apex distinctly caudate, hardly serrulate. Texture membranaceous; surfaces naked; areolae in \pm 3 rows on each side. Sori large, slightly immersed. — Nearly related to *P. persicifolium*, *Desv.*

Mindoro, Java.

* (115b) **P. persicifolium**, *Desv.*, Rac., Flor. Btz., I, 115; *P. cuspidatum*, *Bl.* (not *Don*), Flor. Jav., II, 175, tab. LXXXII; *P. phlebodioides*, *Copel.*, Polypod. Philipp., 123; Philipp. Journ., I, Suppl., 162; *Goniophlebium*

cuspidatum, Pr., Bedd., Ferns Br.I., I, tab. LXXIX; — var. *grandidens*; *P. grandidens*, Kze, Bot. Zeit., 1846, 425.

Rhizome creeping, clothed with lanceolate to linear, dark-brown, \pm ciliate scales, not cretaceous. Stipes 15—45 c.M. long, firm, naked, stramineous or subebeneous. Fronds 50—300 c.M. long, 20—40 c.M. broad, often subpendulous. Pinnae articulate, distant, short-stalked, 1—3½ c.M. broad, the apex acuminate or gradually narrowed, the edge crenate, serrate or slightly toothed, the base rather acutely cuneate or gradually narrowed. Texture membranaceous to subcoriaceous; rachis and surfaces naked; areolae in \pm 3 rows on each side. Sori in 1 row on each side, not deeply immersed (often apparently superficial in dried material).

Var. grandidens: Edges deeply toothed.

Sumatra, Java, Celebes, Philippines.

* (116a) **P. Koningsbergeri**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 21.

Near *P. verrucosum*, Wall. Rhizome scales indistinctly ciliato-dentate, rather entire. Stipes distant, slender, soon naked, \pm 10—15 c.M. long. Fronds small, 15—20 c.M. long, \pm 12—15 c.M. broad. Pinnae paucijugate (\pm 2—3 on each side below the terminal one; terminal pinna \pm 10—12½ c.M. long, the lateral ones similar, sessile, \pm opposite, articulate, erecto-patent, 7—8 c.M. long, \pm 1½ c.M. broad near the base, narrowed gradually towards the acuminate apex, the margin indistinctly crenate or crenato-serrate, the base broadly, abruptly cuneate. Texture subcoriaceous; surfaces naked except a few scattered, deciduous, fimbriate, minute, lanceolate scales on the costae beneath, \pm hairy in the younger forms; areolae in 3 rows on each side. Sori confined to the costal areolae, slightly immersed, not very prominent on the upper surface; paraphyses yellow, indistinctly incised at the apex. — This species bears the name of Dr. J. C. Koningsberger, Subdirector of the Agricultural Department in Netherlands India.

Java (Preanger Regencies, v. A. v. R.).

* (116b) **P. verrucosum**, Wall., Hk Bk., Syn. Fil., 344; Hk., Gard. Ferns, tab. XLI; *Goniophlebium verrucosum*, J. Sm., Bedd., Ferns Br.I., II, tab. CCLVII.

Rhizome firm, creeping, not cretaceous; scales lanceolate-oblong, the base peltate, the edge ciliate, the apex acuminate, erect, subadpressed. Stipes scattered, 30—60 c.M. long, copiously clothed like the rachis with caducous scales,

at length quite naked, the scales linear-acuminate, subulate, ciliate, recurved or spreading. Fronds subpendulous, 45—120 c.M. long, 15—30 c.M. broad. Pinnae numerous, subsessile, articulate to the rachis, erecto-patent or spreading, linear, to 20 c.M. long, 1—4 c.M. broad, the edges mostly \pm parallel, entire or crenato-dentate, the base obtusely cuneate or nearly rounded, the upper half narrowed gradually to the point, or the apex abruptly narrowed, short-cuspidate; terminal pinna simple, similar to the lateral ones but not rarely very elongated and strap-shaped. Texture papyraceous or subcoriaceous; surfaces naked or villose, often conspicuously hairy in the younger forms; costae whether or not with some few scattered fimbriate scales beneath; areolae in 4—5 rows on each side of the costae. Sori in 1 row on each side, confined to the costal areolae, immersed in deep cavities, which form very distinct warts or tubercles on the upper surface; paraphyses dark-brown, oblong and mostly irregularly, pinnately or stellately incised at the apex.

Malaya; Trop. Australia.

(117a) **P. molliculum**, *Copel.*, Polypod. Phiipp., 124; *P. tomentellum*, *C. Chr.*, Ind. Fil.

Rhizome wide-creeping, the scales small, subulate, ferruginous. Stipes \pm 10 c.M. long, suberect, naked or nearly so. Fronds 15—20 c.M. long, \pm 8 c.M. broad, pinnate. Pinnae narrowly lanceolate, \pm 6—8 m.M. broad, acute, serrate or crenate towards the base, short-auriculate, the base \pm truncate, sessile but not adnate. Texture herbaceous; surfaces clothed with a short, soft tomentum; veins free beyond a single row of areolae on each side. Sori scarcely immersed.

Philippines.

* (117b) **P. pallens**, *Bl.*, Flor. Jav., II, 178, tab. LXXXIV, A; *P. subauriculatum*, *Bl.*, var., *C. Chr.*, Ind. Fil.

Rhizome creeping, pruinose, clothed with few, thin, linear scales, at length \pm naked. Stipes subremote, 5—15 c.M. long, glabrescent, pale-green, here and there pale-fuscous. Fronds 10—40 c.M. long, 5—10 c.M. broad, oblong, the rachis dirty-stramineous, hirsute, the hairs intermixed with small, adpressed or curved, narrow-lanceolate or subulate, fimbriato-denticulate, black-fuscous, deciduous scales. Pinnae \pm 10—15-jugate; terminal pinna \pm pinnatifid at the base, the lateral ones opposite or alternate, sessile, straight or subfalcate, the base broadly rounded to cordato-truncate, the edge crenato-serrulate, the apex bluntish or acute. Texture rather herbaceous to coriaceous;

surfaces more or less densely clothed with \pm deciduous, whitish hairs; veins anastomosing commonly in 1 row of large, costal areolae. Sori in 1 row on each side of the costae, immersed.

Java.

(118) **P. Beddomei**, *Bk.*, Hk. Bk., Syn. Fil., 344; *Goniophlebium molle*, *Bedd.*, Handb. Ind. Ferns, 322; Ferns Br.I., II, tab. CCVI.

Rhizome creeping, densely clothed with reddish-brown, subulate scales. Fronds \pm 45 c.M. long, 17—20 c.M. broad, oblong, acuminate. Pinnae numerous, 1—1 $\frac{1}{2}$ c.M. broad, sinuate-crenate, superior base often subauricled; superior pinnae decurrent on the rachis, remainder sessile from a broad base, not adnate, or inferior ones subpetiolate. Texture membranaceous; rachis and both surfaces covered with soft, weak, whitish hairs; veins forming 2—3 rows of areolae. Sori in 1 row on each side of the costae.

Malacca; Burmah.

(119a) **P. argutum**, *Woll.*, Hk. Bk., Syn. Fil., 511; *Goniophlebium argutum*, *J. Sm.*, *Bedd.*, Ferns Br.I., I, tab. VI.

Rhizome firm, wide-creeping, the scales minute, linear-subulate, dark-brown. Stipes 7—15 c.M. long, naked, castaneo-stramineous. Fronds suberect, lanceolate-oblong, 30—60 c.M. long, 20—30 c.M. broad. Pinnae 10—20-jugate, ligulate-acuminate, faintly inciso-crenate, 1 $\frac{1}{4}$ —2 c.M. broad, many free, distant, uniformly narrowed from the base to the apex, lower subpetiolate, lowest not reduced. Texture membranaceous; surfaces bright-green, naked; areolae 1-serial. Sori 1-serial, nearer the midribs than the edge, superficial.

Luzon; North. India.

(119b) **P. pseudoconnatum**, *Copel.*, Philipp. Journ., I, Suppl., 161, tab. XXII.

Rhizome woody, cretaceous, densely clothed with deciduous, lanceolate, acuminate, brown scales. Stipes \pm 20 c.M. long or longer, strong, glabrescent. Fronds \pm 50 c.M. long, 30—50 c.M. broad, acuminate, the rachis sordido-paleaceous towards the apex. Pinnae often opposite, sessile, lowest largest, \pm 20 c.M. long, to \pm 2 c.M. broad, cordate at the base, the edge coarsely serrate, the apex acuminate. Texture herbaceous; surfaces nearly naked; veins forming 1 row of large costal areolae and sometimes 1 or more series of smaller areolae outside the costal ones. Sori large, superficial.

Luzon.

(120) **P. lampongense**, *Zoll.*, Syst. Verz., 37, 48.

Fronds scattered, pinnate. Pinnae linear-lanceolate, the higher and fertile ones narrower than the barren, subensiform, sessile, cordate, subauriculate at the base. Rachis angular, sparingly paleaceo-villose; veins slender, naked. Sori in 3 rows on each side.

Java.

(121a) **P. benguetense**, *Copel.*, Philipp. Journ., I, Suppl., 256; *P. mengtzeense*, *Copel.* (not *Christ*), l. c., 161, tab. XXI.

Rhizome creeping, densely clothed with lanceolate, acuminate, black-fulvous scales. Stipes \pm 15 c.M. long, often stramineous and glabrescent like the rachis. Fronds 20—25 c.M. long, 12—20 c.M. broad. Pinnae \pm 1—2 c.M. apart, often opposite, sessile, the base subcordate or hastato-dilatate, the edge acutely serrate, the apex acuminate; higher pinnae distinct, not decurrent, nor broadly adnate. Texture membranaceous; surfaces naked; costae prominent; veins forming 1 row of costal areolae and sometimes also a second row of smaller areolae outside the costal ones. Sori subimmersed.

Luzon.

* (121b) **P. subauriculatum**, *Bl.*, Rac., Flor. Btz., I, 116; *Bl.*, Flor. Jav., II, 177, tab. LXXXIII; *P. Reinwardtii*, *Kze*, Linn., XXIII, 285; *Goniophlebium Reinwardtii*, *de Vr.*, Ned. Kruidk. Arch., I, 257; *G. subauriculatum*, *Pr.*, Bedd., Ferns Br.I, I, tab. LXXVIII; — var. *serratifolium*; *G. serratifolium*, *Brack.*, Expl. Exp., XVI, 35.

Rhizome wide-creeping, pruinose, scaly, at length \pm naked, the scales dark-brown, linear-lanceolate or subulate, entire or ciliato-dentate, finely acuminate, erecto-patent or spreading. Stipes scattered, firm, 15—50 c.M. long, glossy, clothed like the rachis with densely crowded, \pm spreading or recurved, brownish, deciduous, finely acuminate, ciliate scales, afterwards naked. Fronds subpendulous, \pm 50—500 c.M. long, 8—50 c.M. broad, oblong-lanceolate, pinnate. Terminal pinna mostly pinnatifid at the base; lateral pinnae numerous, subspreading, linear-lanceolate, $\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, articulate to the rachis, sessile, with \pm their own breadth between them, the base cordate, \pm truncate or rounded, often \pm auricled on one or both sides, the edge entire, slightly crenate or faintly toothed, the apex narrowed \pm gradually, acute or acuminate. Texture herbaceous to subcoriaceous; surfaces naked, the lower covered with scattered, minute, roundish, irregularly ciliate, deciduous, peltate scales when young; rachis at length naked;

areolae in 2—3 rows on each side. Sori in 1 row on each side, distinctly immersed.

Var. serratifolium: Pinnae coarsely toothed.

Malaya; Samoa, Trop. Australia, New Caledonia.

§ 6. PLEOPELTIS. a. **Myrmecophila.** *. **Aspidopodium.**

1. *Fronds simple or lobed.* (122) *P. sinuosum.*

1.1. *Fronds deeply pinnatifid or pinnate.*

2. *Rhizome forming a single, bulbous or wide-spreading crust.*

3. *Fronds pinnatifid at the base; sori subcostular.*

(125) *P. lomarioides.*

3.3. *Fronds pinnate at the base. Sori medial.* (124) *P. sarcopus.*

2.2. *Rhizome forming a row of united, subconvex, oblong or roundish crusts.*

(125) *P. imbricatum.*

* (122) **P. sinuosum**, *Wall.*, *Rac. Flor. Btz.*, I, 103; *Hk.*, *Spec. Fil.*, V, tab. CCLXXXIV; *Pleopeltis sinuosa*, *Bedd.*, *Ferns Br.I.*, I, tab. VIII.

Rhizome wide-creeping, thick, densely clothed with adpressed, peltate, roundish scales, dark in the centre and pale round the edge. Stipes $\frac{1}{2}$ —5 c.M. long, firm, naked, arising from large, prominent, conical protuberances. Fronds dimorphous; barren fronds 7—17 $\frac{1}{2}$ c.M. long, 2—2 $\frac{1}{2}$ c.M. broad, the apex blunt or rounded, the edge entire, narrowed to the base, fertile ones \pm 20—40 c.M. long, 1—2 c.M. broad, narrowed to the base, the edge \pm sinuate. Texture coriaceous; surfaces naked; veins obscure. Sori large, immersed, in 1 row on each side, nearer the edge than the midrib, 1 to each lobe.

Malaya.

* (123) **P. lomarioides**, *Kze.*, *Hk. Bk.*, *Syn. Fil.*, 365; *Mett.*, *Farngr.*, I, *Polypod.*, tab. II, fig. 18—19; *Drynaria lomarioides*, *J. Sm.*, *Journ. of Bot.*, III, 397.

Rhizome forming a thick, wide-spreading crust densely clothed with small peltate scales, ferruginous in the centre, with a broad, scariose border. Stipes arising from conical protuberances, 5—10 c.M. long, firm, erect. Fronds 15—45 c.M. long, 7—15 c.M. broad, the barren ones cut down within 5—7 m.M. of the costa into close, horizontal, oblong, entire, blunt

lobes \pm 2 c.M. broad; the fertile ones cut nearly to the costa with more numerous lobes 6—20 m.M. broad. Texture coriaceous; surfaces naked; veins immersed, irregular; areolae fine, with free veinlets. Sori immersed, round or oblong, forming 2 close rows near the central ribs of the lobes.

Malaya; Formosa.

(124) **P. sarcopus**, *de Vr. & Teysm.*, Hk. Bk., Syn. Fil., 514; *P. myrmecophilum*, *Christ*, Fil. Saras., I, 24; III, tab. III, fig. 21—24.

Rhizome inflated, greyish, 6—10 c.M. thick (high), \pm 15 c.M. broad (each way), the scales dense, small, roundish, peltate, black in the centre, with a whitish border. Stipes 10—30 c.M. long, stramineous, thick, arising from conical protuberances or with the base immersed in the rhizome. Fronds \pm 40 c.M. long, with 30—35 pinnae on each side. Pinnae entire, 7—9 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad, the point blunt; higher pinnae close, connected, the lower ones free. Texture moderately coriaceous, rather succulent; veins fine, not very conspicuous; main veins 6—8 on a side, not quite reaching the margin; areolae irregular, with few free veinlets. Sori in 1 row on each side of the costae, medial, very deeply immersed, forming very prominent papillae on the upper surface of the pinnae, the mouth of the pits surrounded by a distinctly raised border.

Celebes; South. China.

(125) **P. imbricatum**, *Karst.*, Ann. Btz., XII, 168, tab. XV, fig. 29—30, tab. XIX; *P. mirabile*, *C. Chr.*, Ind. Fil.

Rhizome climbing, consisting of several to numerous divisions separated by distinct constrictions 4—8 c.M. broad, each internode roundish or oblong, convex, 8—16 c.M. long, 7—15 c.M. broad, \pm 8 m.M. thick in the centre, concave beneath, \pm 2 m.M. thick at the margin, clothed with black, roundish, peltate scales, which are scattered on the upper, but densely crowded on the under side. Fronds 4—9 to each division, ranged in 2 approximate rows (1 row on each side of the longitudinal axis), 30—50 c.M. long, deeply pinnatifid in the fully developed plant, the stipes placed on conical protuberances. Lobes more or less numerous, erecto-patent. Sori deeply immersed, prominent on the upper surface, in 1 row on each side of the costulae.

Moluccas.

§ 6. PLEOPELTIS. a. **Myrmecophila.** **. **Lecanopteris.**

- a.** Soriferous teeth or lobes turned back on the upper surface of the frond.
- b.** Sori immersed in the outgrowths.
- c.** Outgrowths not twisted, the mouth of the cavities facing upwards.
- d.** Sori oblong, rather distant. (126) *P. lecanopteris*.
- d.d.** Sori round, rather close. (127) *P. Nieuwenhuisii*.
- c.c.** Outgrowths twisted, the mouth of the cavities facing towards the apex of the pinna. (128) *P. deparoides*.
- b.b.** Sori superficial on the outgrowths.
- *** Outgrowths twisted. (129) *P. Macleayi*.
- *.*** Outgrowths not twisted. (130) *P. barisanicum*.
- a.a.** Soriferous teeth or lobes erect or erecto-patent, not turned back on the upper surface of the frond. (131) *P. naviculare*.

* (126) ***P. lecanopteris***, *Mett.*, Ann. Mus. Bot. L. B., II, 224; *P. carnosum*, *Christ*, Ann. Btz., XV, 160; *P. patelliferum*, *Burck.*, Ann. Btz., IV, 96, tab. VII; *Lecanopteris carnosa*, *Bl.*, Flor. Jav., II, tab. XCIV, A; *Copel.*, Polypod. Philipp., 133; *C. Chr.*, Ind. Fil.; — *var pumilum*; *L. pumila*, *Bl.*, l. c., tab. XCIV, B; *Copel.*, l. c., 154; *C. Chr.*, l. c.

Rhizome thick, naked, yellowish, inflated, forming a semiglobose mass \pm 15 c.M. broad, \pm 9 c.M. high, covered with several conical or semiglobose protuberances and subulate, simple or branched, scaly prickles (probably reduced fronds). Stipes on the protuberances, 7—15 c.M. long, brown, naked, glossy. Fronds dimorphous, pinnatifid nearly or quite to the costa. Segments many to numerous, $2\frac{1}{2}$ —5 c.M. long, \pm 1—2 c.M. broad, the sterile ones oblong-lanceolate, blunt, with the margin entire or repand, the fertile ovate-lanceolate, acute, with the margin lobed with soriferous projections, the lobes (projections) \pm oblong, oblique, rather distant, obtuse, narrowed at the base, turned back on the upper surface of the frond, with the deeply immersed sori facing upwards. Texture coriaceous; surfaces naked; veins indistinct. Sori rather large, 3—6 m.M. long, \pm oblong, immersed in cavities of the same shape, solitary on and occupying nearly the whole of the lobe-like projections.

Var. pumilum: Segments blunt or bluntish; texture firm but pellucid; veins distinct; sori smaller.

Malaya, especially Celebes.

* (127) **P. Nieuwenhuisii**, v. A. v. R.; *Lecanopteris Nieuwenhuisii*, *Christ*, Ann. Btz., XX, 127, tab. III, fig. 1; C. Chr., Ind. Fil.

Rhizome naked, inflated, thick. Stipes 4—6 c.M. long, firm, naked, stramineous like the rachis. Fronds linear-lanceolate, \pm 40 c.M. long, \pm 5½ c.M. broad, acuminate, narrowed gradually towards the base, the edge deeply pinnatifid nearly to the costa. Lobes \pm 25 on each side below the long terminal one; lower lobes oblong, entire, barren, the central and higher ones fertile, sinuato-lobate, the central ones \pm 3 c.M. long, \pm 1½—2 c.M. broad. Soriferous outgrowths rather close, semiorbicular (the apex and sinuses rounded), subspreading, turned back on the upper surface of the frond, with the deeply immersed sori facing upwards. Texture firm, pellucid when dry; surfaces naked; costulae and main veins visible; areolae fine, with free veinlets. Sori large, 1 to each outgrowth, \pm 5—6 on each side, deeply immersed so as to form semiglobose, \pm cucullate, mamillary papillae on the upper surface.

Borneo.

* (128) **P. deparoides**, *Christ*, Farnkr. d. Erde, 116; *Davallia deparoides*, *Ces.*, Fel. di Bor., 15; *Lecanopteris deparoides*, *Bk.*, Journ. of Bot., XIX, 566; C. Chr., Ind. Fil.; *Cyathea deparoides*, *Ces.*, l. c., tab. IV.

Rhizome \pm 3 c.M. thick, creeping, tortuous, thinly pubescent, afterwards pulverous, at length naked, tuberculous, yellowish-brown. Stipes stramineous, naked. Fronds erect, to \pm 65 c.M. long. Pinnae to 40-jugate, opposite, linear, obtuse, lobato-dentate, the lower ones free, \pm 4½ c.M. long, the following \pm 5 c.M. or more long, 6—8 m.M. broad, the higher ones growing gradually shorter, connected by a narrow wing. Texture firm; surfaces naked; costae and main veins distinct; areolae with free veinlets. Sori on the lobe-like teeth, deeply immersed, 1 to each tooth, the teeth turned back on the upper surface of the frond, induplicate, twisted, the cavities becoming oblong with the opening facing towards the apex of the pinna so as to form spurious iudusia resembling those of *Davallia*. — *Cesati's* drawing does not agree with *Teysmann's* *Karimata* plant (*Becc.*, Mal., II, 244—245), which has the soriferous teeth distinctly inflexed and twisted.

Borneo, Sumatra.

(129) **P. Macleayi**, v. A. v. R.; *Lecanopteris Macleayi*, *Bk.*, *Becc.*, Mal., II, 244; C. Chr., Ind. Fil.

Like *P. deparoides*, *Christ*, but the sori terminal on the teeth, not in cavities.
Java.

(130) **P. barisanicum**, v. A. v. R.; *Lecanopteris incurvata*, Bk., Ann. of Bot., VIII, 123; C. Chr., Ind. Fil.

Stipes short, erect, firm, yellow-grey to brownish, naked. Fronds oblong-lanceolate, caudate, simply pinnate, 25—40 c.M. long. Pinnae multijugate, 3—5 c.M. long, confluent at the base or connected by a narrow wing, the barren ones linear-oblong, entire, $1\frac{1}{2}$ —2 c.M. broad, obtuse, the fertile ones narrower, to 1 c.M. broad, lanceolate, deeply crenate (toothed?). Texture membranaceous; under surface glaucescent; areolae hexagonal, copious, with copious free veinlets. Sori solitary and terminal on the teeth, which are induplicate, boat-shaped, and turned back on the upper surface of the frond.

Sumatra.

* (151) **P. naviculare**, v. A. v. R.; *Lecanopteris Curtisii*, Bk., Journ. of Bot., XIX, 366; Hk., Icon. Pl., XVII, tab. MDCVII; C. Chr., Ind. Fil.

Stipes 5 c.M. long or longer, shining, pale-brown like the rachis. Fronds lanceolate, simply pinnate or deeply pinnatifid, 35—45 c.M. long, 5—12 c.M. broad. Pinnae \pm 10—30 pairs below the terminal one, ligulate, obtuse, narrowed near the base, $2\frac{1}{2}$ —6 c.M. long, 6—12 m.M. broad, all fertile except the 2—7 lower pairs, which are rather shorter or \pm reduced. Texture chartaceous; surfaces naked, the lower glaucous; main veins distinct; areolae fine. Sori small, 1 to each tooth; soriferous teeth minute, \pm 7—8 on each side of the pinnae, erect or erecto-patent, twisted, not turned back on the upper surface of the frond, each with a laterally compressed, boat-shaped cavity, with the mouth facing towards the apex or margin of the pinna.

Sumatra, Lingga, Borneo, Celebes.

§ 6. PLEOPELTIS. b. **Atactosia**. *. **Lepisorus**.

a. Substance of the fronds thick; all the veins \pm obscure.

b. Sori scattered irregularly.

§. Fronds roundish or oblong, nearly as broad as long, under 2 c.M. long. (152) *P. holophyllum*.

§.§. Fronds elongate, more than 2 c.M. long.

*. Rhizome wide-creeping or scandent.

×. Fronds narrow, to 2 c.M. broad.

+ . Fronds \pm 7—15 c.M. long, distinctly stalked.

(135) *P. subsparsum*.

- +.+. Fronds \pm 20—45 c.M. long, sessile or short-stalked. (154) *P. tenuilore.*
- ×.×. Fronds broader, \pm 2½—5 c.M. broad when fully developed.
- +. Rhizome naked. Sori slightly immersed. (135) *P. sarawakense.*
- +.+. Rhizome scaly. Sori superficial.
- △. Rhizome scales brown, lanceolate. (136) *P. superficiale.*
- △.△. Rhizome scales black, roundish, the margin brown. (137) *P. peltatum.*
- *.*. Rhizome short-creeping.
- ×. Fronds short and broad, spathulate, entire. (138) *P. antropoides.*
- ×.×. Fronds moderately long and narrow, lanceolate, crenatodentate. (139) *P. curticens.*
- b.b.** Sori in 1 or sometimes 2 rows on each side of the costa.
- c.** Fronds more or less dimorphous; fertile fronds longer and narrower than the barren ones, sometimes much contracted. (*)
- d.** Fertile fronds stipitiform, the lamina contracted to a mere midrib. (140) *P. Bakeri.*
- d.d.** Fertile fronds not stipitiform, the lamina distinctly developed.
- e.** Sori superficial.
- f.** Fertile fronds entire or slightly crenate.
- g.** Barren fronds narrow, mostly under 1 c.M. broad, the fertile ones not narrowed in the soriferous part.
- h.** Barren fronds under 5 c.M. long. (141) *P. Wrayi.*
- h.h.** Barren fronds \pm 6—10 c.M. long. (142) *P. Beccarii.*
- g.g.** Barren fronds mostly broader, and then often 1½—3 c.M. broad, the fertile ones narrowed in the soriferous part.
- ×. Barren fronds more than 2 c.M. long. (143a) *P. accedens.*

(*) See also *P. neglectum*, Bl.

x.x. Barren fronds under 2 c.M. long.

(143b) *P. damunense*.

f.f. Fertile fronds repand or slightly lobed.

*. Repand. Sori on the lobes.

(144) *P. nummularium*.

.. Lobed. Sori on the anterior side of the lobes.

(145) *P. hammatisorum*.

e.e. Sori distinctly immersed.

‡. Fertile fronds lanceolate in the lower, linear in the
soriiferous upper half. (146) *P. stenopteris*.

‡.‡. Fertile fronds linear-lanceolate. (147) *P. soridens*.

c.c. Fronds uniform or nearly so, the fertile ones sometimes narrower
and longer stalked.

1. Sori superficial or slightly immersed.

2. Sori in 1 row on each side of the midrib.

3. Fronds small, under 3 c.M. long.

*. Fertile fronds cuneate at the base.

(148) *P. neglectum*.

.. Fertile fronds rounded at the base.

(149) *P. rudimentum*.

3.3. Fronds larger, 5—30 c.M. long.

4. Sori submarginal. (150) *P. subecostatum*.

4.4. Sori distant from the margin.

5. Fronds lanceolate-oblong, rostrate, 5—10
c.M. long. (151) *P. rostratum*.

5.5. Fronds linear-lanceolate, not rostrate, 10—
30 c.M. long. (152) *P. lineare*.

2.2. Sori in 2 rows on each side of the midrib.

*. Fronds lanceolate-oblong, under 15 c.M. long, short-
stalked. (153) *P. Forbesii*.

.. Fronds linear-lanceolate, more than 15 c.M. long,
long-stalked. (154) *P. induratum*.

1.1. Sori distinctly immersed.

α. Sori nearer the edge than the midrib.

β. Sori round. (155) *P. stenophyllum*.

β.β. Sori oblong. (156a) *P. revolutum*.

(156b) *P. productum*.

α.α. Sori distant from the margin.

- ×. *Sori round.* (157) *P. subgeminatum.*
 × ×. *Sori oblong.* (158) *P. papuanum.*
a.a. *Substance of the fronds rather thin; veins distinct.* (1)
b. *Sori in 1—2 regular or irregular rows on each side of the midrib.*
c. *Fronds dimorphous.*
 †. *Fertile fronds orbicular and barren in the lower, long-linear in the soriferous upper part.* (159) *P. Annabellae.*
 †.†. *Fertile fronds ligulate, cuneately narrowed at the base.* (160) *P. lycopodioides.*
c'.c'. *Fronds uniform, the fertile sometimes narrower than the barren.*
d. *Fronds rather small, under 20 c.M. long.*
e. *Areolae with free veinlets.* (161) *P. Raapii.*
e'.e'. *Areolae without free veinlets.* (162) *P. zosteriforme.*
d'.d'. *Fronds larger, ± 30—60 c.M. long.* (163) *P. normale.*
b'.b'. *Sori in several rows on each side, or scattered irregularly.* (2)
1. *Sori superficial.*
2. *Rhizome scales blackish or dark-brown, not squarrose.* (164a) *P. myriocarpum.*
 (164b) *P. sablanianum.*
2'.2'. *Rhizome scales brown, squarrose.* (165) *P. glossipes.*
1'.1'. *Sori immersed.* (166) *P. linguaeforme.*

(152) **P. holophyllum**, *Bk.*, Journ. of Bot., XVII, 45 (not XXVI, 325); (? *P. flabellivenium*, *Bk.*, C. Chr., Ind. Fil.).

Rhizome wide-creeping, the scales dense, erecto-patent, minute, linear-subulate. Stipes subdistant, brown, polished, slender, naked, 2½—5 c.M. long. Fronds roundish or oblong, 8—15 m.M. each way, crenulate round the edge except towards the base. Texture rigid; surfaces naked; no distinct midrib; veins flabellate, forming copious irregular areolae without free veinlets. Sori few, round, superficial, scattered irregularly.

Borneo.

(153) **P. subsparsum**, *Bk.*, Journ. of Bot., XVIII, 215.

Rhizome slender, wide-climbing, firm, flexuose, thinly clothed with small, adpressed, bright-brown, lanceolate scales. Stipes distant, stiffly erect, naked,

(1) See also **P. revolutum**, *C. Chr.*

(2) See also **P. normale**, *Don.*

2½—8 c.M. long. Fronds linear-lanceolate, entire, 7—15 c.M. long, ½—2 c.M. broad below the middle, narrowed gradually to both ends. Texture rigidly coriaceous; both surfaces naked; veins obscure, the main ones erecto-patent, enclosing copious, small, irregular areolae. Sori superficial, large, round, crowded in the narrowed upper half of the frond, not in regular rows.

Sumatra.

(154) **P. tenuilore**, *Kze*, Hk. Bk., Syn. Fil., 354; Mett., Farngr., I, Polypod., tab. I, fig. 28; *Diblemma samarensis*, *J. Sm.*, Bau. & Hk., Gen. Fil., tab. CIX, B; *Drynaria tenuiloris*, *J. Sm.*, Journ. of Bot., III, 397.

Rhizome slender, woody, scandent. Fronds 20—45 c.M. long, 1—1½ c.M. broad, the apex acuminate, the edge entire, the lower part narrowed very gradually into a short stipe or the base. Texture rigidly coriaceous; surfaces naked; no main veins; areolae fine, hidden, with free veinlets. Sori small, copious, scattered, sometimes confluent especially along the edge.

Philippines.

(155) **P. sarawakense**, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 228.

Rhizome slender, wide-creeping, branched, naked. Fronds few, distant, erect, simple, oblong-lanceolate, acuminate, with the base cuneately and gradually narrowed, 10—15 c.M. long, 5—4 c.M. broad at the middle, on stipes 3½—8 c.M. long. Texture subcoriaceous; surfaces naked; veins fine, immersed, hidden; areolae hexagonal. Sori large, irregularly scattered all over the under surface of the frond except the apex, slightly immersed.

Borneo, New Guinea.

* (156) **P. superficiale**, *Bl.*, Rac., Flor. Btz., I, 108; *Bl.*, Flor. Jav., II, 136, tab. LVI, fig. 1; *Pleopeltis superficialis*, *Bedd.*, Ferns Br.I., I, tab. LXXV; *P. normale*, *Don*, var. *sumatranum*, *Bk.*, Journ. of Bot., XVIII, 215.

Rhizome wide-creeping, blackish; scales brown, lanceolate, peltate, the base acute, round or cordate, adpressed, the apex recurved or erect. Stipes 5—15 c.M. long, firm, naked, slightly scaly when young. Fronds linear-lanceolate, 15—40 c.M. long, 2½—5 c.M. broad, narrowed gradually towards both ends, the edge entire. Texture coriaceous, commonly weak; surfaces naked; veins hidden or moderately visible; no main veins; areolae fine, with free veinlets. Sori roundish, sometimes slightly irregular, numerous, scattered irregularly.

Malaya; Queensland, Japan, Korea, China, North. India.

(137) **P. peltatum**, v. **A. v. R.**; *Pleopeltis peltata*, *Scort.*, MS.; *Pl. superficialis*, *Bedd.*, var. *latifrons*, *Bedd.*, *Journ. of Bot.*, XXXI, 226.

Near *P. superficiale*, *Bl.* but the scales of the rhizome roundish, \pm adpressed, black with a brown edge. Stipes shorter; fronds shorter and broader.

Malacca.

* (138) **P. antrophyoides**, v. **A. v. R.**, *Bull. Dép. Agr. I. N.*, 1908, XVIII, 22.

Rhizome short-creeping, the scales brownish, linear-lanceolate, entire or slightly toothed. Fronds sessile or short-stalked, 8—15 c.M. long, the upper $\frac{1}{2}$ — $\frac{2}{3}$ ovate-lanceolate, 3—5 c.M. broad, the apex short-acuminate, the edge entire, recurved, the base narrowed into the narrow, linear-cuneate, decurrent lower part the frond. Texture coriaceous; surfaces naked; costa prominent in the lower part, becoming gradually finer and indistinct upwards, nearly wanting towards the apex; veins hidden; areolae very irregular, numerous, with copious free veinlets. Sori scattered irregularly, moderately large.

Sumatra (Forbes, No. 3119).

* (139) **P. curtidens**, *Christ*, *Ann. Botz.*, XX, 122.

Rhizome short-creeping, the scales not dense, lanceolate-subulate, firm, brown. Stipes short, to \pm 3 c.M. long, tufted, slender. Fronds \pm 15—20 c.M. long, \pm 1 $\frac{1}{2}$ c.M. broad, lanceolate, acute, the edge slightly undulate and broadly crenate with shallow, tooth-like, decumbent crenations, the base narrowed very gradually into the stipe. Texture coriaceous but not very thick; surfaces naked; costa prominent; no distinct main veins; areolae irregular, with few free veinlets. Sori numerous, small, placed in 3—4 irregular rows on each side, leaving a space from the margin, hardly immersed.

Borneo.

* (140) **P. Bakeri**, *Lueres.*, *Abh. Nat. Ver. Brem.*, VII, 48, *P. torulosum*, *Bk.*, *Journ. of Bot.*, XVIII, 215 (not *Journ. of Linn. Soc., Bot.*, XVI, 204).

Rhizome slender, wide-creeping, densely clothed with linear, acuminate, ciliate, reddish-brown scales. Fronds very dimorphous; sterile fronds on stiffly erect, naked stipes 2 $\frac{1}{2}$ —4 c.M. long, the lamina simple, oblong or linear-oblong, entire or slightly and broadly crenate, blunt or acute, 2 $\frac{1}{2}$ —7 c.M. long, 1 $\frac{1}{2}$ —2 $\frac{1}{2}$ c.M. broad at or below the middle, the base cuneate. Texture rigidly coriaceous; surfaces naked; veins immersed, obscure; no mains veins; areolae copious, small, irregular, with free veinlets. Fertile fronds on much

longer stipes (10—15 c.M. long), the lamina 10—15 c.M. long, reduced to a mere midrib, from which the large, round or oblong sori bulge out at intervals on each side.

Sumatra.

* (141) **P. Wrayi**, *Bk.*, Journ. of Bot., XXV, 206.

Rhizome slender, wide-creeping, the scales dense, lanceolate, acuminate, membranaceous, ferruginous, ciliate. Stipes of barren fronds $\pm 1\frac{1}{2}$ —4 c.M. long. Barren fronds oblanceolate, $2\frac{1}{2}$ —5 c.M. long, 6—9 m.M. broad, obtuse. Texture thick, coriaceous; surfaces naked like the stipe; veins reticulate, immersed, obscure. Fertile fronds ± 5 —13 c.M. long, 3—5 m.M. broad, linear, obtuse, with a longer stipe. Sori round, superficial, in 1 row on each side of the midrib; receptacle round or oblong.

Malacca, Sumatra.

* (142) **P. Beccarii**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 22.

Rhizome creeping, firm, the scales lanceolate-oblong, with the apex acute and the base rounded or acute. Stipes scattered, 1—5 c.M. long, erect, naked. Sterile fronds ± 6 —10 c.M. long, 7—8 m.M. broad, linear-lanceolate, narrowed gradually towards both ends, the apex more or less blunt, the edge indistinctly crenate. Texture coriaceous; surfaces naked; costa prominent beneath; no main veins; areolae hidden, in ± 1 —2 rows on each side, with free veinlets. Fertile fronds on the longest stalks, linear-lanceolate, 12—15 c.M. long, 5—7 m.M. broad, narrowed gradually towards both ends. Sori in 1 row on each side, medial or nearer the midrib than the edge, distant, superficial, occupying the upper $\frac{1}{2}$ — $\frac{2}{3}$ of the frond; receptacle \pm oblong. — Baker determined this plant as *P. congener*, *Hk.* (Journ. of Bot., XVIII, 215), from which it differs by the quite naked fronds and the anastomosing veins.

Sumatra (Beccari, No. 442).

* (143a) **P. accedens**, *Bl.*, Rac, Flor. Btz., I, 103; *Pleopeltis accedens*, *Moore*, *Bedd.*, Ferns Br.I., II, tab. CCXV.

Rhizome slender, wide-creeping, sparingly clothed with narrow, lanceolate, ciliate scales. Stipes none or to 1 c.M. long, $1\frac{1}{2}$ —3 c.M. apart. Fronds dimorphous. Barren fronds oblong to lanceolate, $2\frac{1}{2}$ —8 c.M. long, $1\frac{1}{2}$ —3 c.M. broad, the apex bluntish, the base cuneate or rounded, the edge entire, not thickened. Texture coriaceous; surfaces naked or sparingly covered with scattered, deciduous, stellate scales; no main veins; veins hidden; areolae

irregular; free veinlets often directed towards the midrib. Fertile fronds narrower and longer than the barren ones, the upper part soriferous, narrowed gradually or linearly. Sori in 1 row on each side of the costa, superficial.

Malaya; Polynesia.

(143b) **P. damunense**, *Rst.*, Fed., Rep., V, 42.

Rhizome wide-creeping, filiform, sparingly clothed with small, thin, lanceolate, ciliate scales. Fronds sessile, ± 1 c.M. apart, dimorphous. Barren fronds roundish-ovate or ovate-lanceolate, $1-1\frac{1}{2}$ c.M. long, $\pm \frac{3}{4}$ c.M. broad, cuneate at the base. Texture subcoriaceous; surfaces glabrescent; costa prominent on both sides; areolae immersed, in 2 rows between the costa and edge. Fertile fronds narrowly linear or more generally elongate-ovate, narrowed at the apex. Sori round, in 2 rows, occupying the narrowed apex of the fronds, intermixed with peltate scales.

New Guinea.

* (144) **P. nummularium**, *Mett.*, Hk. Bk., Syn. Fil., 340; *Mett.*, *Farn.*, I, Polypod. tab. I, fig. 44-45; *P. pyrolifolium*, *Goldm.*, C. Chr., Ind. Fil.; *Marginaria nummularia* *Pr.*, (oldest name).

Rhizome creeping, slender, clothed with linear-lanceolate, acuminate, ferruginous, subadpressed scales. Fronds distant, the barren ones mostly short-stalked, suborbicular, oval or ovate, $\pm 1-4$ c.M. long, $\pm 1-2$ c.M. broad when fully developed, the edge entire or shallowly and broadly crenate. Texture coriaceous; surfaces naked; veins immersed, obscure, forming 3-4 rows of areolae on each side, with free veinlets. Fertile fronds linear, 7-10 c.M. long, $\pm 5-4$ m.M. broad, narrowed gradually towards both ends, on stipes nearly as long as the fronds, the edge repand, the stipes slender, the veins forming 1-2 rows of longitudinally elongated areolae. Sori in 1 row on each side near the costa, 1 to each lobe, superficial, but the position marked on the upper surface of the frond.

Philippines.

(145) **P. hammatisorum**, *Harr.*, Copel., Polypod. Philipp., 125; (? *P. nummularium*, *Mett.*).

Rhizome long, slender, the scales thin, membranaceous, long, lanceolate, light-brown. Stipes scattered, $\frac{1}{2}-4$ c.M. long, slender, erect, naked. Sterile fronds elliptical, 1-4 c.M. long, $\frac{1}{2}-2\frac{1}{2}$ c.M. broad, obtuse, coarsely and shallowly crenate except at the narrowed base. Texture coriaceous; surfaces

naked; veins immersed; midrib distinct. Fertile fronds linear, 5—13 c.M. long, \pm 5 m.M. broad, on stipes $2\frac{1}{2}$ —8 c.M. long, with a lobe under each sorus. Sori \pm 20 in a row on each side of the midrib, large, giving the frond a knotted appearance.

Philippines.

* (146) **P. stenopteris**, *Bk.*, Journ. of Bot., XVII, 43; Christ, Ann. Btz., XX, tab. III, fig. 4.

Rhizome wide-creeping; scales dense, squarrose, brown, the base roundish, peltate, the point long, fine, linear. Stipes nearly none or to 4 c.M. long, the fertile fronds on the longest stipes. Fronds dimorphous. Barren fronds lanceolate-spathulate, 3—5 c.M. long, $\frac{1}{2}$ —1 c.M. broad, the apex bluntly rounded, the edge entire, thickened, the base gradually narrowed, \pm cuneate. Texture very rigidly coriaceous; costa conspicuous; veins hidden. Fertile fronds 20—30 c.M. long, the lower half narrowly lanceolate, the edge entire, the base cuneately narrowed, the apex narrowed gradually into the narrow, linear, repand upper half of the frond. Sori immersed, rather distant, confined to the linear upper half of the frond, placed in the projections of the waves, 1 to each lobe, prominent on the upper surface.

Borneo.

* (147) **P. soridens**, *Hk.*, Hk. Bk., Syn. Fil., 355; Hk., Spec. Fil., V, tab. CCLXXXIII, B.

Rhizome wide-creeping; scales squarrose, the base roundish, peltate, the apex long, fine, linear. Stipes $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. long, slender, naked, scattered. Fronds linear-lanceolate, 10—30 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, narrowed very gradually to both ends, the barren ones with the edge entire or repand; fertile fronds toothed with the teeth distant and rounded, or deeply and very irregularly incised with the lobes spreading, oblong, bluntly rounded, 1—50 m.M. apart, the sinuses broadly rounded. Texture rigid, coriaceous; surfaces naked; no main veins; areolae fine; free veinlets rare. Sori submarginal, terminal on the lobes, 1 to each tooth or lobe, deeply immersed, forming very distinct papillae on the upper surface.

Borneo.

* (148) **P. neglectum**, *Bl.*, Flor. Jav., II, 135, tab. LIV, fig. 1.

Rhizome moderately long, creeping, dirty-brown; scales dense, small, lanceolate-subulate, recurved, the younger ones reddish-brown, ereso-ciliate, villo-

se near the base, the others blackish-brown, adpressed, naked. Fronds subdimorphous, the barren ones on stalks under 8 m.M. long, the fertile on stalks $\pm 2\frac{1}{2}$ c.M. long. Barren fronds distant, 8—25 m.M. long, 6—15 m.M. broad, obovate, the apex rounded, the edge broadly crenate, slightly reflexed, the base narrowed suddenly. Texture coriaceous; surfaces naked; costa distinct beneath in the lower half. Fertile fronds lanceolate-spathulate, ± 2 c.M. long, 5—8 m.M. broad, the apex rounded, the base cuneate. Sori in 1 row on each side, ± 3 —5 to a row, occupying the upper $\frac{1}{2}$ — $\frac{3}{4}$ of the frond.

Java, Luzon.

(149) **P. rudimentum**, *Copel.*, Polypod. Philipp., 125; (*P. oodes*, *Kze*, *Copel.*, Philipp. Journ., III, 35).

Rhizome wide-creeping, slender, the scales linear-subulate, brownish. Stipes $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, erect, filiform, naked, stramineous. Fronds orbicular-ovate, the fertile ones rather narrower, 1—2 c.M. long, rounded at both ends, entire or finely serrate. Texture coriaceous; surfaces naked; costa evanishing near the apex; veins obscure, anastomosing irregularly, or sometimes free but hamate. Sori few, subimmersed, round, in 1 row on each side, \pm medial between the costa and margin.

Philippines.

(150) **P. subecostatum**, *Hk.*, *Hk. Bk.*, Syn. Fil., 353; *Hk.*, Spec. Fil., V, tab. CCLXXXIII, A.

Rhizome creeping, the scales brown, fibrillose, squarrose. Fronds sessile, 10—20 c.M. long, 1— $1\frac{1}{2}$ c.M. broad, narrowed gradually to both ends, the edge entire. Texture thick, coriaceous; colour pale-green; surfaces naked; midrib upwards indistinct; no main veins; areolae obscure. Sori in a single submarginal row.

Borneo.

* (151) **P. rostratum**, *Hk.*, *Hk. Bk.*, Syn. Fil., 353; *Hk.*, Icon. Pl., X, tab. CMLIII; *P. subrostratum*, *C. Chr.*, Ind. Fil.; *Pleopeltis rostrata*, *Bedd.*, Ferns Br.I., II, tab. CLIX; — var. *trifurcatum*, *Christ*, Bull. Herb. Boiss., 2^e Série, I, 449.

Rhizome filiform, wide-creeping, the scales small, fibrillose. Stipes $2\frac{1}{2}$ —5 c.M. long, firm, naked. Fronds 5—10 c.M. long, 1— $2\frac{1}{2}$ c.M. broad, narrowed gradually to both ends, the edge entire, the apex rostrate. Texture

coriaceous; surfaces naked; no distinct main veins; veins immersed; areolae fine, with free veinlets. Sori large, in single rows not far from the midrib.

• **Var. trifurcatum**: Fronds 2—3-furcate, with acute lobes.

Luzon; North. India, South. China; — var. *trifurcatum*: *New Guinea*.

* (152) **P. lineare**, *Thb.*, Hk. Bk., Syn. Fil., 354; *P. loriforme*, *Hk.*, Gard. Ferns, tab. XIV; *P. contiguum*, *Wall.*, List, No. 285; *Pleopeltis nuda*, *Hk.*, Bau. & Hk., Gen. Fil., tab. XVIII; *Pl. Wightiana*, *Bedd.*, Ferns S.I., tab. CLXXX; *Pl. linearis*, *Moore*, *Bedd.*, Handb. Ind. Ferns, 346.

Rhizome wide-creeping, woody, the scales lanceolate, dull-brown. Stipes varying from \pm none to 5 c.M. long. Fronds 10—30 c.M. or more long, 1—5 c.M. broad, entire, narrowed gradually to both ends. Texture subcoriaceous; surfaces nearly naked; areolae fine, with copious free veinlets. Sori hardly immersed, large, \pm oblong, in 1 row on each side, medial or nearer the midrib than the edge, distant, covered with copious peltate scales when young.

Malaya; Hawaii, Japan, China, North. India, Ceylon, South. Africa.

* (153) **P. Forbesii**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 23.

Rhizome wide-creeping, firm, the scales dense, strongly adpressed, ovate or oblong, rounded or acute at both ends, peltate, brown, with a blackish centre. Stipes distant, 5 c.M. or more apart, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, erect, firm, naked, not very slender (with regard to the rhizome). Fronds broadly lanceolate, 10—12 c.M. long, $2\frac{1}{2}$ —4 c.M. broad below the middle, narrowed gradually to an acuminate apex and base. Texture coriaceous, not thick; surfaces naked; veins fine, irregularly anastomosing; areolae with free veinlets. Sori moderately large, distant, in 2 rows on each side of the midrib, the outer row medial between the midrib and edge, the apex and base of the fronds destitute of sori.

Sumatra (Forbes, No. 2333).

(154) **P. induratum**, *Bk.*, Ann. of Bot., V, 475.

Rhizome slender, wide-creeping. Stipes wiry, naked, 7—10 c.M. long. Fronds simple, linear, 20—25 c.M. long, 1 — $1\frac{1}{2}$ c.M. broad at the middle, narrowed gradually to both ends. Texture rigidly coriaceous; surfaces naked; veins quite hidden. Sori semiglobose, superficial, forming a single lax row

on each side of the costa, with a few others forming an outer row at the middle of the frond.

New Guinea.

* (155) **P. stenophyllum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 104; *Bl. Flor. Jav.* II, 134, tab. LV, fig. 1, a—b; *Pleopeltis stenophylla*, *Moore*, *Bedd.*, *Ferns Br.I*, II, tab. CCXXXIV.

Rhizome wide-creeping, the scales dense, lanceolate-subulate, acuminate, entire, brownish. Stipes $\frac{1}{2}$ —5 c.M. apart, 1—2 $\frac{1}{2}$ c.M. or more long, firm, erect. Fronds 8—25 c.M. long, 5—15 c.M. broad, narrowed gradually to both ends, the apex varying from acute to rounded, the edge entire. Texture very coriaceous; surfaces naked; no distinct main veins; veins hidden; areolae with few free veinlets. Sori deeply immersed, very prominent on the upper surface, placed in 1 row on each side near the edge. — The form with the bluntly rounded fronds is *var. obtusum*, *Christ* (*Ann. Btz.*, XV, 153).

Malaya; Fiji.

* (156a) **P. revolutum**, *C. Chr.*, *Ind. Fil.*; *P. longifolium*, *Mett.*, *Rac.*, *Flor. Btz.*, I, 104; *P. contiguum*, *Hk.* (not *Wall.*), *Fil. Exot.*, tab. XX; *Icon. Pl.*, X, tab. CMLXXXVII; *Pleopeltis longifolia*, *Bedd.*, *Ferns Br.I*, I, tab. VII; *Grammitis longifolia*, *Bl.*, *Enum.*, 119; *Drynaria revoluta*, *J. Sm.*, *Journ. of Bot.*, III, 421.

Rhizome creeping, woody, the scales linear-lanceolate, brown or black, not quite spreading. Stipes short, firm. Fronds 30—70 c.M. long, 1—4 c.M. broad, \pm ligulate, narrowed gradually to both ends, the apex blunt or bluntish, the edge entire, often revolute. Texture coriaceous; under surface naked or slightly furfuraceous; no main veins; areolae fine, hidden, with free veinlets. Sori oblong, 2—5 \times as long as broad, immersed, parallel to the edge, confined to the upper $\frac{1}{3}$ — $\frac{2}{3}$ part of the frond, placed in a line nearly close to the edge; young capsules mixed with small peltate scales. — The young plants have the texture thin, membranaceous to herbaceous, the veins conspicuous. A form found in Celebes has the sori round, large.

Malaya; New Caledonia.

(156b) **P. productum**, *Christ*, *Philipp. Journ.*, II, 178.

Rhizome short-creeping, covered with small, scattered, setiform scales. Stipes approximate, \pm 10 c.M. long, stramineous, naked. Fronds linear, \pm 63 c.M. long, \pm 1—1 $\frac{1}{4}$ c.M. broad, the apex acutely long-acuminate, the

base narrowed gradually into the stipe, the edge narrowly revolute. Texture rigidly coriaceous; surfaces naked; colour yellow-greenish; costa prominent, stramineous; veins hidden. Sori oblong, rather large, submarginal, impressed, with a space less than their own length between them, the higher ones protruded beyond the margin.

Mindanao.

(157) **P. subgeminatum**, *Christ*, Schum. & Laut., Flor. deut. Sch.geb., Suppl., 47.

Rhizome wide-creeping, very strong, the scales short, furfuraceous. Fronds subsessile, usually in distant pairs, uniform, entire, $\pm 7\frac{1}{2}$ —15 c.M. long, 1—2 $\frac{1}{2}$ c.M. broad, narrowed towards both ends, the point often blunt. Texture coriaceous; surfaces naked; areolae copious, small, with numerous included veinlets. Sori large, in 1 row on each side, medial between the midrib and edge, much immersed.

New Guinea.

(158) **P. papuanum**, *Blk.*, Becc., Mal., III, 48.

Rhizome wide-creeping. Stipes strong, rigid, naked, erect, 2 $\frac{1}{2}$ —5 c.M. long. Fronds simple, lanceolate-oblong, ± 15 c.M. long, 2 $\frac{1}{2}$ —4 c.M. broad, the base cuneate, the edge entire, the apex acuminate. Texture firm; surfaces naked; veins immersed, obscure. Sori oblong, in 1 row on each side of the midrib, immersed in the substance of the frond.

New Guinea.

(159) **P. Annabellae**, *Forb.*, Journ., of Bot., XXVI, 33, tab. CCLXXX; ? *P. cyclobasis*, *Blk.*, Kew Bull., 1896, 42.

Rhizome creeping, slender, clothed with scattered, brown, membranaceous, lanceolate, acuminate scales. Fronds sessile, remote, imbricating, simple. Barren fronds suborbicular, entire, cordate at the base. Texture membranaceous; veins anastomosing irregularly; areolae large, enclosing smaller ones with few free veinlets; costa disappearing \pm towards the centre of the frond. Fertile fronds similar but the apex suddenly produced into a long, narrow, linear-lanceolate fertile part several times longer but much narrower than the barren part, with the costa distinct quite to the point. Sori \pm oblong, in 1—2 rows on each side of the costa.

New Guinea.

(160) **P. lycopodioides**, L., Hk. Bk., Syn. Fil., 357; Christ, Farnkr. d. Erde, fig. 280; *P. lagopodioides*, Desv., Berl. Mag., V, 313.

Rhizome wide-creeping, the scales lanceolate-subulate, imbricate, reddish when young, at length ferruginous or decolored. Stipes none or short, distant. Fronds dimorphous. Barren fronds 5—10 c.M. long, 1—3 c.M. broad, ovate-oblong, the apex commonly bluntly rounded, the base narrowed, the edge entire. Texture coriaceous; surfaces naked or the costa slightly scaly; veins moderately visible, forming on each side a first row of narrow, longitudinal costal areolae, a second row of large transversal areolae with included, free or anastomosing veinlets, which are directed towards the edge, and near the latter 1 or more rows of small areolae with or without free veinlets. Fertile fronds longer and narrower than the barren ones, narrowed gradually below. Sori large, in 1 row on each side, solitary in the largest areolae (those of the 2nd row), on the included veinlets.

Java; Hawaii, Trop. America & Africa, Mascarenes.

* (161) **P. Raapii**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 23, tab. IV, fig. 1, a—b.

Rhizome creeping, the scales linear-lanceolate, deciduous. Stipes scattered, 1—5 c.M. long, naked. Fronds lanceolate, \pm 10—12 c.M. long, $\frac{3}{4}$ —1 $\frac{1}{2}$ c.M. broad at or a little below the middle, the fertile ones the narrowest, narrowed gradually towards both ends, the apex blunt, acute or acuminate, the edge entire. Texture thinly herbaceous or membranaceous; surfaces naked; costa distinct; veins visible, forming 1—5 rows of irregular areolae on each side of the costa; areolae with free veinlets. Sori distant, partially round, partially oblong and oblique, placed in 1 row on each side nearer the margin than the midrib, sometimes accompanied by some few (1—5) scattered ones. — In swamps and ditches.

Another (submerged) water-fern, found in Lombok by Dr. Ernst, has the stalks shorter, the fronds a little broader and slightly scaly on the costa beneath, and the sori placed in 1—2 rows nearer the midrib than the edge, whether or not accompanied by a few others placed in a 5th, outer row (v. A. v. R., l. c., tab. IV, fig. 2, a—b).

Batu Islands (Raap, No. 671).

(162) **P. zosteriforme**, Wall., Hk. Bk., Syn. Fil., 357; Mett., Farnng., I, Polypod., tab. I, fig. 26—27; *Pleopeltis zosteriformis*, Bedd., Ferns Br.I., I, tab. CXXIII.

Rhizome wide-creeping, slender, scaleless. Stipes 1—2½ c.M. long, slender. Fronds 10—15 c.M. long, ± 5 m.M. broad, the apex scarcely acuminate, the edge entire, the base gradually narrowed. Texture herbaceous; colour dark-green; surfaces naked; no main veins; veins distinct; areolae in 2—3 irregular rows, without free veinlets. Sori distant, round or ± oblong, in 1 regular or irregular row on each side nearer the midrib than the edge.

Malacca.

(163) **P. normale**, *Don.*, Hk. Bk., Syn. Fil., 558; *Pleopeltis normalis*, *Moore*, *Bedd.*, *Ferns Br.I.*, I, tab. X; — var. *longifrons*; *Pol. longifrons*, *Wall.*, *List*, No. 274; *Hk. & Grev.*, *Ic. Fil.*, tab. LXV.

Rhizome woody, scandent, clothed with deciduous, fibrillose, ovate or lanceolate, almost hair-pointed scales. Stipes approximate, 2½—8 c.M. long, stramineous or reddish. Fronds linear-lanceolate, 50—60 c.M. long, 2½—5 c.M. broad, narrowed gradually towards both ends, the point acute, the edge entire or slightly wavy. Texture membranaceous to slightly coriaceous; surfaces naked, or a few scattered, closely adpressed scales present on the stipes and near the base of the costa beneath; no main veins; areolae moderately large, with free veinlets. Sori large, superficial, in 1 regular or more commonly 1—2 irregular, ± wavy rows on each side of the midrib. — Near *P. lineare*, *Thb.*, but the capsules not intermixed with peltate scales.

Var. longifrons: Sori copious, scattered irregularly.

Trop. Asia, China, Madagascar.

* (164a) **P. myriocarpum**, *Mett.*, Hk. Bk., Syn. Fil., 558; *Mett.*, *Farngr.*, I, *Polypod.*, tab. I, fig. 38; *Phymatodes myriocarpa*, *Pr.*, *Tent. Pterid.*, tab. VIII, fig. 12.

Rhizome firm, woody, the scales subadpressed, ovate or lanceolate, peltate, blackish, the edge brownish, toothed, erose or slightly fimbriate. Stipes distant, 1—5 c.M. long. Fronds lanceolate, 15—90 c.M. long, 2½—10 c.M. broad, acuminate, entire, narrowed gradually to the base. Texture papyraceo-herbaceous; surfaces naked or the under side pubescent; primary veins forming a series of large, costal primary areolae, a second series of stout veins forms smaller but still very conspicuous primary areolae outside the costal ones, both including numerous finer ones resembling those between the second series and the margin, with included free veinlets. Sori copious, scattered irregularly, sometimes confluent.

New Guinea, Amboyna, Borneo, Philippines; Cochinchina.

FILICES.

(164b) **P. sablanianum**, *Christ*, Philipp. Journ., II, 177.

Near *P. myriocarpum*, *Mett.*, but: Rhizome scandent, brown, clothed with stiff, subulate, dark-brown scales. Fronds to ± 75 c.M. long, to $\pm 5\frac{1}{2}$ c.M. broad, the apex long-caudate, the margin undulate, the base long-decurrent. Texture membranaceous; colour bright-green; costa light-yellow; venation slender, forming a single row of very large, square primary areolae and another row of very narrow, obscure ones.

Luzon.

(165) **P. glossipes**, *Bk.*, Ann. of Bot., V, 476.

Rhizome wide-creeping, clothed with few, small, squarrose, linear, brown scales. Stipes $3\frac{1}{2}$ —4 c.M. long. Fronds lanceolate-oblong, acute or obtuse, narrowed very gradually at the base, 15—20 c.M. long, $3\frac{1}{2}$ — $6\frac{1}{2}$ c.M. broad. Texture chartaceous; surfaces naked; venation as in *P. myriocarpum*, *Mett.* Sori globose, superficial, scattered all over the frond.

New Guinea; Cambodja.

(166) **P. linguaeforme**, *Mett.*, Hk. Bk., Syn. Fil., 558.

Rhizome stout, short-creeping. Fronds sessile, 20—45 c.M. long, 6—8 c.M. broad, tongue-shaped, narrowed from the centre to $\frac{3}{4}$ of the way down, and then dilated again to a broadly-rounded base, the edge entire. Texture papyraceous; both surfaces naked; veins very distinctly prominent, the main ones irregular; areolae subquadrangular, with a single, distinctly immersed sorus on free or joined veinlets in the centre of each.

Luzon, East. Malaya; Polynesia, Melanesia.

§ 6. PLEOPELTIS. b. **Atactosia**. ** **Pleuridium**.

a. Sori in 1 row on each side of the midrib, solitary between the main veins, sometimes confluent.

b. Sori superficial, round, rarely oblong.

c. Barren fronds distinctly shorter than the fertile ones, and much broader, under 10 c.M. long.

d. Fertile fronds under 6 m.M. broad. (167) *P. costulatum*.

d.d. Fertile fronds broader, 7—20 m.M. broad.

e. Rhizome scales lanceolate, whitish. (168) *P. Hellwigii*.

e.e. Rhizome scales fibrillose, ferruginous.

(169) *P. rynchophyllum*.

c.c. Barren fronds hardly shorter than the fertile ones, more than 10 c.M. long. (170) *P. Griffithianum*.

b.b. Sori \pm immersed, round. (171) *P. Whitfordi*.

b.b.b. Sori \pm immersed, oblong.

*. Sori oblique to the costa. (172a) *P. Treubii*.

.. Sori parallel to the costa. (172b) *P. Valetonianum*.

a.a. Sori in 1 row between the main veins, free or confluent. ⁽¹⁾

1. Upper surface of the fronds copiously covered with white dots.

(175) *P. platyphyllum*.

1.1. Upper surface of the fronds without white dots.

2. Texture membranaceous. (174) *P. campyloneuroides*.

2.2. Texture rather coriaceous.

3. Fronds small, under $12\frac{1}{2}$ c.M. long. Main veins erecto-patent. (175) *P. phlebiscopum*.

3.3. Fronds larger, more than $12\frac{1}{2}$ c.M. long.

*. Main veins \pm horizontal. (176a) *P. hemionitideum*.

.. Main veins oblique. (176b) *P. Werneri*.

a.a.a. Sori in 2 rows between the main veins, 1 to several in a row.

b. Texture membranaceous or herbaceous to papyraceous.

*. Stipes distant. (177) *P. Zippelii*.

.. Stipes approximate. (178) *P. membranaceum*.

b'.b'. Texture chartaceous to coriaceous.

c. Upper surface of the fronds covered with white dots.

(179) *P. leucophorum*.

c'.c'. Upper surface of the fronds without white dots.

d. Rhizome scales blunt at the apex, pale-ochraceous.

(180) *P. triquetrum*.

d'.d'. Rhizome scales acuminate or aristate at the apex.

e. Rhizome scales dark-brown, sometimes white.

f. Sori 4—8 in a row. (181a) *P. rupestre*.

f'.f'. Sori 1—3 in a row. (181b) *P. luzonicum*.

e'.e'. Rhizome scales pale-coloured, yellowish. (182) *P. albulum*.

a.a.a.a. Sori scattered irregularly, free or sometimes confluent. ⁽²⁾

⁽¹⁾ See also § 7, **Selliguea**.

⁽²⁾ See also **P. membranaceum**, *Don*, **Zippelii**, *Bl.* & **leucophorum**, *Bl.*

- 1.** *Fronde small, under 15 c.M. long.*
- *. *Rhizome scales fibrillose, yellowish. Sori rather large, superficial.*
(183) P. oodes.
- *.*. *Rhizome scales lanceolate, pale-brown. Sori rather small, subimmersed.*
(184) P. dulitense.
- 1.1.** *Fully developed fronds more than 15 c.M. long.*
- 2.** *Texture membranaceous or herbaceous.*
- §. *Rhizome naked.*
- *. *Fronde short-stalked.* (185a) P. suboppositum.
- *.*. *Fronde long-stalked.* (185b) P. Scortechinii.
- §.§. *Rhizome scaly.*
- ×. *Sori irregularly 1—3-serial.* (186a) P. ovatum.
- ×.×. *Sori numerous.* (186b) P. Zollingerianum.
- 2.2.** *Texture chartaceous to coriaceous.*
- 3.** *Fronde commonly $\pm 2\frac{1}{2}$ — $6\frac{1}{2}$ c.M. broad. (*)*
- 4.** *Stipes stout, narrowly winged by the gradually decurrent base of the fronds.* (187) P. millisorum.
- 4.4.** *Stipes short or wanting, not winged. Fronde \pm blunt at the base.* (188) P. punctatum.
- 3.3.** *Fronde commonly broader, $\pm 6\frac{1}{2}$ —20 c.M. broad.*
- §. *Fronde cordate or rounded at the base.*
(189) P. musifolium.
- §.§. *Fronde gradually narrowed at the base.*
- *. *Rhizome short, thick.* (190a) P. validum.
- *.*. *Rhizome scandent, slender.*
(190b) P. monstrosum.

* (167) **P. costulatum**, *Bk.*, Journ. of Bot., XVIII, 215; *Acrostichum costulatum*, *Ces.*, Fel. Becc. Polin., 5 & 8; *Drymoglossum rigidum*, *Hk.*, var. *longistipitatum*, Christ, MS. in Herb. Bog.

Rhizome wide-creeping, slender, woody; scales erecto-patent, lanceolate-subulate, brownish or bleached, the base roundish, dilated, peltate. Stipes $2\frac{1}{2}$ —15 c.M. long, those of the fertile fronds the longest, scattered, naked, erect. Fronde dimorphous. Barren fronds ovate or deltoid-oblong, obtuse,

(*) See also **P. monstrosum**, *Copel.*

2½—7 c.M. long, 2—4 c.M. broad, the base broadly cuneate or subtruncate, the edge thickened, broadly and shallowly crenate. Texture rigidly coriaceous; surfaces naked; main veins prominent beneath, distinct and straight nearly to the edge, erecto-patent or the lower spreading; veins quite hidden. Fertile fronds linear, 6—10 c.M. long, 3—5 m.M. broad, narrowed gradually towards both ends; midrib very prominent beneath; edges recurved. Sori ± contiguous, round or oblong, sometimes confluent, medial, in 1 row on each side parallel to the costa, occupying the space between the midrib and edge.

Sumatra, Borneo, New Guinea.

(168) **P. Hellwigii**, *Diels*, Schum. & Laut., Flor. deut. Sch.geb., 140, tab. III, A—B.

Rhizome wide-creeping, slender, the scales copious, large, whitish, lanceolate, long-acuminate. Stipes 1—4 c.M. long, the fertile fronds on the longest stipes. Fronds dimorphous. Barren fronds subobovate, 1½—2½ c.M. long, ± 1¼—1½ c.M. broad, the apex blunt, the base cuneate. Texture coriaceous; the margin entire, subrevolute; surfaces naked; costa prominent; main veins erecto-patent, ± 4 on each side, prominent; areolae hidden. Fertile fronds elongate, 4—5 c.M. long, ± ¾—1 c.M. broad; main veins ± 8—9 on each side. Sori medial, in 1 row on each side of the costa.

New Guinea.

* (169) **P. rhynchophyllum**, *Hk.*, Hk. Bk., Syn. Fil., 359; Hk., Icon. Pl., X, tab. CMLIV; *Pleopeltis rhynchophylla*, *Moore*, Bedd., Ferns Br.I., I, tab. IX.

Rhizome firm, wide-creeping, the scales copious, peltate, lanceolate, acuminate, ferruginous. Stipes 2½—10 c.M. long, the fertile fronds on the longest stipes. Fronds dimorphous. Barren fronds ovate, 2½—4 c.M. long. Texture coriaceous; surfaces naked; main veins distinct to the edge; areolae copious, fine, with free veinlets. Fertile fronds 7—15 c.M. long, 1—2 c.M. broad, narrowed very gradually upwards, the base cuneate, the edge obscurely toothed or rotundato-crenate. Sori 1-serial on each side, confined to the upper ¼—½ of the frond.

Philippines; North. India.

* (170) **P. Griffithianum**, *Hk.*, Hk. Bk., Syn. Fil., 359; Hk., Icon. Pl., X, tab. CMLI; *Pleopeltis Griffithiana*, *Moore*, Bedd. Ferns. Br.I., II, tab. CLVIII; — var. *borneense*, *Christ*, Ann. Btz., XX, 122.

Rhizome wide-creeping, the scales dense, linear, squarrose, dull-brown. Stipes distant, 7—15 c.M. long, firm, erect. Fronds 15—26 c.M. long, 5½—5 c.M. broad, the point acute or acuminate, the edge repand, the base rounded or rather narrowed and auricled. Texture coriaceous; surfaces naked; main veins very distinct to the edge; areolae fine, hidden, with copious free veinlets. Sori large, in 1 row on each side near the midrib, solitary between the main veins.

Var. borneense: Sori more distant from the midrib, smaller, more numerous; rhizome scales linear-lanceolate, pale-brown.

South. China, North. India; — var. borneense: *Borneo.*

(171) **P. Whitfordi**, *Copel.*, Philipp. Journ., I, Suppl., 256, tab. IV, B.

Rhizome creeping, the scales lanceolate, apiculate, persistent, pale-ferruginous when young. Stipes distant, filiform, 1—4½ c.M. long, the fertile fronds on the longest stipes. Fronds dimorphous. Barren fronds 1—3½ c.M. long, the smallest ones orbicular, the largest oblong, obtuse, the margin cartilagineous, broadly and shallowly crenate. Texture coriaceous; surfaces naked; costa black beneath; main veins straight, reaching nearly to the margin; areolae quadrangular, in ± 3—4 rows on each side, with few included, excurrent free veinlets. Fertile fronds 20—60 c.M. long, 6—9 m.M. broad, the smallest ones entire, obtuse, the largest crenate, acute, cuneate at the base. Sori in 1 row on each side, medial between the costa and margin, sub-immersed, prominent on the upper surface, not restricted to the highest part of the frond.

Luzon.

* (172a) **P. Treubii**, *Christ.*, Ann. Bot., XX, 121; *P. dolichosorum*, *Christ.*, MS. in Herb. Bog.

Rhizome wide-creeping, slender; scales firm, the base roundish, peltate, the point long, fine, hair-like, spreading, reddish-brown. Stipes slender, firm, 5—9 c.M. long, naked. Fronds subdimorphous. Barren fronds lanceolate-ovate, 5—7 c.M. long, ± 1½—2 c.M. broad, the apex subacute, the margin entire, the base cuneate. Texture ± coriaceous; surfaces naked; main veins fine but visible, slightly prominent, oblique; areolae hidden, with free veinlets. Fertile fronds on the longest stipes, narrowly lanceolate, longer and narrower than the barren ones, to 11 c.M. long, to 1 c.M. broad; main veins hidden. Sori in 1 row on each side, to ± 20 to a row, oblong, medial between the mid-

rib and edge, parallel to the main veins, immersed, prominent on the upper surface. — Scarcely different from a small form of *P. heterocarpum*, *Mett.*
Borneo.

* (172b) ***P. Valetonianum***, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 25, tab. V.

Rhizome creeping, stout, green, at length naked, the scales thin-membranaceous, yellowish, lanceolate, acuminate, erect, slightly but distinctly crisped, with the base rounded and peltate. Stipes \pm remote, naked, \pm terete, firm, erect, reddish or brownish-green in fresh material, 5—15 c.M. long, the fertile fronds on the longest stipes. Fronds dimorphous. Barren fronds ovate-oblong or sublanceolate, \pm 9—11 c.M. long, \pm 4—5 c.M. broad below the middle, narrowed towards the blunt or bluntish apex, the edge thickened and shallowly crenate, the crenations broad, 1 to each main vein, straight or more commonly slightly convex, the base rounded or cuneate, slightly decurrent upon the stipe. Texture very coriaceous; surfaces naked; colour bright-green above, paler beneath in fresh material; costa prominent; main veins distinct, erecto-patent, not quite reaching the margin; areolae hidden, with free veinlets. Fertile fronds linear, \pm 15 c.M. long, \pm 1 c.M. broad below the middle, narrowed gradually towards the bluntish apex, the base suddenly cuneate; main veins horizontally spreading, except the lowest pair, which is distinctly erecto-patent. Sori large, medial, solitary between the main veins, oblong, perfectly parallel to the costa, the highest ones nearly round, the others gradually becoming longer, the lowest \pm 4—6 \times as long as broad, \pm $\frac{1}{2}$ c.M. long, all deeply immersed so as to form large, conspicuous papillae on the upper surface of the frond. — The species bears the name of Dr. Th. Valeton, Director of the Buitenzorg Herbarium.

Java (Preanger Regencies, v. A. v. R.).

* (175) ***P. platyphyllum***, *Sw.*, Rac., Flor. Btz., I, 104; *P. crassinervium*, *Bl.*, Flor. Jav., II, 145, tab. LXI; *Pleopeltis platyphylla*, *Bedd.*, Handb. Ind. Ferns, Suppl., 94.

Rhizome creeping, \pm woody, the scales dense, lanceolate, subulate-acuminate, dark-brown or blackish. Stipes 2—3 c.M. apart, 5—15 c.M. long, strong, erect, canaliculate. Fronds 15—50 c.M. long, 4—12 c.M. broad, narrowed to both ends, the point acute, the edge entire. Texture rigid, coriaceous, very thick and firm; both surfaces naked, the upper densely covered with small cretaceous dots; midrib prominent; main veins parallel, very distinct to

the edge, glossy. Sori immersed, in single rows between the main veins, 2—12 in a row, confined to the upper $\frac{1}{3}$ — $\frac{2}{3}$ of the frond or rarely occupying the whole lamina.

Java, Borneo, Malacca.

(174) **P. campyloneuroides**, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 229.

Rhizome slender, wide-creeping, the scales small, lanceolate, adpressed. Stipes erect, naked, 1—4 c.M. long. Fronds simple, 10—15 c.M. long, 1—2 c.M. broad, oblanceolate, obtuse, narrowed gradually from the middle to the base. Texture membranaceous; surfaces naked; main veins fine, parallel, erecto-patent, indistinct; transversal veins forming subquadrangular areolae. Sori in 1 row between the main veins, 3—4 in a row.

Borneo.

(175) **P. phlebiscopum**, *Bk.*, Journ. of Linn. Soc., Bot., XV, 110.

Rhizome woody, flexuose, short-creeping, the scales membranaceous, erect, lanceolate, acuminate, pale-brown, $\frac{1}{2}$ c.M. long. Stipes $2\frac{1}{2}$ —8 c.M. long, rigid, naked, stramineous, canaliculate. Fronds simple, oblong or ovate-oblong, entire, $3\frac{1}{2}$ —8 c.M. long, 2— $3\frac{1}{2}$ c.M. broad at the middle, blunt or subacute, deltoid at the base. Texture rigidly coriaceous; surfaces naked; main veins erecto-patent, distinct to the edge; areolae fine, immersed. Sori superficial, in single rows between the main veins, 5—4 in a row, often confluent and then forming an oblong mass 5—8 m.M. broad.

Ternate.

* (176a) **P. hemionitideum**, *Wall.*, Hk. Bk., Syn. Fil., 360; *Pleopeltis hemionitidea*, *Moore*, Bedd., Ferns S.I., tab. CLXXXII.

Rhizome creeping, the scales lanceolate. Stipes $2\frac{1}{2}$ —15 c.M. long, erect, firm, scaly. Fronds 20—45 c.M. long, 5—8 c.M. broad, narrowed towards both ends, the edge entire. Texture thinly coriaceous; surfaces naked; costa slightly scaly; veins distinctly prominent; main veins not reaching the edge, blackish, \pm horizontal, including rather large areolae; areolae in \pm 4 rows, subquadrangular, with many free veinlets. Sori on the anastomosing veins, moderately large, round or oblong, often confluent, in 1 row between the main veins, the rows rather irregular.

Malaya; North. India, South. China.

(176b) **P. Wernerii**, *Rst.*, Fed., Rep., V, 43.

Rhizome wide-creeping, whitish-glaucous, clothed with lanceolate-triangular, acuminate, entire, rufous-brown scales. Stipes slender, \pm 10 c.M. long, 5—10 c.M. apart, stramineous, naked. Barren fronds broadly ovate-lanceolate, \pm 18 c.M. long, \pm 6 c.M. broad near the base, the apex acuminate, the acumen \pm 4 c.M. long, narrow, the edge entire, the base cuneate or subrotund, slightly decurrent; fertile fronds rather similar but narrower, 3—4 c.M. broad. Texture chartaceous; surfaces naked; costa prominent beneath; main veins erecto-patent, nearly straight, prominent on both sides; areolae in 8—9 rows on each side of the costa, with many free veinlets. Sori in 1 row between the main veins, in pairs, subrotund or (principally the lower ones) oblique-oblong or rarely conduplicate, slightly immersed.

New Guinea.

* (177) **P. Zippelii**, *Bl.*, Rac., Flor. Btz., I, 105; *Bl.*, Flor. Jav., II, 172, tab. LXXX; *P. oxyphyllum*, *Kze.*, Bot. Zeit., 1848, 116; *Pleopeltis Zippelii*, *Moore*, Bedd., Handb. Ind. Ferns, 357; — var. *sparsisorum*.

Rhizome creeping, dark-coloured, the scales lanceolate, with the base peltate, the margin ciliate or shortly toothed, the point acute. Stipes distant, 1—10 c.M. long, firm, erect, naked. Fronds lanceolate, 25—70 c.M. long, 3—10 c.M. broad, the edge entire or repand, both ends narrowed, the apex acute, the base gradually decurrent upon the stipe. Texture herbaceous but firm; surfaces naked; costa conspicuous, sometimes with a few scales beneath in the lower part; main veins distinct nearly to the edge; areolae large, with included smaller ones, with numerous free veinlets. Sori distant, in 2 rows near the main veins, 2—4—6 in a row.

Var. sparsisorum: Sori scattered irregularly.

Malacca, Java, Borneo, Philippines; North. India

* (178) **P. membranaceum**, *Don*, Hk. Bk., Syn. Fil., 360; *Pleopeltis membranacea*, *Moore*, Bedd., Ferns S.I., tab. CLXXVII; — var. *grandifolium*; *P. grandifolium*, *Wall.*, List, No. 282.

Rhizome firm, the scales ovate, acuminate, spreading, membranaceous, lurid-green. Stipes 0—12½ c.M. long, firm, erect, approximate or subtufted. Fronds 30—90 c.M. long, 2—15 c.M. broad, lanceolate, narrowed towards both ends, the base decurrent, the edge entire or more or less sinuate. Texture thinly herbaceous or membranaceous; surfaces naked; main veins distinct; areolae large, distinct, copious, with finer secondary areolae and free

veinlets. Sori on the anastomosing veins, in 2 rows between the main veins, round or oblong, \pm 4—8 in a row.

Var. grandifolium: Fronds larger. Sori more copious, scattered irregularly.

Celebes; Philippines; South. China, North. India, Ceylon.

* (179) **P. leucophorum**, *Bl.*, Journ. of Linn. Soc., Bot., XXII, 229.

Rhizome slender, glaucous, creeping, the scales shining, filiform, brown, nearly spreading, deciduous, to $\frac{1}{2}$ c.M. long. Stipes 1—2 c.M. long, erect. Fronds oblong-lanceolate, acuminate, 5—25 c.M. long, 2—3 $\frac{1}{2}$ c.M. broad at the middle, with the margin recurved. Texture coriaceous; upper surface shining, covered with whitish cretaceous dots, many of which are arranged in a close row placed at or near the margin of the frond; under side thinly villose; main veins fine, straight, erecto-patent; veins anastomosing irregularly. Sori small, superficial, in 2 irregular rows between the main veins or scattered irregularly.

Borneo.

* (180) **P. triquetrum**, *Bl.*, Rac., Flor. Btz., I, 105; *Bl.*, Flor. Jav., II, 141, tab. LIX.

Rhizome long, woody, the scales dense, pale, rather large, broad, elliptical or deltoid-ovate, with the base rounded and peltate, the apex bluntish, the edge entire or erose to lacerate, membranaceous, afterwards deciduous. Stipes distant, firm, erect, naked, glossy, 6—20 c.M. long, the fertile fronds often on the longest stipes. Fronds 10—25 c.M. long, 5 $\frac{1}{2}$ —7 $\frac{1}{2}$ c.M. broad, varying from ovate-oblong to lanceolate, the apex obtuse to acuminate, the base cuneate or rounded, \pm slightly decurrent upon the stipe, the edge subentire or \pm repand, bounded by a narrow, often recurved margin. Texture very coriaceous; surfaces naked, the upper covered with minute, indistinct, scattered black dots; costa and main veins distinct, prominent and subtriquetrous beneath, the latter distinct to the edge. Fertile fronds like the barren ones or longer and narrower. Sori superficial, in 2 rows between the main veins, 4—8 in a row.

Malaya; Samoa.

* (181a) **P. rupestre**, *Bl.*, Rac., Flor. Btz., I, 106; *Bl.*, Flor. Jav., II, 142, tab. LV, fig. 2, tab. LX; *P. saxatile*, *Mett.*, Farng., I, Polypod., 116; *P.*

triquetrum, *Bl.*, var., *C. Chr.*, *Ind. Fil.*; *Pleopeltis rupestris*, *Moore*, *Bedd.*, *Handb. Ind. Ferns*, *Suppl.*, 94; — var. *taeniopsis*; *P. taeniopsis*, *Christ*, *Fil. Saras.*, IV, 443; *P. triquetrum*, *Bl.*, var. *taeniopsis*, *Christ*, *Ann. Btz.*, XV, 157, tab. XVI, fig. 24, a—d; — var. *leucolepis*, *Rst.*, *Fed.*, V, 43.

Rhizome long, woody; scales dull-brown, rather small, narrow, elongate-lanceolate, rounded or cordate at the base, the edge slightly denticulate, the apex spreading, acuminate, at length deciduous. Stipes scattered, firm, erect, naked, glossy, 5—25 c.M. long. Fronds 10—20 c.M. long, 1—5 c.M. broad, lanceolate, acuminate, entire or repand, the edge callous, recurved, the base cuneate or rounded. Texture coriaceous; surfaces naked, the upper side covered with minute, indistinct, scattered black dots; costa keeled beneath; main veins distinct. Fertile fronds similar to the barren ones or longer and narrower. Sori superficial, in 2 rows between the main veins, 4—8 in a row.

Var. taeniopsis: Rhizome scales ferruginous; stipes distant, solitary or in pairs, \pm 3 c.M. long; fertile fronds linear or linear-lanceolate, 4—10 m.M. broad, narrowed gradually towards both ends; main veins not very distinct; sori in 1 row on each side, or in 2 irregular rows but then rather marginal.

Var. leucolepis: Rhizome scales white, narrower, minutely ciliatodentate; stipes slender; texture chartaceous; areolae more conspicuous.

Malaya; — var. *taeniopsis*; *Celebes*; — var. *leucolepis*: *New Guinea*.

(181b) **P. luzonicum**, *Copel.*, *Philipp. Journ.*, I, *Suppl.*, 162, tab. XXIII.

Rhizome scandent, clothed with black-brown, lanceolate, long-acuminate, minutely ciliate scales. Stipes 1—5 c.M. long, stramineous, clothed with deciduous, brown scales. Fronds lanceolate or oblanceolate, 20—30 c.M. long, to 4 c.M. broad, entire, acuminate, the base narrowed gradually into the stipe. Texture subcoriaceous; surfaces naked; main veins distinct nearly to the edge; areolae hidden. Sori large, in 2 rows between the main veins, forming 1—3 rows on each side of the costa, superficial.

Luzon.

(182) **P. albulum**, *Christ*, *C. Chr.*, *Ind. Fil.*; *P. lagopodioides*, *Christ*, *Ann. Btz.*, XIX, 57.

Like *P. rupestre*, *Bl.* but smaller; fronds narrower, the point finer; scales of the rhizome yellowish-white, papyraceous, lanceolate, aristate.

Celebes.

(183) **P. oodes**, *Kze*, Hk. Bk., Syn. Fil., 354.

Rhizome slender, the scales small, fibrillose, yellowish. Stipes distant, slender, naked, 7—10 c.M. long. Fronds $5\frac{1}{2}$ — $6\frac{1}{2}$ c.M. long, $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, ovate, entire, the base rounded, hardly decurrent. Texture coriaceous; both surfaces naked; main veins distinct nearly to the edge; areolae fine, with free veinlets. Sori rather large, few, scattered.

Philippines, Borneo.

(184) **P. dulitense**, *Bk.*, Kew Bull., 1893, 211.

Rhizome slender, wide-creeping, the scales lanceolate, membranaceous, pale-brown, erecto-patent. Stipes firm, 7—13 c.M. long, naked, stramineous. Fronds ovate, 5—8 c.M. long, 2 — $5\frac{1}{2}$ c.M. broad, simple, acute or acuminate, crenate in the lower half, the base rounded or deltoid. Texture chartaceous; surfaces naked, glossy; main veins erecto-patent, parallel, distant; areolae hexagonal, with free veinlets. Sori small, globose, subimmersed.

Borneo.

(185a) **P. suboppositum**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 995.

Rhizome creeping, naked. Stipes approximate, \pm 3 c.M. long, naked, winged nearly to the base by the gradually decurrent base of the fronds. Fronds lanceolate, acuminate, 35—55 c.M. long, \pm 6 c.M. broad at the middle. Texture herbaceous, flaccid; surfaces naked; colour pale-green; costa distinct; main veins horizontally spreading, flexuose, distinct nearly to the edge; primary areolae hidden, longitudinal, in 4—5 rows between the midrib and margin, divided into numerous smaller secondary ones, with included free veinlets. Sori irregular, numerous, \pm 25 to each primary areola.

Luzon.

(185b) **P. Scortechinii**, *Bk.*, Ann. of Bot., V, 477; *Pleopeltis Scortechinii*, *Bedd.*, Handb. Ind. Ferns, Suppl., 95.

Rhizome woody, wide-creeping, flexuose, calvate. Stipes brown, naked, 35—40 c.M. long. Fronds oblanceolate, acuminate, 40—45 c.M. long, 7—9 c.M. broad, obscurely repand, narrowed gradually from the middle to the base. Texture membranaceous; surfaces naked; main veins distinct to the edge. Sori minute, superficial, scattered, often confluent.

Malacca.

(186a) **P. ovatum**, *Wall.*, Hk. Bk., Syn. Fil., 359; Hk. & Grev., Ic. Fil., tab. XLI; *P. phyllomanes*, *Christ*, var. *ovatum*, *Christ*, Bull. Ac. Int. Bot., 1902, 210, tab. page 213 (right hand figures); C. Chr., Ind. Fil.; *Pleopeltis ovata*, *Moore*, *Bedd.*, Ferns, Br.I., II, tab. CLVII.

Rhizome wide-creeping, woody, the scales lanceolate, dull-brown, deciduous. Stipes 10—20 c.M. long, erect, slightly scaly below. Fronds 15—30 c.M. long, 5—8 c.M. broad, the point acute or acuminate, the edge entire, the base rounded to decurrent. Texture firm-herbaceous; both surfaces naked; main veins distinct; areolae copious, with free veinlets. Sori irregular, 1—3-serial.

Luzon; China, North. India.

* (186b) **P. Zollingerianum**, *Kze*, Bot. Zeit., 1846, 422; *P. heterocarpum*, *Bl.*, Rac., Flor. Btz., I, 107; *Bl.*, Flor. Jav., II, 167, tab. LXXV.

Rhizome creeping, blackish when dry, the scales adpressed, lanceolate, brown, not dense. Stipes 5—20 m.M. apart, 7—40 c.M. long, the unwinged portion often short, \pm scaly below, those of the fertile fronds the longest. Fronds lanceolate, entire, 20—60 c.M. long, 4—12 c.M. broad, the apex acute, the base gradually or rather suddenly decurrent, so as to form a long, narrow wing to the stipe. Texture thin, herbaceous; surfaces naked; costa conspicuous; main veins distinct to the edge; veins forming copious areolae with free veinlets. Sori scattered irregularly, roundish or elliptical, often confluent.

Malaya.

(187) **P. millisorum**, *Bk.*, Journ. of Linn. Soc., Bot., XV, 109.

Stipes stout, with a narrow wing extending nearly to the base. Fronds linear-ligulate, 60—75 c.M. long, 5—6 $\frac{1}{2}$ c.M. broad at the middle, narrowed very gradually to both ends. Texture rigidly coriaceous; surfaces naked; main veins zigzag, reaching $\frac{3}{4}$ on the way to the edge; veins fine, hidden; areolae copious, minute, with numerous free veinlets. Sori round, superficial, crowded, filling up the upper $\frac{1}{2}$ or $\frac{2}{3}$ of the frond, distinctly visible on the upper surface.

Kei Islands.

* (188) **P. punctatum**, *Sw.*, *Christ*, Farnkr. d. Erde, 106; *P. irioides*, *Poir.*, Rac., Flor. Btz., I, 106; *Bl.*, Flor. Jav., II, 169, tab. LXXVII; *Hk.*, Fil. Exot., tab. IV; *Pleopeltis irioides*, *Moore*, *Bedd.*, Ferns S.I. tab. CLXXVIII; *Pl. punctata*, *Bedd.*, Handb. Ind. Ferns, 357; *Acrostichum punctatum*, *L.*, (oldest

name); — var. *mindanense*, *subirideum* & *subdrynariaceum*; *P. mindanense*, *subirideum* & *subdrynariaceum*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 994.

Rhizome creeping, stout, the scales deltoid-oblong to ovate-lanceolate, blunt or acute, ± dark-coloured. Stipes rather approximate, 0—1 c.M. long. Fronds linear-lanceolate, ± 30—90 c.M. long, 2½—6½ c.M. broad, narrowed gradually towards both ends, the edge entire, narrowly subrevolute, the apex ± acute or acuminate, sometimes bluntish, the very base bluntish or slightly rounded. Texture subcoriaceous, rather fleshy; surfaces naked; costa prominent on both sides; main veins hidden in fresh material, not coloured, fine, slightly prominent when dry, reaching ± the edge; areolae fine, copious, with free veinlets. Sori small, very abundant, often nearly close, scattered irregularly, occupying the upper ⅓—⅔ of the frond.

Var. *mindanense*: Fronds considerably dilated at the base; texture papyraceous; veins prominent; areolae hidden; sori larger.

Var. *subirideum*: Fronds 20—50 c.M. long, 2½—3 c.M. broad, the apex often obtuse, the base often dilated; texture fleshy; sori near the apex, larger.

Var. *subdrynariaceum*: Like var. *mindanense*, but the venation more conspicuous and the base of the fronds scarioso; sori very numerous, small.

Trop. Asia & Australia, Polynesia, Trop. Africa, Mascarenes; — the varieties in *Luzon*.

* (189) ***P. musifolium*, Bl.**, Rac. Flor. Btz., I, 107; Bl., Flor. Jav., II, 171, tab. LXXIX; *P. Schumannianum*, *Diels*, Schum. & Laut., Flor. deut. Sch. geb., 159, tab. III, C—D; *Pleopeltis musaefolia*, *Moore*, Bedd., Ferns Br.I., II, tab. CCCXVII.

Rhizome woody, branched, creeping, firm, the scales ovate-lanceolate, rather dark-brown. Fronds sessile, approximate or distant, 50—100 c.M. long, 7½—20 c.M. broad, linear or ligulate, the apex bluntish or acute, the edge entire, the base rounded or cordate. Texture thinly coriaceous, firm; surfaces naked; costa and main veins distinct; main veins straight, rather dark-coloured in fresh material, connected by transversal, arched veins; included areolae copious, with free veinlets. Sori small, very abundant, sometimes occupying the whole under surface. — *P. Schumannianum*, *Diels* from New Guinea seems to be a local derivative of this species, with smaller fronds, which are ovate-oblong, or panduriform with a narrowed part ± 1—1½ c.M. broad, and with the base cordate, rounded or cuneate.

Malaya.

(190a) **P. validum**, *Copel.*, Polypod. Philipp., 126.

Rhizome short, thick, the scales ovate, adpressed; obtuse, cordate or peltate. Fronds tufted, nearly sessile, \pm 70 c.M. long, 10 c.M. or more broad, lanceolate, entire, subacute, narrowed gradually to the base, the lower $\frac{1}{3}$ part wing-like, the margin entire. Texture carnosu-coriaceous, fragile; surfaces naked; main veins fine, reaching nearly to the margin; transversal veins parallel to the costa, forming \pm 7 rows of divided areolae on each side, with free veinlets. Sori numerous, scattered irregularly.

Philippines.

(190b) **P. monstrosum**, *Copel.*, Leaf., I, 78; — var. *leucophlebium* & *integrius*, *Copel.*, l. c.

Rhizome scandent, slender, clothed with dark-grey, lanceolate, acuminate scales, which are peltate at the base. Stipes $\frac{1}{2}$ —1 c.M. long, naked. Fronds 50—50 c.M. long, 4—7 c.M. or more broad, narrowed gradually towards both ends, acuminate, the edge coarsely crenate or lobed, with the central lobes whether or not much produced. Texture subcoriaceous; surfaces naked; costa prominent, beneath acute, above 2-angular in the upper part; main veins conspicuous, black, reaching nearly the margin; transversal veins \pm 6, inconspicuous; veinlets hidden. Sori crowded, scattered irregularly.

Var. leucophlebium: Fronds often narrowed and rather entire; veins beneath white.

Var. integrius: Hardly different from var. *leucophlebium*, the fronds crenato-sinuate.

Luzon, Mindoro.

Species of an indefinite systematical place. Sori unknown.

(191) **P. Weinlandii**, *Christ*, Bull. Herb. Bois., 2^e Série, I, 448, with figure.

Rhizome long; creeping, the bristles subulate, brown, dense, subadpressed. Stipes distant, 5—6 c.M. long, firm, densely covered like the whole frond with simple, white, diaphanous, spreading hairs. Fronds ovate-elongate, 7—10 c.M. long, \pm 4 c.M. broad, shortly acute or obtuse, the base broadly cuneate, the margin undulate or repand. Texture coriaceous; colour pale-brownish; veins very conspicuous; costa prominent; main veins oblique, reaching to the margin; areolae subquadrangular, large, 5—4 between each pair of main

veins, with included free veinlets; upper surface covered with numerous, scattered, cretaceous dots. Sori unknown.

New Guinea.

§ 6. PLEOPELTIS. c. **Chrysopteris**. *. **Phymatodes**.

a. *Sori superficial.*

b. *Sori in 1—4 rows on each side of the costulae.*

c. *Texture herbaceous to papyraceous.*

d. *Lobes entire or repand.*

e. *Fronds rather small, under 25 c.M. long.*

f. *Lateral segments in \pm 2—4 pairs.*

(192) *P. paucijugum.*

f.f. *Lateral segments in \pm 8—9 pairs.*

(193) *P. sundense.*

e.e. *Fully developed fronds more than 25 c.M. long.*

‡. *Rhizome \pm naked.*

×. *Stipes not winged.*

(194a) *P. alatum.*

×.×. *Stipes winged.*

(194b) *P. flaccidum.*

‡.‡. *Rhizome scaly. Stipes winged.* (195) *P. commutatum.*

d.d. *Lobes serrulate or toothed.*

(196) *P. ebenipes.*

c.c. *Texture chartaceous to coriaceous.*

§. *Under surface bluish-glaucous.*

(197) *P. glaucum.*

§.§. *Under surface green, not glaucous.*

‡. *Sori nearer the costulae than the margin.* (1)

(198) *P. hastatum.*

‡.‡. *Lower sori medial, the higher ones nearer the margin than the costulae.* (199) *P. quinquefidum.*

b.b. *Sori scattered irregularly.* (2)

1. *Fronds dimorphous.*

(200) *P. anomalum.*

1.1. *Fronds uniform.*

2. *Lower segments much reduced.*

(201) *P. grandidentatum.*

2.2. *Lower segments hardly, if at all, reduced.*

3. *Stipes \pm short-winged. Fronds typically 3-fid, but sometimes simple or 5-fid.* (3)

(202a) *P. pteropus.*

(1) See also **P. rostratum**, *Hk.*

(2) See also **P. monstrosum**, *Copel.*

(3) See also **P. rivulare**, *Copel.*

3.3. *Stipes short-winged. Fronds as in 3.3.3.* •

(202b) *P. dolichopterum.*

3.3.3. *Stipes long-winged. Fronds typically pinnatifid with 3 or more pairs of lateral segments below the terminal one.*

4. *Fronds small or middle-sized, less than 40 c.M. long above the wing of the stipe. Rhizome scales linear or lanceolate.*

5. *Rhizome scales sparse.*

6. *Rhizome fleshy.* (205a) *P. rivulare.*

6.6. *Rhizome woody.* (205b) *P. pentaphyllum.*

5.5. *Rhizome scales crowded.* (204) *P. insigne.*

4.4. *Fully developed fronds large, more than 40 c.M. long.* (205) *P. dilatatum.*

a.a. *Sori more or less deeply immersed. (1)*

b. *Fronds dimorphous. Barren fronds with the segments lanceolate-oblong, fertile with the segments narrowly linear.* (206) *P. incurvatum.*

b.b. *Fronds uniform, rarely subdimorphous.*

c. *Fronds ± as broad as long, the lower lobes deflexed.*

(207) *P. macrochasmum.*

c.c. *Fronds distinctly longer than broad.*

d. *Sori shallowly immersed. (2)*

*. *Fronds under 1 M. long.* (208a) *P. phymatodes.*

.. *Fronds more than 1 M. long.*

(208b) *P. phanerophlebium.*

d.d. *Sori deeply immersed.*

e. *Lateral segments in ± 2—10 pairs.*

×. *Under surface pale-violescent. Fronds ± 15 c.M. long.* (209) *P. violascens.*

×.×. *Under surface green. Fronds ± 20—50 c.M. long.* (210) *P. nigrescens.*

e.e. *Lateral segments in many or numerous pairs.*

f. *Texture thinly papyraceous; segments acuminate.*

(211) *P. Schneideri.*

f.f. *Texture firm; segments acute to blunt.*

(212) *P. longissimum.*

(1) See also *P. commutatum*, Bl.

(2) See also *P. longissimum*, Bl.

* (192) **P. paucijugum**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 24.

Rhizome slender, creeping, the scales lanceolate. Stipes slender, erect, scaly, 4—10 c.M. long below the winged portion. Fronds 10—20 c.M. long, with an entire terminal lobe 5—10 c.M. long, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad, and 2—4 similar lobes on each side, which are suberect or erecto-patent, narrowed gradually towards an acute or bluntish point, connected by a costal wing 5—5 m.M. broad, the lowest gradually decurrent upon the stipe. Texture thin but firm; under surface minutely hairy; costa and costulae \pm scaly beneath, prominent like the primary veins, which stop short and unite at $\pm \frac{2}{3}$ of the way to the margin, forming 1 series of large primary areolae on each side, each with commonly 1 sorus in the narrower and mostly 2 sori in the broader segments; secondary areolae in and beyond the primary ones fine, indistinct, with few free veinlets. Sori small, rather oblong.

Borneo (Teuscher).

(195) **P. sundense**, *C. Chr.*, Ind. Fil.; *P. sumatranum*, *Blk.*, Ann. of Bot., VIII, 151 (not Journ. of Bot., XVIII, 214).

Stipes moderately long, naked, stramineous. Fronds lanceolate-oblong, 17—20 c.M. long, 7—10 c.M. broad, deeply pinnatifid, the base hardly narrowed. Segments 8—9-jugate, contiguous, lanceolate, entire, acute, confluent at the base, $1\frac{1}{4}$ — $1\frac{1}{2}$ c.M. broad. Texture moderately firm; surfaces naked; areolae hexagonal, copious, with free veinlets. Sori large, semiglobose, superficial, in 1—2 rows on each side of the costulae.

Sumatra.

(194a) **P. alatum**, *Hk.*, Hk. Bk., Syn. Fil., 365; *P. Wilkesii*, *C. Chr.*, Ind. Fil.; *Drynaria alata*, *Brack.*, Expl. Exp., XVI, 48, tab. VI, fig. 1.

Rhizome woody, scaleless. Stipes 50—45 c.M. long, firm, erect. Fronds 45—60 c.M. long or shorter, 25—30 c.M. or more broad, cut down nearly to the rachis below (within $1\frac{1}{2}$ c.M. above) into few or numerous (4—50), rather distant or more close, erecto-patent, repand, linear-oblong, acuminate lobes $2\frac{1}{2}$ c.M. or more broad. Texture papyraceo-herbaceous; both surfaces naked; main veins distinct to the edge; areolae copious, with free veinlets. Sori in rows of 5—4 near the costulae.

Celebes; Samoa, Fiji.

(194b) **P. flaccidum**, *Christ*, Philipp. Journ., II, 178.

Rhizome creeping, grey-brown. Stipes slender, stramineous, 6—8 c.M. long, naked, or provided with blackish setulae downwards, winged nearly or quite to the base. Fronds oval, \pm 40 c.M. long, \pm 18 c.M. broad, deeply pinnatifid. Terminal lobe \pm 18 c.M. long, caudate; lateral lobes \pm 4—5 on each side, \pm 2—5 c.M. apart, \pm 10 c.M. long, \pm 1 $\frac{1}{2}$ c.M. broad, caudate, connected by a costal wing \pm 1 $\frac{1}{4}$ c.M. broad, the sinuses broadly rounded, the lowest lobes decurrent into the wing. Texture flaccido-papyraceous, hyaline; colour pale-green; surfaces naked; costa and costulae distinct, stramineous; main veins slender, curved, indistinct near the edge; areolae numerous, with included free veinlets. Sori numerous, rather small, roundish, irregularly 2—3-seriate, not immersed in pits.

Luzon.

* (195) **P. commutatum**, *Bl.*, Flor. Jav., II, 165, tab. LXXIII; *P. affine*, *Bl.*, Rac., Flor. Btz., I, 115; *Bl.*, l. c., 160, tab. LXIX.

Rhizome woody, the scales roundish, oblong when young, entire, adpressed, dark-brown, deciduous. Stipes \pm 20—45 c.M. long, firm, glossy. Fronds 50—120 c.M. long, 15—30 c.M. or more broad, cut down into few or several spreading or erecto-patent lobes on each side. Lobes linear, acuminate to caudate, \pm 1—4 c.M. broad, connected at the base by a costal wing $\frac{1}{2}$ —2 c.M. broad, the sinuses rounded; lowest lobes decurrent so as to form a narrow wing to the stipe. Texture papyraceous; surfaces naked; primary veins flexuose, becoming gradually finer towards the margin, uniting several times so as to form \pm 3 series of alternate, \pm hexagonal primary areolae; secondary areolae numerous, fine, indistinct, with copious free veinlets. Sori superficial or immersed, in 2—5 rows on each side of the central ribs of the laciniae.

Malaya.

* (196) **P. ebenipes**, *Hk.*, Hk. Bk., Syn. Fil., 565; *Pleopeltis ebenipes*, *Bedd.*, Ferns Br.I., I, tab. CXXXVIII.

Rhizome stout, creeping, the scales large, lanceolate, glossy, \pm black, subcoriaceous, not hair-pointed. Stipes 15—25 c.M. long, naked, glossy, on short branches of the rhizome. Fronds 15—45 c.M. long, 20—50 c.M. broad, cut down to a broadly winged costa into 6—20 pairs of segments. Segments 1—5 c.M. broad, a little serrulate, acuminate, the lowest pair generally deflexed and either a little decurrent on the stipe or scooped out and subcordate at the lower base, sometimes quite spreading as are the other seg-

ments. Texture firm-papyraceous; surfaces naked; costa puberulous; main veins distinct to the edge; areolae fine, with free veinlets. Sori in single rows near the costulae.

Borneo; Yunnan, North. India.

(197) **P. glaucum**, *Hæ*, Hk. Bk., Syn. Fil., 365; Mett., Farng., I, Polypod., tab. II, fig. 12—13; *P. glauco-pruinatum*, *C. Chr.*, Ind. Fil.

Rhizome short, creeping, the scales lanceolate-subulate, dark-brown, rigid. Stipes 7—50 c.M. long, firm, erect, glossy. Fronds oblong, 20—40 c.M. long, deeply pinnatifid. Lobes \pm 4—6 on each side, 7—15 c.M. long, \pm 1 c.M. broad, erecto-patent, linear, acuminate, the edge entire or distantly crenato-serrulate, the lowest lobes slightly decurrent on the stipe, broad, sterile, a little abbreviate, the higher ones narrow, fertile. Texture coriaceous; under surface glauco-pruinose, upper often covered with calcareous scales; no main veins; areolae hidden. Sori superficial, in 1 row on each side of the costulae.

Philippines.

* (198) **P. hastatum**, *Thb.*, Hk. Bk., Syn. Fil., 361; Kze, Farnkr., I, tab. LXXXIII; *Pleopeltis oxyloba*, *Bedd.*, Ferns S.I., tab. CLXXV; *Pl. hastata*, *Moore*, *Bedd.*, Handb. Ind. Ferns, 362; — var. *trifidum*; *Pol. trifidum*, *Don*, Hk. Bk., l. c., 365.

Rhizome firm, short-creeping, the scales linear-lanceolate, squarrose, brownish. Stipes scattered, 5—15 c.M. long, erect, firm, glossy. Fronds 10—50 c.M. long, either simple, lanceolate, 1—4 c.M. broad, or 3-fid with the lateral lobes erecto-patent, similar to the terminal one but smaller, or the fronds pinnatifid with 2—4 pairs of lateral lobes below the terminal one; lobes elongate, 7—15 c.M. long, 1—2½ c.M. broad, narrowed gradually towards the acute or acuminate point, the margin subentire or slightly crenate to repand; lowest lobes sometimes forked, slightly decurrent on the stipe. Texture thinly coriaceous; surfaces naked, the lower pale-green; main veins distinct, straight, parallel; areolae with free veinlets. Sori round, large, superficial, in 1 row on each side, nearer the costulae than the margin, solitary between the main veins.

Var. trifidum: Larger form, with more than 4 pairs of lateral lobes. *Philippines, Malacca; Ceylon, North. India, China, Japan, Formosa.*

(199) **P. quinquesfidum**, *Bk.*, Journ. of Bot., XVIII, 216.

Rhizome woody, the scales brown, linear. Stipes distant, naked, erect,

rigid, pale-brown, 12—25 c.M. long. Fronds deltoid, 15—20 c.M. long, 20—25 c.M. broad, cut down nearly or quite to the costa. Segments 5—7, lanceolate, acuminate, 10—15 c.M. long, 1—2 c.M. broad, the edge obscurely repand. Texture coriaceous; main veins of the segments raised, erecto-patent, distinct and straight to the edge; areolae small, copious, hexagonal. Sori large, round, superficial, solitary between the main veins, the lower medial between the costulae and margin, the higher nearer the margin.

Sumatra.

(200) **P. anomalum**, *Christ*, Bull. Herb. Boiss., VI, 201, tab. III, fig. 5, a—d; *P. heterolobum*, *C. Chr.*, Ind. Fil.

Rhizome creeping, the scales brown, ovate-subulate, crisped, patent. Stipes scattered, 3—4 c.M. long, naked like the whole plant, stramineous. Fronds \pm 20 c.M. long, the base gradually decurrent; sterile fronds to \pm 8 c.M. broad, irregularly pinnatifid \pm $\frac{1}{2}$ -way down to the costa into lobes different in size and shape; fertile fronds lanceolate-ligulate, \pm $1\frac{1}{2}$ c.M. broad, the margin irregularly repand. Texture membranaceous; costa distinct; main veins conspicuous; areolae large, principally in \pm 4 rows on each side of the costa, with free veinlets. Sori numerous, irregularly scattered.

Philippines.

(201) **P. grandidentatum**, *Bk.*, Ann. of Bot., V, 479; *P. dilatatum*, *Wall.*, var. *grandidentatum*, *Ces.*, Fel. di Bor., 27.

Stipes 50 c.M. or more long below the wing, the wing \pm 15 c.M. long to the lowest segment. Fronds oblong, \pm 90 c.M. long, deeply pinnatifid. Segments lanceolate, the central ones 20—25 c.M. long, $2\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, sharply toothed, the lower ones much dwarfed. Texture membranaceous; surfaces naked; main veins of the segments erecto-patent, distinct to the edge. Sori copious, minute, scattered.

Borneo.

* (202a) **P. pteropus**, *Bl.*, Rac., Flor. Btz., I, 110; *Bl.*, Flor. Jav., II, 168, tab. LXXVI; *P. tridactylum*, *Wall.*, Hk. & Grev., Ic. Fil., tab. CCIX; *Pleopeltis tridactyla*, *Moore*, Bedd., Ferns Br.I., I, tab. XI; *Pl. pteropus*, *Moore*, Bedd., Handb. Ind. Ferns, 559.

Rhizome creeping, the scales lanceolate-subulate, blackish or brownish. Stipes scattered, 5—20 c.M. long, naked or sparingly scaly. Fronds lanceolate, simple to pinnatifid with a large terminal lobe 10—50 c.M. long, $2\frac{1}{2}$ —5 c.M.

broad and 1—2 smaller ones on each side. Lobes lanceolate, acute or acuminate, the edges entire or irregularly crenate; lowest lobes (or the base of the simple frond) gradually decurrent so as to form a narrow wing to the upper part of the stipe. Texture herbaceous; surfaces naked or the lower \pm scaly on the veins; areolae fine, copious, with free veinlets; main veins not reaching the edge. Sori small, not abundant, scattered irregularly, 1—7 between the main veins.

Trop. Asia.

(202b) **P. dolichopterum**, *Copel.*, Philipp Journ., I, Suppl., 162.

Rhizome creeping, the extremity clothed with black, lanceolate scales. Stipes seriate, 10—15 c.M. long, stramineous, naked, or sparingly scaly towards the base. Fronds deltoid, pinnatifid. Lobes divaricating, connected at the base by a costal wing 2—5 m.M. broad, the lowest lanceolate, 10—12 c.M. long, 1—1 $\frac{1}{4}$ c.M. broad, entire or undulate, acuminate, the base decurrent on the stipe so as to form a short wing to 5 c.M. long. Texture membranaceous; surfaces naked; veins laxly reticulate; areolae with few free veinlets. Sori scattered or irregularly seriate, superficial.

Philippines.

(203a) **P. rivulare**, *Copel.*, Philipp. Journ., I, Suppl., 165.

Rhizome creeping, green, fleshy, sparingly clothed with black, lanceolate, acuminate scales. Stipes 5—10 c.M. long, glabrescent, stramineous, narrowly winged, the unwinged basal part 1—5 c.M. long. Fronds 10—16 c.M. long, broadly ovate, pinnatifid, the costal wing \pm as broad as the lobes. Lobes 1—4-jugate, to 7 c.M. long, $\frac{1}{2}$ —1 $\frac{1}{2}$ c.M. broad, acute, entire or undulate, the terminal one often longer. Texture membranaceous; surfaces naked; lateral veins inconspicuous, not visible by reflected light. Sori scattered irregularly.

Luzon.

(203b) **P. pentaphyllum**, *Bk.*, Ann. of Bot., V, 478.

Rhizome wide-creeping, woody, the scales small, sparse, lanceolate, acuminate. Stipes naked, \pm 2 $\frac{1}{2}$ c.M. long below the wing. Fronds \pm 15—50 c.M. long, deeply pinnatifid. Lobes distant, linear-lanceolate, \pm $\frac{3}{4}$ —1 c.M. broad, the lowest ones decurrent so as to form a mere wing to the stipe \pm 15 c.M. long. Texture membranaceous; surfaces naked; veins fine, copious. Sori minute, scattered.

Philippines.

* (204) **P. insigne**, *Bl.*, *Rac.*, *Flor.*, *Btz.*, I, 113; *Bl.*, *Flor. Jav.*, II, 166, tab. LXXIV; *P. tridactylon*, *Mett.*, (not *Wall.*), *Farnng.*, I, *Polypod.*, 104, tab. I, fig. 39—40; *Pleopeltis insignis*, *Bedd.*, *Ferns Br.I.*, II, tab. CCXIV; *Drynaria decurrens*, *Brack.*, *Expl. Exp.*, XVI, 48.

Rhizome creeping, the scales narrow, lanceolate-linear, spreading. Stipes scattered or approximate, erect, naked, \pm 2—5 c.M. long below the wing. Fronds 15—50 c.M. long, 10—20 c.M. broad, with an entire, lanceolate-acuminate terminal lobe 5—10 c.M. long, 1—1½ c.M. broad or broader and 2—6 or more similar ones on each side, which are connected by a costal wing 5—15 m.M. broad, the lowest lobes decurrent so as to form a narrow wing to the stipe \pm 15—50 c.M. long. Texture herbaceous, firm; surfaces naked; veins fine; areolae with free veinlets. Sori small, superficial, roundish or irregularly oblong, often confluent, scattered irregularly.

Malacca, Java, Philippines.

* (205) **P. dilatatum**, *Wall.*, *Hk. Bk.*, *Syn. Fil.*, 565; *P. eurynphyllum*, *C. Chr.*, *Ind. Fil.*; *Pleopeltis dilatata*, *Bedd.*, *Ferns Br.I.*, I, tab. CXXII.

Rhizome woody, the scales ovate, large, brown. Stipes up to 30—40 c.M. long, with a narrow wing reaching down nearly to the base. Fronds 45—90 c.M. long, \pm 50 c.M. or more broad, cut down within ½—2½ c.M. of the costa into 8—16 erecto-patent lobes on each side. Lobes \pm 20 c.M. long, 1—4 c.M. broad, entire, acuminate. Texture membranaceo-papyraceous; surfaces naked; main veins fine, flexuose, not quite reaching the margin; areolae copious, fine, with free veinlets. Sori small, scattered irregularly, often confluent.

Malaya; Ceylon, North. India, South. China, Samoa.

* (206) **P. incurvatum**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 110; *Bl.*, *Flor. Jav.*, II, 151, tab. LXV; *Pleopeltis incurvata*, *Moore*, *Bedd.*, *Ferns Br.I.*, I, tab. CXXIV.

Rhizome creeping, woody; scales ovate-lanceolate, the base roundish, pel-tate, adpressed. Stipes distant, brown, \pm 25 c.M. long. Fronds dimorphous. Barren fronds 15—25 c.M. each way, rarely simple, more generally hastato-trifid or pinnatifid with 2—8 pairs of lateral lobes below the terminal one; lobes approximate, ovate-deltoid or lanceolate-triangular, the edge entire, thickened, the point acute. Texture firm, coriaceous; surfaces naked; main veins distinct to the edge; veins hidden; areolae with free veinlets. Fertile fronds cut down nearly or quite to the costa into erecto-patent, linear or linear-lanceolate lobes, the lobes \pm 5—7 on each side, ½—1 c.M. broad,

distant, decurrent, the lowest sometimes forked. Sori round, or oblong and parallel to the central ribs of the lobes, in 1 row on each side, quite immersed in deep cavities, which form conical, truncate tubercles on the upper surface.

Malaya.

* (207) **P. macrochasmum**, *Bk.*, Journ. of Bot., XVIII, 216; *Hk.*, Icon. Pl., XVII, tab. MDCLXXV; *P. laciniatum*, *Bl.*, Rac., Flor. Btz., I, 111; *Bl.*, Flor. Jav., II, 149, tab. LXIII; *Pleopeltis laciniata*, *Bedd.*, Handb. Ind. Ferns, Suppl., 97.

Rhizome woody, wide-creeping, the scales dense, linear-lanceolate, acuminate, ciliate, squarrose. Stipes distant, 10—50 c.M. long, firm, erect, naked. Fronds deltoid-ovate, 20—50 c.M. each way, cut down nearly (to a narrow wing) to the costa into \pm 6—18 linear-lanceolate, acuminate lobes. Lobes to 25 c.M. long, 1—1½ c.M. broad, with the edge entire or crenulate to slightly undulate, the lowest ones deflexed. Texture coriaceous; surfaces naked; main veins prominent below, distinct and straight to the edge; areolae with free veinlets. Sori in 1 row on each side of the costulae, nearer the costulae than the edge, immersed, prominent on the upper surface.

Malacca, Sumatra, Java, Borneo, Celebes.

* (208a) **P. phymatodes**, *L.*, Rac., Flor. Btz., I, 114; *P. trifidum*, *Ces.* (not *Don*), Fel. Becc., Polin., 5; *Pleopeltis Phymatodes*, *Moore*, *Bedd.*, Ferns S.I. tab. CLXXIII; — var. *dimorphum*, *Christ.*, Fil. Saras., I, 23.

Rhizome wide-creeping, the scales ovate-lanceolate, acuminate, peltate, adpressed, deciduous. Stipes scattered, 7—50 c.M. long, firm, erect, naked, glossy. Fronds varying from simple (oblong-lanceolate, 10—20 c.M. long, entire or irregularly crenate or slightly lobed) to deeply pinnatifid (20—90 c.M. long, 15—60 c.M. broad) with 1 to numerous entire, erecto-patent lobes on each side. Lateral lobes lanceolate-oblong to linear, similar to the terminal one, 10—56 c.M. long, 1—4 c.M. broad, the apex acuminate, the bases connected by a costal wing \pm ½—1 c.M. broad, the sinuses \pm rounded. Texture \pm coriaceous; surfaces naked; no distinct main veins; costal areolae of the fertile fronds large, 1-serial, secondary ones copious, fine, with free veinlets. Sori large, shallowly immersed, round or oblong, placed in 1—2 regular or irregular rows in the smaller forms of the species, or scattered irregularly in the larger.

Var. dimorphum: Barren fronds shallowly 3-lobed, the lobes

very broad. Fertile fronds with ± 4 pairs of lobes ± 12 c.M. long, ± 3 c.M. broad. Sori numerous, copious, scattered.

Trop. and Subtrop. Asia, Africa & Australia, Ppynesia; — var. *dimorphum*: *Celebes*.

(208b) **P. phanerophlebium**, *Copel.*, Philipp. Journ., I, Suppl., 165, tab. XXIV.

Rhizome scandent, clothed with ovate or orbicular, peltate, entire, imbricate, persistent scales. Stipes naked, 40—80 c.M. long. Fronds $\pm 1\frac{1}{2}$ M. long, ± 60 c.M. broad. Lobes ± 55 c.M. long, $\pm 4\frac{1}{2}$ c.M. broad, entire, acuminate, connected at the base by a costal wing 1—1 $\frac{1}{2}$ c.M. broad. Texture herbaceous or subcoriaceous; surfaces naked; venation as in *P. commutatum*, *Bk.*, the primary veins very conspicuous. Sori large, hardly immersed, often in 2 rows on each side of the costulae, 1 in each primary areola.

Mindanao.

(209) **P. violascens**, *Mett.*, Ann. Mus. Bot. L. B., II, 227.

Rhizome creeping, the scales rufo-ferruginous, nearly glossy, lanceolate-acuminate, hardly ciliate. Stipes 7—8 c.M. long. Fronds ± 15 c.M. long, oblong, pinnatisect. Segments approximate, ± 6 on each side, spreading, lanceolate, acute, the margin callous, serrate; lower segments nearly free, with the base narrowed, adnate, the higher ones adnate or decurrent. Texture coriaceous; under surface pale-violescent; costulae prominent; areolae immersed. Sori in 1 row on each side of the higher costulae, deeply immersed.

Java.

* (210) **P. nigrescens**, *Bl.*, Rac., Flor. Btz., I, 111; *Bl.*, Flor. Jav., II, 161, tab. LXX; *Hk.*, Fil. Exot., tab. XXII; *Pleopeltis nigrescens*, *Carr.*, *Bedd.*, Handb. Ind. Ferns, 367; *Pl. longissima*, *Bedd.*, Ferns S.I. tab. CLXXVI.

Rhizome stout, creeping, the scales ovate, adpressed, thin, brown. Stipes 20—90 c.M. long, firm, erect, pale, glossy. Fronds 50—90 c.M. long, 20—50 c.M. broad, cut within $\frac{1}{2}$ —2 $\frac{1}{2}$ c.M. of the costa into ± 2 —10 linear, spreading or erecto-patent, entire, acuminate to caudate lobes on each side to 30 c.M. long, 1—5 c.M. broad, the sinuses as broad or narrower. Texture thinly coriaceous; surfaces naked; colour often blackish when dry; main veins prominent, flexuose, forming 1 row of large costal areolae; areolae in and

beyond the costal ones fine; free veinlets abundant. Sori 1 in each costal areola, sunk in deep cavities, very prominent on the upper surface. — The young plants have the fronds simple or 3-fid.

Malaya; Polynesia, Trop. Australia, Ceylon, South. India.

(211) **P. Schneideri**, *Christ*, Bull. Herb. Boiss., VI, 855.

Rhizome wide-creeping, the scales ovate, acute, pellucid, with the margin toothed, pale-brown. Stipes distant, the base broad, very stout, firm, \pm 60 c.M. long, terete, naked, brownish in the lower part, the upper part yellowish to purpureous like the costa. Fronds \pm 100 c.M. long, to \pm 18 c.M. broad, linear-lanceolate. Segments \pm 24 on each side, linear-lanceolate, erecto-patent, alternate, the higher ones subopposite, \pm $1\frac{1}{2}$ c.M. broad, \pm 2 c.M. apart, \pm 12 c.M. long, acuminate, with the margin entire or undulate, rarely denticulate at the apex, and the bases decurrent, joined so as to form a narrow wing $1\frac{1}{2}$ c.M. broad, with rounded sinuses; lower lobes moderately obtuse, \pm 8 c.M. long. Texture thinly papyraceous; surfaces naked; costa very prominent; costulae yellowish, very prominent; veins conspicuous, fine, prominent, dark-coloured, spreading, straight, forked; areolae subpentagonal, including smaller areolae with free veinlets. Sori large, roundish, occupying the upper part of the frond, 1-serial on each side of the costulae, nearer the costulae than the margin and commonly close to the former, 12—15 in a row, remote, immersed, forming papillae on the upper surface of the frond.

Sumatra.

* (212) **P. longissimum**, *Bl.*, Rac., Flor. Btz., I, 112; *Bl.*, Flor. Jav., II, 159, tab. LXVIII; *Pleopeltis longissima*, *Moore*, Bedd., Ferns S. Br.I., Suppl., tab. CCCLXXXVIII; — var. *batjanense*, *Christ*, in Warb., Mons., I, 62.

Rhizome wide-creeping, the scales ovate or broadly lanceolate, blunt, thin, adpressed. Stipes scattered, 15—120 c.M. long, firm, glossy, naked. Fronds elongate, 50—120 c.M. long, 10—50 c.M. broad, cut down nearly to the costa into numerous erecto-patent lobes. Lobes linear, bluntish or acute, entire or nearly so, 7—20 c.M. long, $\frac{1}{2}$ —2 c.M. broad, the bases connected by a costal wing \pm 1—2 m.M. broad, the sinuses rounded. Texture subcoriaceous; surfaces naked; veins forming large, distinct costular areolae and in and beyond these irregular smaller ones with free veinlets; main veins indistinct, not reaching the margin. Sori deeply immersed, round, in single rows on each side near the costulae.

Var. batjanense: Fronds smaller; lobes abbreviate; texture firmer; sori hardly immersed.

Malaya; Polynesia; — var. batjanense: Batjan Island.

§ 6. PLEOPELTIS. c. **Chrysopteris.** **. **Phymatopsis.**

a. *Fronds dimorphous.* (215) *P. lagunense.*

a.a. *Fronds uniform.*

b. *Sori deeply immersed.*

*. *Pinnae acute.* (214) *P. sculpturatum.*

.. *Pinnae obtuse.* (215) *P. Sauvinieri.*

b.b. *Sori superficial or shallowly immersed.*

c. *Upper surface covered with submarginal or scattered white dots.*

d. *Pinnae under 1 c.M. broad; veins hidden; surfaces pale-green.*

(216) *P. subaquatile.*

d.d. *Pinnae 1—2 c.M. broad or narrower, but then the veins distinct; upper surface bright-green.* (217) *P. albidosquamatum.*

c.c. *Upper surface without white dots.*

✧. *Pinnae dilatato-adsnate at the base.* (218) *P. Moseleyi.*

✧.✧. *At least the lowest pinnae narrowed at the base and not rarely ± stalked.*

*. *Sori superficial.* (219a) *P. palmatum.*

.. *Sori slightly immersed.* (219b) *P. proteus.*

(213) **P. lagunense**, *Christ*, Bull. Herb. Boiss., VI, 201, tab. III, fig. 4, a—d.

Rhizome wide-creeping, the scales dense, ovate, acute at the apex, umbonate towards the base, adpressed, reddish-brown, leaving behind roundish, prominent cicatrices when they fall. Stipes scattered, naked, firm, 10—15 c.M. long. Fronds dimorphous, 15—20 c.M. long, pinnate. Barren fronds with 6—7 pinnae on each side below the pinnatifid apex. Pinnae distant, ovate, ± 4 c.M. long, ± 2 c.M. broad, sessile or short-stalked, the base narrowed, the edge crenate or slightly toothed, the apex acute or elongated into a long, serrate, linear cauda. Surfaces naked; costae conspicuous; main veins less distinct, reaching the edge; areolae indistinct. Fertile fronds with the pinnae linear-lanceolate or ligulate, ± 12 c.M. long, 3—10 m.M. broad, erecto-patent,

distinctly crenate; costae conspicuous. Sori numerous, in 1 row on each side of the costae, 1 to each crenation, large, immersed, medial.

Philippines.

(214) **P. sculpturatum**, *Bk.*, Hk. Bk., Syn. Fil., 514.

Stipes grey, glossy, naked. Fronds oblong, 55—45 c.M. long, 20—25 c.M. broad. Pinnae 15—25-jugate, erecto-patent, subdistant, adnate, much dilated at the base, narrowed to a point, slightly repand, $\frac{1}{2}$ —1 c.M. broad at the middle, lowest largest. Texture rigidly subcoriaceous; surfaces naked; veins distinctly raised in the dried plant. Sori in 1 row on each side of the midribs, midway between these and the edge, 20—55 in a row, deeply immersed, the pits without bordered edges, prominent on the upper surface.

Java.

(215) **P. Sauvinieri**, *Bk.*, Ann. of Bot., V, 480.

Stipes naked, stramineous, 10—13 c.M. long. Fronds lanceolate-oblong, \pm 45 c.M. long, 10—15 c.M. broad at the middle, narrowed to the base. Pinnae 30—40-jugate, linear, obtuse, 6—7 m.M. broad, adnate by a dilated base. Texture firm; surfaces naked; veins hidden. Sori in 1 row on each side of the central ribs, medial, sunk in deep pits forming papillae on the upper surface, \pm 20 in a row in the central pinnae.

Celebes.

* (216) **P. subaquatile**, *Christ*, Ann. Bot., XX, 124.

Rhizome creeping, the scales dense, subulate, brown, \pm 5 m.M. long. Stipes \pm 55 c.M. long, glossy, brownish-grey. Fronds \pm 65 c.M. long, \pm 20 c.M. broad. Pinnae \pm 13 on each side, distant, nearly erect, linear-lanceolate, narrowed gradually towards both ends, the terminal one \pm 25 c.M. long, 6—7 m.M. broad, with the edge \pm undulate. Texture flaccid, subcoriaceous; surfaces pale-coloured, glaucous, opaque, naked, the upper covered with small, submarginal or rarely scattered, cretaceous dots; veins irregular, hidden; areolae oblong, with few free veinlets. Sori in 1 row on each side of the costae, \pm 10—12 in a row, medial, superficial.

Borneo.

* (217) **P. albidosquamatum**, *Bl.*, Rac, Flor. Bot., I, 115; *Bl.*, Flor. Jav., II, 157, tab. LVII; *Hk.*, Gard. Ferns, tab. XLVII; — var. *varians*; *P. varians*, *Bl.*, Mett., Farng., I, Polypod., 108; *Bl.*, l. c., tab. LVIII.

Rhizome woody, creeping, the scales dense, spreading, filiform-subulate, fine, brown, to 15 m.M. long. Stipes 10—50 c.M. long, firm, erect, naked, glossy. Fronds sometimes simple, commonly pinnate, to 50—60 c.M. long, 15—40 c.M. broad, with several (to 17) \pm distant, linear-lanceolate, sessile or stalked, erecto-patent pinnae on each side, which are 10—25 c.M. long, 1—2 c.M. broad, the apex much acuminate, the edges entire or slightly repand towards the point, the base narrowed. Texture coriaceous; upper surface covered with small, white, cretaceous dots all over or only near the edge; veins usually indistinct. Sori in a single row on each side between the midrib and edge.

Var. varians: Pinnae narrower; main veins distinct.

Malaya.

(218) **P. Moseleyi**, *Bk.*, Journ. of Linn. Soc., Bot., XV, 110.

Stipes 15—25 c.M. long, stramineous, naked. Fronds quadrangular or rhomboidal, simply pinnate, 20—40 c.M. long, 20—30 c.M. broad. Pinnae 4—6-jugate, linear, acuminate, all dilatato-adnate at the base, erecto-patent or subpatulous, 1—2 c.M. broad, the terminal one like the rest but often larger, the margin horny, irregularly repand. Texture subcoriaceous; surfaces naked; costae and main veins glossy and stramineous; areolae immersed, rather large; free veinlets copious, knobbed. Sori round, slightly immersed, in 1 row on each side nearer the margin than the costa.

Moluccas, New Guinea.

* (219a) **P. palmatum**, *Bl.*, Rac., Flor. Btz., I, 114; *Bl.*, Flor. Jav., II, 150, tab. LXIV; *P. crenulatum*, *Kze*, Mett., Farng., I, Polypod., 110; *Pleopeltis palmata*, *Moore*, Bedd., Handb. Ind. Ferns, 568; Ferns Br.I., II, tab. CLVI; — var. *angustatum*; *P. angustatum*, *Bl.*, Mett., l. c., 108; *Bl.*, l. c., 148, tab., LXII; — var. *obtusum*.

Rhizome stout, creeping; scales dense, conspicuous, linear-lanceolate or lanceolate, thin, pale-brown, the base broad, rounded, peltate, the point acute, the edge obscurely toothed or ciliate. Stipes distant, 7—50 c.M. long, erect, naked, glossy. Fronds 15—45 c.M. long, 15—30 c.M. broad or broader, pinnate. Terminal pinna linear or linear-oblong, entire or slightly repand, crenate or toothed; lateral pinnae similar, 1—10 on each side, erecto-patent, 10—25 c.M. long, $\frac{1}{2}$ —4 c.M. broad, lanceolate, acuminate or caudate, narrowed to the base and then dilated again, broadly adnate to the rachis; lower pinnae not rarely \pm stalked, and the barren ones often the broadest. Tex-

ture subcoriaceous; surfaces naked; main veins distinct nearly or quite to the edge: areolae fine, hidden. Sori in 1 row on each side of the costae, \pm medial, superficial. — Young plants with the fronds simple and yet often fructiferous.

Var. angustatum: Pinnae \pm cuneate at the base, the lower ones \pm stalked, the higher sessile, narrowly or hardly adnate to the rachis.

Var. obtusum: Like var. angustatum, but the pinnae \pm blunt, very thick in texture.

Malaya.

(219b) **P. proteus**, *Copel.*, Philipp. Journ., I, Suppl., 164, tab. XXV.

Rhizome creeping, the scales subulate, short-ciliate, with the base peltate, rounded or sometimes hastate. Stipes $\frac{1}{2}$ —17 c.M. long, naked. Fronds polymorphous, simple, 2—3-fid or pinnate with 1—4 pinnae on each side below the terminal one. Simple fronds $\frac{1}{2}$ —5 c.M. long, oblong or lanceolate, obtuse at both ends. The 2—3-fid fronds are intermediate forms between the simple and pinnate ones. Pinnate fronds to 20 c.M. long, nearly as broad. Pinnae lanceolate, acute or acuminate, entire or undulate, sessile, adnate or the lower short-stalked, articulate to the rachis. Texture coriaceous; surfaces naked; main veins distinct nearly to the edge; veins immersed. Sori in 1 row on each side of the costa in the simple fronds and in 1—3 rows on each side in the pinnate forms, slightly immersed.

Luzon.

§ 6. PLEOPELTIS. c. **Chrysopteris**. ***. **Drynariopsis**.

*. Sori in 1 row between the main veins. (220) *P. coronans*.

.. Sori scattered irregularly. (221) *P. heracleum*.

* (220) **P. coronans**, *Wall.*, Mett., Farng., I, Polypod., 121; Hk., Fil. Exot., tab. XCI; *P. conjugatum*, *Bk.* (not *Lam.*), Hk. Bk., Syn. Fil., 566; *Drynaria coronans*, *J. Sm.*, Bedd., Ferns Br.I., I, tab. XIII.

Rhizome very thick, the scales dense, long, linear, crisped, brown. Fronds sessile, arranged in a circle, 60—120 c.M. long, the lower part 10—20 c.M. broad, the base dilated, cordate, the edge narrowed, lobed, passing gradually into the upper part of the frond, which is 50—45 c.M. broad and cut down nearly to the costa throughout into entire, erecto-patent segments $2\frac{1}{2}$ —4 c.M.

broad, linear, acute or acuminate. Texture rigid; surfaces naked; main veins of the segments distinct to the edge, united by transversing veins forming \pm 6 quadrangular areolae on each side of the costulae. Sori in single rows between the main veins near their anterior sides, sometimes confluent.

Malacca; North. India, South. China, Formosa.

* (221) **P. heracleum**, *Kze*, Rac. Flor. Btz., I, 116; Hk., Gard. Ferns, tab. I; *Drynaria Heracleum*, *Moore*, Bedd., Handb. Ind. Ferns, Suppl., 93.

Rhizome stout, fleshy, succulent, creeping, the scales narrow, linear or lanceolate, brown. Fronds sessile, 90—550 c.M. long, the lower part 7—15 c.M. broad, moderately sinuato-lobate, rounded or cordate at the base, the upper part much dilated, 50—100 c.M. broad, cut down to a broadly winged costa into linear-lanceolate. entire, acute lobes $5\frac{1}{2}$ —10 c.M. broad at the base. Texture coriaceous, firm; surfaces naked; veins distinct, with 6—8 \pm irregular, quadrangular areolae on each side of the costulae, enclosing smaller ones with free veinlets. Sori small, copious, scattered irregularly, slightly immersed, often confluent.

Malaya.

§ 6. PLEOPELTIS. c. **Chrysopteris**. ****. **Aglaomorpha**.

(222) **P. Meyenianum**, *Hk.* (not *Schott*), Hk. Bk., Syn. Fil., 366; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 166, A—C; *Aglaomorpha Meyeniana*, *Schott*, Bau. & Hk., Gen. Fil., tab. XCI.

Rhizome very stout, the scales dense, linear, crisped, bright-ferruginous, 1—1 $\frac{1}{2}$ c.M. long. Fronds 60—100 c.M. long, the basal part barren, \pm 15 c.M. broad, shallowly lobed, the central part 20—50 c.M. broad, sterile, cut down nearly to the costa into erecto-patent, linear-oblong, blunt, entire lobes 7—15 c.M. long, 1—3 c.M. broad, the upper part fertile, pinnate, the pinnae 10—20 c.M. long, narrowly linear, sinuous, consisting only of a firm midrib with a row of small, \pm semiorbicular lobes on each side, each with a single sorus. Texture coriaceous; surfaces naked; main veins reaching the margin; areolae copious, with free veinlets.

Philippines.

§ 7. SELLIGUEA. (*)

a. *Fronds simple.*

b. *Texture herbaceous to papyraceous; main veins commonly fine.*

1. *Rhizome scales blackish.*

S. *Fronds narrow, 1—3½ c.M. broad. Sori occupying the whole frond.*

‡. *Areolae regular.* (225a) *P. selliguea.*

‡.‡. *Areolae irregular.* (225b) *P. Bolsteri.*

S.S. *Fronds broader, 3½—5 c.M. broad. Sori confined to the upper ⅓—½ of the frond.* (224) *P. interruptum.*

1.1. *Rhizome scales brown.*

2. *Primary areolae ± regularly subquadrate.*

3. *Fertile fronds ± ½ ×, as long as the barren ones.* (225) *P. regulare.*

3.3. *Fertile fronds not or hardly shorter than the barren ones.*

4. *Fronds cuspidate, short-stalked, ± 20—25 c.M. long.* (226) *P. Hosei.*

4.4. *Fronds acute, rather long-stalked, ± 25—45 c.M. long.* (227) *P. macrophyllum.*

2.2. *Primary areolae irregular.* (228) *P. spurium.*

b.b. *Texture coriaceous or subcoriaceous; main veins mostly prominent.*

c. *Sori superficial.*

d. *Lower part of the frond suddenly narrowed into a decurrent base.* (229) *P. pedunculatum.*

d.d. *Lower part of the frond suddenly or gradually narrowed, if suddenly, the base not or hardly decurrent.*

e. *Barren fronds obtuse, truncate at the base, fertile ones acute at both ends.* (230) *P. Elmeri.*

e.e. *Both barren and fertile fronds ± acute at both ends.* (251a) *P. Feei.*

(251b) *P. calophlebium.*

(252) *P. heterocarpum.*

c.c. *Sori immersed.*

a.a. *Fronds deeply pinnatifid or pinnate.*

‡. *Segments decurrent.* (253) *P. ellipticum.*

‡.‡. *Segments not decurrent.* (254) *P. Maingayi.*

(*) See also § 6, b**, *Pleuridium*, under **a.a.**

* (225a) **P. selliguea**, *Mett.*, Christ, Farnkr. d. Erde, 108; *Gymnogramme membranacea*, *Hk.*, Hk. Bk., Syn. Fil., 588; *Selliguea membranacea*, *Bl.*, Flor. Jav., II, 126, tab. LII, fig. 2; *S. Selliguea*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 992.

Rhizome woody, creeping, the scales small, linear, nearly black. Stipes 1—20 c.M. long, nearly naked. Fronds 10—50 c.M. long, 1—5½ c.M. broad, the point acuminate, the edge entire, the lower part narrowed very gradually. Texture papyraceo-herbaceous; surfaces naked; main veins distinct to the edge, zigzag, very slender; areolae hexagonal. Fertile fronds on the longest stipes. Sori in oblique, single, parallel, often interrupted lines between the main veins.

Malaya; Queensland.

(225b) **P. Bolsteri**, *Copel.*, Philipp. Journ., I, Suppl., 257, tab. IV, A.

Rhizome wide-creeping, stout, clothed with black, aculeate-ovate; deciduous scales. Fronds scattered, sessile or short-stalked, 20—50 c.M. long, 1—2 c.M. broad, narrowed gradually towards both ends. Texture membranaceous; surfaces naked; main veins flexuose, hardly reaching the margin; areolae fine, irregular, with free veinlets. Sori oblong, irregular, or in single rows between the main veins.

Mindanao.

(224) **P. interruptum**, *C. Chr.*, Ind. Fil.; *Gymnogramme acuminata*, *Bk.*, Journ. of Bot., XXVI, 526.

Rhizome wide-creeping, the scales minute, linear, black. Stipes remote, slender, naked, 10—15 c.M. long. Fronds simple, oblong-lanceolate, very acuminate, tapering gradually to the base, 15—20 c.M. long, 5½—5 c.M. broad at the middle. Texture membranaceous; surfaces naked; main veins fine, parallel, erecto-patent, reaching nearly to the edge; areolae with free veinlets. Sori linear, much interrupted, confined to the upper ⅓—½ of the frond.

Borneo.

(225) **P. regulare**, *Mett.*, Ann. Mus. Bot. L. B., II, 225; *Gymnogramme regularis*, *Bk.*, Hk. Bk., Syn. Fil., 588.

Rhizome creeping, the scales ovate to lanceolate, subferruginous or dark-brown. Stipes 0—15 c.M. long. Fronds 15—40 c.M. long, 4—7 c.M. broad, the fertile ones the shortest and narrowest, oblong-elongate, narrowed gra-

dually to both ends, the edge entire. Texture papyraceous; surfaces naked; main veins parallel; areolae copious, subquadrate, divided into smaller, distinct secondary ones with free veinlets. Sori in single, continuous or interrupted lines between the main veins.

Borneo.

* (226) **P. Hosei**, **C. Chr.**, Ind. Fil.; *Gymnogramme campyloneuroides*, *Bk.*, Journ. of Linn. Soc., Bot., XXIV, 261; *Selliguea campyloneuroides*, *Bedd.*, Handb. Ind. Ferns, Suppl., 101.

Rhizome woody, wide-creeping, the scales small, lanceolate, dark-brown. Stipes scattered, short. Fronds simple, 20—25 c.M. long, 6—9 c.M. broad at the middle, oblong, cuspidate, narrowed very gradually to the base. Texture moderately firm; surfaces naked; main veins very distinct, erecto-patent, produced regularly from the midrib to the margin; transverse veins regularly parallel; areolae with free veinlets. Sori in single, interrupted rows reaching from the midrib to the margin.

Borneo, Malacca.

* (227) **P. macrophyllum**, **Reinv.** (not *Mett.*), *Mett.*, Farng., I, Polypod., 111; *Gymnogramme macrophylla*, *Hk.*, *Hk. Bk.*, Syn. Fil., 588.; *Selliguea macrophylla*, *Bl.*, Flor. Jav., II, 127, tab. LIII; *Grammitis macrophylla*, *Bl.*, Enum., 119.

Rhizome creeping, slender, the scales lanceolate, brown. Stipes 5—20 c.M. long, naked, distant. Fronds 25—45 c.M. long, 5—10 c.M. broad, acute, narrowed to both ends, the edge entire. Texture thin, papyraceous; surfaces naked; main veins distinct, slender, zigzag; areolae subquadrangular, with free veinlets. Fertile fronds often longer stalked than the barren ones. Sori in single, continuous or interrupted, parallel lines between the main veins.

Malaya; South. China.

(228) **P. spurium**, **Mett.**, Ann. Mus. Bot. L. B., II, 226; *Gymnogramme spuria*, *Bk.*, *Hk. Bk.*, Syn. Fil., 588.

Rhizome creeping, the scales ovate-acuminate, brown. Stipes \pm 20—25 c.M. long, reddish-brown, margined upwards. Fronds \pm 50 c.M. long, \pm 7—8 c.M. broad, ovate-oblong, the base cuneate, the point bluntish, the edge subrepand. Texture papyraceous; surfaces naked; main veins straight, prominent; areolae irregular, large, divided into smaller ones, and those again so-

metimes subdivided, with free veinlets. Sori in single, interrupted rows between the main veins.

Celebes.

* (229) **P. pedunculatum**, *Mett.*, C. Chr., Ind. Fil.; *P. Hamiltonianum*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 316; *Gymnogramme Hamiltoniana*, *Hk.*, Hk. Bk., Syn. Fil., 589; *Selliguea Hamiltoniana*, *Bedd.*, Ferns Br.I., II, tab. CCXXXIX; *Ceterach pedunculata*, *Hk. & Grev.*, Ic. Fil., tab. V.

Rhizome wide-creeping, woody, the scales linear, dull-brown. Fronds dimorphous. Barren fronds \pm 15—50 c.M. long, \pm 6—10 c.M. broad, elliptical-oblong, entire, acuminate, the lower part suddenly narrowed into a gradually decurrent base, the stipe 5—12 c.M. long. Texture rather thinly coriaceous; surfaces naked; main veins distinct to the edge; areolae copious, with free veinlets. Fertile fronds 10—20 c.M. long, 2—4 c.M. broad, entire or repand, on naked, slender stipes more than 50 c.M. long. Sori in broad, continuous rows, solitary between and parallel to the main veins.

Malacca; North. India, South. China.

(230) **P. Elmeri**, *Copel.*, Polypod. Philipp., 152; — var. *separatum*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 992.

Rhizome creeping, firm; scales ovate, acuminate, the base peltate, the margin scariose. Stipes firm, naked, brownish or blackish, 5—6 c.M. long. Sterile fronds deltoid-oblong, \pm 10 c.M. long, 6—8 c.M. broad, obtuse, entire or subsinuate, the base truncate. Texture firmly coriaceous; surfaces naked; main veins parallel, conspicuous, reaching the margin; veins hidden. Fertile fronds \pm 10 c.M. long, \pm 4 c.M. broad, acute, subentire, cuneate at the base, on stipes 15—20 c.M. long; texture coriaceous. Sori narrow, continuous, reaching the costa and the margin.

Var. separatum: Sori not continuous, interrupted, forming 2 rows of round sori on each side of the costa, one costal, the other marginal.

Luzon.

* (251a) **P. Feei**, *Mett.*, *Farngr.*, I, Polypod., 110; *Gymnogramme Feei*, *Hk.*, *Rac.*, *Flor. Btz.*, I, 74; *Selliguea Feei*, *Bory*, *Bl.*, *Flor. Jav.*, II, 125, tab. LI; *Bedd.*, *Ferns Br.I.*, II, tab. CLI; — var. *vulcanicum*; *P. vulcanicum*, *Bl.*, *Mett.*, l. c., 111; *Bl.*, l. c., 144, tab. LVI, fig. 2; — var. *caudiforme*;

P. caudiforme, Bl., Mett., l. c., 110; Bl., l. c., 146, tab. LIV, fig. 2; *Selliguea caudiformis*, J. Sm., Bedd., Handb. Ind. Ferns, 590.

Rhizome creeping, the scales lanceolate-linear, light-brown. Stipes $2\frac{1}{2}$ —40 c.M. long, firm, erect, naked, more or less glossy, pale or brown, those of the fertile fronds the longest. Fronds subdimorphous, 5—25 c.M. long, 2—10 c.M. broad, ovate-oblong, the fertile ones commonly the longest and narrowest, often much the narrowest, acute, entire, narrowed at the base. Texture coriaceous; surfaces naked; main veins distinct; areolae hidden. Sori *not immersed*, linear, continuous, solitary between and parallel to the main veins. not reaching either the midrib or the edge, distant or contiguous.

Var. vulcanicum: Stipes of the fertile fronds not always longer than those of the barren ones. Fronds smaller, \pm ovate, bluntish, uniform, or the fertile ones narrower. Sori interrupted so as to form several parallel rows of round, oblong or linear-oblong sori.

Var. caudiforme: Fertile fronds much narrowed, long-acuminate, subcaudate. Sori large, round, solitary between the main veins.

Malaya, Polynesia.

(251b) **P. calophlebium**, Copel., Philipp. Journ., II, 140, tab. III, A.

Rhizome wide-creeping, clothed with squarrose, broadly lanceolate, fulvo-ferruginous scales, which are peltate and castaneous at the base. Stipes erect, naked, 5—10 c.M. long, those of the fertile fronds the longest. Barren fronds ovate, \pm 14 c.M. long, \pm 5 c.M. broad, long-acuminate, the edge brown, rigid, the base rounded (acute in Copeland's table). Texture coriaceous; surfaces naked; main veins stout, straight, reaching the margin, erecto-patent; veins hidden beneath, visible above. Fertile fronds contracted, linear, \pm 16 c.M. long, \pm 7—11 m.M. broad, broadest near the base, narrowed gradually towards the apex, the base cuneate, the edge entire or crenulate (undulato-crenate in Copeland's drawing). Sori 1—2 between the oblique main veins, the pairs often confluent.

Mindoro.

* (252) **P. heterocarpum**, Mett., Farng., I, Polypod., 110; *Gymnogramme heterocarpa*, Hk. (not Bl.), Rac., Flor. Btz., I, 74; *Grammitis heterocarpa*, Bl., Enum., 118; *Selliguea heterocarpa*, Bl., Flor. Jav., II, 125, tab. LIII, fig. 1; — var. *interruptum*; *P. vulcanicum*, Christ (not Bl.), Farnkr. d. Erde, 108; — var. *abbreviatum*; *P. caudiforme*, Christ (not Bl.), l. c.

Rhizome creeping, woody, the scales dense, linear-lanceolate, brown. Stipes 4—25 c.M. long, firm, erect, naked, those of the fertile fronds often the longest. Fronds 7—30 c.M. long, $1\frac{1}{2}$ —5 c.M. broad, the fertile ones not much narrower than the barren, lanceolate, acute or acuminate, the edge entire, the base more or less gradually narrowed. Texture coriaceous; surfaces naked, glossy; main veins distinct; veins hidden. Sori *immersed*, long, linear, solitary between and parallel to the main veins, continuous or slightly interrupted.

Var. interruptum: Sori distinctly interrupted, forming rows of roundish or oblong sori.

Var. abbreviatum: Fertile fronds much narrowed, subcaudate. Sori roundish or oblong, 1—2 between the main veins.

Malacca, Banca, Java, Celebes.

* (255) **P. ellipticum**, *Thb.*, Christ, Farnkr. d. Erde., 107; *Gymnogramme elliptica*, *Bk.*, Hk. Bk., Syn. Fil., 589; *G. decurrens*, Hk., Spec. Fil., V, 161; *Selliguea elliptica*, *Bedd.*, Handb. Ind. Ferns, 592; *S. decurrens*, *Pr.*, *Bedd.*, Ferns Br.I., I, tab. CL; *Grammitis decurrens*, *Wall.*, Hk. & Grev, Ic. Fil., tab. VI.

Rhizome woody, creeping, firm, the scales adpressed. Stipes scattered, 50—40 c.M. long, firm, erect, naked, pale. Fronds 50—50 c.M. long, 15—25 c.M. broad, cut down nearly or quite to the rachis into 4—6 linear-oblong or lanceolate, distant, sessile, opposite, acute segments narrowed towards the base, mostly connected by the decurrent, oblique bases, which form a wing to the rachis; longest segments 1—4 c.M. broad. Texture herbaceous; rachis and surfaces naked; costal areolae large, with smaller, irregular included ones with free veinlets. Sori linear, oblique, nearer the midribs than the edge.

Malacca, Philippines; Formosa, Japan, North. China, North. India, Queensland.

(254) **P. Maingayi**, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 518; *Gymnogramme Maingayi*, *Bk.*, Hk. Bk., Syn. Fil., 517; *Selliguea Maingayi*, *Bedd.*, Handb. Ind. Ferns, 592.

Stipes \pm 45 c.M. long, dull-brown, naked. Fronds deltoid, to 30 c.M. long, with \pm 5 oblong-lanceolate pinnae. Largest pinnae \pm 15 c.M. long, $4\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, slightly repand, narrowed gradually at base, the lowest pair distant. Texture membranaceous; surfaces naked; main veins distinct to the edge; areolae copious, small, with free veinlets. Sori abundant, irregular, punctiform or confluent in lines or curves.

Malacca.

72. CYCLOPHORUS, *Desvaux.*

Sori round, rarely oblong, and then oblique or sometimes subparallel to the costa. No indusium.

Rhizome creeping. Stipes articulate to the rhizome. Fronds simple, \pm entire in malayan material; surfaces covered with scattered or more or less densely crowded or interlaced stellate hairs, which are fastened by a central point, those of the upper surface often caducous; veins anastomosing, hidden; areolae with free included veinlets. Sori sometimes quite hidden among the tomentum. Hk. Bk., Syn. Fil., tab. V, fig. 48, k—l; Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 168. (*Niphobolus*).

Arrangement of the sections.

- § 1. **NIPHOBOLUS.** Venation as in *Campyloneuron*, i. e. areolae \pm regularly subquadrangular, with 2 or more included, excurrent, simple or forked free veinlets. Diels, l. c., fig. 165, H.
- § 2. **NIPHOPSIS.** Venation as in *Pleopeltis*, i. e. areolae \pm irregular, the free veinlets spreading in various directions. Diels, l. c., fig. 164, B—E.

§ 1. **NIPHOBOLUS.**

a. *Tomentum adpressed, often very thin.*

b. *Fronds dimorphous, the fertile ones mostly narrower and often longer than the barren.*

‡. *Fertile region of the frond with the tomentum as thin as in the barren part.*

*. *Rhizome scales black. Stellate hairs of the under surface scattered.* (1) *C. pachydermus*?

.. *Rhizome scales ferruginous. Stellate hairs of the under surface crowded.* (2) *C. rupestris.*

‡.‡. *Fertile region of the frond with the tomentum thicker and looser than in the barren part.*

×. *Rhizome scales adpressed. Fertile fronds to 1 c.M. broad.* (3) *C. adnascens.*

×.×. *Rhizome scales spreading at the point. Fertile fronds 1½—2 c.M. broad.* (4) *C. varius.*

b.b. *Fronds uniform or nearly so.*

- c.** *The stellate hairs of the under surface scattered.*
- +**. *Main veins hidden. Sori in 5—6 rows between the midrib and margin.* (5) *C. anetioides.*
 - ++.** *Main veins distinct. Sori in 10—20 rows between the midrib and margin.* (6) *C. sphaerostichus.*
- c.c.** *The stellate hairs of the under surface crowded or interlaced, but adpressed.*
- d.** *Main veins hidden.*
 - ‡.** *Fronds contracted in the fertile region. Sori immersed in the tomentum.* (7) *C. macropodus.*
 - ‡‡.** *Fronds hardly contracted in the fertile region. Sori not immersed in the tomentum.* (8) *C. acrostichoides.*
 - d.d.** *Main veins distinct.*
 - e.** *Fronds small, under 30 c.M. long.*
 - f.** *Rhizome scales adpressed, obscurely peltate at the base, rather sessile.* (9) *C. pannosus.*
 - f.f.** *Rhizome scales spreading, distinctly peltate at the base.* (10) *C. heteractis.*
 - e.e.** *Fronds larger, 60—90 c.M. long.* (11) *C. princeps.*
- a.a.** *Tomentum loose, woolly, thick.*
- 1.** *Fronds small, under 6 c.M. long. (*)*
 - ×**. *Barren fronds suborbicular, cordate or rounded at the base, on stipes 1—2 m.M. long.* (12) *C. nummularifolius.*
 - ×.×**. *Barren fronds obovate, cuneate at the base, on stipes ± 10 m.M. long.* (15) *C. obovatus.*
 - 1.1.** *Fronds larger, more than 6 c.M. long when fully developed.*
 - 2.** *Texture thin or weak.*
 - +**. *Fronds linear, ± ½ c.M. broad.* (14) *C. lanuginosus.*
 - ++.** *Fronds lanceolate, 2—8 c.M. broad.* (15) *C. penangianus.*
 - 2.2.** *Texture thick, firm.*
 - 3.** *Fronds mostly 8—30 c.M. long.*
 - 4.** *Fronds narrow, to ± 5 m.M. broad. Sori in 1—3 rows on each side of the costa.* (16) *C. rasamalae.*
 - 4.4.** *Fully developed fronds ± 1 c.M. broad, or broader.*
 - 5.** *Sori in 2—4 rows on each side of the costa.*
 - ‡.** *Upper surface without depressed dots and minute blackish scales.* (17) *C. flocciger.*

(*) See also under **C. rasamalae**, *C. Chr.*

‡.‡. Upper surface purveyed with depressed dots, which are covered with minute, deciduous, blackish, calcareous scales. (18) *C. porosus*.

5.5. Sori in 5—8 rows on each side of the costa.

6. Upper surface purveyed with depressed dots and minute, deciduous, blackish, calcareous scales. (19) *C. sticticus*.

6.6. Upper surface not purveyed with depressed dots and minute blackish scales. (20) *C. mollis*.

3.3. Fully developed fronds more than 30 c.M. long. Sori in 5—8 rows on each side of the costa. (21) *C. albicans*.

(1) **C. pachydermus**, *C. Chr.*, Ind. Fil.; *Polypodium pachydermum*, *Bk.*, Journ. of Linn. Soc., Bot., XV, 108.

Rhizome wide-creeping, woody, flexuose, the scales minute, adpressed, lanceolate, black, mixed with a brown powder. Stipes naked, 1—2½ c.M. long. Fronds dimorphous, the barren ones oblanceolate-oblong, obtuse, 2½—5 c.M. long, 1—2 c.M. broad, cuneate at the base, the fertile ones ligulate, 7—10 c.M. long. Texture thick, coriaceous; upper surface naked, lower clothed with a thin tomentum visible by magnifying power, the hairs with a distinct bare space round each; veins immersed. Sori occupying the upper ½ or ⅔ of the frond, superficial, 6—8 in erecto-patent rows between midrib and edge.

Kei Islands, Moluccas.

* (2) **C. rupestris**, *C. Chr.*, Ind. Fil.; *Polypodium rupestre*, *R. Br.*, *Mett.*, *Farng.*, I, *Polypod.*, 125; *P. confluens*, *Hk.* (not *R. Br.*), *Hk. Bk.*, *Syn. Fil.*, 549; *Niphobolus rupestris*, *Spr.*, *Gies.*, *Niph.*, 169; *Hk. & Grev.*, *Ic. Fil.*, tab. XCIII.

Rhizome wide-creeping, slender, rather filiform, branched, densely clothed with pale-ferruginous, linear-subulate, hair-pointed scales, peltate at the base. Fronds scattered, dimorphous. Barren fronds short-stalked or sessile, orbicular, oblong, rhomboidal or spatulate, ± 1—2 c.M. long, ± ½—1 c.M. broad, mostly broadest at or above the middle, the apex rounded, the base cuneate or rounded. Texture very thick; upper surface often soon naked, lower clothed with dense but adpressed, moderately woolly, brownish tomentum; veins quite hidden. Fertile fronds linear or ligulate, 2—15 c.M. long,

2—5 m.M. broad, the apex blunt or rounded, the lower $\frac{1}{3}$ — $\frac{1}{2}$ cuneate, narrowed gradually into the short stipe; upper surface soon naked, subinduplicato-canaliculate when dry, lower densely clothed with subadpressed, ferruginous tomentum. Sori rather large, in 1—2 irregular rows on each side of the costa, immersed, but elevated above the tomentum, occupying the upper $\frac{1}{2}$ — $\frac{2}{3}$ of the fertile frond, but sometimes covering the whole under surface. — Plants gathered in New Guinea by Dr. Versteeg, which may be a form of this species, have the barren and fertile fronds sessile, the barren oblong, rounded at the apex, cuneate, rounded or cordate at the base; fertile fronds with the young sori deeply immersed, and the cavities closed by a small, roundish, lid-like, deciduous lateral outgrowth of leaf tissue covered with large, globose tufts of densely crowded, very long-stalked, ferruginous stellate hairs, the tufts \pm 2—5 m.M. each way.

? *New Guinea; Australia, New Zealand.*

* (3) **C. adnascens**, *Desv.*, Berl. Mag., V, 500; *Polypodium adnascens*, Sw., Rac., Flor. Btz., I, 98; *P. caudatum*, Mett., Farng., I, Polypod., 126; *P. carnosum*, Mett., l. c., 124; *P. Koenigii*, Bl., Flor. Jav., II, 50; *Niphobolus adnascens*, Klf., Hk., Gard. Ferns, tab. XIX; Gies., Niph., 196; *N. caudatus*, Klf., Bl., l. c., 56, tab. XXII; *N. carnosus*, Bl., l. c., 50, tab. XIX, *N. elongatus*, Bl., l. c., 52, tab. XX; — var. minor, Christ, MS. in Herb. Bog.

Rhizome wide-creeping, the scales dense, peltate, dark-coloured, ovate or lanceolate, with the base rounded, the apex blunt, acute or acuminate, the edge ciliate, pale-brown or whitish. Stipes 1—5 c.M. or more apart, $\frac{1}{2}$ —5 c.M. long, firm. Fronds dimorphous. Barren fronds oblong, linear or spatulate, 2—12 c.M. long, 1—2 c.M. broad, the apex blunt, the edge entire, the base cuneate or gradually narrowed. Texture rigidly coriaceous, thick; costa distinct; veins hidden; upper surface naked or nearly so, lower thinly tomentose, the stellate hairs abundant but not crowded, caducous. Fertile fronds linear, longer than the barren ones, 8—50 c.M. long, 4—10 m.M. broad, with the edges more or less recurved, on stipes 1—8 c.M. long, the tomentum thicker, woolly (the hairs densely crowded) in the soriferous part of the frond. Sori small, often crateriform, occupying the upper $\frac{1}{2}$ — $\frac{2}{3}$ of the frond, immersed, hidden in the tomentum when young. — Ferns with the habit of *Drymoglossum heterophyllum*, C. Chr. in the smaller forms.

Var. minor: Fertile fronds sessile, \pm 6 c.M. long, 3—6 m.M. broad in the fertile region.

Trop. Asia, Polynesia; — var. minor: *Batu Islands (Raap).*

* (4) **C. varius**, *Gaud.*, Freyc., Voy. Ur. Phys., Bot., 564; *Polypodium varium*, *Mett.*, Rac., Flor. Btz., I, 100; *Niphobolus varius*, *Klf.*, Gies., Niph., 208; *Bl.*, Flor. Jav., II, 54, tab. XXI.

Rhizome wide-creeping, slightly branched, the scales oblong-lanceolate, brown, with the base peltate, the apex acuminate and \pm spreading, the margin ciliate. Stipes scattered, 1—2 c.M. long. Fronds dimorphous. Barren fronds \pm 8—12 c.M. long, 2—4 c.M. broad, lanceolate, obtuse. Texture coriaceous; upper surface naked, lower covered with scattered, adpressed stellate hairs; costa conspicuous; main veins hardly visible. Fertile fronds \pm 15—25 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, the soriferous region often narrowed and the under surface densely covered there with deciduous, \pm loose and woolly tomentum. Sori \pm immersed in shallow, indistinct cavities, \pm crateriform, closely placed in 6—8 rows between the midrib and edge, occupying the upper part of the frond, hidden in the tomentum when young.

Malaya; Polynesia.

(5) **C. anetioides**, *C. Chr.*, Ind. Fil.; *Niphobolus anetioides*, *Christ*, Ann. Btz., XIX, 58.

Rhizome wide-creeping, the scales subulate-acuminate, yellow-brown. Stipes distant, $\frac{1}{2}$ —1 c.M. long. Fronds ovate-oblong to lanceolate, 6—10 c.M. long, to $\frac{2}{2}$ c.M. broad, obtuse to acuminate, cuneately narrowed at the base, often lobed there with 1—3 round lobes. Texture papyraceo-coriaceous; upper surface bright-green, the stellate hairs short, whitish; under surface pale, the hairs sparingly scattered, reddish-brown, adpressed; costa ebeneous, conspicuous; main veins hidden, erecto-patent, 8—15 on each side, with 5—6 rows of longitudinal areolae on each side of the costa. Fertile fronds narrower. Sori not immersed in the tomentum, occupying the upper part of the frond, covering the whole space between the midrib and edge, placed in 2 rows between the main veins, with 5—6 sori in a row.

Celebes.

(6) **C. spaerostichus**, *C. Chr.*, Ind. Fil.; *Polypodium sphaerostichum*, *Mett.*, Hk. Bk., Syn. Fil., 550; *Niphobolus sphaerostichus*, *Copel.*, Polypod. Philipp., 114.

Rhizome woody, wide-creeping, the scales ovate, acuminate, brown, crisped. Stipes 5—8 c.M. long, firm, erect. Fronds uniform, 10—13 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, oblong-lanceolate, acuminate, the base narrowed suddenly.

Texture coriaceous; under surface very thinly matted: main veins distinct to the edge. Sori confluent, covering the whole under surface, placed in 10—20 rows between the midrib and edge.

Celebes, Luzon.

(7) **C. macropodus**, *C. Chr.*, Ind. Fil.; *Polypodium macropodum*, *Bk.*, Journ. of Linn. Soc., Bot., XV, 108.

Rhizome woody, short-creeping, the scales minute, adpressed, deltoid, black, mixed with a brown powder. Phyllopoies 7—10 m.M. long, scaly like the rhizome. Stipes bordered very nearly to the base. Fronds uniform, ligulate, 10—20 c.M. long, 1—2 c.M. broad, the fertile region contracted, the point acute, the base narrowed very gradually. Texture coriaceous, thick; upper surface naked, lower clothed with thin tomentum; veins hidden. Sori occupying the upper half of the frond, minute, immersed in a pale-brown tomentum, placed in erecto-patent rows between midrib and edge, and each showing distinctly on the upper surface of the frond, 6—8 in a row.

Aru Islands.

* (8) **C. acrostichoides**, *Pr.*, Epim., 130; *Polypodium acrostichoides*, *Forst.*, Rac. Flor. Btz., I, 98; *Niphobolus acrostichoides*, *Rich.* (not *J. Sm.*), *Copel.*, Polypod. Philipp., 113; *Gies.*, Niph., 213; *Bedd.*, Ferns Br.I., I, tab. LXXXI; *N. puberulus*, *Bl.*, Flor. Jav., II, 57, tab. XXIII; *N. fissus*, *Bl.*, l. c., 58, tab. XXIV (abnormal form); — var. *gracilis*, *Copel.*, Leaflet, I, 234.

Rhizome creeping; scales roundish or oblong, rarely acute, peltate, adpressed or subadpressed, dark-coloured, the margin paler or whitish when young, ± entire. Stipes 5—7 c.M. long, deciduously tomentose with adpressed stellate hairs. Fronds uniform, 20—60 c.M. long, 1—3 c.M. broad, ligulate or linear, the apex bluntish or subacute, the edge entire, the base narrowed gradually. Texture coriaceous; upper surface naked or nearly so, lower tomentose, the stellate hairs greyish or brownish, often white when young, crowded but adpressed; costa conspicuous; veins hidden. Sori small, close, numerous, crateriform, occupying the upper $\frac{1}{3}$ — $\frac{2}{3}$ of the frond, slightly immersed in shallow cavities, which are not or obscurely prominent on the upper surface, not hidden in the tomentum.

Var. gracilis: Stipes shorter; fronds hardly 1 c.M. broad, less coriaceous; sori in 3—4 rows.

Malaya; Ceylon, Polynesia, Queensland.

* (9) **C. pannosus**, *C. Chr.*, Ind. Fil.; *Polypodium pannosum*, *Mett.*,

Hk. Bk., Syn. Fil., 512: *Niphobolus pannosus*, Bedd., Gies., Niph., 105; *N. lingua*, Bedd., Ferns S.I., tab. CCXL.

Rhizome wide-creeping, the scales lanceolate, acuminate, \pm toothed, adpressed, obscurely peltate at the base, rather sessile. Stipes $2\frac{1}{2}$ —15 c.M. long, those of the sterile fronds often the longest. Fronds nearly uniform, 5—15 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, narrowed towards both ends. Texture subcoriaceous; upper surface at length naked, lower tomentose, the stellate hairs close, adpressed, pale-brown, with long branches; costa and main-veins distinct. Sori densely crowded, 5—6 in a row between the main veins, ceasing a space from the edge. — Fertile fronds very small in Beddome's drawing.

? Java; Tenasserim, Ceylon, Burmah.

* (10) **C. heteractis**, C. Chr., Ind. Fil.; *Polypodium Heteractis*, Mett., Linn., XXXVI, 140; *Niphobolus Heteractis*, J. Sm., Gies., Niph., 153; *N. Lingua*, Spr. (not J. Sm.), Copel., Polypod. Philipp., 113, p. p.; Bedd., Ferns S.Br.I., Suppl., tab. CCCLXXXV. (*)

Rhizome wide-creeping, the scales dense, spreading, lanceolate, ferrugineous, with the base peltate, \pm lobed, the edge ciliate towards the apex. Stipes \pm 4—10 c.M. long, firm, erect. Fronds 8—16 c.M. long, $2\frac{1}{2}$ — $4\frac{1}{2}$ c.M. broad, ovate or oblong-lanceolate, the apex often cuspidate, the base cuneate or rounded. Texture coriaceous; upper surface at length naked, lower matted with adpressed, ferrugineous tomentum; branches of the stellate hairs mostly short, some of them long, the longer ones interlaced; costa and main veins distinct. Sori large, rather prominent, immersed among the tomentum when young, placed in close, rather irregular rows of 4—6 between the main veins.

Philippines; North. India.

(11) **C. princeps**, C. Chr., Ind. Fil.; *Polypodium princeps*, Mett., Ann. Mus. Bot. L. B., II, 232; *Niphobolus princeps*, Gies., Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 325.

Rhizome short; scales lanceolate-subulate, with the base dilated, the margin subentire, the apex divided into 3—4 ciliae. Fronds approximate, subsessile, 60—90 c.M. long, 5—10 c.M. broad, spatulate-lanceolate, short-acuminate, the margin entire, subcallous. Texture coriaceous; upper surface covered with small, cretaceous dots, lower thinly tomentose, the stellate hairs much adpressed, their branches short, or long and interlaced; costa prominent;

(*) See Gies., Niph., 157—158.

main veins very distinct, distant, oblique. Sori numerous, contiguous, occupying the upper $\frac{1}{3}$ — $\frac{1}{2}$ of the frond or more.

New Guinea.

* (12) **C. nummularifolius**, *C. Chr.*, Ind. Fil.; *Polypodium nummulariaefolium*, *Mett.*, *Rac.*, *Flor. Btz.*, I, 101; *Acrostichum nummularifolium*, *Sw.*, *Bl.*, *Flor. Jav.*, II, 53, tab. XI, fig. 1—2; *Niphobolus nummulariaefolius*, *J. Sm.*, *Gies.*, *Niph.*, 179; *Bedd.*, *Ferns Br.I.*, II, tab. CCCXX; — var. *rufus*, *v. A. v. R.*, *Bull. Dép. Agr. I. N.*, 1908, XXI, 8.

Rhizome slender, creeping, the scales roundish and peltate at the base, long, linear, acuminate at the point, \pm ciliate, erect or erecto-patent, white or afterwards brownish or blackish. Fronds dimorphous. Barren fronds on stipes 1—2 m.M. long, placed closely in 2 opposite rows, \pm adpressed to the substratum, suborbicular, $\frac{1}{2}$ —2 c.M. each way, the base obliquely cordate, truncate or rounded. Texture fleshy; stipe and upper surface covered with scattered, deciduous or persistent stellate hairs, lower side densely clothed with white or brownish tomentum. Fertile fronds oblong, $1\frac{1}{2}$ —4 c.M. long, 5—7 m.M. broad, on stipes 2—20 m.M. long, scattered, the apex rounded, the base cuneate. Sori in 3—4 rows on each side, \pm confluent, occupying the whole under surface, immersed among the tomentum.

Var. rufus: Scales of the rhizome bright-reddish-brown, copiously ciliate; tomentum bright-reddish-brown, dense.

Malaya; North. India; — var. rufus: Batu Islands (Raap).

(15) **C. obovatus**, *v. A. v. R.*; *C. nummularifolius*, *C. Chr.*, var., *C. Chr.*, Ind. Fil.; *Polypodium obovatum*, *Mett.*, *Rac.*, *Flor. Btz.*, I, 101; *Acrostichum obovatum*, *Bl.*, *Flor. Jav.*, II, 55, tab. XI, fig. 3; *Niphobolus obovatus*, *Kze*, *Bot. Zeit.*, 1848, 120.

Very near *C. nummularifolius*, *C. Chr.*, but the sterile fronds obovate, to $1\frac{1}{2}$ c.M. long, to 7 m.M. broad, cuneate at the base, on stipes \pm 1 c.M. long; fertile fronds linear-elliptical, to 5 c.M. long, under $5\frac{1}{2}$ m.M. broad, on stipes to 3 c.M. long.

Java.

(14) **C. lanuginosus**, *C. Chr.*, Ind. Fil.; *Niphobolus lanuginosus*, *Gies.*, *Niph.*, 190.

Rhizome wide-creeping; scales greyish-brown, the base peltate, oblong, the apex long, finely acuminate. Stipes scattered, 2—3 c.M. long. Fronds

linear, to \pm 50 c.M. long, \pm $\frac{1}{2}$ c.M. broad, the apex bluntish or acuminate the edge entire, recurved, the base narrowed. Texture weak; stipe and upper surface densely tomentose when young, but at length naked; under surface densely tomentose; midrib prominent beneath, but covered by the tomentum; main veins fine. Sori small, hidden among the tomentum when young.

Philippines.

* (15) **C. penangianus**, *C. Chr.*, Ind. Fil.; *Polypodium penangianum*, *Hk.*, Hk. Bk., Syn. Fil., 352; *Niphobolus penangianus*, *Hk.*, Bedd., Handb. Ind. Ferns, 352; *Hk.*, Icon. Pl., III, tab. CCIII; Bedd., Ferns Br.I., I, tab. CXXI; Bau. & Hk., Gen. Fil., tab. LXXXIII; Gies., Niph., 97.

Rhizome short-creeping, the scales lanceolate, long-acuminate, brown. Stipes none or short, approximate. Fronds oblong-lanceolate, 20—70 c.M. long, 2—8 c.M. broad, the apex acuminate, the base very gradually narrowed, the edge commonly irregularly undulate and repand. Texture papyraceous or flaccid; upper surface at length naked, lower clothed with fine, silky tomentum; main veins oblique, connected by nearly straight veins forming 10—12 rows of narrow, longitudinal, subquadrangular areolae between the midrib and edge; areolae with 3—6 sori in each, the rows close, forming a broad mass confined to the upper half of the frond, not reaching the margin; capsules commonly mixed with stellate hairs.

Java, Sumatra, Malacca; Burmah.

* (16) **C. rasamalae**, *C. Chr.*, Ind. Fil.; *Polypodium Rasamalae*, *Rac.*, Flor. Btz., I, 99; *Niphobolus Rasamalae*, *Gies.*, Niph., 187.

Rhizome wide-creeping, black, with the scales pale-brown or ferruginous, ovate or lanceolate, erecto-patent or recurved, finely long-acuminate, the base peltate, rounded. Stipes 1—10 c.M. apart, \pm 1—5 c.M. long, brown, clothed with deciduous tomentum, at length naked. Fronds linear, 8—50 c.M. long, to \pm 5 m.M. broad, the apex bluntish, the edge recurved, the base narrowed gradually into the stipe. Texture coriaceous, thick; upper surface at length naked, lower densely tomentose with short-stalked (long-branched) and long-stalked (short-branched) hairs; veins hidden; costa prominent, but covered by the tomentum. Sori roundish, small, close or confluent, in 1—5 rows between the midrib and edge, 1-serial in the narrowest, 2—3-serial in the broadest forms, immersed in the tomentum when young.

Forma minor: Fronds very short, $2\frac{1}{2}$ —5 c.M. long, short-stalked.

Java, Sumatra.

* (17) **C. flocciger**, *Pr.*, Epim., 151; *Polypodium floccigerum*, *Mett.*, *Rac.*, *Flor. Btz.*, I, 101; *Antrophyum niphoboloides*, *Kze.*, *Hk.*, *Spec. Fil.*, V, 174; *Niphobolus flocciger*, *Bl.*, *Copel.*, *Polypod. Philipp.*, 115; *Bl.*, *Flor. Jav.*, II, 61, tab. XXVI.

Rhizome wide-creeping; scales narrowly lanceolate, ferrugineous, the base adpressed, the apex acuminate, spreading, entire. Stipes 0—5 c.M. long, those of the fertile fronds often the longest. Fronds 15—30 c.M. long, 1—2½ c.M. broad, narrowed gradually towards both ends, the edge entire. Texture coriaceous; upper surface soon naked or nearly so, lower densely covered with cinnamon or yellow-brown, loose tomentum; veins hidden. Sori close, numerous, placed in 2—4 rows on each side of the costa, immersed in the tomentum, at length confluent. — Too near *C. albicans*, *Pr.*

Malaya; North. India.

* (18) **C. porosus**, *Pr.*, Epim., 150; *Polypodium porosum*, *Wall.*, *Mett.*, *Farn.*, I, *Polypod.*, 128; *P. fissum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 351, p. p.; *Niphobolus fissus*, *Bedd.* (not *Bl.*), *Copel.*, *Polypod. Philipp.*, 115, p. p.; *Bedd.*, *Handb. Ind. Ferns*, 550; *N. porosus*, *Pr.*, *Tent. Pterid.*, 202; *Bedd.*, *Ferns S. I.*, tab. CLXXXIII; *N. Mannii*, *Gies.*, *Niph.*, 107.

Rhizome short, creeping, the scales ferrugineous, narrowly lanceolate, long-acuminate, entire, subspreading. Stipes 0—2 c.M. long below the winged part, subapproximate. Fronds 15—35 c.M. long, 1—4 c.M. broad, lanceolate, linear-lanceolate or subspathulate, bluntish to acuminate, narrowed very gradually towards the base so as to form a wing to the stipe. the edge entire, irregularly wavy or purveyed with short or long, subulate, lobe-like outgrowths. Texture coriaceous; surfaces at first wholly tomentose with ferrugineous or greyish, densely crowded stellate hairs, the upper side at length naked, purveyed with small depressed dots, which are covered with deciduous, blackish, calcareous scales, the stellate hairs purveyed with an erect central branch at the point of union of the spreading ones; venation quite sunk; areolae with 2—5 soriferous, excurrent veinlets. Sori rather large, confined to the upper half or occupying the whole under surface of the frond, placed in 2—4 ± irregular rows on each side of the costa, at length scarcely elevated above the tomentum.

Malacca, Philippines; China, North. India.

* (19) **C. sticticus**, *C. Chr.*, *Ind. Fil.*; *Polypodium sticticum*, *Mett.*, *Farn.*, I, *Polypod.*, 128; *P. fissum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 351, p. p.; *Ni-*

phobolus sticticus, Kze, Gies., Niph., 155; *N. fissus*, Bedd. (not Bl.), Copel., Polypod. Philipp., 115, p. p.

Rhizome creeping; scales rigid, dark-brown, lanceolate, suberect, acuminate, the margin pale and more or less ciliate. Fronds approximate, spatulate, lanceolate or linear-lanceolate, 15—50 c.M. long, 1—2 c.M. broad, the apex varying from blunt to acuminate, the base narrowed very gradually so as to form a winged stipe \pm 6 c.M. long, the edge entire. Texture coriaceous; veins hidden; upper surface naked or provided with scattered, adpressed, greyish or brownish hairs and small depressed dots, which are covered with deciduous, blackish, calcareous scales; under surface densely tomentose; areolae with 5—4 soriferous, excurrent veinlets. Sori rather large, confined to the upper half of the frond, placed in 5—8 rows on each side of the costa, immersed among the tomentum, but at length distinct. — Clearly distinguished from *C. porosus*, Pr. by the rhizome scales.

Philippines; South. China, Ceylon.

(20) **C. mollis**, Pr., Epim., 151; *Polypodium molle*, Mett., Farng., I, Polypod., 128; *Niphobolus mollis*, Kze, Bot. Zeit., 1848, 121.

Rhizome creeping, the scales moderately large, membranaceous, reddish-brown, not closely adpressed, narrow, lanceolate, long-acuminate. Stipes \pm 1½ c.M. long. Fronds 20—50 c.M. long, 2—2½ c.M. broad, lanceolate, acuminate, the base narrowed gradually. Texture \pm coriaceous; upper surface covered with dirty-white, the lower with reddish-brown tomentum; costa prominent beneath, at length naked, flexuose; main veins slightly visible. Sori superficial, crowded in the upper part of the frond, not reaching the margin and the apex, placed in 6—8 rows.

Java, Bali, Timor; Irawaddi River.

* (21) **C. albicans**, Pr., Epim., 151; *Polypodium albicans*, Mett., Rac., Flor. Btz., I, 101; *Niphobolus Blumeanus*, Kze, Bot. Zeit., 1848, 120; *N. albicans*, Bl., Flor. Jav., II, 60, tab. XXV; *N. glaber*, Klf., Bl., l. c., 62, tab. XXVII.

Rhizome wide-creeping, the scales dense, pale-brown or ferruginous, lanceolate, sometimes crisped, erecto-patent or spreading, peltate and adpressed at the base. Stipes 2—4 c.M. apart, 4—15 c.M. long, covered with white or brown, often deciduous tomentum. Fronds 50—80 c.M. long, 1—5 c.M. broad, narrowed towards both ends, the edge entire, \pm recurved. Texture very coriaceous, upper surface at length naked, lower densely clothed with white or brownish tomentum; costa less densely tomentose, like the stipe;

main veins often distinct to the edge, venation not rarely partially irregular. Sori in 4—6 close rows between the main veins so as to form 5—8 rows between the midrib and edge, sunk in the tomentum when young.

Java.

§ 2. NIPHOPSIS.

- a.** Sori in 1 row on each side of the costa.
- b.** Fronds small, the barren ones under 3 c.M. long, under $\frac{3}{4}$ c.M. broad. (22) *C. Lauterbachii*.
- b.b.** Fronds larger, the barren ones 3—60 c.M. long, $\frac{3}{4}$ —5 c.M. broad.
- c.** Rhizome scales ciliate. (25) *C. confluens*.
- c.c.** Rhizome scales entire.
- d.** Fronds 4—30 c.M. long, 2—5 c.M. broad; rhizome scales decoloured at the apex; sori slightly immersed in shallow pits. (24a) *C. angustatus*.
- d.d.** Fronds 30—60 c.M. long, 1—2 c.M. broad; rhizome scales pale-brown at the apex; sori superficial. (24b) *C. samarensis*.
- a.a.** Sori not in 1 row on each side of the costa, often densely crowded. (*)
- 1.** Barren fronds generally under 20 c.M. long.
- 2.** Fronds blunt or rounded at the apex. (25) *C. abbreviatus*.
- 2.2.** Fronds acute to caudate at the apex.
- 3.** Barren fronds 5—8 c.M. long, on stipes 1—4 c.M. long. (26) *C. Warburgii*.
- 3.3.** Barren fronds 11—18 c.M. long, on stipes 7—17 c.M. long. (27) *C. Sarasinorum*.
- 1.1.** Fully developed barren fronds more than 20 c.M. long.
- α.** Barren fronds matted beneath with scattered stellate hairs. (28) *C. Christii*.
- α.α.** Barren fronds clothed beneath with densely crowded or interlaced stellate hairs.
- β.** Sori large.
- +. .** Fertile fronds much exceeding the barren ones. (29) *C. distichocarpus*.
- +.+. .** Fertile fronds not much exceeding the barren ones. (30) *C. asterosorus*.

(*) See also *C. albicans*, Pr.

$\beta.\beta.$ *Sori small or minute.*

$\gamma.$ *Rhizome wide-creeping; fronds distant, solitary.*

(51) *C. argyrolepis.*

$\gamma.\gamma.$ *Rhizome short-creeping; fronds not solitary, aggregate or approximate.*

$\delta.$ *Stellate hairs without a spine-like central branch; leaf-tissue with palisade cells (*).*

$\epsilon.$ *Fronds stalked; stipes to 30 c.M. long, subterete or laterally compressed, not angled, grooved in front.*

(52) *C. stigmosus.*

$\epsilon.\epsilon.$ *Fronds sessile or stalked; stipes not subterete, angled, not grooved, flattened in front.*

(53) *C. Beddomeanus.*

$\delta.\delta.$ *Stellate hairs with a very long, erect, spine-like central branch; leaf-tissue without palisade cells.*

(54) *C. splendens.*

* (22) **C. Lauterbachii**, *C. Chr.*, Ind. Fil.; *Niphobolus Lauterbachii*, *Christ, Schum. & Laut., Flor. deut. Sch.ggeb., 142; Gies., Niph., 221, fig. 20.*

Rhizome wide-creeping, filiform, the scales subulate, peltate, pale-coloured, thin, scattered. Fronds scattered, dimorphous. Barren fronds spatulate, ovate or lanceolate, $\pm 1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long, ± 5 m.M. broad, obtuse, entire, on stipes 1—2 c.M. long. Texture rigidly coriaceous; under surface with the stellate hairs small, adpressed, scarcely visible, greyish-green; veins inconspicuous; areolae oblong. Fertile fronds linear-ligulate, 6—8 c.M. long, 2—3 m.M. broad, acute to caudate, sinuato-crenate, on stipes 1—2 c.M. long. Sori submarginal, occupying the crenations, the stalks of the capsules immersed in deep cavities.

New Guinea.

(23) **C. confluens**, *C. Chr.*, Ind. Fil.; *Polypodium confluens*, *R. Br., Warb., Mons., I, 61 (not Hk. Bk., Syn. Fil., 349); P. glabrum, Mett., Hk. Bk., Syn. Fil., 356; Mett., Farnng., I, Polypod., tab. III, fig. 6—8; Niphobolus confluens, Spr., Gies., Niph., 172.*

(*) See *Gies., Niph., 78—79, fig. 15.*

Rhizome filiform, wide-creeping, branched, the scales lanceolate, acuminate, ciliate, ferruginous, peltate at the base. Stipes $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. long. Fronds subdimorphous. Sterile fronds lanceolate or oblong, 5—9 c.M. long, 8—12 m.M. broad, the apex blunt. Texture coriaceous; upper surface naked, lower densely clothed with adpressed, subferruginous stellate hairs, at length not rarely quite naked; veins hidden. Fertile fronds \pm linear-lanceolate or linear, longer than the barren ones. Sori large, not hidden in the tomentum, commonly confined to the apex or the upper $\frac{1}{5}$ — $\frac{1}{2}$ of the frond, in 1 row on each side, often confluent so as to form a ridge on both sides of the costa. — Near *C. angustatus*, *Desv.*, but the sori more confluent.

New Guinea; Australia, Norfolk Island, Lord Howe's Island, New Caledonia, New Zealand.

* (24a) **C. angustatus**, *Desv.*, Berl. Mag., V, 300; *Polypodium angustatum*, *Sw.*, Rac., Flor. Btz., I, 100; *Niphobolus angustatus*, *Spr.* (not *J. Sm.*), Copel., Polypod. Philipp., 115; *Gies.*, Niph., 195; *Hk.*, Gard. Ferns, tab. XX; *Bedd.*, Ferns S.I., tab. CLXXXV; *N. sphaerocephalus*, *Hk. & Grev.*, Ic. Fil., tab. XCIV; *Pleopeltis angustata*, *Pr.*, *Bedd.*, Handb. Ind. Ferns, 551.

Rhizome firm, woody, wide-creeping, branched, the scales with a roundish, peltate, adpressed, brown or black, coriaceous base and a very long, linear, finely acuminate, whitish, decoloured, membranaceous, entire, spreading, deciduous or persistent point. Stipes scattered, 1—10 c.M. long, covered like the upper surface of the frond with fugacious tomentum. Fronds lanceolate, 4—30 c.M. long, 2—5 c.M. broad, narrowed towards both ends, the apex bluntish, the edge entire. Texture coriaceous; under surface densely clothed with adpressed, white or subferruginous tomentum; costa conspicuous; main veins hidden, slightly visible when dry. Fertile fronds commonly longer than the barren ones, \pm 20—50 c.M. long, 1—2 c.M. broad in the fertile region, the barren part often broader. Sori large, \pm immersed, round or oblong, oblique or subparallel to the costa, in 1 row on each side, nearer the \pm recurved edge than the midrib, sometimes confluent, slightly prominent on the upper surface, occupying the upper $\frac{1}{6}$ — $\frac{5}{6}$ of the frond.

Malaya; Polynesia, North. India.

(24b) **C. samarensis**, *C. Chr.*, Ind. Fil.; *Polypodium samarense*, *Mett.*, *Hk. Bk.*, Syn. Fil., 556; *Niphobolus samarensis*, *Fée*, Copel., Polypod. Philipp., 115; *Gies.*, Niph., 191; *Gyrosorium samarense*, *Pr.*, (oldest name).

Rhizome firm, wide-creeping; scales ovate-lanceolate, entire, the base pel-

tate, with a fuscous centre, the point long, linear, acuminate, pale-brown (not white), spreading. Stipes firm, erect, $2\frac{1}{2}$ —10 c.M. long. Fronds linear-lanceolate, 50—60 c.M. long, 1—2 c.M. broad, the soriferous upper part linear, contracted, the base narrowed. Texture coriaceous; upper surface naked or slightly tomentose when young, lower densely coated with dirty-white or greyish-ferruginous tomentum; veins hidden. Sori large, oblong, placed in 1 row on each side of the midrib, confined to the narrowed upper $\frac{1}{3}$ — $\frac{1}{2}$ of the frond, confluent.

Philippines.

* (25) **C. abbreviatus**, *C. Chr.*, Ind. Fil.; *Polypodium abbreviatum*, *Mett.*, *Rac.*, *Flor. Btz.*, I, 98; *Niphobolus abbreviatus*, *Zoll.*, *Gies.*, *Niph.*, 160; *N. Lingua*, *Spr.*, p. p. (*)

Rhizome wide-creeping; scales peltate, oblong-lanceolate, acuminate, brown, the margin pale, ciliate, the ciliae fine, crisped, deciduous. Stipes 14—20 c.M. long, densely covered with adpressed stellate hairs. Fronds dimorphous. Barren fronds roundish or ovate, 7—10 c.M. long, 3—7 c.M. broad, the apex blunt or rounded, the base suddenly narrowed into a short, decurrent wing. Texture coriaceous; upper surface covered with scattered, fugacious, adpressed stellate hairs and often also with white cretaceous dots; under side densely tomentose, the tomentum adpressed; costa and main veins conspicuous; areolae irregular, with few free veinlets. Fertile fronds more elongate, $\pm \frac{1}{3} \times$ as broad as long, narrowed gradually to the stipe and the bluntish apex. Sori densely crowded, at length confluent, occupying the upper $\frac{2}{3}$ of the frond. — The scales at the bases of the stipes are pale, blunt, deciduously ciliate, rather large.

Java, Sumatra.

(26) **C. Warburgii**, *C. Chr.*, Ind. Fil.; *Niphobolus Warburgii*, *Gies.*, *Niph.*, 165; *N. Lingua*, *Spr.*, p. p. (*)

Rhizome wide-creeping, branched; scales peltate and rounded or emarginate at the base, the point long-acuminate, the margin long-ciliate. Stipes 1—4 c.M. long, 1—5 c.M. apart. Sterile fronds commonly 5—8 c.M. long and 2—5 c.M. broad, oblong, the apex acuminate, the base subcuneate; fertile fronds longer and broader, on longer stipes. Texture coriaceous; upper surface naked; under side covered with adpressed, whitish or brownish to-

(*) See *Gies. Niph.*, 157—158.

mentum; costa conspicuous beneath; main veins hidden; areolae irregular, with free veinlets. Sori numerous, densely crowded, occupying the whole or nearly the whole under surface of the frond.

Celebes.

(27) **C. Sarasinorum**, *C. Chr.*, Ind. Fil.; *Niphobolus Sarasinorum*, *Gies.*, *Niph.*, 162; *N. Lingua*, *Spr.*, p. p. (*)

Rhizome wide-creeping; scales dense, black-brown, adpressed, the younger ones narrow, lanceolate, long-acuminate, with the apex crisped, the margin long-ciliate. Sterile fronds oblong, 11—18 c.M. long, \pm 5 c.M. broad, the apex acuminate to caudate, the base suddenly narrowed and slightly decurrent, the stipe 7—17 c.M. long. Fertile fronds sometimes longer and narrower. Texture coriaceous; upper surface naked, lower covered with light-brown, adpressed tomentum; costa and main veins distinct; areolae irregular, with free veinlets. Sori small, dense, occupying the whole under surface or only the upper half of it.

Celebes.

* (28) **C. Christii**, *C. Chr.*, Ind. Fil.; *Niphobolus Christii*, *Gies.*, *Niph.*, 140; *N. Lingua*, *Spr.*, p. p. (*)

Rhizome wide-creeping, the scales brown, obovate and peltate at the base, narrowed to a long, lanceolate-acuminate, spreading, often capillary, crisped point, the margin pale and slightly ciliated towards the point. Barren fronds ovate or oblong, 10—12 c.M. long, to 5 c.M. broad below the middle, on stipes 3—10 c.M. long, narrowed gradually from the broadest part to a long-acuminate point, the base cuneate or rounded; fertile fronds lanceolate, narrowed gradually to both ends, 10—12 c.M. long, to $2\frac{1}{2}$ c.M. broad, on stipes \pm 15 c.M. long. Texture coriaceous; costa and main veins distinct; upper surface naked, lower covered with scattered, adpressed stellate hairs in the barren, but more woolly in the fertile fronds. Sori large, crowded, at length confluent, occupying the whole under surface.

Borneo.

* (29) **C. distichocarpus**, *C. Chr.*, Ind. Fil.; *Polypodium distichocarpum*, *Mett.*, *Hk. Bk.*, *Syn. Fil.*, 352; *Niphobolus distichocarpus*, *Gies.*, *Diels*, in *Engl. & Prantl*, *Nat. Pfl. Fam.*, I⁴, 325.

(*) See *Gies.*, *Niph.*, 157—158.

Rhizome creeping, scaly. Barren fronds linear-oblong, 20—30 c.M. long, 3—5 c.M. broad, narrowed towards both ends, on stipes to 12½ c.M. long; fertile fronds linear-lanceolate, 55—65 c.M. long, 2—4 c.M. broad, on stipes to 5 c.M. long. Texture coriaceous; upper surface soon naked, lower densely clothed with soft, brown tomentum; costa prominent; main veins slightly prominent; areolae hidden, with the included veinlets simple or forked, free or anastomosing. Sori large, in 2 rows between the main veins, occupying the upper ¼—½ of the frond, forming 3—6 rows on each side of the costa in the broader part of the frond; capsules mixed with stellate hairs, especially when young. — Baker and Diels placed this species in *Niphobolus*, but the specimens from Java (Leg. de Vriese) and Sumatra (Leg. Korthals) determined by Burck (as *P. distichocarpum*, Mett.) and a plant sent from Malacca and cultivated in the Buitenzorg gardens have the venation decidedly *irregular*.

Java, Sumatra, Malacca.

* (50) **C. asterosorus**, *C. Chr.*, Ind. Fil.; *Polypodium asterosorum*, *Bk.*, Journ. of Bot., XVIII, 214.

Rhizome slender, wide-trailing, clothed with large, lanceolate, pale-reddish-brown, membranaceous scales. Stipes scattered, naked, pale-brown, 5—8 c.M. long. Fronds lanceolate, entire, 20—50 c.M. long, 2—2½ c.M. broad above the middle, acute, narrowed gradually towards the base. Texture coriaceous; upper surface naked, lower whitish, matted with loose, soft, reddish-brown tomentum; main veins obscure, erecto-patent, enclosing copious minute areolae; free veinlets simple or forked. Sori large, round, superficial, densely pilose, in erecto-patent rows of 5—6 between the midrib and margin, filling up the whole of the upper part of the frond and mixed with copious soft, brown stellate hairs. — Near *C. distichocarpus*, *C. Chr.*

Sumatra.

(31) **C. argyrolepis**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 991.

Rhizome wide-creeping, string-shaped, branched, black-purple, densely clothed with subulate, aristate, scariose, squarrose, glistening-white scales. Stipes distant, solitary, ± 5 c.M. long, flattened, slender, white-woolly, soon naked. Fronds linear-lanceolate, 18—40 c.M. long, 2—2½ c.M. broad, the apex elongate, acuminate, the edge entire, involute when dried, the base ovate, not or hardly decurrent. Texture carnosu-coriaceous, flaccid; costa distinct; veins rather occult; areolae elongate, irregular, with free veinlets; upper surface

rather naked, lower densely covered with adpressed, white, minute stellate hairs intermixed with larger, scattered, fulvous ones. Sori occupying the narrowed apical part of the frond, confluent, also covering the costa, forming a dark-brown mass 5—18 c.M. long.

Luzon, Mindoro.

* (52) **C. stigmatosus**, *Desv.*, Berl. Mag., V, 501; *Polypodium stigmatosum*, *Sw.*, Rac., Flor. Btz., I, 102; Hk. Bk., Syn. Fil. 550, p. p.; *Niphobolus venosus*, *Bl.*, Flor. Jav., II, 65, tab. XXVIII, fig. 1; *N. stigmatosus*, *Moore*, Gies., Niph., 115; *Bedd.*, Handb. Ind. Ferns, 528, p. p.

Rhizome short-creeping; scales long, narrow-lanceolate, finely acuminate, the margin entire or with few short teeth, the base not peltate. Stipes always present, not rarely to 20—50 c.M. long, not angled, subterete or laterally compressed especially in dried material, not angled, tomentose, the upper (inner) side furrowed, not flattened. Fronds lanceolate, 20—60 c.M. long, 3—7 c.M. broad, the apex acute or acuminate, the base cuneately narrowed. Texture coriaceous; costa and main veins distinct beneath; areolae hidden, in 10—16 rows on each side of the costa; upper surface densely tomentose, afterwards naked, often purveyed with numerous depressed dots, which are commonly covered with deciduous, white, dot-like cretaceous scales; under surface clothed with interlaced, long- and short-branched stellate hairs forming a dense coat. Sori numerous, small, close, placed in irregular rows, hidden among the tomentum when young, occupying the upper $\frac{2}{3}$ — $\frac{4}{5}$ of the frond.

Malaya; Ceylon, South. India.

* (53) **C. Beddomeanus**, *C. Chr.*, Ind. Fil.; *Polypodium stigmatosum*, *Sw.*, Hk. Bk., Syn. Fil., 550, p. p.; *Niphobolus stigmatosus*, *Moore*, *Bedd.*, Handb. Ind. Ferns, 528, p. p.; *N. Beddomeanus*, *Gies.*, Niph., 101; *N. costatus*, *Pr.*, *Bedd.*, Ferns Br.I., I, tab. CXX.

Rhizome short-creeping, scaly at the extremity, the scales dark-brown, narrowed from a broad base into a long, acuminate, hair-like point, the base not peltate. Stipes 0—18 c.M. long, deciduously tomentose, not laterally compressed, the upper (inner) side flat, angled, not furrowed, the lower (outer) side convex or angular. Fronds lanceolate, 30—90 c.M. long, 5—7 c.M. broad or broader, the apex acuminate, the lower part narrowed very gradually to the base. Texture coriaceous, rather thin; costa and main veins distinct beneath; areolae hidden, in 8—14 rows on each side of the costa; upper surface covered with long-branched, interlaced stellate hairs,

afterwards naked, not rarely purveyed with minute depressed dots, which are covered with deciduous, white, dot-like cretaceous scales; under surface tomentose like the upper, but the hairs intermixed with short-branched ones, the tomentum pale-brown, or white when young, forming a close coat. Sori numerous, minute, immersed among the tomentum when young, occupying \pm the upper half of the frond, reaching to 1—3 m.M. or more from the margin, placed in irregular, contiguous rows.

Malaya; South. China, North. India.

(54) **C. splendens**, *C. Chr.*, Ind. Fil.; *Polypodium splendens*, *Hk.*, Spec. Fil., V, 52 (not V, 95); *P. nitens*, *Bk.*, *Hk. Bk.*, Syn. Fil., 553; *Nipholobus splendens*, *J. Sm.*, *Gies.*, *Niph.*, 99.

Rhizome short-creeping, the scales narrowed from a sessile, truncate or rounded, broad base into a subulate point. Fronds approximate, sessile or subsessile, 45—90 c.M. long, 6—10 c.M. broad, broadest in the upper half, narrowed very gradually from there to the base, the edge entire. Texture subcoriaceous; upper surface white-cobwebby, at length naked, often purveyed with white, dot-like cretaceous scales; under surface densely clothed with long- and short-branched, brown stellate hairs, the short-branched ones purveyed with a long, erect, spine-like central branch (*Gies.*, *Niph.*, fig. 8, E); costa and main veins distinct. Sori small, numerous, close, immersed among the tomentum, not reaching the margin. — Among the long- and the short-branched hairs there are many intermediate forms with both short and long branches, but without central spines.

Philippines.

73. DRYNARIA, *J. Smith.*,

Sori round or punctiform, placed on the points of union of the anastomosing veins. No indusium.

Rhizome creeping or ascending. Fronds articulate to the rhizome, dimorphous. Barren fronds (cup-leaves) rigid, mostly oakleaf-like, sessile, soon becoming brown, commonly cordate at the base, collecting quantities of humus into which the roots grow; fertile fronds large, stalked, green, pinnatifid or pinnate, the segments mostly articulate; veins anastomosing copiously; primary areolae often quadrangular, secondary with free included veinlets spreading in various directions. *Diels*, in *Engl. & Prantl*, *Nat. Pfl.Farm.*, I⁴, fig. 171, A.

Trop. Asia, Australia, & Polynesia.

Arrangement of the sections.

- § 1. **EUDRYNARIA.** Cup-leaves rather flat, at best slightly convex. Fertile fronds pinnatifid, or pinnate with the segments contiguous and adnate with a broad base.
- § 2. **PORONEMA.** Cup-leaves as in § 1. Fertile fronds pinnate, the segments stalked, or sessile with a cuneate or acute base.
- § 3. **THAYERIA.** Cup-leaves induplicate or convolute.

 § 1. **EUDRYNARIA.**

- a.* Sori in 1 row on each side of the costulae (central ribs of the segments).
- b.* Segments quite entire. (1) *D. pleuridioides.*
- b.b.* Segments crenate or slightly toothed. (2) *D. propinqua.*
- a.a.* Sori not as in *a.*
- 1.* Sori in 2 rows between the main veins of the segments.
- 2.* Segments connected by a broad wing. (3) *D. descensa.*
- 2.2.* Segments connected by a narrow wing. (4) *D. quercifolia.*
- 1.1.* Sori scattered irregularly. (5) *D. sparsisora.*

* (1) ***D. pleuridioides*, Pr.** (not *J. Sm.*), Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 530; *Polypodium pleuridioides*, Mett., Rac., Flor. Btz., I, 117; *P. Willdenowii*, Bl. (not *Bory*), Flor. Jav., II, 156, tab. LXVI; — var. *amboinensis*.

Rhizome thick, the scales dense, narrow, linear or lanceolate. ciliate. Cup-leaves 10—30 c.M. long, 7—20 c.M. broad, lobed \pm $\frac{1}{2}$ -way down, the lobes blunt, linear-oblong. Fertile fronds 45—100 c.M. long, 20—40 c.M. broad, with a distinct stipe, the lobes reaching down nearly or quite to the costa, 1—2 c.M. broad, blunt, acute or acuminate, quite entire; lower lobes decurrent upon the stipe. Texture subcoriaceous; surfaces naked; areolae copious, with free veinlets. Sori in 1 row on each side, very near the midribs, more or less immersed.

Var. amboinensis: Cup-leaves shallowly lobed; lobes of fertile fronds \pm 5 c.M. broad, united at the base by a broad wing.

Java; — var. *amboinensis*: *Amboyna* (Boerlage).

* (2) **D. propinqua**, *J. Sm.*, *Bedd., Handb. Ind. Ferns*, 339; *Ferns Br.I.*, II, tab. CLX; *Polypodium propinquum*, *Wall., Hk. Bk., Syn. Fil.*, 367; *P. Willdenowii*, *Hk. (not Bory)*, *Gard. Ferns*, tab. XXXV; *Pleopeltis Parishii*, *Bedd., l. c.*, I, tab. CXXV; — var. *sumatrana*.

Rhizome stout, creeping, the scales linear-lanceolate, reddish. Cup-leaves 10—25 c.M. long, 7—10 c.M. broad, cut $\frac{1}{2}$ — $\frac{3}{4}$ -way down into bluntish or acute lobes. Fertile fronds 40—90 c.M. long, often 50 c.M. broad, with a distinct stipe, the lobes 10—15 c.M. long, 1—2 c.M. broad, reaching down nearly or quite to the costa, crenate, slightly toothed or undulato-crenate; lowest lobes decurrent. Texture subcoriaceous; surfaces naked; areolae copious, with free veinlets. Sori in a single row on each side, near the midribs. — Very near *D. pleuridioides*, *J. Sm.*

Var. sumatrana: Sori very large, nearly medial.

Java; North. India, South. China, Fernando Poo; — var. *sumatrana*: *Sumatra* (Leg. . . . ?).

(3) **D. descensa**, *Copel.*, *Philipp. Journ.*, III, 36.

Near *D. quercifolia*, *J. Sm.* Rhizome 5—10 c.M. thick, the apex mammiform, the scales peltate, aristate, caducous. Cup-leaves 5—10 c.M. long, dry, hard, ovate, slightly crenato-lobate, brown, or grey beneath, the base cordate with the basal lobes overlapping. Stipes of fertile fronds winged, the wing consisting of the lower, reduced lobes. Fertile fronds 40—60 c.M. long, deeply lobed or pinnatifid. Lobes to 10—20 c.M. long, $\pm 1\frac{1}{4}$ — $2\frac{1}{2}$ c.M. broad, acuminate. crenate, slightly narrowed at the base, connected by a wing ± 4 c.M. broad. Texture papyraceous; surfaces naked.

Luzon.

* (4) **D. quercifolia**, *J. Sm.*, *Bedd., Handb. Ind. Ferns*, 341; *Diels, in Engl. & Prantl, Nat Pfl.Fam.*, I⁴, fig. 171, A; *Bedd., Ferns S.I.*, tab. CLXXXVII; *Polypodium quercifolium*, *L., Rac., Flor. Btz.*, I, 118.

Rhizome stout, ascending, densely clothed with narrow, linear-subulate, eroso-ciliate scales, which are narrowed gradually and rather regularly from the rounded, peltate base to the point. Cup-leaves $7\frac{1}{2}$ —30 c.M. long, 5—15 c.M. broad, rigid, bluntly lobed often $\frac{1}{2}$ -way down. Fertile fronds 60—150 c.M. long, 50—40 c.M. broad, deeply pinnatifid nearly to the costa, on stipes to 30 c.M. long. Lobes entire, erecto-patent, linear-lanceolate, blunt, acute or acuminate, 1—4 c.M. broad, the lowest not rarely short, often suddenly decurrent so as to form a wing reaching not rarely to the base of the stipe.

Texture rigidly coriaceous; surfaces naked; main veins (of the lobes) distinct \pm to the edge; primary areolae \pm quadrangular, in 4—6 rows, enclosing copious finer secondary ones with few included free veinlets. Sori rather large, in 2 rows between the main veins.

Trop. Asia & Australia, South. China, Fiji.

* (5) **D. sparsisora**, *Moore*, C. Chr., Ind. Fil.; *D. Linnaei*, *Bedd.*, Handb. Ind. Ferns, 543; Ferns Br.I., II, tab. CCCXV; *Polypodium Linnei* (*Linnaei*), *Bory*, Rac., Flor. Btz., I, 118; *P. sparsisorum*, *Desv.*, (oldest name).

Rhizome stout, the scales ovate, short, adpressed, black or brown, peltate, eroso-ciliate, with a suddenly narrowed, long, linear-subulate, often recurved, fugacious apex. Cup-leaves bluntly lobed, \pm similar to those of *D. quercifolia*, *J. Sm.* Fertile fronds 45—90 c.M. long, 15—35 c.M. broad, stipitate, cut down nearly to the costa into erecto-patent, lanceolate to linear, entire lobes; lower lobes decurrent so as to form a shorter or longer wing to the stipe. Texture rigidly coriaceous; surfaces naked; main veins distinct to the edge; areolae large, \pm quadrangular, divided into numerous finer ones. Sori small, scattered irregularly.

Malaya; Ceylon, Trop. Australia, Fiji.

§ 2. PORONEMA.

* (6) **D. rigidula**, *Bedd.*, Handb. Ind. Ferns, 344; Ferns Br.I., II, tab. CCCXIV; *Polypodium rigidulum*, *Sw.*, Rac., Flor. Btz., I, 117; *P. diversifolium*, *R. Br.*, Hk., Gard. Ferns, tab. V; *P. Gaudichaudii*, *Bory*, Bl., Flor. Jav., II, 158, tab. LXVII.

Rhizome stout, creeping, the scales linear-lanceolate, narrow, ciliate, erecto-patent, brown, with the base adpressed. Cup-leaves sessile, 15—30 c.M. long, 7—10 c.M. broad, the base sinuate, the edge deeply incised towards the costa into blunt, oblong or linear lobes. Fertile fronds 60—120 c.M. long, 15—45 c.M. broad, pinnate, long-stalked, the stipe and rachis grey or brown, densely clothed with a greyish tomentum consisting of deciduous, copiously ciliate, peltate scales, at length naked, the stipe purveyed with 2 opposite rows of few distant, large glands (rudimentary pinnae). Pinnae to 2½ c.M. or more apart, 4—10 m.M. broad, stalked or sessile, with a cuneate base, articulate to the rachis, the edge entire, crenate or toothed, the apex acuminate. Texture subcoriaceous; surfaces naked; main veins indistinct, rather wanting;

areolae copious; free veinlets few. Sori immersed, in 1 row \pm medial between the midribs and the edge.

Trop. Asia, Australia & Polynesia.

§ 3. THAYERIA.

- *. *Cup-leaves deeply lobed or pinnatifid.* (7) *D. nectarifera.*
 (8) *D. cornucopia.*
 *. *. *Cup-leaves slightly undulate, subentire.* (9) *D. involuta.*

(7) ***D. nectarifera***, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 530; *Polypodium nectariferum*, *Bk.*, *Becc.*, *Mal.*, II, 247, tab. LXV; *Thayeria nectarifera*, *Copel.*, *Philipp. Journ.*, I, Suppl., 166.

Rhizome stout, woody, tortuose, scaly towards the extremity, the scales adpressed, orbicular, peltate. Cup-leaves ovate, pinnatifid, induplicate, the lobes oblong, obtuse. Fertile fronds stipitate, oblong-lanceolate, 45—60 c.M. long, simply pinnate. Pinnae numerous, the lower ones adnate, lanceolate, covered beneath with copious dot-like glands; upper pinnae narrowed or contracted, linear, torulose, soriferous, \pm similar to those of *Polypodium Meyenianum*, *Hk.*

New Guinea, ? Luzon.

(8) ***D. cornucopia***, *v. A. v. R.*, *Bull. Dép. Agr. I. N.*, 1908, XXI, 8; *Thayeria cornucopia*, *Copel.*, *Philipp. Journ.*, I, Suppl., 165, tab. XXVIII.

Rhizome wide-creeping, scandent, often descendent, clothed with small, lanceolate, subpeltate, acuminate, adpressed scales. Cup-leaves to 50 c.M. long, cornet-shaped, i. e. with the large basal lobes convolute and overlapping so as to form an oblique, humus-collecting cup placed on a horizontal branch of the rhizome 1—6 c.M. long, which passes through the sinus between the lobes and ends in the base of the cornet; lower part scariose, brown, upper green, deeply pinnatifid, the segments deciduous. — Fertile fronds unknown.

Mindanao.

* (9) ***D. involuta***, *v. A. v. R.*, *Bull. Dép. Agr. I. N.*, 1908, XXI, 8, tab. IV.

Rhizome long, woody, ascending or climbing, flexuose, densely scaly; scales suberose, the base fuscous, broadly ovate, peltate, the margin paler, scariose,

the apex narrowed suddenly into a long, linear-subulate, recurved, deciduous point. Cup-leaves distant, 6—10 c.M. long, induplicate or convolute, scarioso-coriaceous, with the outer side subopaque, the inner glossy, the base cordate, the apex rounded, the side edges slightly undulate and overlapping so as to form rather short, closed, subtubular organs quite enclosing distant, rooting portions of the rhizome, not solitary on lateral branches of the rhizome. Fertile fronds distant, similar to those of *D. quercifolia*, *J. Sm.*, but often much smaller. Sori placed in 2 rows of 4—7 between the main veins of the lobes.

Borneo (Teysmann, Nos. 8577 & 8590; Teuscher).

74. DRYOSTACHYUM, *J. Smith.*

Sori oblong or subquadrangular, close, confined to the contracted upper pinnae, in 1 row on each side of the midribs. No indusium.

Rhizome stout, creeping. Fronds sessile, articulate to the rhizome, pinnate or subpinnate; lower pinnae broad, barren; veins anastomosing copiously; areolae with free included veinlets. Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 170, A—C.

Malaya.

* ***D. splendens***, *J. Sm.*, Journ. of Bot., III, 399; Bau. & Hk., Gen. Fil., tab. XCV; Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 170, A—C; *Polypodium splendens*, *Hk.*, Hk. Bk., Syn. Fil., 567; *Hk.*, Spec. Fil., V, 95 (not V, 52); *Drynaria splendens*, *Bedd.*, Ferns Br.I., II, tab. CCCXVI; — var. *pilosum*; *Dryost. pilosum*, *J. Sm.*, Kze, Farnkr., I, 159, tab. LXI.

Rhizome stout, creeping, the scales linear-lanceolate, grey or brown. Fronds ± sessile, 60—90 c.M. long, 20—30 c.M. or more broad, the lowest part brown, rigid, narrow, ± 4 c.M. broad, forming a shallowly lobed wing, the central part sterile, cut down nearly or quite to the costa into 4—8 suberecto-patent, linear-oblong, acute or acuminate lobes or pinnae 10—20 c.M. long, 2½—5 c.M. broad, dilated at the base; highest part fertile, pinnate, the pinnae distant, ligulate, 10—20 c.M. long, ± 1 c.M. broad. Texture firmly herbaceous; surfaces naked; main veins distinct; areolae copious, with few free veinlets. Sori oblong or subquadrangular, in 1 row on each side of the costae, filling up the whole under surface of the fertile pinnae except the costa and the main veins.

Var. pilosum: Upper surface of the fertile pinnae covered with brown, hairy, elliptical patches corresponding with the receptacles of the sori. Capsules crowned with \pm 6 bristles.

Malaya; — var. *pilosum*: *Philippines*.

75. DRYMOGLOSSUM, *Presl.*

Sori long, linear, 1 on each side, parallel to the costa, continuous or interrupted. No indusium.

Rhizome creeping. Stipes articulate to the rhizome. Fronds dimorphous, simple, entire; veins anastomosing. Hk. Bk., Syn. Fil., tab. VI, fig. 58; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 160.

Tropical and extra-tropical.

§. *Rhizome scales lanceolate, rhomboidal or ovate, toothed, fimbriate or lacinate.*

‡. *Sori nearer the midrib than the edge.* (1) *D. carnosum.*

‡‡. *Sori nearer the edge than the midrib, often submarginal.*

(2) *D. heterophyllum.*

§.§. *Rhizome scales subulate, entire.*

×. *Barren fronds under 4 c.M. long; fertile fronds long-stalked.*

(3) *D. rigidum.*

×.×. *Barren fronds \pm 8 c.M. long; fertile fronds not long-stalked.*

(4) *D. novo-guineae.*

* (1) ***D. carnosum***, *J. Sm.* (not *Hk.*), Hk. Bk., Syn. Fil., 397 (excl. var.); Bau. & Hk., Gen. Fil., tab. LXXVIII, A; Bedd., Ferns Br.I., I, tab. LV; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 160; *Nothochlaena carnosa*, *Wall.*, (oldest name).

Rhizome creeping, long, filiform, clothed with peltate, lanceolate, toothed scales. Barren fronds 5—6 c.M. long, on stipes 1—2½ c.M. long, lanceolate-oblong, rather obtuse or subacute. Texture coriaceous, thick and fleshy; areolae with free veinlets. Fertile fronds linear-spathulate, \pm 5—8 c.M. long, \pm 6 m.M. broad, subacute, narrowed at the base, on stipes 3—4½ c.M. long. Sori linear, continuous, forming a line nearer the midrib than the

edge, soon filling up nearly the whole under surface of the frond, covered by numerous stalked peltate scales when young.

Amboyna, Philippines; Formosa, North. India.

* (2) **D. heterophyllum**, *C. Chr.*, Ind. Fil.; *D. piloselloides*, *Pr.*, Rac., Flor. Btz., I, 61; *Bedd.*, Ferns S.I., tab. LV; *Hk.*, Gard. Ferns, tab. XLVI; *D. carnosum*, *Rac.* (not *Hk.*), l. c., 62; *Niphobolus nummularifolius*, *Bedd.*, l. c., tab. CLXXXVI; *Acrostichum heterophyllum*, *L.*, (oldest name).

Rhizome wide-creeping, slender, the scales dense, roundish, oblong or lacinate. Fronds dimorphous. Barren fronds roundish or oblong, sessile or short-stalked, 1—5 c.M. long, 1—2 c.M. broad, the apex rounded, the edge entire, the base more or less obliquely truncate or cuneate. Texture coriaceous or fleshy; surfaces covered with scattered, deciduous stellate hairs, especially when young; veins immersed; areolae with free veinlets. Fertile fronds 5—10 c.M. long or longer, 5—8 m.M. broad, on stipes to 2 c.M. long. Sori in a broad, submarginal line, not rarely filling the whole under surface of the fertile fronds in the narrowest forms; capsules mixed a few stellate, pedicellate hairs.

Trop. Asia.

* (5) **D. rigidum**, *Hk.*, *Hk. Bk.*, Syn. Fil., 598; *Hk.*, Icon. Pl., X, tab. CMXCVI.

Rhizome firm, wide-creeping, the scales dense, lanceolate-subulate, brownish. Fronds dimorphous, the barren ones $1\frac{1}{2}$ — $5\frac{1}{2}$ c.M. long, oblong, the apex rounded, the base \pm cuneate, the edge entire, thickened, on firm stipes to \pm $2\frac{1}{2}$ c.M. long; fertile fronds 10—15 c.M. long, nearly semiterete, \pm 5 m.M. broad, narrowed into a stipe \pm 10—15 c.M. long. Texture very thick and rigid; surfaces naked; main veins hidden; areolae without free veinlets. Sori sunk in a deep groove between the midrib and edge.

Borneo.

(4) **D. novo-guineae**, *Christ*, Schum. & Laut., Flor. deut. Sch. geb., 157.

Rhizome wide-creeping, the scales at first white, afterwards reddish-brown, subulate, dense, subadpressed. Sterile fronds scattered, on stipes 3—4 c.M. long, naked, ovate-elongate, \pm 8 c.M. long, \pm 3 c.M. broad, the apex acute, the base narrowed, the margin entire, cartilagineous. Texture thick, rigidly coriaceous; costa conspicuous; veins inconspicuous near the margin;

areolae hidden. Fertile fronds longer, ± 10 c.M. long, linear-ligulate, obtuse, ± 5 m.M. broad, on stipes ± 4 c.M. long; costa conspicuous. Sori linear, parallel to the costa and covering the margin.

New Guinea.

76. CHRISTOPTERIS, *Copeland.*

Sori long, linear, close to the costa, 1 on each side, finally covering the whole under surface of the contracted fertile fronds. No indusium.

Rhizome creeping. Stipes articulate to the rhizome. Fronds dimorphous. Barren fronds sagittate, fertile 5-partite with narrow segments; veins anastomosing; areolae with included free veinlets.

Philippines.

C. sagitta, *Copel.*, Polypod. Philipp., 111; Philipp. Journ., I, Suppl. 157, tab. XIII; *Polypodium Sagitta*, *Christ.*, Bull. Herb. Boiss., VI, 199.

Rhizome creeping or scandent, the younger part densely clothed with spreading, cinnamon-coloured, filiform scales ± 8 m.M. long, which are deciduous, leaving older parts sparsely covered with ferruginous scales ± 4 m.M. long. Stipes fuscous or ebeneous, of sterile fronds ± 10 c.M., of fertile ± 20 c.M. long, naked. Barren fronds 15—20 c.M. long and broad, deltoid, the point acuminate, with a triangular sinus 2 c.M. high and wide where the stipe is attached, with a broad, shallow sinus between the apex and each lateral point. Texture coriaceous; surfaces naked; the 5 main ribs conspicuous, the main veins strictly parallel, subevident. Fertile fronds cleft to within ± 1 c.M. of the base into 5 narrowly linear, somewhat spreading, obtuse segments 8—20 c.M. long, ± 5 m.M. broad.

Philippines.

TRIBE XI. ACROSTICHEAE.

77. PLATYCERIUM, Desvaux.

Sori indefinite, forming large patches on the under surface of the frond, occupying the ultimate segments or the lower sinuses, or placed on transversally or longitudinally oblong, stalked segments near the base of the frond.

Rhizome commonly short-creeping. Fronds dimorphous, articulate into a depression of the rhizome. Barren fronds (basal fronds) roundish-reniform in general outline, closely applied to the substratum, at least at the base, collecting quantities of detritus into which the roots grow, the outer edge entire or more or less deeply lobed; fertile fronds standing out from the substratum, forked or more generally dichotomously branched with stag-horn-like divisions; surfaces often clothed with more or less deciduous stellate hairs; principal veins dichotomous, secondary ones anastomosing; areolae with included free veinlets. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 177. (*)

Paleotropical, and in temperate Australia.

Arrangement of the sections.

§ 1. **EUPLATYCERIUM.** Sori on the ultimate segments or in the lower sinuses of the fertile fronds, but then not as in § 2.

§ 2. **SCUTIGERA.** Sori placed on specialized, stalked, oblong or broadly reniform segments springing from the lower part of the frond, in the lowest sinus.

§ 1. **EUPLATYCERIUM.**

a. Sori 1 to each fertile frond, occupying the primary disk (*i. e.* the cuneate base of the frond), placed against the upper edge. (1) *P. grande.*

a.a. Sori 2 to each fertile frond, occupying the secondary disks (*i. e.* the cuneate base of the primary fork branches), placed against the upper edge.

*. Fronds forked, both fork branches dichotomously branched. (2) *P. Wallichii.*

.. Fronds forked, the outer branch of each frond simple, broadly cuneato-flabellate, 2-cornate, the inner dichotomously branched.

(3) *P. Wilhelminae* Reginae.

a.a.a. Sori several to each fertile frond, occupying the ultimate segments.

(*) Mostly beautiful plants, whose extraordinary appearance has made them universally beloved in cultivation.

- b.** Fertile fronds shorter than the barren ones, with the ultimate segments cuneate, dilated towards the apex. (4) *P. Wandae*.
- b.b.** Fertile fronds longer than the barren ones, with the ultimate segments not cuneate, but rounded, blunt or narrowed at the apex.
- c.** Soriferous segments involute, the sori thickened with a dense tomentum giving them a corky appearance. (5) *P. sumbawense*.
- c.c.** Soriferous segments not involute, the sori not thickened, not corky.
- d.** Barren fronds 1, fertile 3, pendulous. (6) *P. Willinckii*.
- d.d.** Barren fronds 2 or more, fertile 4—7, erect or nutant, but not decidedly pendulous. (7) *P. bifurcatum*.

(1) ***P. grande***, *J. Sm.*, *Rac.*, *Flor. Btz.*, I, 56, p. p.; *Hk.*, *Fil. Exot.*, tab. LXXXVI; *Nich.*, *Dict. d' Hort.*, IV, fig. 210; *Bedd.*, *Ferns Br.I.*, II, tab. CCCXXVI; *Acrostichum grande*, *A. Cunn.*, MS.; *Alcicornium grande*, *Und.*, *Bull. Torr. Cl.*, XXXII, 594.

Basal fronds roundish, 40—90 c.M. each way, convex, deeply incised, the lobes simple or once or more times forked. Fertile fronds in pairs, 100—200 c.M. long, dichotomously branched, the basal part (disk) broadly cuneate, the ultimate segments broad, linear, loriform, bluntly rounded. Texture thinly coriaceous; surfaces naked. Sori only 1 to each frond, occupying the disk only, placed against the upper edge between the 2 primary fork branches.

Malaya; Trop. Australia.

(2) ***P. Wallichii***, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 425; *Hk.*, *Fil. Exot.*, tab. XCVII; *Bedd.*, *Ferns Br.I.*, I, tab. CVIII; *Alcicornium Wallichii*, *Und.*, *Bull. Torr. Cl.*, XXXII, 596.

Near *P. grande*, *J. Sm.*, but the fertile fronds shorter, 30—70 c.M. long. Texture thick; under surface covered with a thin, yellowish or reddish tomentum. Sori 2 to each frond, occupying the secondary disks, i. e. the cuneate base of the primary fork branches, placed against the upper edge, which is much produced in the sinus.

Malacca.

* (3) ***P. Wilhelminae Reginae***, *v. A. v. R.*, *Bull. Dép. Agr. I. N.*, 1908, XVIII, 24, tab. VI—VII; *Alcicornium Wilhelminae Reginae*, *v. A. v. R.*, l. c., 25.

Basal fronds 2 or more, large, to \pm 100 c.M. each way, convex, the

upper edge deeply lobed, the lobes linear, elongate, erect, simple or 1—2 × forked, the base rounded, obliquely cordate, the outer edge sinuate, with the lobes gradually passing into the higher, linear ones. Fertile fronds in pairs, to 200 c.M. long, divided into 2 unequally developed halves. The inner halves (i. e. the 2 contiguous halves) of the fronds pendulous, dichotomously branched, the segments cuneate, widening gradually from the base to the next higher branching; ultimate segments lanceolate or triangular-elongate, narrowed towards the apex. The outer halves of the fronds spreading horizontally in opposite directions, simple, large, to 75 c.M. broad, broadly flabellato-cuneate, the outer edge slightly rounded and flanked by 2 blunt, falcate, incurved horns, the horns horizontal or pendulous. Texture flaccido-coriaceous, firm when dry; surfaces clothed with numerous, scattered but not crowded, stellate hairs. Sori large, 2 to each frond, occupying the secondary disks, i. e. one placed in the lowest sinus of the dichotomously divided half of each frond and the other on the simple half, between the horns, the latter much the largest, to ± 45 c.M. broad, to ± 20 c.M. deep; capsules copiously intermixed with stellate hairs. — The description of this splendid and interesting fern is taken from a specimen cultivated in the Buitenzorg gardens and probably gathered in one of the malayan islands. Raciborski referred it to *P. grande*, *J. Sm.*, which has the segments loriform, naked, and the sori solitary, i. e. 1 sorus to each frond, occupying the primary disk. — This species is dedicated to Her Majesty WILHELMINA, Queen of the Netherlands.

? *Malayan.*

(4) ***P. Wandae***, *Rac.*, Bull. Ac. Crac., 1902, 58.

Basal fronds 150—200 c.M. long, 100 c.M. or more broad, the upper edge deeply incised, the base deeply cordate, the lobes furnished with numerous excrescences, which are very thin, ± 6 c.M. long, several times dichotomously forked and hiding the apex of the scaly rhizome. Fertile fronds pendulous, 2 × forked, smaller than the barren ones, to 120 c.M. long; ultimate segments cuneate, 20—40 c.M. broad at the apex, which is not truncate but repand or incised. Sori covering the under surface of the terminal segments.

New Guinea.

* (5) ***P. sumbawense***, *Christ.*, Warb., Mons., I, 64; *Alcicornium sumbawense*, *Und.*, Bull. Torr. Cl., XXXII, 596.

Basal fronds roundish-reniform, 15—30 c.M. each way, lacinate, the laciniae deeply forked; texture scariosa, fragile, spongy; surfaces ± glossy;

colour ochraceous-fulvous. Fertile fronds 2—5, narrowed gradually towards the base, to 65 c.M. long, repeatedly dichotomous, pale-coloured, flocculent, the tomentum adpressed; ultimate segments rather numerous (\pm 16—20) to each frond, lanceolate, ligulate or elongate-triangular, to 7 c.M. long, narrow subobtusate, rostrate, involute. Sori occupying the whole under surface of the ultimate segments, densely covered with a thick, whitish tomentum when young, but less so at full maturity. — A plant gathered in Timor by Teysmann, very probably a form of this species, has the fertile fronds very coriaceous and rigid, 45—75 c.M. long, with the ultimate segments not rostrate, the sori often reaching from the apex of the ultimate segments to considerably below the highest sinuses.

Sumbawa, Timor.

(6) **P. Willinckii**, *Moore*, Nich., Dict. d' Hort, IV, 175; Gard. Chron., N. S., III, fig. 56; *Alcicornium Willinckii*, *Und.*, Bull. Torr. Cl., XXXII, 596.

Basal fronds single, erect, suborbicular and lobed. Fertile fronds triple, elongate, pendulous; each frond narrowly cuneate, dichotomously forked, the lobes narrow, ensiform. Sori on the under surface of the ultimate lobes.

Java.

* (7) **P. bifurcatum**, *C. Chr.*, Ind. Fil.; *P. alcicorne*, *Desv.*, Rac., Flor. Btz., I, 57; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 177, A; Nich., Dict. d' Hort., IV, fig. 209; *Alcicornium bifurcatum*, *Und.*, Bull. Torr. Cl., XXXII, 594; *Acrostichum bifurcatum*, *Cav.*, (oldest name).

Rhizome thick, creeping, densely scaly, bearing the fronds in more or less distant groups. Basal fronds 2 or more, roundish or cordate, imbricate, convex, entire, sinuate or lobed, 10—20 c.M. each way, afterwards brown, the lobes spreading or erect. Fertile fronds mostly 4—7, clustered, the stipes thick, to \pm 5 c.M. long or longer, \pm erect, the fronds 40—90 c.M. long, cuneate at the base, 2—3 \times dichotomously branched, with about 4—8 ultimate lobes, which are ligulate, flat, 10—20 c.M. long, 2—5 c.M. broad, with the apex blunt or bluntish. Texture thick; colour pale-green; young fronds covered with a thin, whitish or pale-reddish, \pm fugacious, adpressed tomentum; areolae \pm hexagonal, with few free veinlets. Sori occupying \pm the upper $\frac{3}{4}$ of the ultimate lobes.

Much cultivated. — *Orig. loc.:* *Australia, New Caledonia, Lord Howe's Island.*

§ 2. SCUTIGERA.

* (8) **P. coronarium**, *Desc.*, Prod., 215; *P. biforme*, *Bl.*, Rac., Flor. Btz., I, 57; *Bl.*, Flor. Jav., II, 44, tab. XVIII; *Bedd.*, Ferns Br., I, tab. CIX; II, tab. CCXXIV; *Alcicornium coronarium*, *Und.*, Bull. Torr. Cl., XXXII, 594; *Osmunda coronaria*, *Kön.*, (oldest name); — var. *cucullatum*, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 25.

Basal fronds roundish, thick, at length yellow-brown, the upper edge often deeply lobed. Fertile fronds single, double or clustered, to 100—450 c.M. long, stalked, pendulous, repeatedly dichotomous; ultimate lobes ligulate, 2—3 c.M. broad. Texture coriaceous; surfaces not rarely clothed with more or less numerous stellate hairs. Sori placed on a broadly reniform or transversally oblong, mostly convex, sometimes half-cupshaped, stalked segment 10—20 c.M. broad, springing from the lower part of the fertile fronds, in the lowest sinus, 1 to each frond.

Var. cucullatum: Ultimate lobes of fertile fronds dilated towards the obliquely rotundato-truncate and slightly emarginate apex. Soriferous segment longitudinally oblong, subcucullate, rather slipper-shaped, the edge broadly recurved.

Malaya; Siam, Burmah; — var cucullatum: Lingga Islands (Teysmann).

78. ELAPHOGLOSSUM, *Schott.*

Sori indefinite; sporangia densely covering the under surface of the fertile fronds.

Rhizome mostly creeping. Fronds simple, articulate or subarticulate to the rhizome, uniform or nearly so, rarely exceedingly different in shape; veins free or united by a submarginal veinlet or a marginal, cartilagineous or decoloured strand. *Hk. Bk.*, Syn. Fil., tab. VII, fig. 60, a—b; *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam., I⁴, fig. 173 & 174, A, E & H.

Pantropical, but more especially neotropical.

Arrangement of the sections.

§ 1. **ACONIOPTERIS**. Fronds bordered by a submarginal vein or marginal, diaphanous, decoloured, cartilagineous or thickened line or strand.

§ 2. **EU-ELAPHOGLOSSUM**. Fronds not bordered as in § 1.

§ 1. ACONIOPTERIS. (*)

- a.** Veins united by a submarginal veinlet. (1) *E. gorgoneum*.
- a.a.** Veins united by a marginal, thickened, cartilaginous, decoloured or diaphanous line or strand.
- b.** Rhizome scales blackish, jagged or fimbriate. (2) *E. conforme*.
- b.b.** Rhizome scales yellowish. (3) *E. angulatum*.
- b.b.b.** Rhizome scales brown.
- c.** Fronds bluntly rounded at the apex.
- * Veins rather hidden. (4a) *E. decurrens*.
- ** Veins rather visible. (4b) *E. luzonicum*.
- c.c.** Fronds not bluntly rounded at the apex.
- d.** Fronds naked or very sparingly scaly, the scales minute, orbicular.
- e.** Stipes short with regard to the fronds. (5) *E. Beccarianum*.
- e.e.** Stipes long with regard to the fronds. (6) *E. callifolium*.
- d.d.** Fronds minutely scaly, especially beneath, the scales lanceolate-subulate or stellate-fimbriate. (7) *E. laurifolium*.

(1) **E. gorgoneum**, *Brack.*, Expl. Exp., XVI, 74; *Acrostichum gorgoneum*, *Klf.*, Hk. Bk., Syn. Fil., 416; ? *A. pellucido-marginatum*, *Christ*, Fil. Saras., III, 255.

Rhizome short-creeping, firm, the scales brown. Stipes approximate, firm, erect, 2½—5 c.M. long, passing gradually into the barren fronds, which are 15—40 c.M. long, 5—8 c.M. broad, the point acute, the lower ⅔ narrowed very gradually, the edge entire. Texture subcoriaceous; surfaces naked; veins united by a straight, submarginal veinlet. Fertile fronds narrower, on stipes 15 c.M. or more long.

Malaya; Polynesia.

* (2) **E. conforme**, *Schott* (not *Moore*), *Christ*, Farnkr. d. Erde, 33; *Diels*, in *Engl. & Prantl*, Nat. Pfl.Fam., I⁴, fig. 173; *Bedd.*, Ferns S.I., tab. CXCVIII; *Acrostichum conforme*, *Sw.*, Hk. Bk., Syn. Fil., 401; *A. marginatum*, *Wall.*, List., No. 17; *Vittaria acrostichoides*, *Hk. & Grev.*, Ic. Fil., tab. CLXXXVI.

(*) See also **E. hirtum**, *C. Chr.*

Rhizome creeping, woody, the scales blackish, ovate, jagged or fimbriate, not hair-pointed. Stipes firm, approximate, erect, 5—8 c.M. long, blackish at the base, generally clothed with scattered, pale, sheathing scales. Sterile fronds 5—30 c.M. long, 2—3 c.M. broad, lanceolate, narrowed towards both ends, the point acute or acuminate, the edge entire, decoloured or cartilagineous, slightly revolute in age. Fertile fronds the longest and narrowest, longer stalked. Texture coriaceous; both surfaces covered with a few deciduous scales when young, those on the midrib beneath the largest; veins hidden, generally once forked.

Pantropical.

* (3) **E. angulatum**, *Moore*, C. Chr., Ind. Fil.; *Acrostichum angulatum*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 47; *Bl.*, *Flor. Jav.*, II, 25, tab. VI.

Rhizome creeping, the scales yellowish, lanceolate, thinly membranaceous. Stipes firm, erect, stramineous, naked or slightly scaly, 1—3 c.M. apart, 5—20 c.M. long, those of the fertile fronds the longest. Fronds elliptical-lanceolate, 8—40 c.M. long, 2—4 c.M. broad, the fertile ones the smallest, narrowed towards both ends, the apex acute, the margin entire, with a narrow, decoloured line. Texture coriaceous; surfaces naked or slightly scaly on the midrib beneath; veins subparallel, simple or forked.

Java.

* (4a) **E. decurrens**, *Moore*, C. Chr., Ind. Fil.; *Acrostichum decurrens*, *Desv.*, *Rac.*, *Flor. Btz.*, I, 47; *Bl.*, *Flor. Jav.*, II, 32, tab. X; *A. obtusifolium*, *Spr.* (not *Brack.*), *Hk.*, *Spec. Fil.*, V, 204; — var. *Cumingii*; *E. Cumingii*, *Moore*, *Copel.*, *Polypod. Philipp.*, 137; *A. Cumingii*, *Kée*, *Hk. Bk.*, *Syn. Fil.*, 407.

Rhizome creeping, the scales dense, long, linear-lanceolate, brown. Stipes 2½—15 c.M. long, those of the fertile fronds the longest. Barren fronds 15—40 c.M. long, 4—8 c.M. broad, the apex bluntly rounded, the margin cartilagineous or ± pellucid, the lower part narrowed gradually into the stipe. Texture coriaceous; surfaces naked or the under side sparingly and minutely scaly, the edge more or less fringed with minute, ovate-lanceolate, acuminate, thin, brown scales; veins immersed, nearly hidden. Fertile fronds often small and narrow.

Var. Cumingii: Stipes longer; barren fronds 3—4 c.M. broad, fertile ones narrower; upper surface scaly like the lower.

Sumatra, Java, Moluccas, Philippines; — var. *Cumingii*: *Philippines*.

(4b) **E. luzonicum**, *Copel.*, *Leafl.*, I, 255.

Rhizome strong, densely scaly. Stipes approximate, slender, 16—30 c.M. long, scaly, the scales different, the largest ones narrowly ovate. Barren fronds obtuse, 20—25 c.M. long, 4—5½ c.M. broad, cuneate at the base, the margin bounded by a cartilagineous strand and ciliate with an intramarginal row of deciduous, subciliate scales. Texture subcoriaceous; surfaces glabrescent; veins rather visible, at length conspicuous. Fertile fronds smaller, on longer stipes. — Near *E. decurrens*, *Moore*.

Luzon.

(5) **E. Beccarianum**, *C. Chr.*, *Ind. Fil.*; *Acrostichum Beccarianum*, *Bk.*, *Becc.*, *Mal.*, III, 27; *A. Norrisii*, *Ces.* (not *Hk.*), *Fel. di Bor.*, 31.

Rhizome short-creeping, densely clothed with small, lanceolate, brown scales. Barren fronds oblong-lanceolate, firm, naked, to 30—40 c.M. long, to ± 4 c.M. broad, narrowed gradually into a short stipe, the edge entire, thick; veins inconspicuous, erecto-patent, not closely placed. Fertile fronds often much smaller, on stipes to ± 15 c.M. long.

Borneo.

* (6) **E. callifolium**, *Moore*, *C. Chr.*, *Ind. Fil.*; *Acrostichum callaeifolium*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 47; *Bl.*, *Flor. Jav.*, II, 22, tab. IV.

Rhizome thick, creeping, the scales narrow, lanceolate, brown, dense. Stipes ± 15—50 c.M. long, firm, erect, naked or slightly scaly. Barren fronds 10—80 c.M. long, 5—10 c.M. broad, fertile ones shorter and narrower than the barren, elliptical-lanceolate, narrowed towards both ends or rounded at the base, the edge with a narrow, decoloured line. Texture membranaceous to coriaceous; surfaces naked or slightly scaly, the scales microscopical, roundish; veins fine, indistinct, simple or forked.

Java, Mindanao.

* (7) **E. laurifolium**, *Moore*, *C. Chr.*, *Ind. Fil.*; *Bedd.*, *Ferns S. I.*, tab. CC; *E. latifolium*, *Bedd.*, *Handb. Ind. Ferns*, 416; *Acrostichum laurifolium*, *Thouars*, *Flor. Trist. d'Ac.*, 51; *A. commutatum*, *Mett.*, *Ann. Mus. Bot. L. B.*, IV, 292; *A. conforme*, *Bl.* (not *Sw.*), *Flor. Jav.*, II, 23, tab. V.

Rhizome creeping, woody, densely clothed with lanceolate, acuminate, brown scales. Stipes erect, firm, subangular, 5½—7½ c.M. long, those of the fertile fronds the longest, covered with scattered, small, deciduous scales, and purveyed with an elevated, decurrent margin on each side. Sterile fronds

7—15 c.M. long, $1\frac{1}{2}$ —4 c.M. broad, oblong-lanceolate, narrowed towards both ends, but more gradually into the stipe, the point acute, the margin entire or slightly repand, thickened, diaphanous. Texture rigid; both surfaces (especially the lower) clothed with scattered, minute, deciduous, brownish, lanceolate-subulate or stellate-fimbriate scales; veins fine, hardly visible, often forked. Fertile fronds narrower, thinner, subrepand.

Luzon, Celebes, Java; Ceylon, South. India, Mascarenes, Tristan d'Acunha.

§ 2. EU-ELAPHOGLOSSUM.

- 1.** *Under surface of the fronds naked or nearly so.*
- 2.** *Fronds small, under 10 c.M. long.* (8) *E. microphyllum.*
- 2.2.** *Fully developed fronds larger, more than 10 c.M. long.*
- 3.** *Stipes of barren fronds wanting or very short, under 3 c.M. long.*
(9) *E. melanosticticum.*
- 3.3.** *Stipes of barren fronds distinct, 7—15 c.M. long.* (10) *E. Lessonii.*
- 1.1.** *Under surface of the fronds rather densely tomentose or scaly.*
- a.** *Barren fronds suddenly narrowed at the base. Under surface scaly.*
(11) *E. miniatum.*
- a.a.** *Barren fronds gradually narrowed at the base.*
- b.** *Barren fronds clothed beneath with stellate hairs.*
(12) *E. petiolatum.*
- b.b.** *Barren fronds clothed beneath with ciliate or fimbriate scales.*
- c.** *Barren fronds densely scaly above.* (13) *E. hirtum.*
- c.c.** *Barren fronds sparingly scaly above.* (14) *E. Copelandi.*

* (8) **E. microphyllum**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 25.

Rhizome slender, creeping, blackish, the scales linear-lanceolate, brown, entire. Stipes distant, $\frac{1}{2}$ —4 c.M. apart, slender, 1— $2\frac{1}{2}$ c.M. long. Sterile fronds lanceolate, entire, 5—6 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, shortly acuminate but the point bluntish, the base narrowed gradually into the stipe. Texture herbaceous, firm; surfaces naked; veins not close, simple or once or more times forked, not quite reaching the unaltered margin. Fertile fronds shorter, narrower, the edge subentire, the base narrowed more suddenly and

then tapering very gradually into the stipe so as to form a very narrow wing to the upper part of the stipe, which is \pm 7—8 c.M. long.

Java (Ploem).

(9) **E. melanosticticum**, *Moore*, C. Chr., Ind. Fil.; *Acrostichum melanosticticum* (*melanostycton*), *Bl.*, Flor. Jav., II, 26, tab. VII; *A. Norrisii*, *Hk.*, Hk. Bk., Syn. Fil., 401; *E. Norrisii*, *Bedd.*, Handb. Ind. Ferns, 418.

Rhizome creeping, stout, woody, the scales ovate-lanceolate, long-acuminate, suberose, brownish, dense. Stipes wanting or very short, to $2\frac{1}{2}$ c.M. long, approximate. Barren fronds 30—45 c.M. long, 2—4 c.M. broad, broadest \pm $\frac{1}{3}$ on the way to the base, blunt or acute, narrowed gradually from the broadest part towards both ends. Texture coriaceous; surfaces naked or nearly so; veins immersed, indistinct. Fertile fronds much narrower than the barren ones.

Java, Penang.

(10) **E. Lessonii**, *C. Chr.*, Ind. Fil.; *Acrostichum Lessonii*, *Mett.*, *Linn.*, XXXVI, 60.

Rhizome thick, the scales dense, brownish, oblong-lanceolate, with the edge slightly ciliate. Stipes approximate, 7—15 c.M. long, compressed. Sterile fronds \pm $57\frac{1}{2}$ c.M. long, \pm 4 c.M. broad, linear-spathulate, obtuse, narrowed gradually into the stipe. Texture coriaceous; upper surface naked or nearly so, lower hardly covered with punctiform, fugacious scales; veins immersed, repeatedly forked, erecto-patent. Fertile fronds linear-lanceolate, \pm 20 c.M. long, \pm 2 c.M. broad, narrowed gradually towards both ends, on stipes \pm 2 c.M. long; costa prominent.

Sumatra; Vanikoro.

(11) **E. miniatum**, *Christ*, Mon. Elaph., 72, fig. 30; *Acrostichum miniatum*, *Christ*, Ann. Bot., XV, 175; *Fil. Saras.*, III, 254, tab. III, fig. 25—27.

Barren fronds \pm 25 c.M. long, \pm 3 c.M. broad, finely acuminate, suddenly narrowed into the stipe. Texture very thick; veins hidden; upper surface naked, light-green, scaly on the costa; under side covered with regularly scattered, branched, reddish-brown scales, those of the margin peltate, ciliate, densely crowded. Fertile fronds longer, narrowly linear, on longer stipes; upper surface densely covered with cinnamon-coloured, round, stellate hairs.

Celebes.

* (12) **E. petiolatum**, *Urb.*, C. Chr., Ind. Fil.; *E. viscosum*, *Schott*, Christ, Farnkr. d. Erde, 39; *Bedd.*, Ferns S.I., tab CXCVI; *Acrostichum viscosum*, *Sw.*, Rac., Flor. Btz., I, 46; *Hk. & Grev.*, Ic. Fil., tab. LXI; *A. petiolatum*, *Sw.*, (oldest name).

Rhizome woody, short-creeping, the scales dense, linear-lanceolate, dark-brown. Stipes approximate, 7—15 c.M. long, firm, erect, covered with small scales and stellate hairs. Barren fronds 15—30 c.M. long, 1—2½ c.M. broad, the apex acute or acuminate, the lower part narrowed gradually. Texture thinly coriaceous; under surface covered with stellate hairs, upper naked or hairy too; midrib conspicuous; veins visible, fine. Fertile fronds narrower and longer stalked.

Trop. Asia, Africa, America.

* (13) **E. hirtum**, *C. Chr.*, Ind. Fil.; *E. squamosum*, *J. Sm.*, Christ, Farnkr., d. Erde, 56, fig. 83; *Bedd.*, Ferns S.I., tab. CXCVII; *Acrostichum squamosum*, *Sw.*, *Hk. Bk.*, Syn. Fil., 411; *A. paleaceum*, *Hk. & Grev.*, Ic. Fil., tab. CCXXXV; *A. hirtum*, *Sw.*, (oldest name).

Rhizome woody, short-creeping, the scales ovate to lanceolate, ciliate, blackish-brown. Stipes tufted or approximate, 2½—10 c.M. long, densely clothed with lanceolate, ciliate, pale, brown and blackish scales. Fronds 15—30 c.M. long, 2—3½ c.M. broad, elongate-ligulate, acute or short-acuminate at the point, the base narrowed gradually. Texture thick but flaccid; surfaces densely covered with ovate-lanceolate, ciliate, brown scales, the scales on the midrib are longer and those on the under surface intermixed with polished, blackish-brown ones; midrib conspicuous; veins hidden. Fertile fronds narrower and longer stalked than the barren ones, the edge more or less membranaceous.

Sumatra; Ceylon, South. India, East. Africa, Madeira, Azores, Trop. America, Hawaii.

(14) **E. Copelandi**, *Christ*, Philipp. Journ., II, 176.

Rhizome creeping, woody, sparingly clothed with black-brown, subulate setulae. Stipes approximate, to 35 c.M. long, with the fertile fronds on the longest stipes, rufo-stramineous, clothed with black, fimbriato-lacerate squamulae, the base black-brown, provided with scattered, brown, lanceolate scales. Barren fronds linear-lanceolate, to ± 42½ c.M. long, ± 2½ c.M. broad, narrowed towards the acute apex and base, not decurrent on the stipe. Texture firm-chartaceous; colour dark-brown; upper surface sparingly scaly, lower

densely so, the scales peltate, copiously ciliate to fimbriate, partially smaller and decoloured, partially larger and cinnamon-coloured; costa prominent, rufous, rather sparingly scaly; veins hidden, horizontal. Fertile fronds linear, to \pm 35 c.M. long, \pm 1 c.M. broad; upper surface densely scaly, lower densely soriferous.

Mindanao.

79. STENOCHLAENA, J. Smith.

Sori indefinite; sporangia densely crowded over the under surface of the much contracted or narrowed, \pm linear pinnae of the fertile fronds.

Rhizome woody, high-scandent, often reaching the tops of the highest trees. Fronds pinnate, dimorphous or polymorphous, whether or not articulate to the rhizome, the abnormal barren fronds, when present, variously and often finely divided, the fertile fronds Lomaria-like; veins free, simple or forked, those of the fertile pinnae sometimes slightly anastomosing, but always united at the base by a more or less distinct or spurious, \pm costal, transversing vein, those of the barren pinnae sometimes so. Hk. Bk., Syn. Fil., tab. VII, fig. 60, f—g; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 133. (*)

Pantropical.

Arrangement of the sections.

§ 1. **LOMARIOPSIS.** Barren fronds uniform; abnormal fronds wanting.

§ 2. **TERATOPHYLLUM.** Barren fronds dimorphous; abnormal fronds present.

§ 1. **LOMARIOPSIS.**

a. Barren pinnae not articulate to the rachis, cordate at the base.

(1) *S. laurifolia.*

a.a. Barren pinnae articulate to the rachis.

b. Barren pinnae unequal-sided at the base; upper base rather truncate, lower \pm cuneate.

(2) *S. arthropteroides.*

(*) The fertile pinnae have the edges not rarely recurved, \pm modified and indusium-like, and then it is rather difficult to distinguish the species from Lomaria, except when the rhizome is shown.

b.b. Barren pinnae equal-sided at the base.

c. Lateral pinnae few, 1—3 on each side. (5) *S. Smithii*.

c.c. Lateral pinnae more numerous, ± 6 or more on each side.

d. Rhizome naked or nearly so; barren pinnae with a small gland at or near the base on the upper edge.

(4) *S. palustris*.

d.d. Rhizome rather copiously scaly; barren pinnae without a gland at the base.

e. Barren pinnae with the edges not cartilaginous.

(5) *S. sorbifolia*.

e.c. Barren pinnae with the edges cartilaginous.

(6) *S. subtrifoliata*.

(1) **S. laurifolia**, *Pr.* (not *Diels*), *Copel.*, *Polypod. Philipp.*, 88; *Acrostichum laurifolium*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 412.

Stipes firm, erect, naked. Fronds 60—90 c.M. long, 30—45 c.M. broad. Barren pinnae 15—25 c.M. long, $5\frac{1}{2}$ — $4\frac{1}{2}$ c.M. broad, sessile, not articulate to the rachis, narrowed gradually from the cordate base to the apex, the edge sharply but finely toothed, without a gland. Texture coriaceous; rachis and both surfaces naked. Fertile pinnae as long as the barren ones, $\frac{1}{2}$ — $1\frac{1}{2}$ c.M. broad.

Philippines and Celebes to Polynesia.

(2) **S. arthropteroides**, *Christ*, *Bull. Herb. Boiss.*, 2^e Série, VI, 998.

Rhizome wide-excurrent, rugulose, not aculeate, brown, clothed with scattered, adpressed, black, furfuraceous scales. Stipes scattered, fulvo-stramineous, to 5 c.M. long. Fronds ± 25 c.M. long, ± 15 c.M. broad, broadly ovate, hardly narrowed at the base, the rachis slender, minutely winged. Pinnae lanceolate, ± 8 — 10 on each side, spreading, stalked, articulate on distinct, black, calliform protuberances of the rachis, ± 9 c.M. long, ± 2 c.M. broad in the fully developed plants, the upper base auriculate-truncate, the lower cuneate, the edge crenato-serrate. Texture thinly herbaceous; colour bright-green; veins forked, more than 1 m.M. apart, clothed beneath with scattered, black, furfuraceous scales, as is the rachis. Fertile fronds with the pinnae linear, very distant, acuminate, rounded or nearly cordate at the base.

Luzon.

* (3) **S. Smithii**, *v. A. v. R.*; *S. Raciborskii*, *C. Chr.*, Ind. Fil.; *Acrostichum Smithii*, *Rac.*, Bull. Ac. Crac., 1902, 59.

Rhizome wide-creeping (scandent), densely clothed with lanceolate, acuminate, fimbriate scales. Stipes 8—10 c.M. long, green, slightly canaliculate, scaly especially near the base. Fronds pinnate, 30—45 c.M. long, 30—56 c.M. broad. Pinnae 1—3 on each side below the terminal one, alternate, articulate, distant, linear-lanceolate, 13—20 c.M. long, 2—3 c.M. broad, the base narrowed, the margin entire or indistinctly crenate; terminal pinna 25—50 c.M. long, often inequilateral. Texture rigid, fragile; surfaces smooth, naked; veins parallel, spreading or suberecto-patent, simple or often once forked. Fertile pinnae linear, 2½—5 m.M. broad, thinly herbaceous.

Moluccas, Aru Islands.

* (4) **S. palustris**, *Bedd.* (not *Mett.*), Copel., Polypod. Philipp., '88; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 133, C—D; *S. scandens*, *J. Sm.*, *Bedd.*, Ferns S.I., tab. CCI; *Acrostichum scandens*, *Hk.* (not *J. Sm.*), *Rac.*, Flor. Btz., I, 54; *Lomariopsis palustris*, *Kuhn* (not *Mett.*), *Christ*, *Farnkr. d. Erde*, 41; *Polypodium palustre*, *Burm.*, (oldest name); — var. *juglandifolia*; *S. juglandifolia*, *Pr.*, Copel., l. c.

Rhizome woody, wide-scandent, naked or sparingly covered with roundish or ovate, adpressed scales. Stipes 7—20 c.M. long, firm, erect, naked. Fronds 30—200 c.M. long, 20—50 c.M. broad, pinnate. Pinnae articulate, numerous, lanceolate, 2—4 c.M. broad, acuminate, pungently serrate towards the apex, the base cuneately rounded, with a small pulvinate gland on the upper side, which is often black in dried material. Texture firm; surfaces naked; veins simple or forked, united at the base by a costal line or costal arches, otherwise free, but not rarely irregularly and slightly anastomosing in the fertile pinnae. Fertile fronds with the pinnae linear, 2—5 m.M. broad, the edge mostly covering the sori when young.

Var. juglandifolia: Fertile pinnae leafy at the base; stipes mucronate.

Trop. Asia, Australia & Polynesia; — var. *juglandifolia*: *Philippines*.

* (5) **S. sorbifolia**, *J. Sm.*, Copel., Polypod. Philipp., 88; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 133, A—B; *Acrostichum sorbifolium*, *L.*, *Hk. Bk.*, *Syn. Fil.*, 412; *Lomariopsis sorbifolia*, *Fée*, *Christ*, *Farnkr. d. Erde*, 59; *Bedd.*, Ferns Br.I., II, tab. CXCII; — var. *spectabilis*; *A. spectabile*, *Rac.*, *Flor. Btz.*, I, 54; *Lomaria spectabilis*, *Kze.*, *Bot. Zeit.*, 1848, 144.

Rhizome high-climbing, thick, woody, often angled by the decurrent bases of the stipes, clothed with lanceolate-subulate, often ciliate, brownish scales. Stipes firm, slightly scaly, 10—30 c.M. long. Fronds 30—100 c.M. long, 20—60 c.M. broad, pinnate, the rachis often narrowly winged towards the apex. Pinnae distant, articulate, \pm 6—50 on each side, sublanceolate, 2—5 c.M. broad, acute, acuminate or subcaudate, entire, crenate, serrate or dentate, the base \pm cuneately narrowed. Texture firm; surfaces naked; colour blackish-green when dry; veins parallel, distinct quite to the margin, simple or forked. Fertile fronds with the pinnae smaller and narrower, \pm 5—9 m.M. broad.

Var. spectabilis: Fertile pinnae as long as the barren ones, linear, 2—5 m.M. broad.

Pantropical; — var. *spectabilis*: *Malaya*.

(6) **S. subtrifoliata**, *Copel.*, Philipp. Journ., I, Suppl., 152.

Rhizome scandent, angular, densely clothed with rather long, lanceolate or ovate, fuscous scales. Stipes 10—20 c.M. long, firm, densely scaly at the base, subglabrescent towards the frond. Fronds of the fully developed plants pinnate, 45—70 c.M. long, ovate. Pinnae articulate to the rachis, the lower stalked, \pm 25 c.M. long, $2\frac{1}{2}$ —3 c.M. broad, acute or caudate, the edge entire, cartilagineous; higher pinnae sessile, often oblique. Texture coriaceous; surfaces naked. Fertile pinnae \pm 1 c.M. broad.

Mindanao.

Indefinite species.

* (7)? **S. dubia**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 26.

Rhizome scandent, woody, stout, densely scaly like the base of the stipes, blackish-brown, the older parts rough and rather naked. Stipes \pm 35 c.M. long, dark-purple-brown, densely scaly at the base, densely tomentose in front like the rachises, provided with 2 rows of elliptical, flattened, depressed, paler coloured glands resembling scars of fallen pinnae, the scales long, linear-subulate, brown, leaving the stipes asperous when they fall, the tomentum dark-reddish-brown. Fronds 2-pinnate, the rachis dark-purple-brown. Pinnae erecto-patent, not articulate to the rachis. Pinnulae numerous, rather spreading, articulate to the secondary rachises, ovate-oblong, acuminate, \pm 3 c.M. long, \pm $\frac{1}{2}$ c.M. broad near the base, the base rounded or broadly

cuneate, slightly unequal-sided, the edge shallowly crenate but more deeply towards the apex. Texture coriaceous; surfaces naked, upper dark-coloured, lower paler; costae prominent on both sides, with the upper side tomentose towards the base; veins prominent beneath, the higher ones simple or forked, the lower 2 × furcate or subpinnate, all united at the base by a prominent costal line. Fertile fronds not seen. — The description is taken from a specimen determined as *Lomariopsis sorbifolia*, *Fée*.

Amboyna (Teysmann).

§ 2. TERATOPHYLLUM.

* (8) **S. aculeata**, *Hæe*, Bot. Zeit., 1848, 142; *S. gracilis*, *Kze*, l. c.; *Acrostichum aculeatum & gracile*, *Rac.*, Flor. Btz., I, 55; *Teratophyllum aculeatum*, *Mett.*, Ann. Mus. Bot. L. B., IV, 296; *Olfersia aculeata & gracilis*, *Bl.*, Flor. Jav., Suppl., tab. XCV—XCVI; *Davallia achilleaefolia*, *Wall.*, Christ, Farnkr. d. Erde, fig. 96, i; *Hk.*, Spec. Fil., I, tab. LVI, D; *Lomaria aculeata & gracilis*, *Bl.*, (oldest names); — var. *inermis*, *Mett.*, l. c., 297; Ann. Btz., XII, tab. XIV—XV, fig. 25—28; *Bedd.*, Ferns Br.I., II, tab. CCIX.

Rhizome firm, wide-climbing, prickly, varying from slender to 5 m.M. thick. Fronds scattered, the barren ones dimorphous. Normal barren fronds stalked, 40—100 c.M. long, pinnate. Pinnae articulate, acute or acuminate, entire, cuneate or rounded at the base, to 11 c.M. long, to 2½ c.M. broad. Texture firm and green (thin and reddish-purplish in *S. gracilis*, *Kze*); rachis and costae often minutely scaly; surfaces naked. Abnormal barren fronds variously divided, with very small, entire, crenate or incised, roundish, oblong or linear ultimate segments, much smaller than the normal fronds and adpressed to the substratum; texture membranaceous. Fertile fronds pinnate, the pinnae narrowly linear. — The forms of this species are often considered as forms of *S. sorbifolia*, *J. Sm.* or *palustris*, *Bedd.*

Var. inermis: Rhizome unarmed.

Trop. Asia.

80. POLYBOTRYA, *Humb. & Bonpl.*

Sori at first terminal on the veins, but soon occupying ± the whole under surface of the much contracted fertile segments.

Rhizome erect or creeping, but not high-scandent. Fronds not articulate to the rhizome, at least simply pinnate, dimorphous, the fertile ones not Lomaria-like; veins free in malayan material.

Pantropical.

Arrangement of the sections.

- § 1. **EGENOLFIA.** Veins free; leaflets with a tooth or seta in the sinuses of the crenations or lobes. Rhizome short-creeping. Hk. Bk., Syn. Fil., tab. VII, fig. 60, h—j; Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 102.
- § 2. **PSOMIOCARPA.** Veins free; leaflets without a tooth or seta in the sinuses. Rhizome erect or slightly tuberous. Diels, l. c., fig. 105, B.
- § 3. **EUPOLYBOTRYA.** Veins free; leaflets without a tooth or seta in the sinuses. Rhizome wide-creeping, scaly. Hk. Bk., l. c., fig. 60, c—e; Diels, l. c., fig. 104. (Malayan?).
- § 4. **ARTHROBOTRYA.** Like § 3, but the rhizome naked; leaflets articulate to the rachis.
- § 5. **OLFERSIA.** Vein tips united by an intramarginal transversing vein. Hk. Bk., l. c., fig. 60, m—o. Diels, l. c., fig. 103, E. (Not in Malaya).
- § 6. **SOROMANES.** Veins anastomosing \pm as in Nephrodium, but less regularly. Hk. Bk., l. c., fig. 60, r—s; Diels, l. c., fig. 105, C. (Not in Malaya).

Key to the species ().*

- a.** *Pinnae not articulate to the rachis.*
- b.** *Leaflets with a tooth or seta in the sinuses of the crenations or lobes.* (1) *P. appendiculata.*
- b.b.** *Leaflets without a tooth or seta in the sinuses.*
- c.** *Fronds 2-pinnatifid.*
- d.** *Pinnae deciduously scaly at the margin.* (4) *P. stenosemioides.*
- d.d.** *Pinnae not scaly at the margin.* (2) *P. Nieuwenhuisii.*
- c.c.** *Fronds 2—3-pinnate.* (3) *P. apiifolia.*
- a.a.** *Pinnae articulate to the rachis.* (5) *P. articulata.*

(*) The rhizome of one of the species being unknown to me, I don't follow the arrangement of the sections in this key.

§ 1. EGENOLFIA.

* (1) **P. appendiculata**, **J. Sm.** (not *Bedd.*), *Christ, Farnkr. d. Erde*, 43; *Diels, in Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 102; *Bedd., Ferns S.I.*, tab. CXCIV; *P. marginata*, *Bl., Enum.*, 100; *Acrostichum appendiculatum*, *Willd., Rac., Flor. Btz.*, I, 52; — var. *Hamiltoniana*, *Bedd., Handb. Ind. Ferns*, 426; *Ferns Br.I.*, I, tab. CXI (the 3 lower figures); *P. Hamiltoniana*, *Pr., Tent. Pterid.*, 251; *A. Hamiltonianum*, *Wall., List*, No. 29.

Rhizome firm, woody, short-creeping, the scales brown, lanceolate. Stipes approximate, 7—15 c.M. long, erect, slightly scaly like the rachis, the scales linear. Barren fronds simply pinnate, ovate-lanceolate, sometimes rooting at the apex, 15—50 c.M. long, 6—20 c.M. broad. Pinnae short-stalked, ± approximate, 1/2—2 c.M. broad, the sides parallel, the apex blunt, the edges varying from slightly crenate to cut 1/2-way down to the midrib into oblong, blunt lobes, with a seta or bristle-like tooth in the sinuses, the base unequal-sided, the upper base subtruncate, auricled, the lower truncate or cuneate. Texture herbaceous, firm; surfaces naked; veins visible, pinnate. Fertile fronds on the longest stipes, pinnate; pinnae oblong or linear-oblong, much shorter and narrower than the barren ones.

Var. Hamiltoniana: Fertile pinnae contracted. Sori in bead-like clusters without any connecting leafy membrane.

Trop. Asia; — var. *Hamiltoniana*: *Malacca; Chittagong, Assam.*

§ 2. PSOMIOCARPA.

* (2) **P. Nieuwenhuisii** (*Nieuwenhuisii*), **Rac.**, *Bull. Ac. Crac.*, 1902, 57; *Acrostichum Nieuwenhuisii*, *Rac.*, MS. in *Herb. Bog.*

Rhizome erect, scaly at the extremity, the scales yellowish-brown, spreading, lanceolate. Stipes 5—40 c.M. long, clothed towards the base with many scattered, spreading or deflexed, lanceolate-acuminate, brownish scales, the rest glossy, minutely pubescent or short-fibrillose and blackish-purple, as is the rachis. Fronds lanceolate-oblong, acuminate, 10—50 c.M. long, 7—15 c.M. broad near the base, the apex pinnatifid, the rest pinnate. Pinnae linear-oblong, acuminate, pinnatifid, to 2 1/2 c.M. broad, broadest in the upper half, slightly narrowed towards the base, the lowest a little abbreviate and deflexed. Lobes oblong, erecto-patent, rather close, rounded at the apex, the longest 6—10 m.M. long, 3—5 m.M. broad, the lowest spreading and not rarely a little elongated. Texture firm-herbaceous; surfaces naked; costae fibrillose

above; veins 5—7-jugate, simple. Fertile fronds as long or shorter, narrower, often on the longest stipes; pinnae contracted, with shorter and narrower, more distant, spreading lobes; lowest pinnae not deflexed, except when casually barren.

Borneo.

(3) **P. apiifolia**, **J. Sm.** (not *Hk.*), *Copel.*, *Polypod. Philipp.*, 40; *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 103, B; *Kze, Farnkr.*, I, tab. LXII; *Acrostichum apiifolium*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 415.

Rhizome stout, woody, erect. Stipes of barren fronds 5—8 c.M. long, erect, densely fibrillose-tomentose. Barren fronds 10—15 c.M. each way, deltoid, 3-pinnate. Pinnae close, only the lowest pair with pinnate pinnulae; ultimate divisions oblong-romboidal, $\frac{1}{2}$ —1 c.M. long, the base equally cuneate, the outer edge slightly toothed. Texture herbaceous; rachises tomentose. Fertile fronds on slender, rather naked stipes 15—20 c.M. long, paniced, with a few distant, slender, simple or compound, beaded branches.

Philippines.

§ ?

(4) **P. stenosemioides**, **Copel.**, *Polypod. Philipp.*, 40; *Acrostichum stenosemioides*, *Bk.*, *Journ. of Linn. Soc., Bot.*, XXII, 230.

Stipes of barren fronds to \pm 50 c.M. long, castaneous, covered with a few lanceolate scales towards the base. Fronds oblong-lanceolate, 50—45 c.M. long, 16—20 c.M. broad, 2-pinnatifid. Pinnae sessile, multijugate, lanceolate, pinnatifid, the lower ones to 12 c.M. long, $2\frac{1}{2}$ — $3\frac{1}{2}$ c.M. broad, the margin beset with a few hyaline, deciduous scales. Segments oblong, obtuse, 4—6 m.M. broad. Texture submembranaceous; surfaces naked; main veins pinnate in the segments; veins 6—8-jugate, the lower ones forked, the higher simple. Fertile fronds similar, but the pinnae smaller, less dense, the stipes \pm 60 c.M. long.

Borneo, Philippines.

§ 4. ARTHROBOTRYA.

(5) **P. articulata**, **J. Sm.**, *Copel.*, *Polypod. Philipp.*, 40; *Christ, Farnkr. d. Erde*, fig. 101; *Acrostichum articulatum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 415; *Lomagramma articulata*, *Copel.*, *Philipp. Journ.*, III, 52.

Rhizome woody, wide-creeping, naked. Stipes firm, erect, slightly scaly, \pm 50 c.M. long. Barren fronds 50—80 c.M. long, 20—30 c.M. broad, oblong-ovate. Pinnæ numerous, articulate to the rachis, stalked, rather distant, spreading, lanceolate, pinnate. Pinnulae rhomboidal, articulate, unequal-sided, the lower base narrow-cuneate, the upper more broadly cuneate and often \pm auricled, the edge deeply toothed. Texture herbaceous; costae and veins conspicuous, scaly. Fertile fronds with the pinnulae contracted, blunt, often sessile.

Timor, Celebes and the Philippines across Melanesia

81. STENOSEMIA, *Presl.*

Sori indefinite, densely covering the dorsal or both surfaces of the much contracted linear segments of the fertile fronds.

Rhizome short. Stipes not articulate to the rhizome. Fronds dimorphous, 3—5-angular in general outline, the basiscopical side of the lower segments strongly developed; veins anastomosing \pm as in *Pleocnemia*, i. e. forming 1—2 rows of areolae against the costae and costulae. Hk. Bk., Syn. Fil., tab. VII, fig. 60, t—u; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 105, A—B.

Malayan and Polynesian.

-
- | | |
|--|------------------------------|
| §. Barren fronds ternate, \pm 20—30 c.M. long. | (1) <i>S. aurita</i> . |
| §.§. Barren fronds pinnate. | |
| *. Barren fronds \pm 30 c.M. long. | (2) <i>S. pinnata</i> . |
| *.*. Barren fronds 45—90 c.M. long. | (3) <i>S. Teysmanniana</i> . |

* (1) ***S. aurita***, *Pr.*, Copel., Polypod. Philipp., 41; *Pr.*, Tent. Pterid., tab. X, fig. 24; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 105, A—B; *Acrostichum auritum*, Sw., Rac., Flor. Btz., I, 52; *Polybotrya aurita*, Bl., Hk., Fil. Exot., tab. LXXXI; *Gymnopteris aurita*, Keys., Christ. Farnkr. d. Erde, fig. 124; — var. minor, Christ, Fil. Saras., III, 252.

Rhizome short-creeping or erect, firm, the scales blackish, lanceolate. Sterile fronds ternate, 20—30 c.M. each way, the central segment deeply pinnatifid with oblong, blunt or acute, straight or subfalcate, entire or crenate

lobes. Lateral segments unequal-sided, the upper side cut $\pm \frac{1}{2}$ -way down into erecto-patent lobes, the lower lobes shorter than the medial ones, the lower side broader than the upper, deeply pinnatifid into entire, toothed or lobed lobes, the lowest of which are much elongated, pinnatifid, falcate. Texture herbaceous; surfaces naked. Fertile fronds of the same shape, but all the divisions linear, $1-1\frac{1}{2}$ m.M. broad. — Stipes to 45 c.M. long.

Var. minor: Dimensions $\frac{1}{3}$ smaller than those of the type.

Malaya; Salomon Isles; — var. minor: Celebes.

(2) **S. pinnata**, *Copel.*, Philipp. Journ., I, Suppl., 146.

Rhizome stout, erect. Stipes tufted, black-fuscous, or yellowish towards the frond, clothed with a few distant, deciduous scales, 25—40 c.M. long, the fertile fronds on the longest stipes. Barren fronds deltoid-ovate, ± 30 c.M. long, ± 20 c.M. broad, pinnate, the rachis densely short-hairy. Lowest pinnae stalked, subfalcate, ovate, acuminate, pinnatifid to $\frac{2}{3}$ on the way to the costa, with the lobes on the lower side produced, those near the base shorter, the edges serrate, the sinuses provided with minute setae; next following pinnae sessile, higher adnate, highest connected. Texture membranaceous; surfaces nearly naked. Fertile fronds ± 10 c.M. long, 2-pinnate.

Mindanao.

* (5) **S. Teysmanniana**, *Diels* (not *Bk.*), in Engl. & Prantl, Nat. Pfl. Fam., I⁴, 198; *Acrostichum Teysmannianum*, *Bk.*, Becc., Mal., III, 56.

Stipes castaneous like the rachis, naked, ± 45 c.M. long. Sterile fronds oblong-rhomboidal, 45—90 c.M. long, 50—60 c.M. broad, pinnate. Pinnae erecto-patent, the lower ones short-stalked, acuminate, $5\frac{1}{2}-6$ c.M. broad, deeply pinnatifid nearly to the costa into numerous linear-oblong, bluntish, entire or inciso-crenate to toothed lobes 5—8 m.M. broad, the lower lobes shorter. Texture herbaceous, firm; surfaces naked, or the costae and costulae clothed with numerous short, gland-like hairs. Fertile fronds similar, but smaller and narrower; ultimate divisions linear, ± 2 m.M. broad, united by a very narrow costal wing.

Sumba Island.

82. HYMENOLEPIS, *Kaulfuss.*

Sporangia densely covering the under side of the much contracted, linear or spoon-shaped apex of simple fronds, often protected by the recurved edge when young.

Rhizome \pm creeping. Stipes articulate to the rhizome. Sterile part of the fronds lanceolate; veins anastomosing copiously; areolae with free included veinlets. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 161, F—H; Hk. Bk., Syn. Fil., tab. VIII, fig. 60, cc—ee.

§. Fertile region linear.

‡. Fronds stalked.

*. Fronds cartilaginous.

(1) *H. rigidissima*.

.. Fronds \pm coriaceous, sometimes membranaceous.

(2) *H. spicata*.

(3) *H. callifolia*.

‡.‡. Fronds sessile.

(4) *H. brachystachys*.

§.§. Fertile region spoon-shaped.

(5) *H. platyrhynchos*.

(1) ***H. rigidissima*, Christ**, Bull. Herb. Boiss., 2^e Série, VI, 990.

Rhizome firm, black, crowned with minute, brown, strigillose scales. Stipes seriate, short. Fronds 1—2, linear-lanceolate, 8—16 c.M. long, 2—8 m.M. broad, the base narrowed, the sporangia occupying the upper $\frac{1}{3}$ — $\frac{1}{2}$ of the frond. Texture cartilaginous, indurate; colour blackish-ferruginous; costa distinct; veins hidden. Sori fulvo-testaceous.

Luzon, Mindoro.

* (2) ***H. spicata*, Pr.**, Christ, Farnkr. d. Erde, 129, fig. 561; Hk., Fil. Exot., tab. LXXVIII; Bedd., Ferns S.I., tab. XLVI; *H. revoluta*, Bl., Enum., 201; *Acrostichum spicatum*, L.f. (not L.), Rac., Flor. Btz., I, 51; *A. platyrhynchos*, Ces. (not Hk.), Fel. Becc. Polin., 5; — var. *latifrons*, Bedd., Journ. of Bot., XXVI, 6; — var. *tenella*; *Acrostichum revolutum*, Zoll., var. *tenellum*, Zoll., MS. in Herb. Bog.

Rhizome creeping, the scales short, ovate-lanceolate, short-acuminate. Stipes 1—7 c.M. long, firm, often dark-coloured like the costa. Fronds 15—45 c.M. long, 1—5 c.M. broad, linear-lanceolate, the lower part narrowed very gradually, the edge entire, or slightly and irregularly undulate, the upper part more or less suddenly narrowed into a linear, contracted apex 1—15 c.M. long, 2—4 m.M. broad, with the unaltered edge much recurved, quite enclosing the sori when young. Texture rigid, chartaceous; surfaces naked; costa distinct; veins hidden; areolae with free veinlets. Sori broadly linear, occupying the linear apex of the frond; capsules intermixed with peltate scales.

Var. latifrons: Fronds to 75 c.M. long, 6—7 c.M. broad; fertile part 15—20 c.M. long.

Var. tenella: Fronds linear, 20—50 c.M. long, $\pm \frac{1}{2}$ c.M. broad, broadest $\pm \frac{1}{3}$ — $\frac{1}{6}$ from the base, narrowed gradually towards both ends. Texture membranaceous.

Trop. Asia, Australia, Polynesia, Madagascar, Mascaren Islands; — var. *latifrons: Malacca;* — var. *tenella: ? (Zollinger).*

* (3) **H. callifolia**, *Christ*, Ann. Btz., XX, 128.

Stipes ± 7 c.M. long, thick, brown, canaliculate. Fronds 40—60 c.M. long, ± 5 — $5\frac{1}{2}$ c.M. broad, lanceolate, acuminate, the base narrowed, the sori-ferous apex ± 25 c.M. long, ± 5 m.M. broad, with the margin recurved. Texture coriaceous; surfaces opaque, naked; costa prominent; veins fine, oblique, flexuose; primary areolae broad, divided into smaller ones with included free veinlets. — Perhaps identical with *H. spicata*, *Pr.*, var. *latifrons*.

Borneo.

* (4) **H. brachystachys**, *J. Sm.*, Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 305; *H. spicata*, *Pr.*, var. *brachystachys*, *Hk.*, Gard. Ferns, tab. III; *Acrostichum brachystachyum*, *Rac.*, Flor Btz, I, 51; *A. spicatum*, *L.f.*, var. *brachystachys*, *Hk.*, Spec. Fil., V, 280; *A. validinervis*, *Rac.*, MS. in Herb. Bog.

Much resembling *H. spicata*, *Pr.*, but the scales of the rhizome with a very long, linear-subulate, finely acuminate apex. Fronds sessile, less rigid, the costa often green or pale-coloured.

Malaya.

(5) **H. platyrhynchos**, *Kze.*, *Christ*, Farnkr. d. Erde, 129; *Hk.*, Icon. Pl., X, tab. IM; *Acrostichum platyrhynchos*, *Hk.*, *Hk. Bk.*, Syn. Fil., 424; *Gymnopteris platyrhynchos*, *J. Sm.*, (oldest name).

Stipes rather tufted, short. Fronds 50—40 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, the lower part narrowed gradually, the edge entire, the sori-ferous apex spoon-shaped. Texture very coriaceous; surfaces naked; areolae fine, hidden, with copious free veinlets. Sori in a subterminal patch $2\frac{1}{2}$ —5 c.M. long, ± 1 c.M. broad, not reaching to the edge.

Celebes, Philippines.

83. ACROSTICHUM, *Linné*.

Sori indefinite, densely covering the under surface of the higher pinnae of simply pinnate fronds.

Rhizome stout, erect. Stipes not articulate to the rhizome. Fronds commonly large; pinnae not articulate, uniform or nearly so, the fertile ones rarely much contracted; veins anastomosing copiously and moderately regular; areolae obliquely elongato-subhexagonal, without free veinlets. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 174, F & J. (*)

Pantropical, mostly in brackish marshes.

* **A. aureum**, *L.*, Rac., Flor. Btz., I, 50; Bedd., Ferns S.I., tab. CCIV; *A. inaequale*, *Willd.*, Bl., Flor. Jav., II, 40, tab. XVI; *A. speciosum*, *Willd.*, Bl., l. c., 42, tab. XVII; *Chrysodium aureum*, *Mett.*, Christ, Farnkr. d. Erde, 47; — var. *attenuatum*, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XXI, 9; (? *A. praestantissimum*, *Bory*, Hk. Bk., Syn. Fil., 455; *Ch. praestantissimum*, *Mett.*, Christ, l. c.).

Caudex erect, woody, scaly. Stipes tufted, 50—60 c.M. long, strong, erect, naked, glossy, pale-brown when dried. Fronds 60—300 c.M. long, 50—60 c.M. broad. Pinnae numerous, distant, erecto-patent, stalked, \pm 15—45 c.M. long, 1—9 c.M. broad, ligulate-oblong or lanceolate, both sides often parallel, entire; barren pinnae with the base \pm truncate or cuneate, the apex blunt (*A. inaequale*, *Willd.* = *aureum* typicum), acute or acuminate (*A. speciosum*, *Willd.*), if blunt sometimes retuse with a mucro; upper pinnae fertile, similar to the barren ones but mostly smaller and blunt. Texture firm, often rigidly coriaceous; surfaces naked; costae prominent and veins often so; areolae small, copious, moderately regular, without free veinlets. Sori filling up the space between the costae and margin. — The young plants have the fronds simple or ternate, sterile.

Var. attenuatum: Fertile pinnae much contracted, narrow-linear. Both barren and fertile segments resembling those of *Leptochilus Raapii*, v. *A. v. R.*, but longer and narrowed gradually towards both ends.

Pantropical; — var. *attenuatum*: *Karimon Djawa Islands* (Koorders).

(*) The genus *Acrostichum* formerly included all ferns with the sporangia covering the fruiting surface, without evident differentiation of sori (Copel.).

84. PHOTINOPTERIS, *J. Smith.*

Sporangia crowded over the under surface of the much contracted, linear, highest segments of the pinnatipartite or pinnate fronds.

Rhizome creeping. Fronds articulate to the rhizome, deeply pinnatifid or pinnate; segments dimorphous, the barren ones lanceolate, articulate to the rachis, free and stalked or adnate and joined; veins anastomosing copiously; areolae with included free veinlets. Diels, in Engl. & Prantl, Nat. Pfl. Fam., I⁴, fig. 170, D—F.

Malayan.

-
- §. Fronds deeply pinnatifid in the lower, barren part. (1) *Ph. drynarioides*.
 §.§. Fronds pinnate in the lower, barren part. (2) *Ph. speciosa*.
-

* (1) ***Ph. drynarioides*, *Bedd.***, Handb. Ind. Ferns, 442; Ferns Br.I., II, tab. CCCXXV; *Acrostichum drynarioides*, *Hk.*, Hk. Bk., Syn. Fil., 425; *Polypodium drynarioides*, *Christ*, Farnkr. d. Erde, 117; *Dryostachyum drynarioides*, *Kuhn*, Ann. Mus. Bot. L. B., IV, 296; C. Chr., Ind. Fil.

Fronds subsessile, to 90 c.M. long, deeply pinnatifid below with numerous close, entire, lanceolate, acuminate, sterile lobes. Lobes 10—20 c.M. long, 2½—4½ c.M. broad, connected by a narrow wing, articulate though adnate to the costa (rachis), the lower ones passing down gradually into a mere, undulate wing to the costa, which is continued to its very base; above the sterile lobes numerous distant, linear pinnae sometimes 50 c.M. long, ± 2½ m.M. broad. Texture very coriaceous; surfaces naked; main veins prominent, with transverse veins; areolae copious, small, with free veinlets.

Malaya; Salomon Islands.

* (2) ***Ph. speciosa*, *Bl.*** (not *J. Sm.*), Copel., Polypod. Philipp., 154; Diels, in Engl. & Prantl., Nat. Pfl.Fam., I⁴, fig. 170, D—F; *Ph. rigida*, *Bedd.*; Ferns Br.I., II, tab. CCXI; *Polypodium speciosum*, *Christ*, Farnkr. d. Erde, 121, fig. 546; *Acrostichum rigidum*, *Wall.*, Rac., Flor. Btz., I, 50; *Lomaria speciosa*, *Bl.*, (oldest name); — var. *simplex*; *Ph. simplex*, *J. Sm.*, Journ. of Bot., III, 405.

Rhizome woody, creeping, whitish, the scales brown, subulate, mixed with hairs. Stipes short or long, naked, provided with 2 opposite rows of

few distant glands or auricles. Fronds ovate-lanceolate, 60—90 c.M. long, 20—50 c.M. broad. Pinnae numerous, the lower ones sterile, stalked, jointed at the base, alternate, rather distant, 7—15 c.M. long, $3\frac{1}{2}$ —8 c.M. broad, ovate, acuminate, the stalks provided at the base with a stipule- or auricle-like outgrowth. Upper pinnae fertile, distant, 10—15 c.M. long, 5— $5\frac{1}{2}$ m.M. broad, short-stalked. Texture very coriaceous, rigid, thick; colour light-green; rachis naked; upper surface covered with scattered, deciduous, white-cretaeous dots, lower with rather long hairs, soon naked; main veins connected by raised transverse veins; areolae copious, with free veinlets.

Var. simplex: Fertile pinnae reduced to 1, the terminal one.

Malaya.

85. CHEIROPLEURIA, *Presl.*

Sori indefinite, covering the whole under surface of the much contracted, nearly linear fertile fronds.

Rhizome creeping. Fronds scattered, not articulate to the rhizome, dimorphous. Barren fronds simple or more generally 2-fid, the lobes sometimes 2-fid again; main ribs 2 or more in each lobe, uniting towards the apices of the segments; veins anastomosing copiously. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 175.

Australasia.

* **Ch. bicuspis, Pr.**, Christ, Farnkr. d. Erde, 128, fig. 360; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 175; *Acrostichum bicuspe*, Hk., Rac., Flor. Btz., I, 48; *Gymnopteris Vespertilio*, Hk., Lond. Journ. of Bot., V, tab. VII—VIII; *Polypodium bicuspe*, Bl., Flor. Jav., II, 175, tab. LXXVIII, B; — var. *integrifolia*, Hk. Bk., Syn. Fil., 421.

Rhizome woody, creeping or subscandent, densely clothed with silky, yellow hairs. Stipes of barren fronds \pm 50 c.M. long, firm, naked. Barren fronds \pm 20 c.M. long, 10 c.M. or more broad, the base cuneate, rounded or subcordate, the upper part consisting of 2 elongate-deltoid, divaricate, acute or acuminate lobes \pm half as long as the whole frond, sometimes 2-fid again. Texture coriaceous; surfaces naked; main ribs \pm 4—8, springing from the base of the frond, uniting towards the apices of the lobes; areolae copious. Fertile fronds 15—25 c.M. long, 1— $1\frac{1}{2}$ c.M. broad, linear-

lanceolate, simple, narrowed towards both ends, with \pm 3 prominent ribs, on stipes 45 c.M. or more long.

Var. integrifolia: Barren fronds ovate-acuminate, simple.

Sumatra, Java, Borneo, New Guinea, Philippines; Formosa, Liu-Kiu.

86a. LEPTOCHILUS, *Kaulfuss.*

Sporangia densely crowded over the under surface of the contracted fertile fronds or pinnae of fertile fronds. Paraphyses mostly small or wanting. Spores generally bilateral or ellipsoidal, often purveyed with more or less deciduous, elevate, reticulate, entire, plicate or fimbriate membranes.

Rhizome generally creeping. Stipes not articulate to the rhizome. Fronds dimorphous, simple or pinnate; veins anastomosing copiously, \pm as in *Aspidium*, very rarely as in *Nephrodium*; areolae generally with included free veinlets.

Pantropical.

Arrangement of the sections.

- § 1. **POECILOPTERIS** (*Gymnopteris*). Main veins distinct nearly or quite to the edge. Hk. Bk., Syn. Fil., tab. VII, fig. 60, v—x, aa—bb; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 106—107.
- § 2. **CHRYSODIUM**. Main veins wanting or indistinct. Fronds simple or ternate, the segments not articulate to the rachis. Hk. Bk., l. c., fig. 60, y—z.
- § 3. **LOMAGRAMMA**. Like § 2, but the fronds pinnate, the segments articulate to the rachis. (*)

§ 1. POECILOPTERIS

a. Barren fronds typically simple and entire, but some of them sometimes ternate or quinate.

§. Veins anastomosing in costal arches only, the rest free.

(1) *L. oligodictyus*;

§.§. Veins anastomosing copiously.

(*) This section would perhaps better be treated as a distinct genus,

- *. *Barren fronds 1—1½ c.M. broad. Areolae without free veinlets.*
(2) L. Linnaeanus.
- *.*. *Barren fronds 4—10 c.M. broad. Areolae with free veinlets.*
(5) L. decurrens.
- a.a.** *Barren fronds typically 3-fid, but sometimes simple or pinnate.*
- ×. *Segments of the fertile fronds lanceolate-oblong, rather short.*
(4) L. trifidus.
- ×.×. *Segments of the fertile fronds narrowly linear, rather long.*
(5) L. latifolius.
- a.a.a.** *Barren fronds typically 3-foliate or pinnate, but some of them sometimes simple, or very rarely 2-pinnate in No. 10a.*
- b.** *Fronds pedatisect.* (6) L. Hollrungii.
- b.b.** *Barren fronds with 1—6 pairs of lateral pinnae.*
- c.** *Fronds small, under 12½ c.M. long.*
- +. *Lateral pinnae 1-jugate.* (7) L. zeylanicus.
- +.+. *Lateral pinnae 2—6-jugate.* (8) L. inconstans.
- c.c.** *Fully developed fronds larger, more than 12½ c.M. long.*
- d.** *Lateral pinnae truncate at the base, the edge broadly crenate and finely toothed.* (9) L. prolifer.
- d.d.** *Lateral pinnae cuneate or rounded at the base, the edge entire, repand, toothed or sinuate.*
- e.** *Fronds scattered.* (10a) L. heteroclitus.
(10b) L. stolonifer. (*)
- e.e.** *Fronds approximate.* (11) L. diversifolius.
- b.b.b.** *Barren fronds as a rule copiously pinnate.*
- 1.** *Fronds to 30 c.M. long.*
- §. *Pinnae entire or crenate.*
- *. *Pinnae entire, acuminate, narrowed at the base.*
(12) L. Preslianus.
- *.*. *Pinnae crenate, acute, truncate at the base.*
(15) L. exsculptus.
- §.§. *Pinnae lobed.* (14) L. celebicus.
- 1.1.** *Fully developed fronds mostly 30—70 c.M. long.*
- 2.** *Pinnae herbaceous or subchartaceous, under 4 c.M. broad.*
- 3.** *Rachis narrowly winged. Pinnae under 8 c.M. long.*
(15) L. virens.

(*) I don't know whether this species is well-classified here.

3.3. *Rachis* not winged. *Pinnae* \pm 9—15 c.M. long in the barren fronds.

4. *Pinnae* whether or not lobed, the lobes without produced anterior teeth. (16a) *L. cuspidatus*.

4.4. *Pinnae* subpinnatifid, the lobes toothed, with the lower anterior teeth produced. (16b) *L. Naumanni*.

2.2. *Pinnae* subcoriaceous, more than 4 c.M. broad in the fully developed barren fronds.

×. *Pinnae* $3\frac{1}{2}$ —5 c.M. broad. (17) *L. Zollingeri*.

×.×. *Pinnae* 5—8 c.M. broad, tinged with red.

(18) *L. scalpturatus*.

(1) **L. oligodictyus**, *C. Chr.*, Ind. Fil.; *Acrostichum oligodictyon*, *Bk.*, Journ. of Linn. Soc., Bot., XXIV, 261.

Rhizome short-creeping. Stipes 5—8 c.M. long, naked, pale-brown. Fronds lanceolate, 20—25 c.M. long, 1—2 c.M. broad at the middle, narrowed gradually to both ends. Texture rather rigid; both surfaces naked; veining of Nephrodium, paucijugate, only the lowest veins anastomosing. Fertile fronds linear, 5—8 m.M. broad, on stipes 15—20 c.M. long.

Borneo.

* (2) **L. Linnaeanus**, *Fée*, *C. Chr.*, Ind. Fil.; *L. heteroclitus*, *C. Chr.*, var. *Linnaeanus*, *Christ*, *Philipp. Journ.*, II, 160; *Acrostichum Linnaeanum*, *Hk.*, *Hk. Bk.*, Syn. Fil., 417; *Hk.*, 2nd Cent. of Ferns, tab. XXVI; *Gymnopteris Linnaeana*, *Christ*, *Copel.*, *Polypod. Philipp.*, 41.

Rhizome creeping, scaly. Stipes more or less distant, $\frac{1}{2}$ —5 c.M. apart, erect, slender, 5—15 c.M. long, scaly \pm throughout or at the base only. Barren fronds 10—25 c.M. long, 1—1 $\frac{1}{2}$ c.M. broad, linear-lanceolate, entire or repand, narrowed rather gradually towards both ends, simple or sometimes with a terminal pinna similar to the simple fronds and below this 1—2 pairs of mostly very small, sessile or stalked, oblong or linear-lanceolate lateral segments to \pm 12 $\frac{1}{2}$ c.M. long, to \pm $\frac{1}{2}$ c.M. broad. Texture herbaceous to papyraceous; surfaces naked; main veins distinct nearly to the edge, more or less zigzag; areolae copious; no free veinlets. Fertile fronds simple or sometimes pinnate, similar to the barren ones but shorter and narrower, sometimes linear, on stipes 10—20 c.M. long.

Malaya, Annam.

* (3) **L. decurrens**, *Bl.*, Enum., 206; *Acrostichum variabile*, *Hk.*, *Hk. Bk.*, Syn. Fil., 417 (excl. var.); *A. variabile*, *Hk.*, var. *decurrens*, *Rac.*, Flor. Btz., I, 49; *Gymnopteris variabilis*, *Bedd.*, Handb. Ind. Ferns, 429; Ferns Br.I., II, tab. CCLXXII; *G. decurrens*, *Moore* (not *Hk.*), *Hk.*, Gard. Ferns, tab. VI; — var. *rasamalae*, *Rac.*, l. c., 50.

Rhizome wide-creeping, the scales linear-lanceolate. Stipes $\frac{1}{2}$ —2 c.M. apart, 0—10 c.M. long. Sterile fronds ovate-oblong or spatulate, 15—45 c.M. long, 4—10 c.M. broad, acute or acuminate, the edge entire, the base cuneate or rounded, decurrent on the stipe. Texture papery or herbaceous; surfaces naked; midrib prominent; main veins distinct, straight or zigzag, reaching nearly to the edge; areolae with free veinlets. Fertile fronds linear or linear-lanceolate, 15—50 c.M. long, 2—7 m.M. broad, on stipes to 50 c.M. long.

Var. rasamalae: Barren fronds 40—80 c.M. long, the apex acuminate; fertile fronds lanceolate, to 3 c.M. broad, long-stalked.

Trop. Asia; — var. *rasamalae*: *Java*.

* (4) **L. trifidus**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 26.

Rhizome creeping. Stipes approximate, \pm 10—20 c.M. long, those of the fertile fronds the longest, densely clothed with lanceolate-subulate, short, crisped, brown scales, at length less scaly. Fronds simple, lanceolate or more commonly 3-fid with the lateral lobes similar to the central one but shorter. Simple fronds \pm 25 c.M. long, \pm 7 c.M. broad, the apex rather suddenly acuminate, the edge irregularly wavy, the base suddenly narrowed and then short-decurrent. Central lobe of the 3-fid fronds similar to the simple fronds, but joined at the base with the lateral segments, which have the lower base suddenly short-decurrent upon the stipe. Texture herbaceous; surfaces naked; costae \pm scaly beneath, but more distinctly towards the base; main veins conspicuous, nearly straight to the edge, united by strongly arching transversing veins forming \pm 6—7 distinct primary areolae on each side of the costae, divided into finer secondary ones with free veinlets. Fertile fronds quite similar to the barren ones but much smaller, the central lobe \pm 8 c.M. long, \pm 2 c.M. broad, the lateral lobes smaller. — The description is taken from specimens cultivated in the Buitenzorg gardens and probably gathered in one of the malayan islands. Near *L. Harlandii*, *C. Chr.* (= *Gymnopteris decurrens*, *Hk.*, Fil. Exot., tab. XCIV).

? *Malayan*.

(5) **L. latifolius**, *C. Chr.*, Ind. Fil.; *Acrostichum taccaefolium*, *Hk.*, *Hk. Bk.*, Syn. Fil., 418; *Gymnopteris latifolia*, *Pr.*, Tent. Pterid., 244; *G. taccaefolia*, *J. Sm.*, Copel., Polypod. Philipp, 42; *G. trilobata*, *J. Sm.*, *Hk.*, Icon. Pl., X, tab. CMVII; *G. subquinquefida*, *Pr.*, Epim., 151; *Hemigramma latifolia*, *Copel.*, Philipp. Journ., II, 406; III, 51, tab. I; *Polybotrya latifolia*, *Meyen.* (oldest name).

Rhizome short, stout, creeping, the scales copious, dark-reddish-brown linear. Stipes of barren fronds $2\frac{1}{2}$ —10 c.M. long, those of fertile fronds about 50 c.M. long, both clothed below with narrow, spreading scales. Barren fronds to 50 c.M. long or longer, typically 5-fid, sometimes simple and 5—8 c.M. broad or pinnate, sometimes rooting at the apex, the pinnate fronds to 60 c.M. long. Pinnae acute, entire, lanceolate, rachis winged in the more complex forms and the lowest pinnae forked. Texture papyraceous; surfaces naked; main veins distinct nearly to the edge, with regular transverse veinlets and copious areolae with included free veinlets. Fertile fronds typically 5-fid, the central segment \pm 15 c.M. long, the lateral ones \pm 10 c.M. long, both 5—4 m.M. broad, exceptionally simple or pinnate. — Exceedingly variable. Copeland includes this under *Hemigramma latifolia* (*Syngamma Zollingeri*, *Diels.*; probably he is correct in this, and I am inclined to believe that *Hemigramma* (*Syngamma* §) would better be treated as a distinct genus.

Philippines.

(6) **L. Hollrungii**, *C. Chr.*, Ind. Fil.; *Acrostichum Hollrungii*, *Bk.*, Ann. of Bot., V, 495; *Gymnopteris Hollrungii*, *Kuhn*, Schum. & Hollr., Flor. Kais. Wilhl., 8.

Rhizome erect, densely clothed with ferrugineous, long-acuminate, bristle-like scales. Stipes scaly at the base, 40—50 c.M. long, canaliculate. Sterile fronds 25—50 c.M. long, 35—40 c.M. broad, with the base pedatisect, the higher part pinnatifid; lobes lanceolate, 25—50 c.M. long, 5—4 c.M. broad, entire, acuminate. Texture subcoriaceous, surfaces naked. Fertile fronds 25—30 c.M. long, 30—40 c.M. broad, pinnatifid, on stipes 60—65 c.M. long; lobes 15—20 c.M. long, linear, entire or slightly sinuato-repand.

New Guinea.

* (7) **L. zeylanicus**, *C. Chr.*, Ind. Fil.; *Acrostichum quercifolium*, *Retz*, *Hk. Bk.*, Syn. Fil., 418; *Gymnopteris quercifolia*, *Bernh.*, *Hk.*, Fil. Exot., tab. LXXX; *Hk.*, Icon. Pl., X, tab. CMV; *Bedd.*, Ferns S.I., tab. XLVII; *Ophioglossum zeylanicum*, *Houtt.* (oldest name).

FILICES.

Rhizome stout, creeping, scaly. Stipes of barren fronds $2\frac{1}{2}$ —5 c.M. long, clothed with soft, brownish or reddish, spreading hairs. Barren fronds oblong, 7—10 c.M. long, $5\frac{1}{2}$ —6 c.M. broad, with an ovate or oblong, large, blunt, bluntly lobed terminal pinna, the lobes sometimes reaching to $\frac{1}{4}$ -way down to the costa. Below this usually a single pair of small, sessile, roundish or ovate, spreading, auricled segments, blunt at the apex, \pm truncate or subcordate at the base. Texture thinly herbaceous; main veins distinct to the edge; veins hairy beneath; areolae copious, with free veinlets. Fertile fronds ternate, on slender stipes 15—25 c.M. long, naked except at the base; terminal segment linear, $2\frac{1}{2}$ —5 c.M. long, $\pm 2\frac{1}{2}$ m.M. broad, entire; lateral segments erecto-patent, similar but shorter.

Trop. Asia.

(8) **L. inconstans**, *C. Chr.*, Ind. Fil.; *Gymnopteris inconstans*, *Copel.*, Polypod. Philipp., 43; *L. heteroclitus*, *C. Chr.*, var. *inconstans*, *Christ Philipp. Journ.*, II, 160.

Rhizome wide-creeping, scaly. Sterile fronds rarely simple, often pinnate, 4—8 c.M. or sometimes more long, linear, lanceolate, oblanceolate or ovate, on stipes 1—3 c.M. long, the stipes slender, scaly or naked; terminal pinna commonly very long, narrow, proliferous; lateral pinnae 2—6-jugate, erecto-patent, commonly obovate, rarely orbicular or lanceolate, obtuse, entire or rarely serrate, \pm decurrent; veins inconspicuous; areolae few, with free veinlets. Fertile fronds linear, on stipes 5—10 c.M. long, pinnate, the stipes erect, filiform, naked above, the pinnae 2—4-jugate, distant, the lowest sometimes stalked, orbicular or oblong, entire, 1—4 m.M. long; free veinlets few or none.

Luzon.

(9) **L. prolifer**, *C. Chr.*, Ind. Fil.; *Polybotrya prolifera*, *Bory. Bél.*, *Voy. Bot.*, II, 18; *Poecilopteris prolifera*, *Pr.*, *Epim.*, 172.

Stipes naked, subtrigonal, twisted, $\pm 17\frac{1}{2}$ c.M. long. Fronds dimorphous. Barren fronds pinnate with ± 5 pinnae on each side below the often proliferous terminal one, $\pm 22\frac{1}{2}$ c.M. long. Pinnae alternate, 7—10 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, acuminate, ovate-oblong, short-stalked, truncate at the base, the edge broadly and bluntly crenate, the crenations finely toothed. Colour pale-green; veins whitish and very conspicuous beneath. Fertile fronds with the pinnae linear, pale-reddish, slightly twisted.

New Guinea; Br. India.

* (10a) **L. heteroclitus**, *C. Chr.*, Ind. Fil.; *Acrostichum flagelliferum*, *Wall.*, *Rac.*, *Flor. Btz.*, I, 56; *Hk. & Grev.*, *Ic. Fil.*, tab. XXIII; *Bl.*, *Flor. Jav.*, II, 57, tab. XIII; *Gymnopteris flagellifera*, *Bedd.*, *Handb. Ind. Ferns*, 453; *Poecilopteris flagellifera*, *J. Sm.*, *Bedd.*, *Ferns Br.I.*, I, tab. CXII; *Acrostichum heteroclitum*, *Pr.*, (oldest name); — var. *eurybasis*, *Christ*, *Philipp. Journ.*, II, 159; — var. *Foxworthyi*, *Christ*, *l.c.*, 160.

Rhizome creeping, firm, the scales brown, lanceolate. Stipes scattered, 15—25 c.M. long, often scaly. Fronds 15—45 c.M. long, simple or more commonly with an entire, toothed or repand, often long-acuminate and rooting terminal pinna and 1—6 pairs of stalked, lanceolate, similar but not long-acuminate lateral ones; terminal pinna to 40 c.M. long, lateral ones 7—15 c.M. long, 2—5 c.M. broad. Texture herbaceous; surfaces naked; main veins distinct to the edge; areolae copious, with or without free veinlets. Fertile fronds similar, but the pinnae smaller and narrower, sometimes partially covered with meniscioid sori. — Variable.

Var. eurybasis: Differs from the type by its aberrant fronds being in part 2-pinnate, deltoid at the enlarged base, posteriorly deeply incised. Even the fertile fronds are sometimes 2-pinnate at the base.

Var. Foxworthyi: Fronds reduced; lateral pinnae 2, reduced, orbicular or oblong, hardly $\frac{1}{2}$ c.M. long; terminal pinna linear or oblong, linear-lanceolate when fertile, with 2 ovate lateral lobes.

Trop. & Subtrop. Asia; Melanesia; — var. *eurybasis*: *Mindanao*; — var. *Foxworthyi*: *Luzon*.

(10b) **L. stolonifer**, *Christ*, *Bull. Herb. Boiss.*, 2^e Série, VI, 1004.

Rhizome stoloniferous, the stolones bearing rows of simple fronds similar to the pinnae of the normal barren fronds, decurrent at the base, on stipes \pm 6 c.M. long. Stipes of barren fronds stramineous, 20—25 c.M. long, more or less densely clothed with black, subulate or setiform scales. Barren fronds deltoid-ovate, \pm 40 c.M. long, \pm 50 c.M. broad, pinnate. Terminal pinna largest, \pm 25 c.M. long, \pm 4 $\frac{1}{2}$ c.M. broad, oblong, acute, repando-lobate, decurrent at the narrowed base, the lobes irregular, short, obtuse; lateral pinnae distant, alternate, \pm 2—3 on each side, erecto-patent, \pm 20 c.M. long, 2—4 c.M. broad, sessile, the base rather unequal-sided, sometimes subauriculate on the lower side, the edge repando-lobate or undulato-crenate, the apex acute, the lobes \pm 10 on each side, short; lowest pinnae hardly reduced. Texture herbaceous; costae distinct; main veins distinct when seen towards the light, straight from the costa to the edge; areolae with free

veinlets. Fertile fronds $\pm \frac{1}{2} \times$ as long as the barren ones, on stipes \pm 14 c.M. long; pinnae 5—7, much contracted, linear-lanceolate, \pm 5 c.M. long, 7—10 m.M. broad, subacute, the base subcordate.

Luzon.

* (11) **L. diversifolius**, *C. Chr.*, Ind. Fil.; *Acrostichum diversifolium*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 48; *Bl.*, *Flor. Jav.*, II, 56, tab. XII; *Poecilopteris diversifolia*, *Pr.*, *Tent. Pterid.*, 242.

Rhizome woody, creeping, the scales lanceolate. Fronds approximate; barren fronds \pm 50—70 c.M. long, long-stalked, simple or more commonly with a terminal segment to 40 c.M. long, to $12\frac{1}{2}$ c.M. broad, ovate-lanceolate, acuminate, entire or repando-sinuate and below this 1—5 pairs of lateral segments similar to the terminal one, but shorter and narrower, to 20 c.M. long, to 7 c.M. broad, stalked. Texture herbaceous; surfaces naked; areolae with few free veinlets. Fertile fronds with 1—2 pairs of short-stalked, ovate-lanceolate lateral pinnae to 10 c.M. long, to $1\frac{1}{4}$ c.M. broad and a longer terminal one.

Malaya.

(12) **L. Preslianus**, *C. Chr.*, Ind. Fil.; *Acrostichum Preslianum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 420; *Poecilopteris Presliana*, *Moore*, *Bedd.*, *Ferns Br.I.*, II, tab. CCLXIX; *Gymnopteris Presliana*, *J.Sm.*, *Bedd.*, *Handb. Ind. Ferns*, 459; *Heteronevron Preslianum*, *Fée*, (oldest name).

Rhizome stout, creeping, the scales narrowed, lanceolate. Stipes 10—25 c.M. long, more or less scaly like the rachis and the stalks of the pinnae. Fronds 15—50 c.M. long, pinnate with \pm numerous erecto-patent, remote pinnae. Lower pinnae stalked, 5—10 c.M. long, $1-1\frac{1}{2}$ c.M. broad, lanceolate, acuminate, the base narrowed, the edge entire. Texture membranaceous or subcoriaceous; rachis \pm winged, especially towards the apex; surfaces naked; venation as in *Nephrodium*. Fertile fronds generally longer stalked; pinnae much narrowed, lanceolate, 2—5 c.M. long, 5—7 m.M. broad.

Philippines; South. India, Madagascar.

(15) **L. exsculptus** (*exsculptus*), *C. Chr.*, Ind. Fil.; *Acrostichum exsculptum*, *Bk.*, *Journ. of Bot.*, XXVI, 526.

Rhizome creeping. Stipes of barren fronds slender, naked, \pm 15 c.M. long. Barren fronds lanceolate, pinnate, 20—50 c.M. long, 5—7 c.M. broad. Pinnae multijugate, sessile, lanceolate, acute, crenate, truncate at the base, $2\frac{1}{2}-3\frac{1}{2}$ c.M. long, 8—9 m.M. broad, the lower ones not reduced. Texture

rigid; surfaces naked; veins anastomosing as in *Nephrodium*; main veins distinct to the edge; veins 3-jugate. Fertile fronds \pm 50 c.M. long, \pm 5 $\frac{1}{2}$ c.M. broad, on longer stipes; pinnae sessile, remote, 4—5 m.M. broad.

Borneo.

(14) **L. celebicus**, *C. Chr.*, Ind. Fil.; *Acrostichum celebicum*, *Bk.*, *Kew Bull.*, 1901, 145.

Stipes densely tufted, slender, naked, 7—15 c.M. long or longer, those of the fertile fronds the longest. Fronds oblong-lanceolate, the barren ones 12—15 c.M. long, 5—8 c.M. broad, pinnate. Pinnae sessile, lanceolate, 1 $\frac{1}{5}$ —1 $\frac{1}{2}$ c.M. broad at the base, those of the fertile fronds narrower, slightly pinnatifid, the lobes obtuse, the base truncate, auricled on the upper side. Rachis slender, pubescent; under surface obscurely pubescent; main veins pinnate; veins 3—4-jugate, anastomosing at the apex.

Celebes.

* (15) **L. virens**, *C. Chr.*, Ind. Fil.; *Acrostichum virens*, *Wall.*, *Hk.* *Bk.*, *Syn. Fil.*, 420, p. p.; *Hk. & Grev.*, *Ic. Fil.*, tab. CCXXI; *Gymnopteris virens*, *Keys.*, *Pol. Cyath. Bung.*, 35; *G. contaminans*, *Bedd.*, *Handb. Ind. Ferns*, 435; *Poecilopteris repanda*, *Bedd.*, *Ferns S.I.* tab. CCII; *P. semicordata*, *Moore*, *Bedd.*, *Ferns Br.I.*, II, tab. CCLXX.

Rhizome thick, short-creeping. Stipes scaly, 10—20 c.M. long. Fronds 30—60 c.M. long, pinnate; rachis \pm scaly, more or less winged. Pinnae 11—30 on each side, sessile or short-stalked, lanceolate, to 8 c.M. long, to 2 $\frac{1}{2}$ c.M. broad, the point acute, the edge entire, crenate or more or less deeply lobed or repand, with the lobes crenate or toothed; terminal pinna often \pm elongated and proliferous. Texture rather herbaceous; costae prominent; main veins distinct to \pm $\frac{2}{3}$ of the way to the margin; areolae broad, irregular, in 3—4 rows, the costal ones without free veinlets, the others generally with several free or irregularly joining included veinlets. Fertile fronds with the pinnae much contracted.

Malacca, Celebes, Philippines; Burmah, South. India, Ceylon.

* (16a) **L. cuspidatus**, *C. Chr.*, Ind. Fil.; *Acrostichum repandum*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 55; *Bl.*, *Flor. Jav.*, II, 39, tab. XIV—XV; *Gymnopteris repanda*, *Christ*, *Farnkr. d. Erde*, 50; *Poecilopteris repanda*, *Pr.*, *Epim.*, 172; *Nephrodium cuspidatum*, *Pr.*, (oldest name); — var. *Quoyanus*; *A. Quoyanum*, *Gaud.*, *Freye.*, *Voy. Ur. Phys.*, *Bot.*, 506, tab. III.

Rhizome creeping, the scales, short, lanceolate. Stipes of barren fronds 15—50 c.M. or more long, naked or slightly scaly. Barren fronds 30—70 c.M. long, 20—30 c.M. broad, sometimes elongated, with about 10—20 pinnae on each side; lower pinnae 10—15 c.M. long, 2—3 c.M. broad, short-stalked, the edge cut into entire, crenate or serrate, blunt lobes reaching to $\frac{1}{3}$ of the way down, sometimes with a seta in the sinuses. Texture herbaceous; rachis and both surfaces naked, or the rachis scaly; main veins distinct; areolae copious, without free veinlets. Pinnae of fertile fronds 5—10 c.M. long, 5—7 m.M. broad, stalked, entire or more or less deeply repand.

Var. Quoyanus: Barren pinnae lobed $\frac{1}{2}$ -way down or more to the costa, the lobes toothed.

Trop. Asia, Australia, Polynesia, Seychelles; — var. *Quoyanus: Philippines, Moluccas, New Guinea.*

(16b) **L. Naumanni, C. Chr.,** Ind. Fil.; *Heteroneuron Naumanni, Kuhn, Forsch. Gaz., IV, Farne, 5, tab. I; Gymnopteris Naumanni, Diels, Schum. & Laut., Flor. deut. Sch.geb., 117.*

Rhizome creeping, elongate, clothed like the base of the stipes with lanceolate, obtuse, ferruginous scales. Fronds distichous, pinnate, pinnatifid towards the often proliferous apex. Stipes of barren fronds 18—24 c.M. long, stramineous, deeply sulcate, sparingly scaly. Fronds 25—47 c.M. long, 14—22 c.M. broad, oblong, acuminate. Lower pinnae stalked, 9—13 c.M. long, ± 2 c.M. broad, lanceolate-acuminate, subpinnatifid, the apex sharply serrate, the base cuneate; higher pinnae sessile, highest confluent. Lobes oblong, toothed, the lower anterior teeth produced. Texture subchartaceous; surfaces naked; colour greenish-opaque; costulae prominent; areolae distinct, lax, appendiculate. Fertile fronds 50—55 c.M. long, 18—20 c.M. broad, ovate-lanceolate, acuminate, on stipes to ± 42 c.M. long; lower segments subpetiolate, higher ones sessile, the longest 10—15 c.M. long, ± 1 c.M. broad, lobato-pinnatifid, acuminate; lobes ovate or elongate, entire; areolae in 1 row on each side of the costulae.

New Guinea; New Hannover.

(17) **L. Zollingeri, Fée,** Gen., 55; *Acrostichum subrepandum, Hk., Hk. Bk., Syn. Fil., 419; Gymnopteris subrepanda, J. Sm., Copel., Polypod. Philipp., 45; Poecilopteris subrepanda, Pr., Bedd., Ferns Br.I., II, tab. CCCXXXIX; Acrostichum Zollingeri, Kze, (oldest name).*

Rhizome woody, wide-creeping. Stipes of barren fronds stout, erect,

nearly naked. Barren fronds varying from simple, \pm 30 c.M. long, \pm 5 $\frac{1}{2}$ —5 c.M. broad, to copiously pinnate, 60 c.M. long, 50 c.M. broad, with several linear-oblong, entire or subrepand pinnae on each side, which are sometimes 15—25 c.M. long, \pm 5 c.M. broad. Texture subcoriaceous; rachis and surfaces naked; main veins distinct nearly to the edge, with copious free veinlets in the areolae. Fertile fronds similar, but smaller.

Malaya; Fiji.

(18) **L. sculpturatus**, **C. Chr.**, Ind. Fil.; *Acrostichum costatum*, Wall., Hk., Spec. Fil., V, 262; *Gymnopteris costata*, Bedd., Copel., Polypod. Philipp., 44; *Poecilopteris costata*, Moore, Ind., 8; *Heteronevron sculpturatum*, Fée, (oldest name).

Rhizome creeping, bearing subulate scales. Stipes 45 c.M. or less long. Fronds 60 c.M. or more long. Pinnae up to 35 c.M. long and 8 c.M. broad, stalked, acuminate, entire, crenate or sinuate. Texture subcoriaceous; surfaces naked; main veins prominent, close; areolae and excurrent included veinlets copious. Pinnae of fertile fronds smaller and more coriaceous.

Philippines; North. India, Burmah.

§ 2. CHRYSODIUM.

- a.** Fertile fronds simple, as are the barren ones.
 - b.** Barren fronds small, under 7 $\frac{1}{2}$ c.M. long.
 - *. Fertile fronds obtuse. (19a) L. minor.
 - *.*. Fertile fronds acuminate. (19b) L. rizalianus.
 - b.b.** Barren fronds larger, more than 7 $\frac{1}{2}$ c.M. long when fully developed.
 - c.** Barren fronds rounded at the base. (20) L. antrophyoides.
 - c.c.** Barren fronds subacute or gradually narrowed at the base.
 - d.** Barren fronds 10—15 c.M. long, to \pm 1 c.M. broad.
 - *. Rhizome short. (21a) L. hydrophyllus.
 - *.*. Rhizome long. (21b) L. modestus.
 - d.d.** Barren fronds 15—45 c.M. long, 1—5 c.M. broad.
 - e.** Rhizome creeping. (22) L. lanceolatus.
 - e.e.** Rhizome scandent. (23) L. axillaris.
 - a.a.** Fertile fronds ternate, barren ones simple or ternate. (24) L. Raapii.
-

* (19a) **L. minor**, *Fée*, C. Chr. Ind. Fil.; *Acrostichum minus*, *Mett.*, Hk. Bk., Syn. Fil., 420; *Gymnopteris minor*, *Hk.* (not *Bedd.*), *Copel.*, Philippod. Philipp., 42; Hk., 2nd Cent. of Ferns, tab. LXXVIII (the description, not the figure, which is *Asplenium*); *G. minus*, *Bedd.*, Handb. Ind. Ferns, 450; Ferns Br.I., I, tab. CXVI.

Rhizome wide-creeping, firm, scaly towards the apex, the scales lanceolate. Stipes remote, slender, those of the barren fronds 2½—5 c.M. long. Sterile fronds 3½—5 c.M. long, ± 1½—2½ c.M. broad, ovate-oblong, the point bluntish, the edge quite entire, the base tapering gradually into the stipe. Texture membranaceous or herbaceous; surfaces naked; costa distinct; areolae with free veinlets, bounded by a ± wavy intramarginal line. Fertile fronds 3½—5 c.M. long, ± 5 m.M. broad, linear-lanceolate, obtuse, on stipes 7½—10 c.M. long.

Philippines, Malacca; North. India.

(19b) **L. rizalianus**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1004; *L. normalis*, *Copel.*, Philipp. Journ., III, 31, p. p.

Rhizome short-creeping, slender, crowned with flaccid, brown, subulate scales. Fronds scattered or approximate. Barren fronds rather sessile, spatulate, cuneate-oblong, ± 4½ c.M. long, ± 1 c.M. broad, obtuse, the base narrowed. Texture herbaceous; costa distinct; areolae hidden, in 1 or rarely 2—5 rows on each side, with free included veinlets. Fertile fronds linear, ± 6 c.M. long, ± 1½ c.M. broad, acuminate, on stipes ± 6 c.M. long.

Luzon.

(20) **L. antrophyoides**, *C. Chr.*, Ind. Fil.; *Acrostichum antrophyoides*, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 231.

Rhizome short-creeping. Fronds sessile, oblanceolate, 15—20 c.M. long, 2—2½ c.M. broad, the point acute, the base rounded. Texture subcoriaceous; surfaces naked; veins obscure, immersed; areolae small. Fertile fronds linear, rigid, 20—25 c.M. long, 5—5 m.M. broad, bearing the sori in the upper half, the lower half narrowed to a naked wing to the costa.

Borneo.

(21a) **L. hydrophyllus**, *Copel.*, Philipp. Journ., I, Suppl., 146.

Rhizome short-creeping, clothed at the extremity with short, fuscous scales. Stipes 1—7 c.M. long, those of the fertile fronds the longest. Barren fronds linear-lanceolate, acute at the apex, subacute at the base, 1 c.M.

broad, \pm 10 c.M. long and entire, rarely 20 c.M. long and undulate. Texture fleshy, papyraceous when dry; surfaces \pm naked; veins hidden, inconspicuous; areolae with few free veinlets. Fertile fronds linear, \pm 7 c.M. long, \pm 3 m.M. broad, obtuse, narrowed gradually into the stipe.

Mindanao.

(21b) **L. modestus**, *C. Chr.*, Ind. Fil.; *Acrostichum modestum*, *Bk.*, Journ. of Linn. Soc., Bot., XXII, 231.

Rhizome slender, wide-creeping, the scales small, lanceolate. Stipes distant, slender, 5—8 c.M. long. Fronds simple, 10—15 c.M. long, 5—9 m.M. broad at the middle, tapering gradually to the base and acuminate apex, the margin entire, the surfaces naked. Areolae hexagonal. Fertile fronds 5—8 c.M. long, 4—4½ m.M. broad.

Borneo.

* (22) **L. lanceolatus**, *Fée*, *C. Chr.*, Ind. Fil.; *Acrostichum lanceolatum*, *Hk.*, *Hk. Bk.*, Syn. Fil., 420; *Gymnopteris lanceolata*, *Bedd.* (not *Hk.*), *Copel.*, Polypod. Philipp., 42; *G. Feei*, *Moore*, *Bedd.*, Ferns S.I., tab. XLVIII; *G. variabilis*, *Bedd.*, var. *lanceolata*, *Bedd.*, Handb. Ind. Ferns, 429; — var. *normalis*, *Hk.*, Spec. Fil., V, 276; *G. normale*, *J. Sm.*, Journ. of Bot., III, 405; *L. normalis*, *Copel.*, Philipp. Journ., III, 51, p. p.

Rhizome woody, creeping. Stipes of barren fronds 2½—15 c.M. long. Barren fronds 15—30 c.M. long, 1—5 c.M. broad, the point blunt or acute, the edge entire, the lower part narrowed very gradually. Texture herbaceous; surfaces naked; main veins short or indistinct; areolae copious, with free veinlets. Fertile fronds 15—30 c.M. long, 3—7 m.M. broad, the stipes as long.

Var. normalis: Barren fronds sessile.

Trop: Asia.

* (23) **L. axillaris**, *Klf.*, Enum., 147, tab. I, fig. 10; *Acrostichum axillare*, *Cav.*, *Hk. Bk.*, Syn. Fil., 420; *A. variabile*, *Hk.*, var. *axillare*, *Rac.*, Flor. Btz., I, 49; *Gymnopteris variabilis*, *Bedd.*, var. *axillaris*, *Bedd.*, Handb. Ind. Ferns, 430; *G. axillaris*, *Pr.*, *Copel.*, Polypod. Philipp., 42; *Bedd.*, Ferns Br.I., II, tab. CCLXXI.

Rhizome wide-scandent, slender, much branched. Barren fronds 2—7 c.M. apart, 15—45 c.M. long, 1—4 c.M. broad, the point bluntish or nearly so, the edge entire, the lower half tapering gradually to the base or a short

or rather long stipe. Texture papyraceous; surfaces naked; areolae copious, with abundant free veinlets. Fertile fronds 15—50 c.M. long, $2\frac{1}{2}$ — $7\frac{1}{2}$ m.M. broad, flexuose, on stipes $2\frac{1}{2}$ —15 c.M. long.

Trop. Asia.

* (24) **L. Raapii**, *v. A. v. R.*, Bull. Dép Agr. I. N., 1908, XVIII, 27, tab. VIII.

Rhizome erect, clothed with densely crowded, erect, linear-lanceolate scales with a dark-brown longitudinal axis and a pale-brown margin. Stipes densely tufted, 15—50 c.M. long, naked, firm, erect, those of the fertile fronds the longest. Fronds dimorphous. Barren fronds simple or some of them ternate, the simple ones ovate-elongate, \pm entire, 12—20 c.M. long, 3—4 c.M. broad, broadest near the \pm rounded or broadly cuneate base, narrowed gradually towards the short-acuminate point, the ternate ones with the segments similar to the simple fronds, but smaller. Texture flaccido-coriaceous; surfaces naked; veins prominent, reticulate as in *Acrostichum aureum*, L. Fertile fronds ternate, the segments stalked, linear, \pm 4—6 c.M. long, \pm 5 m.M. broad, acute. Sori occupying the whole under surface except the midribs and edges.

Batu Islands (Raap).

§ 3. LOMAGRAMMA.

*. *Texture membranaceous. Pinnae serrate.* (25) *L. lomarioides.*

.. *Texture subcoriaceous. Pinnae entire or obscurely crenate.*

(26) *L. perakensis.*

* (25) **L. lomarioides**, *Bl.*, Enum., 206; *Acrostichum Blumeum*, *Hk.*, Rac., Flor. Btz., I, 55, p. p.; *A. lomarioides*, *Christ*, Ann. Btz., XV, 179; *Lomagramme lomarioides*, *J. Sm.*, Bedd., Handb. Ind. Ferns, Suppl., 106; *Gymnopteris lomarioides*, *Christ*, Bull. Herb. Boiss., 2^e Série, I, 456; — var. *pteroides*; *Lom. pteroides*, *J. Sm.*, Journ. of Bot., III, 402; *Bau. & Hk.*, Gen. Fil., tab. XCVIII; — var. *subcoriacea*, *Copel.*, Philipp. Journ., III, 52.

Rhizome wide-scandent, the scales lanceolate. Stipes 15—50 c.M. long, deciduously scaly. Barren fronds 50—100 c.M. long. Pinnae numerous, sessile, articulate, rather falcate, 6—20 c.M. long, $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. broad, the base

truncate, rounded or broadly cuneate, the upper base not rarely broader than the lower, the edge serrate or toothed, but more prominently, towards the apex, the apex acuminate. Texture thinly membranaceous, transparent, pelucid-dotted; rachis and costae \pm scaly beneath, the former more or less winged, especially towards the apex; veins reticulate; areolae without free veinlets. Fertile pinnae \pm 6 m.M. broad, normally soriferous throughout beneath, sometimes partially contracted and soriferous.

Var. pteroides: Sori confined to the margin of the fertile pinnae.

Var. subcoriacea: Texture subcoriaceous, reddish when dry; veins very conspicuous beneath.

Malaya; Assam, Polynesia; — var. *pteroides: Celebes, Philippines;* — var. *subcoriacea: Mindanao.*

(26) **L. perakensis, C. Chr.,** Ind. Fil.; *Acrostichum Blumeianum, Hk., Rac., Flor. Btz., I, 55, p. p.; Lomagramme perakensis, Bedd., Handb. Ind. Ferns, Suppl., 107.*

Like *L. lomarioides, Bl.*, but the texture subcoriaceous, the pinnae entire or obscurely crenate, the veins less conspicuous, the rachis not winged. Fertile pinnae 1—4 m.M. broad.

Java, Malacca.

86b. PLATYTAENIA, Kuhn.

Like *Leptochilus*, but the paraphyses with a large terminal cell and the spores globoso-tetrahedral.

Fronds pinnate, with the vegetative characters of *Taenitis* by the appearance of *Leptochilus*.

Malayan and Polynesian.

P. Requiniana, Kuhn, Copel., Polypod. Philipp, 111; *Acrostichum Requinianum, Gaud., Hk., Spec. Fil., V, 269; Freyc, Voy. Ur. Phys., Bot., 504, tab. IV.*

Rhizome short, hairy. Stipes approximate, 20—30 c.M. long. Barren fronds ternate or pinnate. Pinnae linear-lanceolate, 12—18 c.M. long, \pm 3—4 c.M. broad, narrowed gradually towards the apex, the base narrowed. Venation \pm as in *Acrostichum aureum, L.* Fertile fronds pinnate; pinnae

linear, short-stalked, 10—20 c.M. long, \pm 5 m.M. broad, acute, the base narrowed. Sori filling up the whole under surface of the fertile pinnae, except the costa.

Singapore, Philippines, Moluccas, New Guinea; West. Polynesia.

FAMILY VII. PARKERIACEAE.

1941

VII. PARKERIACEAE.

Sporangia dorsal, not gathered in sori, scattered irregularly on the few longitudinal veins of the fertile fronds. Indusium wanting, but the margin of the fertile fronds broadly recurved, hardly changed in texture, the edges meeting against the midribs, quite enclosing the capsules when young. Ring vertical, broad, incomplete, short or interrupted by the very short, nearly wanting stalk of the capsule, wanting in the american form. Stomium transversal.

Rhizome short, erect. Fronds divided, rather dimorphous; veins anastomosing, but more copiously in the barren than in the fertile fronds.

Pantropical; in quiet, shallow waters, or even dry ground in humid regions or during the rains.

No. 87. **Ceratopteris.**

87. CERATOPTERIS, *Brongniart.*

Characters of the family.

* **C. thalictroides**, *Brong.*, *Rac.*, *Flor. Btz.*, I, 163; *Hk. Bk.*, *Syn. Fil.*, tab. III, fig. 32; *Diels*, in *Engl. & Prantl*, *Nat. Pfl. Fam.*, 1⁴, fig. 178—179; *Bau. & Hk.*, *Gen. Fil.*, tab. XII; *Bedd.*, *Ferns S.I.*, tab. LXXV; *Parkeria pteridioides*, *Hk.*, *Exot. Flor.*, II, tab. CXLVII; *Hk. & Grev.*, *Ic. Fil.*, tab. XCVII; *Acrostichum thalictroides*, *L.*, (oldest name).

Rhizome erect. Stipes tufted, thick, succulent, filled with large air-cells, naked, 5—20 c.M. long. Barren fronds floating or erect, 2—3-pinnatisect with linear or lanceolate ultimate segments 1—7 c.M. long; fronds of the young plants simple or slightly incised. Texture flaccid, succulent; surfaces naked; veins anastomosing copiously and irregularly. Fertile fronds with the segments narrow-linear, pod-like, the veins few, distantly anastomosing, running down the frond longitudinally. — *Parkeria pteridioides*, *Hk.* is the american form with exannulate capsules.

Pantropical.

1870

1871

1872

1873

1874

1875

1876

1877

FAMILY VIII. OSMUNDACEAE.

VIII. OSMUNDACEAE.

Sori dorsal (on the under surface of normal segments) or lateral (on the margin of much contracted fertile segments). Sporangia with a very short, incomplete, horizontal ring often consisting of a small group of thickened cells only. Stomium vertical.

Caudex thick, rarely arborescent. Stipes dilated at the base, not articulate to the caudex. Fronds large, at least simply pinnate, covered with a tomentum of long, woolly hairs when young, at length naked; veins free.

Generally inhabiting temperate climates.

Key to the genera.

- ✧. Fertile segments much contracted, articulate to the rachis. Sori globose, placed in opposite, marginal clusters. Texture of the fronds herbaceous or coriaceous. No. 88. **Osmunda**.
- ✧✧. Fertile segments not contracted, not articulate to the rachis. Sori \pm flat, dorsal. Texture of the fronds membranaceous. No. 89. **Leptopteris**.

88. OSMUNDA, *Linné*.

Sori subglobose, placed in opposite, marginal, thyrsoïd clusters. No indusium.

Fronds large, at least simply pinnate, herbaceous or coriaceous; leaflets dimorphous, articulate to the rachis, the fertile ones much contracted; veins free. Hk. Bk., Syn. Fil., tab. VIII, fig. 62; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 205.

Pantropical, and extra-tropical in the northern hemisphere.

Arrangement of the sections.

- § 1. **PLENASIUM**. Pinnae at best lobed.
 - § 2. **OSMUNDASTRUM**. Pinnae deeply pinnatifid. (Not in Malaya).
 - § 3. **EU-OSMUNDA**. Pinnae pinnate. (Not in Malaya).
-

§ 1. PLENASIUM.

* (1) **O. javanica**, *Bl.*, *Rac.*, *Flor. Btz.*, I, 5; *Bedd. Ferns S.I.*, tab. LXXVII; *O. Banksiaefolia*, *Kuhn*, *Ann. Mus. Bot. L. B.*, IV, 299; *O. Presliana*, *J. Sm.*, *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 205, A; *O. bromeliaefolia*, *Copel.*, *Philipp. Journ.*, II, 147.

Caudex stout, erect, 30—50 c.M. high, 10—20 c.M. or more thick. Stipes 15—30 c.M. long, erect, firm, naked. Fronds to 3 M. long, 30—60 c.M. broad, simply pinnate, the lower or central pinnae fertile. Barren pinnae erecto-patent, 2—2½ c.M. broad, lanceolate, the point acute, the edge entire, crenate, serrulate or toothed, the base cuneate. Texture coriaceous; rachis and surfaces naked; veins distinct, pinnate, subspreading (erecto-patent in *O. Banksiaefolia*, *Kuhn*).

Trop. Asia, Japan, Kamschatka.

(2) **O. blechnoides**, *Zoll.*, *Syst. Verz.*, 49.

Very near *O. javanica*, *Bl.*, but broader and thicker. — A dubious species. *Java.*

89. LEPTOPTERIS, *Presl.*

Sori roundish-oblong, dorsal, occupying the lower part of the veins.

Fronds at least 2-pinnate, thinly membranaceous (\pm as in *Hymenophyllum*); pinnae not articulate to the rachis, the fertile ones not contracted; veins free. *Diels*, in *Engl. & Prantl, Nat. Pfl.Fam.*, I⁴, fig. 204, B—C.

Almost confined to the south temperate zone.

*. *Fronds 2-pinnate. Pinnulae deeply toothed.*

(1) *L. Wilkesiana.*

.. *Fronds 3-pinnatifid. Tertiary segments entire.*

(2) *L. alpina.*

(1) **L. Wilkesiana**, *Christ*, *Farnkr. d. Erde*, 534; *Todea Wilkesiana*, *Brack.*, *Wilk.*, *Expl. Exp.*, XVI, tab. XLIII; *T. Fraseri*, *Hk. & Grev.*, var. *Wilkesiana*, *Hk. Bk.*, *Syn. Fil.*, 427.

Caudex erect, arborescent, to 2—3 M. high. Stipes 15—25 c.M. long, firm, naked. Fronds 1—2 M. long, 50—80 c.M. broad, ovate, 2-pinnate. Pin-

nae close, lanceolate, $2\frac{1}{2}$ —3 c.M. broad, connected by a narrow wing, the lower ones not much shorter, deflexed. Pinnulae linear-oblong, \pm 5 m.M. broad, the edge sharply and deeply toothed. Texture thin; rachises villose.

New Guinea; Polynesia.

(2) **L. alpina**, *C. Chr.*, Ind. Fil.; *Todea alpina*, *Bk.*, Kew Bull., 1899, 121.

Stipes short, naked. Fronds oblong-lanceolate, 45—60 c.M. long, 20—25 c.M. broad at the middle, narrowed gradually towards the base. Pinnae lanceolate, sessile, \pm $2\frac{1}{2}$ c.M. broad, the rachis narrowly winged; lower pinnae deflexed. Pinnulae lanceolate, sessile, erecto-patent, deeply pinnatifid. Ultimate segments linear, entire, 1-nerved. Texture membranaceous; rachis and surfaces naked. Sori few, scattered irregularly.

New Guinea.



FAMILY IX. MARATTIACEAE.

IX. MARATTIACEAE.

Sori on the under surface of normal fronds, round or oblong, consisting of \pm 6—24 free or connected capsules placed in a whorl or in 2 close rows; indusium wanting or minute, consisting of 2 opposite, flat, entire, or deeply fimbriate, inconspicuous, inferior scales, adpressed to the parenchyma; annulus wanting; sporangia opening by a slit down the inner side or by a pore at the apex.

Rhizome erect or creeping. Stipes with 2 large, stout stipules. Fronds *either* large and pinnately divided *or* small and simple or digitate; leaflets often articulate; veins free or anastomosing.

Tropical, and extra-tropical in the southern hemisphere.

Key to the genera.

- a.** Sporangia 2-serial, contiguous but not concrete. Fronds large, pinnately divided; veins free. No. 90. **Angiopteris.**
- a.a.** Sporangia concrete, joined in 2-valved or circular synangia.
- b.** Synangia 2-valved. Fronds large, pinnately divided; veins free. No. 91. **Marattia.**
- b.b.** Synangia circular. Fronds small, simple, ternate or quinate; veins anastomosing. No. 92. **Christensenia.**

90. **ANGIOPTERIS**, *Hoffmann.*

Capsules sessile, not concrete, closely placed in 2 contiguous rows, 2—10 in a row, forming roundish or oblong, submarginal sori, opening by a slit down the inner side. Indusium minute, squamiform or deeply fimbriate.

Caudex short, thick, often semiglobose. Stipes flanked by 2 large, firm stipulae at the base. Fronds large, 1—3-pinnate; veins free, simple or forked; not rarely accompanied by intermediate spurious veins (recurrent veinlets) running from the margin in the direction of the costa. Hk. Bk., Syn. Fil., tab. IX, fig. 69; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 258, A & 259, A—B.

✧. *Fronde simply pinnate.* (1) *A. Smithii*.

✧✧. *Fronde 2—3-pinnate.*

*. *Pinnulae pectinato-serrate, the teeth cartilagineous.* (2) *A. cartilagens.*

.. *Pinnulae entire, crenate, serrate or toothed, the teeth not cartilagineous.*
(5) *A. evecata.*

* (1) ***A. Smithii*, Rac.**, Bull. Ac. Crac., 1902, 54.

Caudex very short, ± 10 c.M. thick. Fronds large, $\pm 1\frac{1}{2}$ — $2\frac{1}{2}$ M. high (incl. the stipe), simply pinnate. Pinnae 5—10 or more on each side, short-stalked, articulate, ovate-oblong, 20—40 c.M. long, 5—10 c.M. broad in the lower half, the base obliquely or unequally rounded or subcordate, the edge entire, the upper half narrowed gradually to the acuminate apex. Texture flaccido-coriaceous; upper surface naked, dark-green, lower pale-green, minutely and sparingly scaly, nearly naked; costae prominent on both sides; veins numerous, generally forked, upcurved, rather spreading at the base, oblique towards the apex; recurrent veinlets wanting. Sori close, submarginal, the largest consisting of ± 20 —30 capsules.

? *Malayan.*

(2) ***A. cartilagens*, Christ**, Bull. Herb. Boiss., VI, 207; *A. cartilaginea*, Christ, l. c., tab. IV, fig. 8, a—d.

Like *A. evecata*, Hoffm., but the fronds larger, to 5 M. long; rachis dark-purple, covered with oblong, subulate, spinuloso-ciliate scales, the ciliae geminate. Pinnulae close, sessile or short-stalked, linear-lanceolate, 10 c.M. or more long, ± 1 c.M. broad, the margin pectinato-serrate with the teeth oblique or spreading, $\frac{1}{2}$ —1 m.M. long, the apex acuminate, the base equally rounded or cuneate on both sides or the upper base truncate. Texture very coriaceous, the teeth cartilagineous; surfaces naked; costae and veins covered beneath with small, rough, bullate scales; veins very numerous, conspicuous, spreading, with spurious veins. Sori not contiguous, small.

Luzon.

* (5) ***A. evecata*, Hoffm.**, Rac., Flor. Btz., I, 1; Bedd., Ferns S.I., tab. LXXVIII; Hk., Fil. Exot., tab. LXXV; *Polypodium evecatum*, Forst., (oldest name).

Caudex erect, semiglobose, to ± 50 c.M. thick. Stipes long, articulate at the base, flanked by 2 large, persistent, leathery stipulae. Fronds large,

to $4\frac{1}{2}$ M. long, 2—4-pinnate. Pinnae to 1 M. long, spreading, the lowest the largest. Leaflets 10—40 c.M. long, 1—4 c.M. broad, linear-lanceolate or linear, sessile or short-stalked, acute, acuminate or cuspidate, the edge entire, crenate, serrate or toothed, often conspicuously serrato-dentate towards the apex, the base cuneate, rounded or truncate, articulate. Texture more or less coriaceous, rarely thin; surfaces often naked; veins parallel, simple or forked, whether or not pellucid; recurrent veinlets mostly present. Sori in 1 row on each side, near the margin, consisting of \pm 8—18 capsules. — A very variable plant with numerous forms.

Trop. Asia & Africa, Tahiti.

91. MARATTIA, Swartz.

Capsules sessile, concrete in oblong sori consisting of 2 opposite, boat-shaped, \pm 5—15-celled synangia, opening by slits down the inner side, \pm medial or subterminal on the lateral veins, rarely occupying the central ribs. Indusium wanting or present, but then minute, inconspicuous, fimbriate, often persistent.

Fronde large, 2—3-pinnate, stipitate, the stipes provided with 2 distinct stipulae at the base; veins free, simple or forked; recurrent veinlets wanting. Hk. Bk., Syn. Fil., tab. IX, fig. 70; Diels in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 239, C & 239, E—F.

Tropical, and subtropical in the southern hemisphere.

Arrangement of the sections.

- § 1. **EUMARATTIA.** Sori occupying the lateral veins of the ultimate segments. Indusium present.
- § 2. **GYMNOTHECA.** Sori as in § 1. Indusium wanting.
- a. **Eugymnotheca.** Sori sessile.
- b. **Eupodium.** Sori stalked. (Not in Malaya).
- § 5. **MESOCARPUS** (*Mesosorus*). Sori occupying the central ribs of the ultimate segments.

§ 1. EUMARATTIA.

a. *Rachis densely scaly.*

(1) *M. vestita.*

a.a. *Rachis not densely scaly.*

- b.** *Ultimate pinnulae often acutely cuneate at the base.*
 *. *Stipes tuberculate; main rachis \pm asperulous.* (2) *M. sambucina.*
 .. *Stipes not tuberculate, main rachis smooth or nearly so.*
 (5) *M. silvatica.*
- b.b.** *Ultimate pinnulae mostly abruptly cuneate, truncate or rounded at the base.*
- c.** *Texture rather thin, membranaceous or herbaceous.*
 +. *Veins nearly always simple. Texture herbaceous.*
 (4) *M. pellucida.*
 +.+ *Veins partially simple, partially forked. Texture membranaceous.*
 (5) *M. fraxinea.*
- c.c.** *Texture rather thick, chartaceous or coriaceous.*
- d.** *Veins beneath naked. Synangia \pm 8—10-celled.* (6) *M. ternatea.*
d.d. *Veins beneath deciduously dilacerato-paleaceous. Synangia \pm 15-celled.*
 (7) *M. Smithii.*

(1) **M. vestita**, *Christ*, Philipp. Journ., II, 185.

Large. Stipes thick, reddish-brown, opaque, densely pustulous and clothed with ovate, flaccid scales, which are intermixed with narrower and smaller paleae, the main rachis rufo-fuscous below, ochraceo-rufous above, densely clothed like the lateral rachises with brown, lanceolate scales and fibrils. Pinnae oblong, \pm 70 c.M. long, \pm 30 c.M. broad, petiolate, narrowed at the base, the apex acuminate. Pinnulae petiolate, \pm 10 c.M. apart, 20 c.M. or more long, \pm 15 c.M. broad, narrowed at the base. Tertiary segments \pm 2½ c.M. apart, in several pairs below the terminal one, lanceolate, \pm 9 c.M. long, \pm 1¼ c.M. broad, alternate, articulate, acuminate, cuneate at the base, toothed with open, spreading teeth, the lowest very short-stalked. Texture thinly but firmly chartaceous; upper surface dark-green, lower paler, opaque; veins conspicuous, simple, remote. Sori minute, remote; synangia \pm 5-celled.
Mindanao.

* (2) **M. sambucina**, *Bl.*, Rac., Flor. Btz., I, 2; de Vr., Mon. Maratt., 6; *Christ*, Philipp. Journ., II, 184; *M. sambucifolia*, *Christ* (not *Bl.*), Ann. Btz., XIX, 43.

Stipes to \pm 50 c.M. long, tuberculate or asperous, the rachis naked, asperulous. Fronds 2—5-pinnate, \pm 60—120 c.M. long, the ultimate rachises narrowly winged. Leaflets lanceolate, the higher ones longer than the lower,

the central ones to ± 5 c.M. long, to $\pm \frac{3}{4}$ — $1\frac{1}{2}$ c.M. broad, with the apex acuminate, the edge serrate, the base cuneate. Texture membranaceous to subcoriaceous; surfaces naked; veins simple, pellucid. Sori distant from the margin; synangia ± 5 —8-celled.

Malaya.

* (3) **M. silvatica**, *Bl.*, de Vr., Mon. Maratt., 6, tab. III, fig. 23; Christ, Philipp. Journ., II, 184.

Stipes naked. Fronds large, 3-pinnate. Ultimate leaflets opposite or alternate, to ± 10 c.M. long, to $\pm 1\frac{1}{3}$ c.M. broad, the higher ones larger than the lower, linear-lanceolate, acute, serrate, the base cuneate. Texture firm-chartaceous; surfaces naked, upper dark-green, lower paler, minutely albopunctulate; ultimate rachises narrowly winged, rather marginate; veins simple, rather spreading. Sori intramarginal, or nearly medial on the veins; synangia ± 12 —15-celled.

Java, Luzon, New Guinea.

* (4) **M. pellucida**, *Pr.*, de Vr., Mon. Maratt., 6; Christ, Philipp. Journ., II, 184.

Fronds 2-pinnate. Leaflets linear-oblong, subabruptly acuminate, serrate, the base cuneate; highest leaflets the largest, to $\pm 12\frac{1}{2}$ c.M. long, to $\pm 1\frac{1}{2}$ c.M. broad. Texture herbaceous; surfaces naked, upper rather dark-green, lower paler; costae and veins beneath naked or nearly so; veins mostly simple, very rarely forked, straight, subspreading, pellucid. Sori rather remote from the margin, sometimes nearly medial on the veins; synangia ± 6 —8-celled.

Philippines; New Caledonia.

* (5) **M. fraxinea**, *Sm.*, Hk. Bk., Syn. Fil., 440; Bedd., Ferns S.I., tab. LXXIX; *M. calliodus*, de Vr., Mon. Maratt., 6, tab. III, fig. 20.

Stipes 20—60 c.M. long, smooth, deciduously scaly and swollen in the lower part. Fronds rather large, 2—3-pinnate. Highest leaflets largest, elongate-lanceolate, ± 10 —15 c.M. long, $\pm 1\frac{1}{2}$ —2 c.M. broad, acuminate, the edge serrate, the base broadly and obliquely cuneate or slightly rounded. Texture membranaceous; surfaces naked; costae rather naked, or slightly scaly at the very base; veins simple or forked, \pm pellucid. Sori somewhat remote from margin; synangia ± 6 —12-celled.

Trop. Asia, Australia & Africa.

* (6) **M. ternatea**, *de Vr.*, Mon. Maratt., 4, tab. III, fig. 16; Christ, Philipp. Journ., II, 185.

Fronde 5-pinnate. Ultimate leaflets \pm 12—20 c.M. long, \pm 2—2½ c.M. broad, linear-lanceolate, acuminate or acute, the edge serrate, the base ovate-cuneate, broadest on the upper side, the petiole squamulose. Texture coriaceous; surfaces naked, upper dark-green, lower paler; rachis \pm ochraceous; costae and veins fuscous (ochraceous after Christ); veins simple, nearly horizontal, pellucid. Sori not close, somewhat remote from the margin; syngangia \pm 15-celled. — A plant found in Ternate by Mr. Teysmann and determined as *M. ternatea*, *de Vr.* has the leaflets not dark-coloured above, the rachis, costae and veins ochraceous, the costae and veins beneath sparingly provided with very minute squamulae, the veins hardly pellucid.

Luzon, Moluccas; ? New Caledonia.

(7) **M. Smithii**, *Mett.*, Verh. Zool. Bot. Ges., XIX, 584.

Fronde 2-pinnate. Pinnae oblong, the base slightly narrowed. Pinnulae spreading, to \pm 15 c.M. long, to \pm 2½ c.M. broad, linear-oblong, acuminate, the lower base rounded, the upper rotundato-truncate, the edge slightly serrate, but more distinctly at the acuminate apex. Texture coriaceous, opaque; rachis smooth, soon naked; costae and veins covered beneath with lanceolate, ferruginous, lacerato-fimbriate, caducous scales; veins commonly simple, rather spreading, subapproximate. Sori distant from the margin, subapproximate; syngangia oblong, 8—10-celled.

New Guinea; Polynesia.

§ 2. GYMNOTHECA. a. **Eugymnotheca.**

(8a) **M. alata**, *Sm.*, Hk. Bk., Syn. Fil., 442; Christ, Farnkr. d. Erde, fig. 1128.

Stipes 50—60 c.M. long, deciduously scaly. Fronde 90—120 c.M. long, 2—5-pinnate at the base, the ultimate rachises distinctly winged. Lower pinnae largest. Leaflets 1—2½ c.M. long, ½—1 c.M. broad, oblong, blunt or acute, the base cuneate, the edge crenate, serrate or toothed. Texture herbaceous; under surface more or less chaffy. Sori copious, submarginal, 1—2 m.M. long. — A plant found in Celebes is probably a form of this species.

? Celebes; Trop. America.

§ 3. MESOCARPUS.

(9) **M. Wernerii**, *Rst.*, Fed., Rep., V, 44.

Rhizome and stipes unknown. Fronds large, ± 2 M. long, 5-pinnate. Pinnae petiolate, elongate-oblong, to 70 c.M. or more long, ± 25 c.M. broad, broadest at the centre, the rachis adherent to the main one, canaliculate above, brown, clothed with few scattered, minute, narrowly lanceolate scales. Pinnulae short-stalked, suberecto-patent, articulate to the secondary rachises, deciduous, linear, the central ones the longest, ± 14 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, the apex abruptly acuminate, the terminal pinnulae somewhat longer than the highest ones, the internodations of the rachises winged between the pinnulae, the wings nearly $\frac{1}{2}$ m.M. broad at the centre, narrowed towards both ends. Tertiary divisions sessile, articulate at the base, deciduous, elongate-oblong, subfalcate, $\pm 7\frac{1}{2}$ m.M. long, ± 5 m.M. broad, short-acuminate, serrato-dentate at the margin, cuneate at the base, the terminal ones caudiform. Texture coriaceous; surfaces naked, upper surface lucent (opaque when dry), darker in colour than the lower; midribs of the ultimate segments (leaflets) prominent above; veins simple, terminating in the teeth, sterile. Sori 1 to each leaflet, occupying the upper part of the midrib; synangia $\pm 9-11$ -celled; indusium wanting.

New Guinea.

92. CHRISTENSENIA, *Maxon.*

Capsules sessile, concrete in raised, circular, $\pm 10-20$ -celled synangia, which are concave at the top so as to form nearly cupuliform or ring-shaped masses, opening by slits down the inner side or by apical pores.

Rhizome creeping. Stipes scattered or approximate, with 2 fleshy stipulae. Fronds simple, or digitate and 5-5-foliolate: veins anastomosing. Hk. Bk., Syn. Fil, tab. IX, fig. 72; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 258, D & 259, G-H. (*Kaulfussia*).

Australasia.

-
- *. *Fronds simple, or 3-partite with adnate lateral segments. Synangia in 2 rows between the main veins.* (1) *C. Cumingiana.*
- *.*. *Fronds 3-5-foliolate, with the lateral segments \pm stalked. Synangia scattered irregularly.* (2) *C. aesculifolia.*
-

(1) **C. Cumingiana**, *Christ*, Philipp. Journ., II, 186.

Rhizome thick, short, fleshy. Stipes approximate, or subfasciculate in the younger plants, to 14 c.M. long, rufo-stramineous and furfuraceous, as are the costae and main veins. Fronds *either* simple, ovate, subcordate at the base, 8–15 c.M. long, 3–5 c.M. broad, repand or coarsely toothed, shortly acute at the apex, *or* 3-partite with the central segment broadly ovate, \pm 11 c.M. long, \pm 5 c.M. broad, acute at the point, repando-lobate at the edge, narrowly cuneate at the base and the lateral segments adnate, very unequal-sided, cordate on the produced lower, narrowly cuneate on the upper side, \pm 9 c.M. long, \pm 5 c.M. broad. Texture succulent; under surface whitish, provided with numerous roundish pores; main veins distinct, horizontal. Synangia in 2 rows between the main veins, \pm 4 in a row.

Mindanao.

* (2) **C. aesculifolia**, *Max.*, C. Chr., Ind. Fil.; *Kaulfussia aesculifolia*, *Bl.*, *Rac.*, *Flor. Btz.*, I, 2; *Hk. & Grev.*, *Ic. Fil.*, tab. CCXXIX; *K. assamica*, *Griff.*, *Bau. & Hk.*, *Gen. Fil.*, tab. LIX, A; *Bedd.*, *Ferns Br.I.*, II, tab. CLXXXV.

Rhizome creeping. Stipes succulent, 20–45 c.M. long, auricled at the base with 2 fleshy stipulae. Fronds digitate, ternate or quinate. Central segment the largest, ovate-lanceolate, narrowed to both ends, with the edge entire, crenate or slightly toothed, 15–40 c.M. long, 7–12 c.M. broad; lateral segments similar but smaller, falcate, unequal-sided. Texture fleshy-herbaceous; surfaces naked, but the lower side dotted over with copious stomata-like pores; main veins conspicuous, pinnate; veins anastomosing; areolae copious, immersed, with free veinlets. Synangia copious, scattered irregularly.

Malaya; Assam.

FAMILY X. OPHIOGLOSSACEAE.

X. OPHIOGLOSSACEAE.

Sori on much contracted, spicate or paniculate fertile segments opposite to and springing from the base, the centre or the margin of the normal barren segments. Capsules \pm globose, opening across the apex or by a slit down the side; ring wanting.

Rhizome erect, rarely creeping, sometimes tuberous. Stipes not articulate to the rhizome. Barren segment of the fronds simple or variously divided; veins free or anastomosing.

Tropical and extra-tropical.

Key to the genera.

1. Fertile segment spicate.

2. Capsules sessile or immersed, placed in 2 opposite rows. Barren segment mostly simple; veins anastomosing.

No. 93. **Ophioglossum.**

2.2. Capsules placed in numerous small, crested clusters spreading in all directions. Barren segment divided; veins free.

No. 94. **Helmintostachys.**

1.1. Fertile segment paniculate; capsules placed in 2 rows on the branches of the panicle, on the side directed towards the mostly divided barren segment; veins free.

No. 95. **Botrychium.**

93. OPHIOGLOSSUM, Linné.

Capsules sessile or immersed, arranged in 2 opposite rows so as to form a long-stalked, close spike (fertile segment).

Mostly small and erect, rarely long and pendulous plants. Rhizome short, with 1—2 or rarely 3—4 or more naked, stalked fronds; each frond with a barren and 1 or rarely more fertile segments; fertile segments simple or sometimes double or triple, opposite to the barren one, arising from its base, centre or margin; barren segment mostly simple, rarely divided; veins reticulate; costa mostly indistinct.

Tropical and extra-tropical.

Arrangement of the sections.

- § 1. **EU-OPHIOGLOSSUM.** Fertile spike single, mostly simple, rarely 2—5-partite, arising from the base of the barren segment, which is sometimes wanting or rudimentary. Rhizome naked, hypogeous. Hk. Bk., Syn. Fil., tab. IX, fig. 73; Diels, in Engl. & Prantl, Nat. Pfl.Fam., 1⁴, fig. 259, A & C.
- § 2. **OPHIODERMA.** Fertile spike as in § 1, but arising from the centre of the barren segment. Rhizome papillose, epiphytical. Diels, l. c., fig. 265, A.
- § 5. **CHEIROGLOSSA.** Fertile spikes many or numerous, springing from the base and margin of the barren segment. Rhizome long-hairy, epiphytical. Diels, l. c., fig. 265, B—C. (Not in Malaya).

§ 1. **EU-OPHIOGLOSSUM.**

- a.* Barren segment wanting or rudimentary. (1) *O. simplex.*
- a.a.* Barren segment distinctly developed.
- b.* Barren segment linear or linear-lanceolate, more than 5 × as long as broad.
- ×. Fronds ± 4½ c.M. high, the barren segment 1—1½ c.M. long, 1—2 m.M. broad. (2) *O. gramineum.*
- ×.×. Fronds 7—20 c.M. high, the barren segment ± 1½—8 c.M. long, 2—6 m.M. broad. (3) *O. inconspicuum.*
- b.b.* Barren segment lanceolate or oblong, less than 5 × as long as broad.
- c.* Barren segment cuneate at the base.
- d.* Barren segment small, mostly ¼—1 c.M. long.
- §. Barren segment roundish, nearly as broad as long. (4) *O. pumilum.*
- §.§. Barren segment elongate, ± ½ × as broad as long. (5) *O. macrorrhizum.*
- d.d.* Barren segment larger, mostly 1—5 c.M. long.
- e.* Barren segment placed quite at the base of the whole frond. (6) *O. fibrosum.*
- e.e.* Barren segment placed above the base of the frond.
- f.* Veins indistinct.
- +. The whole plant 5—10 c.M. high, with the barren segment ± ovate, broadest at or below the middle. (7) *O. parvifolium.*

++. *The whole plant ± 15 c.M. high, with the barren segment obovate, broadest above the middle.* (8) *O. obovatum.*

f.f. *Veins distinct.*

g. *Barren segment membranaceous.*

(9) *O. timorense.*

g.g. *Barren segment flaccid or subcarnose.*

(10) *O. pedunculosum.*

c.c. *Barren segment cordate, subtruncate or rounded at the base.*

+. *Barren segment with the margin minutely crenulate, i. e. bordered by a row of cellules, which have the outer wall convex (*).* (10) *O. pedunculosum.*

++. *Barren segment with the margin quite continuous, i. e. bordered by a row of cellules, which have the outer wall straight (*).* (11) *O. reticulatum.*

(1) ***O. simplex***, *Ridl.*, Ann. of Bot., XVIII, 205, tab. XV; XXII, 327.

Rhizome short, tuberous. Fronds solitary or in pairs, 10—15 c.M. high, flattened, 3—5½ m.M. broad, the sterile segment absent or rudimentary and represented by a more or less pronounced outgrowth at a short distance below the fertile region, with a slightly increased width of the petiole below that point. Fertile spike ± 2½ c.M. long.

Sumatra.

(2) ***O. gramineum***, *Willd.*, Prantl, Ophiogl., 311, tab. VII, fig. 4.

Rhizome cylindrical, slightly tuberous. Fronds commonly in pairs, ± 4½ c.M. high; sterile segment linear or linear-lanceolate, 1—1½ c.M. long, 1—2 m.M. broad, sessile, placed below the middle, the apex acute. Texture rigid; primary veins 5, the central one stouter, straight, the lateral parallel, reticulate; areolae elongate, rarely with free veinlets. Fertile spike under 1 c.M. long, the peduncle 1—2 × as long as the barren segment.

New Guinea; Br. India, West. Africa.

(*) Only visible in fresh or boiled material, by high (± 100 ×) magnifying power.

* (5) **O. inconspicuum**, v. A. v. R., Bull. Dép. Agr. I. N., 1908, XXI, 9; *O. moluccanum*, Schl., forma *inconspicua*, Rac., Nat. Tydschr. Ned. Ind., LIX, 257, tab. II, fig. 5; — var. *majus*, v. A. v. R., l. c.

Rhizome \pm cylindrical. Fronds 1—2, sometimes more, 7—10 c.M. high, the barren segment sessile, elongate-lanceolate, \pm $1\frac{1}{2}$ —3 c.M. long, 2— $3\frac{1}{2}$ m.M. broad, acute at both ends, placed mostly below the middle. Texture \pm fleshy, often thin when dry; areolae longitudinally elongated. Fertile spike \pm 1—2 c.M. long, the peduncle \pm $1\frac{1}{2}$ —2 \times as long as the barren segment.

Var. majus: Like the type but all the dimensions \pm 2 \times as large, and the fronds more numerous, even to 12—15, tufted.

Java, New Guinea; — var. *majus*: *New Guinea* (Versteeg).

(4) **O. pumilum**, v. A. v. R.; *O. moluccanum*, Schl., forma *pumila*, Rac., Nat. Tydschr. Ned. Ind., LIX, 257, tab. II, fig. 2—3; ? Ann. Btz., XXI, tab. XIX, fig. 154.

Rhizome short. Whole plant 1—5 c.M. high. Barren segment roundish or rotundo-ovate, \pm sessile, placed near the base of the plant, 3—10 m.M. long, nearly as broad. Fertile spike 2—6 m.M. long, with the peduncle to $3\frac{1}{2}$ c.M. long.

Java.

(5) **O. macrorrhizum**, Kze, Prantl, Ophiogl., 522, tab. VIII, fig. 22; Kze, Farnkr., I, tab. XXIX, fig. 1, a—e.

Rhizome short, thick. Whole plant under 5 c.M. high. Sterile segment placed at or below the middle, lanceolate or oblong, 5—10 m.M. long, 2—3 m.M. broad, the apex blunt or acute, not apiculate, the base cuneate, decurrent on the petiole. Texture thin; principal veins 5, fine, the central one stouter; areolae indistinct. Fertile spike \pm as long as the sterile segment; peduncle slender, 1—2 c.M. long.

? *Philippines*; *South. Brasilia, Argentine Republic*.

(6) **O. fibrosum**, Schum., Prantl, Ophiogl., 525, tab. VIII, fig. 26; *O. brevipes*, Bedd., Ferns S.I., tab. LXXII.

Rhizome thick, bulbous. Fronds 1—7. Stipes very short, with the sterile segment \pm close to the base, oblong or lanceolate, 4—7 c.M. long, $1\frac{1}{4}$ —2 c.M. broad, sometimes smaller (\pm $2\frac{1}{2}$ c.M. long and \pm 1 c.M. broad), the apex obtuse or acute, the base cuneate. Texture thick, subcarnose; veins

indistinct (except when dry), the midrib a little stouter. Fertile spike 4—5 c.M. long, on a peduncle 10—12½ c.M. long.

Sumatra; Br. India, Trop. Africa.

* (7) **O. parvifolium**, *Grev. & Hk.*, Bot. Misc., III, 218; Bedd., Ferns S.I., tab. LXXI; *O. nudicaule*, *L.f.*, Hk. Bk., Syn. Fil., 445; Kze, Farnkr., I, tab. XXIX, fig. 5, b.

Rhizome slightly tuberous. Whole plant ± 5—10 c.M. high. Barren segment ovate or oblong, ± 1¼—4 c.M. long, ± ½—1½ c.M. broad, sessile or subsessile, placed considerably below the middle of the whole plant, narrowed towards both ends. Texture thin; no evident costa; veins indistinct. Fertile spike 1—1½ c.M. long, the peduncle often 5—5 c.M. long or even longer. — Besides the fertile fronds there are often some shorter ones without fertile segments.

Malaya; Japan, South. India, West. Africa, Trop. & Subtrop. Americ.

(8) **O. obovatum**, *Miq.*, Ann. Mus. Bot. L. B., IV, 93,

The whole plant ± 15 c.M. high. Sterile segment oblong-obovate, rarely broadly obovate, placed considerably below the middle, ± 3½ c.M. long, ± 1¼—1¾ c.M. broad, with a haft ± 1½ c.M. long, the apex rounded or obtuse, the base broadly cuneate. Texture coriaceous; veins nearly hidden. Fertile spike ± 2½ c.M. long, on a slender peduncle ± 10 c.M. long.

Timor.

(9) **O. timorense**, *Miq.*, Ann. Mus. Bot. L. B., IV, 93.

The whole plant mostly 6—10 c.M. high. Sterile segment often placed considerably below the middle, ovate, 2—3½ c.M. long, the apex obtuse or subacute, the base broadly cuneate, narrowed into a short and broad haft. Texture membranaceous; veins distinct, densely reticulate. Fertile spike 1—2½ c.M. long, on a slender peduncle 2½—5 c.M. long.

Timor.

* (10) **O. pedunculatum**, *Desv.*, Prantl, Ophiogl., 328, tab. VIII, fig. 33—34; *O. moluccanum*, *Schl.*, Rac., Flor. Btz., I, 4; Nat. Tydschr. Ned. Ind., LIX, 236, tab. II, fig. 1; Ann. Btz., XXI, tab. XIX, fig. 155; *O. petiolatum*, *Hk.*, Exot. Flor., I, tab. LVI; — var. *lanceolatum*; *O. moluccanum*, *Schl.*, forma *lanceolata*, Rac., l. c. 257, tab. II, fig. 4.

Rhizome cylindrical. Fronds 1—2, rarely 3—4, the whole plant ±

6—25 c.M. high. Barren segment placed (often considerably) below the middle, ovate, lanceolate, oval or rarely subrhomboidal, $\pm 1\frac{1}{2}$ —9 c.M. long, ± 1 —4 c.M. broad, sessile or with a haft, the broadest part below the middle, the base broadly cuneate, rounded or subtruncate, rarely \pm cordate, the apex obtuse or subacute, whether or not apiculate, the marginal cells with the outer wall convex. Texture flaccid or subcarnose, thin when dry; veins distinct; costa flexuose, somewhat stouter near the base, often indistinct towards the apex. Fertile spike with the peduncle to $\pm 3 \times$ longer than the barren segment.

Var. lanceolatum: Whole plant 10—18 c.M. high. Barren segment lanceolate, 3—4 c.M. long, to 1 c.M. broad, placed a little below the middle, acute at both ends; marginal cells.....?; peduncle ± 3 —5 c.M. long.

Trop. Asia, Australia, New Zealand, New Caledonia, Japan; — var. lanceolatum: Java.

* (11) **O. reticulatum**, *L.*, Hk. Bk., Syn. Fil., 446; Prantl, Ophiogl., 530, tab. VIII, fig. 32; Bedd., Ferns S.I., tab. LXX; Hk. & Grev., Ic. Fil., tab. XX; Christ, Farnkr. d. Erde, fig. 1157.

Rhizome not tuberous. Fronds 1—2, the whole plant 10—30 c.M. high. Barren segment placed at or sometimes considerably below the middle, 2—8 c.M. long, $1\frac{1}{2}$ —7 c.M. broad, with a haft, oblong, rarely reniform, the apex obtuse, often rounded, rarely subacute, often apiculate, the base cordate, rarely subtruncate, the margin quite continuous, i. e. the marginal cells with the outer wall straight. Texture mostly thin, rarely rigid; veins distinct, the midrib somewhat stouter near the base, flexuose, becoming finer towards the apex. Fertile spike $\pm 2\frac{1}{2}$ —4 c.M. long, on a slender peduncle often $3 \times$ as long as the barren segment.

Pantropical.

§ 2. OPHIODERMA.

+ . Fronds erect, 10—20 c.M. long.

(12) *O. intermedium.*

+ . + . Fronds pendulous, 30—150 c.M. long.

(13) *O. pendulum.*

* (12) **O. intermedium**, *Hk.*, Hk. Bk., Syn. Fil., 446; Hk., Icon Pl., X, tab. CMXCV.

Fronds erect, 10—20 c.M. long, the flattened stipe passing gradually into the leafy part, which is \pm 1 c.M. broad not far from the apex. Texture fleshy; veins indistinct. Fertile spike 3—4 c.M. long, on a peduncle 3—4 c.M. long.

Java, Borneo, Mindoro, New Guinea.

* (13) **O. pendulum**, **L.**, Rac., Flor. Btz., I, 3; Prantl, Ophiogl., 351; Hk. & Grex., Ic. Fil., tab. XIX; Bedd., Ferns S.I., tab. CCLXIX; Hk., Gard. Ferns, tab. XXXIII.

Fronds 1—5, pendulous, 50—150 c.M. long, the short, terete stipe passing gradually into the leafy part, which is $2\frac{1}{2}$ — $7\frac{1}{2}$ c.M. broad, simple, forked or dichotomous. Texture fleshy or coriaceous; veins indistinct. Fertile spike 5—20 c.M. long, on a peduncle shorter than itself.

Trop. Asia, Australia & Polynesia.

94. HELMINTOSTACHYS, *Kaulfuss.*

Capsules in crested clusters spreading in all directions, forming a long, loose spike (fertile segment).

Rhizome thick, creeping. Fronds scattered, erect. Each frond with a barren and a fertile segment, placed on the same stipe; the fertile segment simple or rarely double or triple, opposite to the barren one and arising from its base; barren segment digitate or pinnatipartite; veins free; costae distinct. Hk. Bk., Syn. Fil., tab. IX, fig. 74; Diels, in Engl. & Prantl, Nat. Pfl.Fam. I⁴, fig. 260, F.

Tropical Asia and Australia.

* **H. zeylanica**, **Hk.**, Rac., Flor. Btz., I, 4; Hk., Gard. Ferns, tab. XXVIII; Bedd., Ferns S.I., tab. LXIX; Hk., 2nd Cent. of Ferns, tab. XCIV; *H. dulcis*, *Klf.*, Enum., 28, tab. I, fig 1; *Osmunda zeylanica*, *L.*, (oldest name).

Stipes erect, 15—30 c.M. long. Barren segment digitate with commonly 5 primary divisions. Primary divisions stalked and forked or pinnate. Ultimate segments 7—15 c.M. long, 2—4 c.M. broad, lanceolate, acute or acuminate, the edge entire or slightly toothed, the base \pm cuneate. Fertile spike 7—20 c.M. long, on a peduncle of about the same length.

Trop. Asia & Australia, New Caledonia.

95. BOTRYCHIUM, Swartz.

Capsules sessile, placed in 2 rows on the inner face of spikes, which form a more or less compound panicle (fertile segment).

Mostly small, erect plants. Rhizome short, erect. Each frond with a barren and a fertile segment placed on the same stipe, the fertile segment arising from the base or the centre of the barren one, which is once or more times pinnatisect; veins free. Hk. Bk., Syn. Fil., tab. IX, fig. 75; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 259, B.

Ferns inhabiting different zones and climates.

Arrangement of the sections.

§ 1. **EUBOTRYCHIUM.** Fronds always naked. (Not in Malaya).

§ 2. **PHYLLOTRICHIMUM.** Fronds hairy, at least when young.

§ 2. PHYLLOTRICHIMUM.

‡. Fertile segment arising from the base of the barren one. (1) *B. daucifolium*.

‡.‡. Fertile segment arising from the centre of the barren one.

(2) *B. lanuginosum*.

* (1) ***B. daucifolium*, Wall.**, Ræc., Flor. Btz., I, 4; Hk. & Grev., Ic. Fil., tab. CLXI; *B. subcarnosum*, Wall., Bedd., Ferns S.I., tab. LXVIII.

Stipes stout, 15—30 c.M. long. Barren segment 15—30 c.M. long, ± as broad, deltoid, 3-pinnatifid to 3-pinnate, on a petiole 2½—15 c.M. long. Lowest pinnae largest. Ultimate segments ½—1 c.M. broad, finely toothed. Texture herbaceous; surfaces at length naked. Panicle 5—10 c.M. long, rather loose, long-stalked, arising from the base of the barren segment.

Philippines, Java; Ceylon, North. India.

(2) ***B. lanuginosum*, Wall.**, Christ, Farnkr. d. Erde, 366; *B. virginianum*, Sm. (*virginicum*, Willd.), var *lanuginosum*, Bedd., Handb. Ind. Ferns, 471; Ferns S.I. tab LXVII; — var. *nanum*, Christ, Philipp. Journ., II, 187.

Stipes 7½—45 c.M. long. Barren segment 10—40 c.M. each way, deltoid,

3—4-pinnatifid. Lower pinnae much the largest. Pinnulae ovate-oblong, variously lobed and decomposed. Texture subcarnose; main and partial rachises more or less hairy; surfaces slightly hairy or at length naked. Fertile peduncle arising from the centre of the barren segment, which the panicle not or scarcely overtops.

Var. nanum is the form with the smaller fronds

Br. India, Ceylon; — var nanum: also in Luzon.

APPENDIX.

APPENDIX.

This appendix principally treats with the ferns discovered after January 1st 1907, besides the additions which could not be inserted in the first part of the work, when in the press.

The numbers within brackets indicate the sequence in the keys of the handbook, but only approximately for the new or supplementary species (marked *a*, *b*, *c*, &c.).

BUITENZORG, September 1908.

v. A. v. R.

1. CYATHEA, *Smith.*

* (1) **C. moluccana**, *R. Br.*; *C. Brunonis*, *Wall.* — Insert: *Bedd.*, *Ferns Br.I.*, I, tab. LXXXVII.

(2) **C. philippinensis**, *Bl.*, *Christ*, *Bull. Herb. Boiss.*, 2^e Série, VI, 1009.

Trunk 1—2 M. high, slender. Stipes unarmed, angular, rufo-stramineous, ± 10 c.M. long, densely clothed with narrow, subulate, aristate scales provided with a black-brown central line. Fronds 2-pinnate, ± 65 c.M. long, ± 25 c.M. broad, ovate, acuminate, narrowed towards the base; rachises scaly like the stipe, but the scales adpressed, shorter and the main rachis clothed on the upper side with short, grey pubescence. Pinnae close, ± 15 on each side, broadly oblong; central pinnae largest, ± 15 c.M. long, ± 6 c.M. broad, sessile, acuminate. Pinnulae ± 16 on each side, lanceolate, close, sessile, $\pm 2\frac{1}{2}$ c.M. long, ± 1 c.M. broad, cut to $\frac{1}{3}$ -way down (or rarely nearly) to the costa. Lobes triangular, acute, ± 6 on each side, obtuse, entire, ± 3 m.M. each way, the lowest posterior one adnate, the lowest anterior overlapping the rachis. Texture coriaceous; surfaces naked, upper black-green, lower paler; main veins pinnate in the lobes; veins ± 3 on a side, distinct, simple. Sori 2—3 to a lobe; indusium persistent, coriaceous, brown, at length breaking up into irregular pieces. — This

description does not agree with Baker's original diagnosis taken from a cultivated specimen.

Luzon.

* (5) **C. alternans**, *Pr.* — Insert as a synonym: *Polypodium alternans*, *Wall.*, (oldest name).

* (12) **C. cyclodonta**, *v. A. v. R.* — Insert: *Bull. Dép. Agr. I. N.*, 1908, XVIII, 1.

(15a) **C. rufopannosa**, *Christ*, *Philipp. Journ.*, II, 180.

Stipes thick, \pm 55 c.M. long, castaneous, provided at the base with numerous short, pungent verrucae and long, lanceolate, acuminate, falcate, rigid, squarrose, very lucid, castaneous scales and clothed like the rachises with a dense, furfuraceous, fibrillose, brick-red tomentum. Fronds 5-pinnatifid, broadly ovate, to 150 c.M. long, \pm 60 c.M. broad, rather narrowed towards the base, the rachis verrucose. Pinnæ remote, \pm 15 on each side below the pinnatifid apex, central longest, \pm 57 c.M. long, \pm 11 c.M. broad, acuminate, somewhat narrowed at the base, lowest deflexed, \pm 22 c.M. long. Pinnulae \pm 50 on each side, approximate, \pm 5½ c.M. long, \pm 1½ c.M. broad, lanceolate, subacute, cut down to the costa. Segments pectinate, \pm 20 on each side, ligulate, bluntish, crenulate, \pm 5 m.M. long, \pm 2 m.M. broad. Texture herbaceous; surfaces sparingly setulose; costae densely covered beneath with rufous, lanceolate and bullato-umbonate scales; costulae pinnate in the segments; veins forked. Sori small, few, occupying the lower veins; indusium reddish-brown, breaking down irregularly.

Mindanao.

(15a) **C. ferruginea**, *Christ*, *Philipp. Journ.*, II, 181.

Stipes not very thick, and like the rachis somewhat glossy, angular, castaneous, floccose-paleaceous. Fronds 5-pinnatifid. Pinnæ \pm 26 c.M. long, \pm 11½ c.M. broad, stalked, ovate-acuminate, the rachis floccose-paleaceous with rufous, crisped scales. Pinnæ \pm 15 on each side below the lobed apex, \pm 2 c.M. apart, lanceolate, acute, sessile, \pm 6 c.M. long, \pm 11 m.M. broad, cut down to a narrowly winged costa, the lowest somewhat reduced. Lobes \pm 14 on each side, close, ovate, falcate, subacute, \pm 5 m.M. long, \pm 3 m.M. broad, minutely crenulate. Texture herbaceous; colour pale-green, opaque; surfaces nearly naked; costae puberulous above, clothed beneath with grey-

rufous, ovate, acute, adpressed scales; veins ± 6 on each side of the costulae, often forked. Sori small, $\pm 4-5$ on each side, confluent; indusium fugacious, hardly conspicuous.

Palawan.

* (16) **C. Zollingeriana**, *Mett.*

Sori on the lower veins, 1—4 on each side of the costulae, the lowest close to the costae.

(18a) **C. adeno-chlamys**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1008.

Fronds above 1 M. long and 60 c.M. broad, ovate, the rachis rufo- or fulvo-stramineous, pubescent. Pinnae alternate, distant, subpetiolate, articulate to the main rachis, hardly narrowed at the base, to 40 c.M. long, ± 15 c.M. broad, caudate-acuminate. Pinnulae in ± 25 pairs, close, ± 7 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, pinnatifid to the costa, the apex shortly, abruptly acuminate. Segments close, falcate, pectinate, ± 17 on each side, oblong, obtuse or subacute. Texture coriaceous: upper surface black-green, lower paler, but not glaucous: costae and costulae rufo-pubescent on both sides; veins 10—12 on each side, distinct, black, simple, lower forked. Sori $\pm 5-6$ on each side, near the costulae; indusium persistent, rigid, green, covered densely with pale, sessile glands, opening down by a slit.

Luzon.

* (19) **C. orientalis**, *Moore*. — Insert as a synonym: *Disphenia orientalis*, *Kze*, (oldest name), p. p.

* (19a) **C. caudata**, *Copel.*, Philipp. Journ., I, Suppl., 144; *Also-phila caudata*, *J. Sm.*, Hk. Bk., Syn. Fil., 42; v. A. v. R., Mal. Ferns, 37; Hk., Spec. Fil., I, tab. XX, B.

Indusium present, membranaceous, evanishing completely.

(19b) **C. Christii**, *Copel.*, Philipp. Journ., I, Suppl., 144.

Stipes sparingly aculeate with short prickles, minutely and deciduously tomentose, the base clothed with rigid, dirty-brown, setaceous scales. Fronds $1\frac{1}{2}-2$ M. long, 5-pinnatifid, the rachis subsperous below, fuscous and short-pubescent above. Pinnae \pm horizontally spreading, the central ones the largest, ± 50 c.M. long, $\pm 12-14$ c.M. broad, acuminate, the lower diminished,

deflexed. Pinnulae $\pm 1\frac{1}{2}$ c.M. broad, serrato-caudate, cut down nearly to the costa into linear-oblong, remote, denticulate, often obtuse segments. Texture thinly coriaceous; upper surface blackish-green, lower paler, minutely scaly on the veins. Sori costular; indusium fragile.

Luzon, Mindanao.

* (20) **C. runensis**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 1.

(21a) **C. Wernerii**, *Rst.*, Fed., Rep., V, 34.

Fronds 3-pinnatifid, the rachis livid-fuscous, unarmed, strigose above, clothed beneath with linear-lanceolate, ferruginous, copiously and irregularly dilacerate, spinuloso-ciliate scales. Pinnae to 40 c.M. long or longer, to 20 c.M. broad, short-stalked, the petiole articulate to the main rachis, the rachis hirsute and scaly like the main one. Pinnulae 12—15 on each side, sessile, linear, narrowly acuminate, deeply pinnatifid, to 10 c.M. long, ± 2 c.M. broad, slightly narrowed at the base, the terminal one lobed, serrate at the apex. Lobes broadly linear, obtuse, entire, subfalcate, remote. Texture chartaceous; surfaces livid-green, glossy, nearly naked; costae and costulae scaly like the rachises, the scales copiously intermixed with white, acuminate, bullate squamulae; veins 4 on each side of the costulae, forked, conspicuous, brown. Sori \pm medial between the costulae and the margin, ± 7 —8 to each lobe, distant, large, fuscous; indusium globose, pale-fuscous, membranaceous, at length irregularly lacerate, persistent.

New Guinea.

* (28) **C. spinulosa**, *Wall.* — Insert: Bedd., Ferns S.I., tab. LVII.

(29a) **C. negrosiana**, *Christ.* Philipp. Journ., II, 181.

Stipes rather thick, 15—20 c.M. long, the base black, densely clothed with subulate-setaceous, flexuose, lucent, black-brown scales, the rest rufo-castaneous and rough with minute verrucae, as is the rachis. Fronds 5-pinnate, to 1 M. or more long, ± 25 c.M. broad, ovate, acuminate, narrowed gradually towards the base. Pinnae distant, ± 10 —15 on each side, erecto-patent, medial largest, sessile, ± 35 c.M. long, ± 14 c.M. broad, caudato-acuminate, not narrowed at the base. Pinnulae subremote, $\pm 1\frac{1}{2}$ c.M. apart, sessile, ± 20 on each side, cut down to the costa (rachis). Tertiary segments ap-

proximate, falcato-lanceolate, subacute, ± 7 m.M. long, $\pm 2\frac{1}{2}$ m.M. broad, obtusely crenulate. Texture herbaceous; colour black-green, opaque; costae and costulae densely scaly with reddish-brown, inflated scales; veins forked at the base, 8—10 on each side. Sori small, 3—5 on each side, costular; indusium green-brown, soon breaking down irregularly.

Negros.

(51a) **C. Loheri**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1007; — var. *tonglonensis*, *Christ*, Philipp. Journ., II, 180.

Fronde above 1 M. long, pinnate \pm to the base of the stipe. Stipes stout, terete, castaneous, scabrous with minute pustules, clothed densely like the rachises and costae with adpressed, lanceolate, white-scariose, fimbriate scales intermixed with a short, rufous tomentum and small, umbilicate, bullate, rufous scales with a black centre and placed on the pustules. Fronds ovate-oblong, ± 40 c.M. broad, narrowed conspicuously towards the base. Pinnae alternate, short-acuminate, central ones ± 50 c.M. long, ± 12 c.M. broad, ± 10 c.M. apart, the lowest hardly 6 c.M. long. Pinnulae imbricate, ± 22 on each side below the incised apex, ± 7 c.M. long, $\pm 1\frac{1}{4}$ c.M. broad, subfalcate, sessile, acuminate, cut down \pm to the costa. Segments pectinate, falcate, ± 20 on each side, linear-lanceolate, entire or slightly crenulate, subacute, ± 5 m.M. long, $\pm 2\frac{1}{2}$ m.M. broad. Texture subcoriaceous; upper surface black-green, lower paler; costulae rufo-puberulous above, scabrous beneath with small, umbilicate, bullate scales at the base; veins inconspicuous, ± 8 on each side. Sori confluent, at length occupying the whole under surface of the fertile segments; indusium globose, lucid, thin, fulvous, soon opening.

Var. tonglonensis: Stipes rugose and cicatrose, scaly, the scales subulate-setaceous, spreading, flexuose, black-brown, flaccid, verrucose at the base. Costae densely scaly with brown, bullate scales. Segments smaller, densely pectinate.

Luzon.

(51b) **C. callosa**, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1008.

Large. Stipes unknown. Main rachis stout, pale-yellow-brown, covered with numerous minute verrucae and provided with some long, black, scattered setae. Pinnae ± 60 c.M. long, ± 20 c.M. broad, nearly sessile, hardly narrowed at the base, calloso-articulate to the main rachis. Pinnulae close, numerous, sessile, ± 11 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, oblong, acuminate,

caudate, incised to the costa (rachis?), the lowest segments distinct, narrowed at the base, the others pectinate, close, ± 20 , falcate, broadly linear, bluntish, ± 1 c.M. long, ± 4 m.M. broad, crenato-serrulate. Texture coriaceous; upper surface black-green, lower very pale, but not glaucous; secondary and tertiary rachises sparingly clothed with brown, furfuraceous, scattered scales; costulae rufous, densely rufo-puberulous on the upper side; veins black, distinct, 2—3-furcate, ± 11 on each side. Sori ± 4 on a side, costular; indusium pale-green, long-persistent.

Luzon.

* (55) **C. celebica**, *v. A. v. R.* — Insert. Bull. Dép. Agr. I. N., 1908, XVIII, 2.

(54) **C. tripinnata**, *Copel.*, Philipp. Journ., I, Suppl., 251.

Stipes ± 60 c.M. long, black-purple, prickly, densely fulvo-tomentose like the rachis, the base hidden among the long, linear, whitish-stramineous scales of the trunc, the prickles scattered, acute. Fronds ± 2 M. long, $\pm 1\frac{1}{2}$ M. broad, 5-pinnate. Central pinnae largest, ± 80 c.M. long, $\pm 22\frac{1}{2}$ c.M. broad, abruptly acute. Pinnulae 2—2 $\frac{1}{2}$ c.M. broad, abruptly acuminate. Tertiary segments sessile, ± 5 m.M. broad, the base cordate, the edge serrulate, the apex subacute, the higher ones adnate at the base. Texture herbaceous; surfaces naked, but the under side scaly on the costae and some times on the veins. Sori costal, ± 5 on each side, occupying the central veins; indusium whitish-stramineous, thin, soon breaking down into persistent, sericeous fragments. — Near *C. arachnoidea*, *Hk.* and *celebica*, *v. A. v. R.*; the latter has the tertiary segments pinnate at the base.

Luzon, Mindoro.

2. HEMITELIA, *R. Brown.*

(2) **H. capensis**, *R. Br.* — Insert as a synonym: *Polypodium capense*, *L.f.*, (oldest name).

* (4) **H. sumatrana**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 2.

3. ALSOPHILA, *R. Brown.*

* (2) **A. glabra**, *Hk.* — Insert as a synonym: *Gymnosphaera glabra*, *Bl.*, (oldest name).

* (5) **A. squamulata**, *Hk.* — Insert as a synonym: *Gymnosphaera squamulata*, *Bl.*, (oldest name).

(11a) **A. calocoma**, *Christ*, Philipp. Journ., II, 182; — var. *congesta*, *Christ*, l. c.; *A. lepifera*, *J. Sm.*, var. *congesta*, *Christ*, in Bull. Herb. Boiss., VI, 157.

Stipes unarmed, but covered with minute tubercles, clothed at the base with a dense mass of long, scariose, pale, lucent, silvery scales, which are broadly ovate at the base, subulate, caudate, aristate at the apex, the point reddish. Fronds large, 5-pinnatisect, the rachis rufo-ochraceous, opaque, provided with crowded, verrucose tubercles and scattered, furfuraceous scales. Pinnæ \pm 75 c.M. long, \pm 25 c.M. broad, petiolate, caudato-acuminate. Pinnulae close, \pm 20—50 on each side below the pinnatifid apex, nearly sessile, lanceolate, acuminate, \pm 15 c.M. long, \pm 2½ c.M. broad, cut down to the costa, the highest broadly adnate, the lowest not reduced. Tertiary segments ligulate, falcate, bluntish or subacute, \pm 55 on each side, \pm 11 m.M. long, \pm 5 m.M. broad, subentire, close, pectinate, the lowest free, distant, tolerably narrowed. Texture flaccid, herbaceous; upper surface dark-coloured, naked, lower paler, scaly on the costae and costulae with a row of white, lucent, ovate, ciliate, adpressed scales; veins \pm 12 on each side, fine, 2 or more times forked. Sori 8—10 on each side, medial, not confluent.

Var. congesta: Differs from the type in having the ultimate segment shorter, narrower, more falcate and more strongly serrate. The pinnulae do not exceed 8 c.M. in length. — In general appearance different from the type, but with the same essential characters.

Luzon.

* (19) **A. latebrosa**, *Wall.* — Insert as a variety: — var. *major*, *Christ*, Philipp. Journ., II, 185.

A plant found in Mindoro and which is probably this species has the base of the stipes densely clothed: *a*, with long, brown hairs; *b*, with lanceolate, membraneous, almost white scales to 3 c.M. long, with a narrow, black base; *c*, with paleae intermediate in form. — It seems incredible that these

should have escaped description if this is the real *A. latebrosa*, *Wall.* (*Copel.*, *Philipp. Journ.*, II, 145).

Var. major: Main rachis fulvo-stramineous, naked or clothed with minute, furfuraceous scales. Pinnae \pm 28 on each side, \pm 55 c.M. long, \pm 18 c.M. broad, oblong, acuminate, nearly sessile. Pinnulae close, horizontal, sessile, oblong, \pm 2 c.M. broad, caudate, cut down to the slender, black costa. Segments imbricate, oblong, straight, \pm 20 on each side, \pm 2 $\frac{1}{2}$ m.M. broad, obtusely rounded, crenulate. Texture thinly herbaceous; costulae clothed with roundish, peltate, yellow, umbilicate scales; veins 2—3-furcate. — *Mindanao*.

* (20) ***A. saparuensis***, *v. A. v. R.* — Insert: *Bull. Dép. Agr. I. N.*, 1908, XVIII, 2.

(25) ***A. ornata***, *Scott.* — var. *sikkimensis*, *Bedd.*, *Journ. of Bot.*, XXXI, 225; *A. sikkimensis*, *Clarke & Bk.*, *Journ. of Linn. Soc., Bot.*, XXIV, 409.

Var. sikkimensis: Not spinose. Fronds 3-pinnatifid, 1 $\frac{3}{4}$ —3 M. long, the rachis densely clothed with adpressed, brown-castaneous, linear-ligulate scales. Pinnae 45—60 c.M. long, \pm 15 c.M. broad, oblong, the sides parallel. Pinnulae \pm 10 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad. Tertiary segments narrowly oblong, crenate or sometimes slightly incised. Texture thin; rachises ferrugineo-bullate, not short-hairy; veins 2-fid, sometimes 3-fid. Sori 4—10 to a segment, placed in 2 straight rows. — The Malacca fern agrees with the Sikkim plant, except that the stipe and main rachis are more prominently muricate.

Malacca; Sikkim.

* (27) ***A. glauca***, *J. Sm.*; *A. contaminans*, *Wall.* — Insert: *Bedd.*, *Ferns Br.I.*, I, tab. LXXXV.

(27a) ***A. Clementis***, *Copel.*, *Philipp. Journ.*, I, Suppl., 143.

Stipes fuscous in the lower part, furnished with rather long, acute, dense, nearly black scales, the base clothed with linear, whitish-fulvous scales, the upper part stramineous like the main rachis. Fronds 1—1 $\frac{1}{2}$ M. long, ovate, 3-pinnatifid; rachis naked, prickly, the prickles rather short. Central pinnae 30—40 c.M. long, the rachis asperous. Pinnulae 6—7 c.M. long, \pm 1 $\frac{1}{2}$ c.M. broad, cut down nearly to the costa into entire, subfalcate lobes.

Texture coriaceous; surfaces naked, upper blackish-green, lower subglaucous. Sori nude (no spurious indusium?).

Mindanao.

* (32) **A. tomentosa**, *Hk.* — Insert as a synonym: *Gnoophora tomentosa*, *Bl.*, (oldest name), and as a variety: — var. *novo-guineensis*.

Var. novo-guineensis: Spines of the rachis longer; laciniae broader, the secondary ones densely scaly beneath. — *New Guinea.*

* (34) **A. lurida**, *Hk.* — Insert as a synonym: *Gnoophora lurida*, *Bl.*, (oldest name).

(34a) **A. melanorhachis**, *Copel.*, *Philipp. Journ.*, II, 146.

Stipes \pm 60 c.M. long, black, the lower part beautifully brown-dotted, the base clothed with long, lanceolate, rigid, castaneous, pale-margined scales, the rest asperulous, clothed like the main rachis with narrow, crinite, fulvous scales. Fronds to \pm 1½ M. long, broadest below the middle, 5-pinnate. Pinnae \pm 12 on each side, the largest \pm 50 c.M. long, \pm 15 c.M. broad, nearly horizontal, stalked, acuminate, the rachis scaly above, black and glabrescent beneath. Pinnulae distinct, \pm 50 on each side, short-stalked, \pm 7 c.M. long, \pm 1 c.M. broad, the apex acuminate, serrate, the lower part pinnate, the rachis clothed beneath with 2 different kinds of scales, densely short-hairy above. Tertiary divisions obovate, obtuse, adnate, minutely crenato-serrate. Texture thinly coriaceous; upper surface nearly black, naked, lower black-olivaceous, clothed on the costae with ciliate, bullate scales; veins simple, the lower barren ones sometimes forked. Sori costal, filling up the whole under side of the contracted fertile segments.

Mindoro.

4. DICKSONIA, *VHéritier.*

* (1) **D. Blumei**, *Moore.* — Also in the *Philippines.*

(2) **D. Copelandi**, *Christ*, *Philipp. Journ.*, II, 183.

Stipes rufo-stramineous, rather glossy, hairy when young, the base densely clothed with fine, reddish-brown, moderately long hairs. Fronds large. Pinnae deltoid, acuminate, petiolate, \pm 45 c.M. long, \pm 24 c.M. broad, produ-

ced on the upper side at the base. Pinnulae ± 20 on each side, close, lower and central deltoid, lowest ± 15 c.M. long, ± 12 c.M. broad, stalked, acuminate. Tertiary segments deltoid-oblong, acuminate, ± 12 on each side, the lowest ± 6 c.M. long, ± 5 c.M. broad, petiolate, rather unequal-sided. Quaternary segments oblong, acute, the lowest free, $\pm 1\frac{1}{2}$ c.M. long, ± 8 m.M. broad, rather unequal-sided, the margin serrate, the teeth triangular, acute, mucronate. Texture coriaceous; upper surface naked, rather glossy, lower hairy on the veins; colour ochraceous-green; veins forked or pinnate in the ultimate segments, rather obscure. Sori numerous, terminal on the teeth, surpassed by the mucrones, 1 to each tooth, occupying especially the anterior side; indusium coriaceous, globose, irregularly 2-valved.

Luzon.

(5) **D. grandis**, *Rst.*, Fed., Rep., V, 34.

• Fronds large, 4-pinnate. Pinnae subsessile, to nearly 1 M. long, 30 c.M. broad, linear-oblong, the apex acuminate, serrate, the rachis sulcate above. Pinnulae subsessile, linear-lanceolate, acuminate, suberecto-patent or subfalcate, the central ones the longest, to 16 c.M. long, $4\frac{1}{2}$ c.M. broad, the lower not much smaller, the higher gradually reduced. Tertiary divisions short-stalked or sessile, the higher ones sessile with a broad base, the lower and central linear, obtuse, to 2 c.M. long or a little longer, nearly $\frac{3}{4}$ c.M. broad, pinnate towards the base, pinnatifid towards the apex. Ultimate segments short-oblong, narrowed at the base, the lower ones short-stalked, the central sessile, both sterile and fertile deeply toothed with 2–3 teeth on each side. Texture coriaceous; surfaces nearly naked, upper blackish-green, glossy, lower paler; primary rachis unknown, secondary and tertiary ones short-ferrugineo-tomentose above (the tomentum more or less intermixed with longer, rubro-fuscous hairs), the under side deciduously tomentose, glossy, black, verruculose; central ribs of the ultimate segments subpellucid, with 2–3 veins on each side. Sori subglobose, 1 to each ultimate segment, terminal on the lowest anterior vein; indusium 2-valved, the valves similar, subentire.

New Guinea.

5. CIBOTIUM, Kaulfuss.

* **C. barometz**, *J. Sm.*; *Polypodium Barometz*, *L.*, (oldest name).

This species, as construed on page 48, includes several forms: it is not very difficult to recognize the following.

α. Fertile pinnulae rather broad, $1\frac{3}{4}$ — $2\frac{1}{2}$ c.M. broad.

Forma typica (*C. Baranetz*, *J. Sm.*, *Christ*, *Philipp. Journ.*, II, 117): Ultimate segments lanceolate-falcate, acute, subentire or crenato-serrate. Rachis and costae deciduously floccoso-araneous. Sori 1—2 or sometimes 3 or 4 on each side of the costulae, occupying the lower part of the segments, the lowest at a very short distance from the costae, suboblique, but subparallel to the edge, small, not prominent beyond the teeth, \pm 1 m.M. broad; valves of the indusium unequal, the outer one semiglobose, the inner narrower, oblong, yellow-brown, not pruinose. — *Trop. Asia*.

Var. assamicum (*C. Assamicum*, *Hk.*, *Spec. Fil.*, I, 83, tab. XXIX, B; *Christ*, l. c.; *C. glaucum* *J. Sm.*, *Bedd.*, *Ferns Br.I.*, I, tab. LXXXIII): Ultimate segments lanceolate-oblong, acute, subfalcate, distinctly serrate-dentate. Rachis nearly naked; costae and costulae sparingly, deciduously floccoso-araneous. Sori 5—6 on each side in the fully developed fronds, occupying more than the lower half of the segments, prominent beyond the teeth, the lowest remote from the costae, $1-1\frac{1}{2}$ m.M. broad; valves of the indusium unequal, as in the type, brown, not pruinose. — *Assam, Tonkin*.

α.α. Fertile pinnulae rather narrow, $\frac{3}{4}$ — $1\frac{1}{2}$ c.M. broad.

b. Costae and costulae \pm naked or bristly.

Var. sumatranum (*C. Sumatranum*, *Christ*, l. c., 118): Ultimate segments ligulate-oblong, bluntish or acute, crenate-denticulate. Rachis, costae and costulae naked or nearly so. Sori 2 or rarely 3 on each side, occupying the lower part of the segments, the lowest remote from the costae, small, $\frac{1}{2}$ —1 m.M. broad; valves of the indusium nearly equal, rufo-fulvous, not pruinose. — *Sumatra*.

Var. setosum, v. A. v. R., *Bull. Dép. Agr. I. N.*, 1908, XVIII, 5: Like var. sumatranum, but the rachis, costae and costulae densely provided with pale-brown or yellowish bristles spreading in various directions and leaving the rachis subasperous when they fall. Sori 1 or rarely 2 on each side, occupying the base of the segments, remote from the costae; indusium fulvous, not pruinose. — *Malayan?*

b.b. Costae and costulae pubescent or furfuraceous.

Var. Cumingii (*C. Cumingii*, *Kze*, *Christ*, l.c.): Ultimate segments triangular-ovate, acute, decumbent, minutely subserrulate-crenate. Costae and costulae densely pubescent (furfuraceous?, the hairs whitish or rufous. Sori 1 on each side, occupying the base of the segments, adpressed to the costae,

rather large, $1\frac{1}{2}$ —2 m.M. broad, often very oblique and then nearly parallel to the costae; valves of the indusium nearly equal, pale-brown, often pruinose. — A specimen received from Luzon (Leg. A. D. E. Elmer) has the costae and costulae densely but deciduously furfuraceous, the tomentum adpressed, pale-coloured, intermixed with some firm, spreading, pale, rather long, bristle-like hairs. — *Philippines*.

Var. lampongense, v. A. v. R., l.c.: Like var. *Cumingii*, but the ultimate segments falcate, bluntly crenato-serrate. Costae and costulae copiously (not densely), but deciduously furfuraceous, the tomentum loose, pale-brown, intermixed with long, firm, straight or crisped, bristle-like, dark-brown hairs. Sori rather close to the costae, less oblique; indusium brown, not pruinose. — *Sumatra (Lamong Districts, Teysmann)*.

7. GLEICHENIA, *Smith.*

§ 2. EUGLEICHENIA.

* (1) **G. circinata**, *Sw.*; *G. semivestita*, *Lab.* — Insert: *Bedd.*, *Ferns Br.I., II*, tab. CLXXVII.

(2a) **G. dicarpa**, *R. Br.*, *Hk. Bk., Syn. Fil.*, 12; *Hk., Spec. Fil.* I, tab. I, C; *Fil. Exot.*, tab. XL.

Lobes of pinnulae (leaflets) round, subhemispherical, very fornicate. Capsules about 2, concealed within the almost slipper-shaped leaflets and mixed with ferruginous, paleaceous hairs, which often extend to the rachis. — Very near *G. vulcanica*, *Bl.*, and approaching it in some specimens, but larger and the leaflets more close.

Mindoro, Palawan; Australia, Tasmania, New Caledonia.

§ 3. MERTENSIA.

(4) *G. glauca*, *Hk.*, as construed on page 58, includes 2 distinct forms: *G. glauca*, *Hk.* & *gigantea*, *Wall.*

* (4a) **G. glauca**, *Hk.*, *Spec. Fil.*, I, 4, tab. III, B; *G. longissima*, *Bl.*, *Rac., Flor. Btz.*, I, 10, p.p.; *Polypodium glaucum*, *Thb.*, (oldest name): Lobes obtuse, crenulate or serrulate at the apex, very glaucous beneath; costae flattened, not marginate above.

(4b) **G. gigantea**, *Wall.*, Hk., l.c., 5, tab. III, A; *G. longissima*, *Bl.* Rac., l.c., p.p.; *G. excelsa*, *J. Sm.*, Hk., l.c., 5, tab. IV, B: Lobes obtuse, entire, slightly glaucous beneath, provided at the base above with an elevated crest forming a margin on each side of the channelled costa.

* (6) **G. Warburgii**, *Christ.* — Also in *Luzon*.

* (7) **G. linearis**, *Clarke*, var. *normalis*.

An irregularly developed form of this shows a frond with some of the internodations geniculate midway between the accessory branches and the next higher forkings and provided there with 2 simply pinnatifid segments, the segment placed on the outer side provided with an axillary bud at the base on the upper side, and not rarely elongate and growing out to a 5-partite or normally developed forked branch often again geniculate at the middle and provided there with 2 nearly opposite pinnatifid segments. Ultimate lobes irregular, varying from $\frac{1}{2}$ —4 c.M. long.

Billiton (Vorderman), *Banca* (Teysmann, Berkhout), *Penang* (Hunger).

* (7) **G. linearis**, *Clarke*, var. *rigida*. — Insert as synonyms: ? *G. crassifolia*, *Copel.*, Philipp. Journ., I, Suppl., 257 (*G. rigida*, *J. Sm.*, Journ. of Bot., III, 420; *Mertensia crassifolia*, *Pr.*, Epim., 23, tab. XIII).

(7) **G. linearis**, *Clarke*. — Insert as varieties: — var. *tenera*, *Bl.*, Enum., 249; — var. *stipulosa*, *Christ*, Bull. Herb. Boiss; 2^e Série, VI, 1010.

Var. tenera: Texture membranaceous.

Var. stipulosa; Axillary stipulae of all the forkings very long, \pm 20 c.M. long, \pm 4 c.M. broad, developed like the terminal branches, which are shorter, \pm 14 c.M. long, \pm 5 c.M. broad. — *Luzon*.

(7a) **G. laevissima**, *Christ*, Bull. Ac. Int. Bot., 1902, 268.

Fronds to 1 M. high, repeatedly dichotomous, the higher axes interrupted, terminating in a globose coma (tuft) of ovate, acute, ivory-white, squarrose, entire scales. Ultimate fork branches \pm 45 c.M. long, \pm 20 c.M. broad, ovate, narrowed towards both ends. 2-pinnate, the apex caudate. Pinnulae \pm 25 on each side, sessile, the central ones the longest, to 12 c.M. long, the lowest shorter, \pm 6 c.M. long, $\frac{1}{2}$ —2 c.M. broad, caudate. Ultimate segments (leaflets), linear-subulate, falcate, pectinate, \pm 36 on each side, to \pm 1 c.M. long,

the base $\pm 1\frac{1}{2}$ m.M. broad, slightly decurrent, adnate, the margin reflexed. Texture firm; surfaces naked, upper light-green, lower somewhat bluish; ribs distinct, stramineous; veins numerous, forked at the base. Sori medial between the ribs and the margin, few, punctiform; capsules whitish.

Luzon; China.

* (8) **G. laevigata**, *Hk.* — Insert as a synonym: *Mertensia laevigata*, *Willd.*, (oldest name).

(9a) **G. candida**, *Rst.*, *Fed.*, *Rep.*, V, 53.

Stipes subterete, black-brown, or yellow-brown upwards. Fronds large, ± 2 M. long, repeatedly dichotomous, rigid. Lowest internodations 2—5 c.M. long, firm, more or less yellow-brown, naked or clothed with lanceolate, acuminate, brownish scales, not leafy. Secondary and tertiary internodations shorter, of the same colour, deeply pinnatifid, with the lowest segments (lobes) often reduced or wanting; the axillary buds scaly, proliferous. Ultimate branches oblong-lanceolate, acuminate, to 25 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad, straight or curved, deeply pinnatifid; lobes suberecto-patent, the largest to $1\frac{1}{2}$ c.M. long, $5\frac{1}{2}$ m.M. broad, linear-oblong, acute, the edge slightly revolute, serrulate. Texture subcoriaceous; surfaces naked, upper dark-green, lower white or glaucous; costae sparingly scaly; veins pellucid, forked. Sori \pm medial or nearer the margin; sporangia 2—5, yellow.

New Guinea.

(15a) **G. loheri**, *Christ*, *Bull. Herb. Boiss.*, 2^e Série, VI, 1009.

Between *G. hispida*, *Mett.* & *vestita*, *Bl.* Rachises cinnamon-coloured, densely clothed with flaccid, lanceolate and subulate, fimbriato-ciliate, rufous, spreading scales. Ultimate segments triangular-lanceolate, contiguous, $\pm 1\frac{1}{4}$ c.M. long, ± 3 m.M. broad at the base, the edge often reflexed, entire, the apex rather acute; texture subcoriaceous; upper surface black-green, lower bright-green, scaly on the midribs.

Luzon.

* (16) **G. amboinensis**, *v. A. v. B.* — Insert: *Bull. Dép. Agr. I. N.*, 1908, XVIII, 3.

S. HYMENOPHYLLUM, *Smith.*

§ 1. EUHYMENOPHYLLUM.

* (6) **H. australe**, *Willd.* — Insert as a synonym: *H. crispatum*, *Wall.*, *Bedd.*, *Ferns S.I.*, tab. CCVII; *Hk. & Grev.*, *Ic. Fil.*, tab. LXXVII.

* (7) **H. demissum**, *Sw.* — Insert as a synonym: *Trichomanes demissum*, *Forst.*, (oldest name).

(8) *H. subdemissum*, *Christ.* — Omit this as a species. *Christ* considers it now as a form of *H. pycnocarpum*, *V. D. B.*, which he separates from *H. Blumeanum*, *Spr.*

* (14) **H. Blumeanum**, *Spr.*; *H. pycnocarpum*, *V. D. B.*; *H. subdemissum*, *Christ*, *Bull. Herb. Boiss.*, VI, 140; *Philipp. Journ.*, II, 155.

H. pycnocarpum, *V. D. B.*: Smooth. Fronds ovate, 5-pinnatifid, the rachis winged. Sori terminal, paniculate, the valves acutely triangular.

(22a) **H. halconense**, *Copel.*, *Philipp. Journ.*, II, 144.

Rhizome filiform, creeping, hirsute, at length naked. Stipes filiform, 2—5 c.M. long, glabrescent, winged narrowly towards the frond. Fronds 4—9 c.M. long, 1½—2 c.M. broad, 2-pinnatifid, with the secondary segments simple or sometimes forked, the edges entire, the rachis winged throughout. Upper surface naked, lower covered with red-brown hairs on the rachis, costae and veins and the bases of the indusia. Sori sessile, hardly sunk ½-way down, terminal on the lowest, anterior secondary segments.

Mindoro.

§ 2. LEPTOCIONIUM.

* (27) **H. Kurzii**, *Prantl.* — I found in the Buitenzorg Herbarium about 5—4 plants quite similar to those gathered by the Ladies von Grävenitz and Stein; they were labeled with the names of other species. The wings of the secondary rachises are whether or not denticulate.

(29) **H. tunbridgense**, *Sm.* — Insert as a synonym: *Trichomanes tunbridgense*, *L.*, (oldest name).

(29a) **H. campanulatum**, *Christ*, Philipp. Journ., II, 155.

Densely caespitose. Rhizome slender, but firm, branched. Stipes $\pm 1\frac{1}{2}$ c.M. long, slender, black and hispid with rigid hairs, as is the rachis. Fronds ± 4 c.M. long, ± 1 c.M. broad, oblong, narrowed at both ends. Pinnae ± 6 on each side, arcuate-reflexed, ± 1 c.M. long, flabellatipartite with 4—5 linear lobes hardly 1 m.M. broad and sparingly aristate-serrate. Texture rather rigid; colour fuscous. Sori few, placed near the base of the costae, stalked; indusium campanulate, the valves ovate, entire. — General habit of *H. tunbridgense*, *Sm.*

Negros.

(51a) **H. Merrillii**, *Christ*, Philipp. Journ., II, 154.

Rhizome creeping, filiform, caespitose, sparingly clothed with short, reddish-brown hairs, as are the stipes. Stipes filiform, ± 5 c.M. long. Fronds ovate, ± 6 c.M. long, ± 5 c.M. broad, acuminate, narrowed at the base, the rachis sparingly hairy, not winged. Pinnae close, ± 8 on each side, cuneate-ovate, higher acute, sessile, not adnate, lower stalked, ± 6 m.M. broad, deeply pinnatifid. Secondary segments cuneate, obtuse, ± 5 on each side, deeply incised. Lobes lanceolate, flat, ± 2 m.M. broad, serrulate-dentate. Texture rather rigid; colour black-brown. Sori ± 5 —4 on each side, solitary on the pinnae, occupying the lowest anterior laciniae; indusium ovate, the apex 2-valved, the valves serrulate; receptacle much exerted.

Luzon.

* (54) **H. multifidum**, *Sw.* — Insert as a synonym: *Trichomanes multifidum*, *Forst.*, (oldest name).

(55) **H. serrulatum**, *C. Chr.* — Insert as a synonym: *Didymoglossum serrulatum*, *Pr.*, (oldest name).

A form gathered in Johor State (Malacca), which may be a narrow intermediate between this species and *H. multifidum*, *Sw.* has the fronds 5—8 c.M. long, $1\frac{1}{2}$ —2 c.M. broad, the main rachis flexuose, sparingly piloso-paleaceous below, winged above, the wing entire. Pinnae erecto-patent, pinnatifid, the segments linear, simple or forked, the ultimate lobes 2—5 m.M. long, under 1 m.M. broad, serrate, the serratures rather distant. Indusium elliptical, whether or not exerted.

* (42) **H. aculeatum**, *Rac.* — Insert as a synonym: *Trichomanes aculeatum*, *J. Sm.*, (oldest name).

9. TRICHOMANES, *Linné*.

§ 2. EUTRICHOMANES.

* (9) **T. nitidulum**, *V. D. B.* — Also found in *Luzon* and *Mindoro*.

* (15) **T. diffusum**, *Bl.* — Also found in *Luzon* and *Mindanao*.

(22a) **T. Wernerii**, *Rst., Fed., Rep., V, 35.*

Rhizome wide-creeping, filiform, short-tomentose. Stipes slender, ± 5 m.M. long, winged nearly to the base. Fronds ovate, to 3 c.M. long, 2 c.M. broad. Pinnæ to 7-jugate, the central ones the longest, ± 12 m.M. long, 5 m.M. broad, lanceolate-rhomboidal, the lower \pm smaller. Lower secondary segments pinnatifid, higher forked or simple, the laciniae linear, $\frac{4}{1}$, $-\frac{5}{10}$ m.M. broad, simple or the lowest forked, emarginate at the apex, the margin bounded by a diaphanous line. Texture thin; rachises narrowly winged: veins slender. Sori immersed; indusium infundibuliform, the mouth dilated, undulate; receptacle 12--15 \times longer than the indusium ($2\frac{1}{2}$ —3 c.M. long).

New Guinea.

(59) **T. maximum**, *Bl.* is construed as including 2 distinct forms: **T. maximum**, *Bl.* & **millefolium**, *Pr.*

* (59a) **T. maximum**, *Bl.*, Enum., 228; Rac., Flor. Btz., I, 29; *V. D. B.*, Hym. Jav., 25, tab. XVIII.

Stipes not winged, or winged towards the frond only. Ultimate lobes recurved towards the apex. Indusium winged quite to the mouth.

* (59b) **T. millefolium**, *Pr.*, Hym., 16, 43; *V. D. B.*, Hym. Jav., 27, tab. XX; *T. anceps*, *Hk., var. β* , *Hk.*, Spec. Fil; I, tab. XL, C, fig. 3.

Stipes winged \pm throughout. Ultimate lobes flat, straight. Indusium stalked, or winged at the base only.

(41) **T. javanicum**, *Bl.* is construed as including 2 distinct forms: **T. javanicum**, *Bl.* & **rhomboideum**, *J. Sm.*

* (41a) **T. javanicum**, *Bl.*, Enum., 224; *Hk. & Grev.*, Ic. Fil.,

tab. CCXL; Rac., Flor. Btz., I, 27, p. p.; *Cephalomanes javanicum*, Pr., V. D. B., Hym. Jav., 50, tab. XXII; *C. Zollingeri*, V. D. B., l. c., 51, tab. XXIII; — var. *intercalatum*, Christ, Philipp. Journ., II, 156.

Pinnae irregularly toothed, the teeth on the anterior side blunt, those on the posterior side acute or subsetiform, but rather short. Indusium erect, with the mouth slightly contracted or slightly dilated.

Var. intercalatum: Pinnae often dissected, the lower side toothed, the teeth elongate, aristate, conspicuous. Sori numerous, \pm 12 to a pinna, placed on the anterior side, the apex and the upper part of the posterior side, all conspicuously exserted; indusium clavato-campanulate, the mouth not dilated. — *Luzon*.

* (41b) **T. rhomboideum**, *J. Sm.*, Journ. of Bot., III, 417; *T. javanicum*, *Bl.*, Rac., l. c., p. p.; *Cephalomanes rhomboideum*, V. D. B., l. c., 55, tab. XXIV.

Like *T. javanicum*, *Bl.*, but the teeth on the posterior side very long, setiform, often curved. Indusium reflexed, the mouth distinctly dilated.

* (42a) **T. sumatranum**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XVIII, 4.

Near *T. javanicum*, *Bl.* and *aspolenioides*, Pr. Pinnae growing gradually shorter towards the apex of the frond, the lower rather suberect, lobed on the upper side, the lobes toothed at the apex like the apex of the pinnae, the 1—5 lower lobes \pm elongate and auriculiform, free in the lowest pinnae and then not rarely short-stalked; higher pinnae erect, undivided, toothed, the uppermost contracted to mere, narrowly marginate petioles to the sori. Sori confined to the higher part of the frond, terminal on the pinnae, those on the not contracted pinnae sessile; indusium narrowly winged, the limb erect, truncate, upcurved at the sides, provided with 2 short, lateral, tooth-like, erect processes consisting of the slightly produced ends of the wings; receptacle exserted.

Sumatra (Burck).

(47a) **T. Christii**, *Copel.*, Philipp. Journ., I, Suppl., 251.

Rhizome wide-creeping. Stipes slender, \pm 1 c.M. long, sparingly pubescent with a few short hairs. Fronds 5—8 c.M. long, $2\frac{1}{2}$ — $5\frac{1}{2}$ c.M. broad, acute, 5-pinnatifid, the main rachis narrowly winged in the upper part. Pinnae oblong, obtuse, the lowest reduced. Ultimate segments linear, entire.

Indusium infundibuliform-campanulate, narrowly winged, the limb suberect, sub-2-labiate; receptacle included.

Mindoro.

* (48a) **T. Merrillii**, *Copel.*, Philipp. Journ., I, Suppl., 144, tab. I; — var. borneense.

Rhizome erect, strong. Stipes tufted, 10—15 c.M. long, clothed with short, rigid, reddish-black hairs. Fronds 15—20 c.M. long, acute, 2-pinnate, the rachis terete, pilose. Pinnae \pm spreading, 3—3½ c.M. long, obtuse; higher pinnae \pm erecto-patent. Lowest acroscopical pinnulae largest, \pm 9 m.M. long, \pm 5 m.M. broad, deeply incised, the segments setiform. Texture coriaceous; colour blackish-green; surfaces naked. Sori deflexed, 2-angular, placed above the axils of the acroscopical segments of the pinnae; indusium with the mouth slightly dilated. — General aspect of *T. javanicum*, *Bl.*

Var. borneense: Fronds smaller. Pinnulae somewhat shorter and narrower. Sori not deflexed, narrowly 2-marginate; receptacle not conspicuously exerted.

Palawan; — var. borneense: *Borneo* (Teuscher).

* (49a) **T. setaceum**, *V. D. B.*, Ned. Kruidk. Arch., V², 176; *T. cupressoides*, *Desv.*, var., *C. Chr.*, Ind. Fil.

Rhizome short, ascending. Stipes approximate, subtufted, 10—18 c.M. long, clothed with deciduous, fuscous hairs, at length quite naked. Fronds 15—20 c.M. long, 5—7 c.M. broad, oblong, acuminate, 2-pinnate, the rachis densely paleaceo-hirsute. Pinnae \pm spreading, the lower ones often deflexed, the higher erecto-patent, oblong or lanceolate-oblong. Pinnulae erecto-patent, subcontiguous, lanceolate, deeply incised, the segments setiform, long. Texture firm; surfaces naked; colour olive-green. Sori in the axils of the lower anterior segments of the pinnae, \pm deflexed, emerged; indusium not conspicuously marginate, cylindric-infundibuliform, the mouth not dilated; receptacle exerted.

Singapore, Banca, Borneo.

10. LYGODIUM, *Swartz.*

§ 1. EULYGODIUM.

* (2) **L. Teysmannii**, *v. A. v. B.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 5.

FILICES.

(2a) **L. basilanicum**, *Christ*, Philipp. Journ., II, 179.

Main rachis slender, ochraceous. Primary petioles \pm 5 c.M. long, narrowly winged. Pinnulae stalked, 5—4-lobate, the stalk \pm $\frac{1}{2}$ c.M. long, winged, the central part undivided; lobes pedately arranged, \pm 9 c.M. long, \pm 6 m.M. broad, patent, thinly marginate, the basal ones deflexed. Texture coriaceous; costulae slender, distinct, rufo-stramineous; veins prominent, oblique, 2—5-furcate, close. Fertile segments suddenly contracted at or below the middle to a wing \pm 1 m.M. broad. Soriferous spikes close, pectinate, \pm $2\frac{1}{2}$ m.M. long.

Basilan.

* (5) **L. circinatum**, *Sw.* — Insert as a synonym: *Ophioglossum circinatum*, *Burm.*, (oldest name). — After var. *monstruosum* & *cristatum*, insert: v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 5.

Var. cristatum: Marginal strand very indistinct, apparently wanting. By its crested spores this approximates the examined specimens of *L. trifurcatum*. *Bk.*, but the fertile spikes are closely pectinate, not in groups.

* (5) **L. trifurcatum**, *Bk.* — The specimens gathered in New Guinea by Prof. Dr. Treub and those cultivated in the Buitenzorg gardens have the laciniae narrower than in the typical form, the fertile spikes partially in groups of 1—5, and the spores verrucose and crested.

* (7) **L. scandens**, *Sw.* — Insert as a synonym: *Ophioglossum scandens*, *L.*, (oldest name).

* (10) **L. flexuosum**, *Sw.* — Insert as a synonym: *Ophioglossum flexuosum*, *L.*, (oldest name).

* (11) **L. japonicum**, *Sw.* — Insert as a synonym: *Ophioglossum japonicum*, *Thb.*, (oldest name).

(11a) **L. Mearnsii**, *Copel.*, Philipp. Journ., III, 57.

Near *L. flexuosum* & *japonicum*, *Sw.* Primary petioles 5—9 m.M. long. Barren pinnae \pm 15 c.M. long, 12—18 c.M. broad. Pinnulae 2—5 on each side, pinnate, on stalks more than 1 c.M. long. Leaflets in 1 pair below the terminal ones; lateral leaflets stalked, 5-fid with obtuse lobes, terminal ones larger, lanceolate; edges obscurely crenate, the crenations denticulate. Texture

herbaceous; surfaces naked. Fertile pinnae smaller, the leaflets entire, ovate. Spikes short, oblong.

Batan Island.

§ 2. HYDROGLOSSUM.

(12) **L. Matthewi, Copel.**, Philipp. Journ., III, 56.

Fronds high-scandent. Pinnae more than 50 c.M. long. Pinnulae stalked, 1—2 on each side below the terminal one, not or hardly articulate at the base, the stalks $1\frac{1}{2}$ — $2\frac{1}{2}$ c.M. long; terminal pinnulae deeply palmatifid with \pm 6 spreading segments 20—30 c.M. long, 5—4 c.M. broad, the lateral ones 2—3-furcate with the segments \pm 20 c.M. long; edges serrulate. Texture herbaceous; surfaces naked, or the veins pubescent at the base on the under side; veins anastomosing. — Fertile fronds unknown.

Philippines.

(15) **L. Merrilli, Copel.**, Philipp. Journ., II, 146, tab. IV.

Near *L. Matthewi, Copel.*, but the terminal pinnulae forked. Fertile fronds with the spikes not very numerous, not close (conf. Copeland's drawing).

Mindoro.

11. SCHIZAEA, *Smith.*

* (3) **S. dichotoma, Sm.** — Insert as a synonym: *Acrostichum dichotomum, L.*, (oldest name).

* (4) **S. digitata, Sw.** — Insert as a synonym: *Acrostichum digitatum, L.*, (oldest name).

13. DIPLAZIOPSIS, *Christensen.*

Indusium sausage-shaped, quite enclosing the sorus when young, rupturing irregularly when the spores mature, apparently fastened all round the receptacle, i.e., the indusium is fixed by its very broad base to the posterior side of the vein, with the free edge strongly involute and closely adpressed to the anterior side of the vein. — Rac., *Ueber die Farngattung Allantodia*, Wall., Bull. Ac. Crac., 1905, 546—549; Bot. Centralbl., 1906, I, 261.

14. DENNSTAEDTIA, *Bernhardi*.

(5) **D. erythrorachis**, *Diels*. — Also in the *Philippine Islands*.

(6a) **D. dennstaedtioides**, *Copel.*, *Philipp. Journ.*, II, 126; *Microlepia dennstaedtioides*, *Copel.*, l. c., I, Suppl., 148, tab. IV; — var. *arthrotricha*, *Copel.*, l. c., III, 34.

Rhizome creeping, naked or nearly so. Stipes 15—20 c.M. long, brown, naked. Fronds 40—60 c.M. long, 20—30 c.M. broad, 4-pinnatifid. Pinnae lanceolate, acute, the lower smaller, the rachis narrowly winged. Pinnulae ovate-lanceolate, obtuse. Ultimate segments obovate, entire or toothed. Texture herbaceous; surfaces naked. Sori small, marginal or submarginal.

Var. arthrotricha: Fronds larger; rachis and upper surface clothed with short, articulate hairs.

Mindanao, Luzon.

(6b) **D. Merrilli**, *Copel.*, *Philipp. Journ.*, II, 126; (? *Microlepia*).

Rhizome thick, creeping. Stipes erect, firm, \pm 60 c.M. long and like the main rachis castaneous, slightly puberulous, rather glabrescent. Fronds 100—120 c.M. long, \pm 50 c.M. broad, 4-pinnatifid. Largest pinnae \pm 50 c.M. long, \pm 11 c.M. broad, acuminate, with the rachis puberulous below; lower pinnae reduced. Pinnulae lanceolate-deltoid, distinctly acuminate. Tertiary segments lanceolate-deltoid, obtuse, decurrent, subpetiolate, the lowest acroscopical ones the largest, to 2 c.M. long, \pm 1 c.M. broad, cut down on each side to a winged costa into \pm 4 subremote, obtuse or truncate, entire or obscurely serrate lobes. Texture herbaceous; surfaces naked, the lower paler than the upper. Sori \pm marginal, in the sinuses, the margin of the lobe forming with the true indusium an imperfect cup to the sorus.

Mindoro.

(7) **D. samoensis**, *Moore*. — Insert as a synonym: *Sitolobium samoense*, *Brack.*, *Expl. Exp.*, XVI, tab. XXXVIII, fig. 1.

(7a) **D. Williamsi**, *Copel.*, *Philipp. Journ.*, I, Suppl., 148; *Christ*, l. c., II, 168.

Stipes tufted, \pm 2 M. long, rufous-yellowish, densely clothed like the rachis with fulvous, strigose hairs, at length rather naked. Fronds \pm 2 M. long, 4-pinnatifid. Pinnae sessile, \pm 90 c.M. long (\pm 120 c.M. long after

Copel.), to 50—55 c.M. broad, oblong, acuminate, hardly narrowed at the base. Pinnulae \pm 35 on each side, petiolate, oblong, acuminate, lowest distant, \pm 16 c.M. long, 4—7 c.M. broad, broadest at the base, upper base truncate, lower cuneate; higher pinnulae approximate. Tertiary divisions close, rather imbricate, decurrent, \pm 25 on each side, lowest anterior largest, \pm 2 c.M. long, \pm $\frac{1}{2}$ c.M. broad, deeply pinnatifid. Lobes oblong, obtuse, crenate on the upper side, \pm $\frac{1}{2}$ c.M. long, \pm $\frac{1}{4}$ c.M. broad. Texture rigidly coriaceous; colour grey-green; costae and veins clothed with short, adpressed, rufous hairs. Sori in the sinuses, 1 or rarely more on each lobe, marginal.

Mindanao.

(7b) **D. Hooveri**, *Christ*, Philipp. Journ., II, 169.

Rhizome creeping. Stipes $1\frac{1}{2}$ —2 M. long. Fronds \pm 2 M. long, subdeltoid, 4-pinnatifid, the rachises pubescent with short, strigose, rufous hairs. Pinnae \pm 60 c.M. long, \pm 18 c.M. broad, short-stalked, hardly narrowed at the base, acuminate. Pinnulae alternate, approximate, petiolulate, \pm 55 on each side, \pm 9 c.M. long, \pm 5 c.M. broad, the upper base produced, the rachis not winged. Tertiary segments \pm 18 on each side, close, subimbricate, obtusely ovate, \pm 1 c.M. long, \pm $\frac{1}{2}$ c.M. broad, cut down to the costa, unequal-sided, upper base produced, truncate, lower cuneate. Ultimate segments 5—4 on each side, cuneate-oblong, obtuse, \pm 5 m.M. long, \pm $2\frac{1}{2}$ m.M. broad, crenato-lobate in the fertile, elongate and more acutely toothed in the barren fronds. Texture thinly herbaceous; colour dark-green; surfaces pilose with whitish, tortuous hairs; veins pinnate and forked in the ultimate segments. Sori roundish, 1 (or rarely more) on each ultimate segment, placed on the short, obtuse, basal teeth; indusium thin, membranaceous, grey, soon evanished.

Midanao.

* (10) **D. cuneata**, *Moore*. — Insert as a synonym: *Sitolobium cuneatum*, *J. Sm.*, (oldest name).

* (12) **D. flaccida**, *Bernh.* — Insert as a synonym: *Trichomanes flaccidum*, *Forst.*, (oldest name).

(13a) **D. Elmeri**, *Copel.*, Leaflet, I, 233.

Rhizome creeping, black, aculeate. Stipes to 25 c.M. apart, probably

to 2 M. long, aculeate. Fronds to 5 M. long, \pm 1—1½ M. broad, 5-pinnate, the main rachis short-aculeate, castaneous. Pinnæ opposite, apparently articulate to the main rachis, broadly lanceolate, 20—25 c.M. broad, acuminate, the rachis aculeate towards the base. Pinnulae sessile, pseudo-articulate, the lower ones opposite, the higher alternate, acuminate, the rachis slightly pubescent. Leaflets sessile, obtuse, auriculate, to 1½ c.M. long, to 7 m.M. broad, the edge cut more than ½-way down to the costa. Texture subcoriaceous; costae slightly pubescent beneath, the rest naked. Sori to 7 on a leaflet, at the tops of the sinuses, inflexed.

Luzon.

* (14) **D. sumatrana**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 6.

15. OLEANDRA, *Cavanilles.*

* (1) **O. cuspidata**, *Bk.* — A form gathered in New Guinea has the rhizome scales subulate and much deflexed at the apex, the fronds \pm 25—30 c.M. long, \pm 4—5 c.M. broad, the surfaces hairy on the veins, the indusia fugacious. — This may be Copeland's *O. colubrina*, *Copel.*, *var. membranacea*.

* (3) **O. colubrina**, *Copel.*, *var. brachypus*. — Copeland distinguishes 2 forms (Philipp. Journ., III, 32—33):

Var. nitida (? *var. brachypus*): Fronds to 40 c.M. long, \pm 2 c.M. broad, coriaceous, soon naked, glossy.

Var. membranacea (? *O. cuspidata*, *Bk.*). Fronds \pm 20 c.M. long, \pm 2½ c.M. broad, narrowly caudate, pubescent, ciliate, membranaceous. Sori small, in an irregular row \pm medial between the costa and margin.

(4a) **O. Wernerii**, *Rst.*, Fed., Rep., V, 40.

Rhizome long, scandent, subcompressed, branched, scaly; scales dense, lanceolate, deciduously acuminate at both ends, at length ovate, adpressed, peltate, the centre black, the margin rufo-ferrugineous, fimbriate; branches 3—10 c.M. or more apart, simple or branched again, slightly scaly, the ultimate ones to ½—5 c.M. long, bearing the fronds. Stipes \pm 2—4 c.M. long, articulate \pm at the base. Fronds dimorphous. Barren fronds elongate-

lanceolate, to 25 c.M. long, to 5 c.M. broad, narrowed towards both ends, the base cuneate, the edge entire, the point long-cuspidate. Texture papyraceous; surfaces naked, \pm glossy; costa prominent on both sides, sparingly scaly; veins \pm 1—2 \times forked. Fertile fronds linear, to 50 c.M. long, 1½—2 m.M. broad (!). Sori rather distant, large, transversal (not oblique), often protruded beyond the margin of the frond; indusium reniform.

New Guinea.

* (5) **O. musifolia**, *Pr.* — Insert as a synonym: *Aspidium musifolium*, *Bl.*, (oldest name).

17. NEPHROLEPIS, *Schott.*

§ 1. LINDSAYOPSIS.

* (5a) **N. davalliae**, *v. A. v. B.*, Bull. Dép. Agr. I. N., 1908, XXI, 2.

Rhizome short, stoloniferous, densely clothed with bright-brown, subulate scales. Stipes \pm 15 c.M. long, approximate, grey-brown and deciduously furfuraceous like the rachis, the tomentum consisting of fuscous, crisped, fibril-like scales. Fronds \pm 50 c.M. long, narrowed towards the apex. Pinnæ numerous, rather distant, horizontally spreading, falcate, short-stalked, lower sterile, lowest more or less reduced but not auricle-like, oblong, higher fertile, central largest, \pm 4 c.M. long, 5—6 m.M. broad at the rotundato-truncate base, narrowed towards the apex, the margin entire or shallowly crenate in the lower part, more deeply crenate, dentate or dentato-lobate towards the apex, the crenations or teeth oblique, bluntish, the fertile ones truncate. Texture coriaceous; surfaces naked; veins 2—5-furcate. Sori roundish, solitary and terminal on the crenations or teeth, rarely solitary on the veins, mostly uniting the apices of the fork branches of the veins, slightly immersed and prominent on the upper surface; indusium brown, semiorbicular, attached at the rounded base and sides, free at the truncate apex, opening outwardly.

New Guinea.

§ 2. EUNEPHROLEPIS.

* (5) **N. floccigera**, *Moore.* — Insert as a synonym: *Aspidium floccigerum*, *Bl.*, oldest name.

* (8) **N. hirsutula**, *Pr.* — Insert as a synonym: *Polypodium hirsutum*, *Forst.*, (oldest name).

(9a) **N. glabra**, *Copel.*, Philipp. Journ., I, Suppl. 146.

Near *N. cordifolia*, *Pr.* Fronds 7—15 c.M. long, rarely longer, lanceolate, the rachis winged like the stipe, naked or nearly so. Pinnae hardly deciduous, the fertile ones crenato-serrate. Texture herbaceous; veins distinct. — This species is notable for its exposure to extremes of moisture. Instead of resisting desiccation, as do all the other species, this one yields readily and endures it, without losing its pinnae, and promptly recovers (*Copel.*).

Luzon.

* (15) **N. exaltata**, *Schott.* — Insert as a synonym: *Polypodium exaltatum*, *L.*, (oldest name).

18. CYCLOPELTIS, *J. Smith.*

* (1) **C. semicordata**, *J. Sm.* — Insert as a synonym: *Polypodium semicordatum*, *Sw.*, (oldest name).

* (2) **C. Presliana**, *Berk.* — Insert as a synonym: *Lastraea Presliana*, *J. Sm.*, (oldest name). — After var. *biauriculata*, insert; v. A. v. R., Bull. Dép. Agr. I. N., 1908, XVIII, 6.

19. POLYSTICHUM, *Roth.*

* (1) **P. auriculatum**, *Pr.* — Insert: *Bedd.*, Ferns S.I., tab. CXX; and after var. *nervosum*, insert: *P. deltodon*, *Diels*, var. *nervosum*, *Christ*, Philipp. Journ., II, 157.

(1a) **P. Copelandi**, *Christ*, Philipp. Journ., II, 157.

Rhizome caespitose. Stipes fasciculate, slender, $\pm 2\frac{1}{2}$ c.M. long, sparingly clothed with subulate, black scales, as is the rachis. Fronds linear-lanceolate, simply pinnate, not narrowed at the base, to 20 c.M. long and 2 c.M. broad, acuminate. Pinnae densely imbricate, ± 50 on each side, rhomboidal, ± 1 c.M. long, $\pm \frac{1}{2}$ c.M. broad at the base, the apex subacute, the edges parallel; upper base acutely auriculate, truncate, lower cuneate; lower side entire, upper minutely toothed, the teeth not aristate. Texture herba-

ceous; surfaces naked; colour yellow-green, opaque; veins forked. Sori \pm 6 on a pinna, placed in 1 row on the anterior side of the costae, distant from the margin, impressed; indusium minute, fugacious.

Luzon.

(1b) **P. obliquum**, *M. vore*, C. Chr., Ind. Fil.; *P. caespitosum*, Schott., Bedd., Ferns Br.I., I, tab. XX'XIII; *P. auriculatum*, Pr., var. *caespitosum*, Bedd., Handb. Ind. Ferns, 204; *Aspidium auriculatum*, Sw., var. *obliquum*, Hk. Bk., Syn. Fil., 493; *A. obliquum*, Don, (oldest name); — var. *luzonicum*, Christ, Philipp. Journ., II, 157.

Rhizome short, thick, the scales ferrugineous, ovate, often ciliate. Stipes tufted, slender, 5—20 c.M. long, stramineous as well as the rachis. Fronds 10—25 c.M. long or longer, oblong- or linear-lanceolate, rather acute, simply pinnate. Pinnae rather distant, horizontal, rhomboidal-oblong, $1\frac{1}{2}$ — $3\frac{1}{2}$ c.M. long, obtuse or acute, sharply but scarcely mucronato-serrate; upper base truncate and obtusely auricled, lower cuneate. Texture chartaceo-membranaceous; surfaces naked; veins forked, lowest anterior pinnate in the auricle. Sori in 1 row on each side of the costae, medial or nearer the edge than the costae; indusium peltate, ciliate or toothed, at length deciduous.

Var. luzonicum: Pinnae imbricate, acutely auricled, the edge nearly entire, the teeth very faint, scarcely visible but nevertheless aristate. Texture herbaceous; veins hidden.

North. India; — var. *luzonicum*: *Luzon*.

(2a) **P. nudum**, *Copel.*, Philipp. Journ., I, Suppl., 145.

Rhizome ascending, clothed with long, linear, brown scales. Stipes close, 40—50 c.M. long, clothed towards the base with long, brown, often ovate scales and towards the frond with paler, often smaller, crinite ones. Fronds \pm 70 c.M. long, \pm 50 c.M. broad, the rachis clothed with brown, crinite scales. Pinnae close, horizontally spreading, 15—18 c.M. long, \pm $2\frac{1}{2}$ c.M. broad, the lowest somewhat shorter. Pinnulae oblong, acute, inciso-serrate, the teeth aculeate, the auricle nearly free, not imbricating. Texture herbaceous; upper surface naked, lower pale-coloured, provided with whitish scales on the veins. Sori naked.

Mindanao.

* (3) **P. aculeatum**, *Schott.* — Insert as a synonym: *Polypodium aculeatum*, L., (oldest name).

(4a) **P. blepharistegium**, *Copel.*, Philipp. Journ., I, Suppl., 145.

Stipes close, 25—40 c.M. long, clothed with rather long, polymorphous scales. Fronds to \pm 40 c.M. long, to \pm 25 c.M. broad, the rachis densely clothed with narrow, pale-fulvous, silky hairs and scales. Lower pinnae deflexed, hardly shorter than the following which are \pm horizontally spreading, \pm 15 c.M. long, 2—2 $\frac{1}{2}$ c.M. broad, acuminate, pinnate. Lowest acroscopical pinnulae cut down nearly to the costa, the others not or hardly incised, broadly auriculate, obscurely toothed, the teeth often provided with short setae, the auricle imbricating. Texture papyraceous; upper surface nearly naked, lower clothed with fulvous, stellate hairs. Indusium whitish, silky, ciliate.

Luzon.

(6) **P. lobatum**, *Pr.* — Insert as a synonym: *Polypodium lobatum*, *Huds.*, (oldest name).

(8) **P. platypus**, *Diels.* — Insert as synonyms: *Aspidium platypus*, *Kze*, Bot. Zeit., 1848, 262; *Lastraea platypus*, *Moore*, Bedd., Ferns Br.I., II, tab. CCXXVIII.

* (10) **P. aristatum**, *Pr.* — Insert as a synonym: *Polypodium aristatum*, *Forst.*, (oldest name). — Omit: var. *Hamiltonii*, it being not malayan.

21. PLEOCNEMIA, *Presl.*

(1) **P. heterophylla**, *v. A. v. R.* — Insert as a synonym: *Haplodictyum heterophyllum*, *Pr.*, (oldest name).

* (4) **P. Leuzeana**, *Pr.* — Insert: *Tectaria leuzeana*, *Copel.*, Philipp. Journ., II, 417; *Polypodium Leuzeanum*, *Gaud.*, (oldest name); — var. *angilogensis*; *Aspidium Angilogense*, *Christ*, Bull. Herb. Boiss., 2^e Série, VI, 1005.

Var. angilogensis: Very large, the adult plant very probably arborescent. Main rachis fulvo-castaneous, naked or furfuraceous, sublucid. Pinnae petiolate, \pm 60 c.M. long, \pm 18 c.M. broad, acuminate. Pinnulae horizontally spreading, \pm 35 on each side, short-stalked, \pm 3 c.M. broad, oblong, acute, deeply pinnatifid. Lobes pectinate, \pm 20 on each side, broadly

linear, subacute, 5 m.M. broad, separate, the sinuses rounded, the higher lobes close, the edges lobato-crenate, the teeth obtuse, \pm 8 on each side, \pm 1 m.M. each way. Costae and veins subfurfuraceous or naked. Sori near the costulae, in the forkings of the veins, 6—8 on a side. — See Copeland's information, l. c. — *Luzon*.

* (5) **P. gigantea**, *Pr.* — Omit as synonyms: *P. Trimeni*, *Bedd.* and *Sagenia gigantea*, *Moore*. — *P. Trimeni*, *Bedd.* is a distinct species and *Sagenia gigantea*, *Moore* its synonym.

(7) *P. Bakeri*, *v. A. v. R.* — Omit this name, it should be *P. Trimeni*, *Bedd.*

* (7) **P. Trimeni**, *Bedd.*, *Handb. Ind. Ferns*, 223; *Nephrodium giganteum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 503; *Sagenia gigantea*, *Bedd.*, *Ferns S.I.*, tab. LXXX; *Aspidium giganteum*, *Copel.* (not *Bl.*), *Polypod. Philipp.*, 38; (*Bl.*, *C. Chr.*, *Ind. Fil.*, p.p.).

Stipes tufted, 50—60 c.M. long, grey or brown; basal scales linear. Fronds 50—90 c.M. long, deltoid or deltoid-lanceolate, pinnatifid at the apex, below pinnate with \pm 5—6 lateral pinnae on each side, which are more or less stalked and all pinnatifid more than $\frac{1}{2}$ -way down to the costa. Lowest pinnae largest, pinnate at or towards the base, with the lower (lowest) basis-copical pinnulae prolonged and deeply pinnatifid. Ultimate lobes bluntish, slightly crenate or nearly entire. Texture rather thin, both surfaces naked; costae and veins \pm rusty above, less so or rather naked beneath; veins forming 1—2 rows of areolae near the midribs, the rest free, no included free veinlets. Sori in 1 row on each side of the costulae; indusium subcordate, reniform. — A form found in Sumatra by Dr. Burck has the rhizome erect, densely clothed with long, linear-subulate, fuscous scales, the stipes and rachises sparingly fibrilloso-paleaceous, the ultimate lobes blunt to subacute, entire to broadly toothed.

? *Philippines*, *Sumatra*; *Ceylon*, *South. India*.

* (8) **P. devexa**, *v. A. v. R.*; *P. membranacea*, *Bedd.*, *Handb. Ind. Ferns*, 225; *Aspidium membranaceum*, *Hk.*, *Hk. Bk.*, *Syn. Fil.*, 259; *A. devexum*, *Kze.*, *Bot. Zeit.*, 1848, 259; *C. Chr.*, *Ind. Fil.*; *Tectaria devexa*, *Copel.*, *Philipp. Journ.*, II, 415; — var. *minor*; *A. giganteum*, *Bl.*, var. *minor*, *Hk.*, *Spec. Fil.*, IV, 50; *Sagenia gigantea*, *Bedd.*, var. *minor*, *Bedd.*, *Ferns S.I.*, tab. CCXLIII.

Stipes tufted, 20—30 c.M. long, nearly naked, with a few spreading, li-

near scales below. Fronds 55—60 c.M. long, deltoid, acuminate, pinnatifid at the apex. Lateral pinnae \pm 4—7 on each side, the lowest much the largest, subdeltoid, long-stalked, pinnate, with the lower pinnulae on the lower side prolonged and again pinnate at the base, the tertiary segments being stalked and deeply pinnatifid; higher pinnae gradually less compound. Ultimate lobes entire or crenulate to toothed. Texture thin, herbaceous or membranaceous; upper surface somewhat hairy towards the margins, lower slightly puberulous; costae and veins rusty or puberulous; veins anastomosing principally in costal arches, but not rarely also as in *Aspidium*. Sori copious, commonly submarginal and terminal on free veinlets; indusium orbicular or reniform.

Var. minor: A small and less divided form, divided \pm as in *P. Trimeni*, *Bedd.*, or with the tertiary leaflets not deeply pinnatifid. — A plant gathered in Borneo and cultivated in the Buitenzorg gardens, which may be a form of this, has the fronds inconspicuously glanduloso-pilose beneath and the costae and veins \pm naked.

Malaya; Formosa, South. China, Ceylon.

(9) **P. profereoides**, *v. A. v. B.*; *Aspidium profereoides*, *Christ*, *Philipp. Journ.*, II, 158; *Tectaria ambigua*, *Copel.*, l. c. 415, p.p.

Stipes \pm 40—75 c.M. long, castaneous, sulcate and sparingly scaly and bristly like the rachis, the scales brown, subulate. Fronds dimorphous, the fertile ones on the longest stipes. Barren fronds more than 50 c.M. long, 52 c.M. broad, deltoid-oblong, pinnate. Pinnae distant, horizontally spreading, \pm 15 on each side below the acuminate, inciso-lobate apex, sessile, broadly lanceolate, \pm 18 c.M. long, \pm 3 $\frac{1}{2}$ c.M. broad, acute, cut $\frac{1}{2}$ -way down or more to the costa, the lowest not reduced, not forked. Lobes \pm 18 on each side, subentire, ovate-obtuse, \pm 1 $\frac{1}{4}$ c.M. long, \pm 1 c.M. broad, separated by narrow, obtuse sinuses, the lowest a little larger. Texture flaccido-herbaceous; colour dark-green, opaque; costae furfuraceo-puberulous above; main veins not prominent, pinnate in the lobes; veins forming 1 row of narrow costal areolae, 1—5 rows of areolae near the costulae and 3—4 areolae between the costal ones and the sinuses. Fertile fronds contracted, \pm 50 c.M. long, the pinnae distant, \pm 10 c.M. long, \pm 2 $\frac{1}{2}$ c.M. broad, deeply incised, the lobes linear-lanceolate, falcate, \pm 4 m.M. broad. Sori round; indusium flaccid, soon corrugate, probably reniform.

Mindanao.

22. DRYOPTERIS, *Adanson.*

§ 1. LASTRAEA.

* (9) **D. Teuscheri**, *v. A. v. R.*, — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 6.

(10) **D. pectinata**, *v. A. v. R.* — Insert as a variety: — var. *Webbiana*: *Nephrodium Webbianum*, *Hk.*, Spec. Fil., IV, 85; *D. orientalis*, *C. Chr.*, var. *Webbiana*, Christ, Philipp. Journ., II, 207.

Var. Webbiana: Differs from the type in having the fronds quite naked, without calcareous dots on the upper surface, and the lobes more horizontal and more angular. — *Mindanao, Amboyna, New Guinea; Fiji.*

(11) **D. ligulata**, *O. Ktze.* — Insert as synonyms: *D. immersa*, *O. Ktze*, var. *ligulata*, Christ, Philipp. Journ., II, 208; *Lastrea ligulata*, *J. Sm.*, (oldest name).

* (15) **D. attenuata**, *O. Ktze.* — Insert as a synonym: *Lastrea attenuata*, *J. Sm.*, (oldest name).

* (18) **D. viscosa**, *O. Ktze*, Christ, Philipp. Journ., II, 211.

Detailed description. — Rhizome thick, erect, blackish, provided with many old stipes. Stipes densely tufted, 12—20 c.M. long, firm, erect, densely tomentose like the rachis with ochraceous-grey hairs and clothed with ovate, black-brown scales. Fronds 20—55 c.M. long, 8—15 c.M. broad, oblong, acuminate. Pinnae spreading, 20—25 on each side, 1—1½ c.M. broad, short-acuminate, sessile, cut down nearly to the costa into oblong, obtuse, subcrenate segments separated by acute sinuses; lower pinnae rather distant, somewhat shorter, deflexed. Texture somewhat fleshy; costae, costulae and especially the under surface densely pubescent with white, rigid hairs (glands?); veins \pm 6 on each side, simple. Sori medial; indusium persistent, brown, coriaceous.

* (25) **D. Peekeli**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 7.

* (24b) **D. diversifolia**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 7.

(25) **D. patens**, **O. Ktze.** — Insert as a synonym; *Polypodium patens*, Sw., (oldest name).

(26) **D. exigua**, **O. Ktze.** — Insert as a synonym: *Lastraea exigua*, J. Sm., (oldest name).

(27a) **D. Foxii**, **Christ**, Philipp. Journ., II, 208: *Nephrodium Foxii*, Copel., MS.

Rhizome short, creeping or ascending. Stipes close, 5—10 c.M. long, stramineous, naked. Fronds lanceolate, 20—30 c.M. long, 6—9 c.M. broad, narrowed at both ends. Pinnæ lanceolate, acuminate, ascending, the lowest remote, minute. Lower pinnulae \pm 8 m.M. long, 1—1½ m.M. broad, acute, obscurely toothed, adnate, the others confluent, the higher tapering gradually into a subserrate tail. Texture membranaceous; upper surface naked, lower sparingly pubescent, the hairs whitish, short; veins \pm 5 on each side. Sori medial; indusium naked, subpersistent. — Fronds sometimes reaching a size 4 \times as large as indicated above.

Luzon, Mindanao.

* (54) **D. filix mas**, **Schott.** — Insert as a synonym: *Polypodium Filix mas*, L., (oldest name). — Insert as a variety: — var. *marginata*; *Nephrodium Filix-Mas*, Rich., var. *marginata*, Clarke, Trans. Linn. Soc., Bot., 2nd Series, I, 521, tab. LXXI; *Aspidium marginatum*, Wall., List, No. 591 (not 366); ? *Lastraea Filix-Mas*, Pr., var. *elongata*, Bedd., Handb. Ind. Ferns, 250; ? *L. elongata*, Pr., Bedd., Ferns S.I. tab. CXII; ? *Nephrodium elongatum*, Hk. & Grev., Ic. Fil., tab. CCXXXIV; *D. marginata*, Christ, Philipp. Journ., II, 212.

Var. marginata: Fronds large, oblong or ovate-lanceolate, not narrowed at the base; lowest pinnæ falcate, often 2—3-pinnate; main and partial rachises nearly naked; tertiary segments oblong, obtuse, serrate or pinnatifid sometimes nearly to the costa. — *Luzon; Centr. China, North. India.*

Var. parallelogramma: — Also found in *Luzon*.

(54a) **D. heleopteroides**, **Christ**, Philipp. Journ., II, 212.

Rhizome short, thick. Stipes fasciculate, \pm 6—17 c.M. long, those of the barren fronds the shortest, brown at the base, rufo-stramineous, clothed with diaphanous, subulate, pale-brown scales, those of the fertile fronds the longest, fibrillose. Fronds dimorphous, the barren ones deltoid, \pm 16 c.M.

long, nearly as broad, pinnate, the apex lobed. Pinnae ± 10 , approximate, lowest largest, petiolulate, ± 8 c.M. long, $\pm 5\frac{1}{2}$ c.M. broad, oblong, obtuse, deeply pinnatifid, higher coarsely lobed. Lobes 8—10, ovate-rhomboidal, obtuse, lowest largest, ± 2 c.M. long, ± 1 c.M. broad, coarsely and irregularly crenato-serrulate. Texture subcoriaceous; colour pale-green, opaque; rachis and surfaces naked; veins pinnate and forked in the lobes. Fertile fronds ± 10 c.M. long, ± 7 c.M. broad, pinnate; pinnae ± 8 on each side, distant, the segments rhomboidal, obtuse, equal, subentire, ± 1 c.M. long, $\pm \frac{1}{2}$ c.M. broad. Sori submarginal, close, ± 4 on each side in the lobes; indusium pale, bullate, subpersistent.

Luzon.

* (55) **D. syrmatia**, *O. Ktze.* — Insert as a variety: var. *petiolosa*, Christ, Philipp. Journ., II, 210.

Var. petiolosa: Pinnae long-stalked, ± 16 c.M. long, $\pm 5\frac{1}{2}$ c.M. broad, on stalks 1 c.M. or more long. Sori medial, minute; indusium grey. — *Luzon, Mindanao.*

(55a) **D. Metteniana**, *Hier.*, Christ, Philipp. Journ., II, 210.

Near *D. syrmatia*, *O. Ktze.* but the pinnae cut to $\frac{1}{2}$ -way down to the costa, to 4 c.M. broad or broader, the lowest pinnae 2-pinnatifid, conspicuously produced on the lower side, with the lowest posterior segment deflexed and deeply lobed. Lobes serrato-crenate, subacute. Texture thinly membranaceous; veins 2—5-furcate. Sori medial, minute; indusium rudimentary.

Luzon, Mindanao, Celebes.

(56) **D. erythrosora**, *O. Ktze.* — Insert as a synonym: *Aspidium erythrosorum*, *Eat.*, (oldest name).

* (40) **D. flaccida**, *O. Ktze.* — Insert as a synonym: *Lastraea flaccida*, *Moore*, *Bedd.*, *Ferns S.I.*, tab. CCL.

* (44) **D. sparsa**, *O. Ktze.* — Insert as a synonym: *Polypodium sparsum*, *Ham.*, (oldest name).

(44a) **D. rizalensis**, *Christ*, *Bull. Herb. Boiss.*, 2^e Série, VI, 1001; ? *Philipp. Journ.*, II, 216.

Rhizome short. Stipes tufted, to ± 40 c.M. long, slender, stramineous,

rigid, clothed with short, grey pubescence and long, black hairs. Fronds elongate-deltoid, ± 30 c.M. long, ± 20 c.M. broad at the base, 2-pinnatifid. Terminal pinna incised, the lobes obtuse, subentire. Lateral pinnae $\pm 7-8$ on each side, oblong, bluntish, lobed nearly to the costa at the base, the lowest ones elongate-deltoid, with the lowest lobes on the lower side produced and deeply lobed. Texture herbaceous, flaccid; under surface paler than the upper; costae and costulae distinct, black, the latter pinnate and forked in the lobes, flexuose. Sori several on each lobe, terminal on the veins; indusium grey, flaccid, fugacious. — Near *D. sparsa*, *O. Ktze.*

Luzon, Mindanao.

* (45) **D. Raciborskii**, *v. A. v. R.* — This species has a distinct sagenioid habit, with the veins sometimes pleocnemioid, forming here and there a single costal or costular areola.

* (46) **D. recedens**, *O. Ktze.* — Insert as a synonym: *Polypodium recedens*, *J. Sm.*, (oldest name).

* (51) **D. intermedia**, *O. Ktze.* — Insert as varieties: — var. *Mannii*, *Christ*, *Philipp. Journ.*, II, 215; *Nephrodium Mannii*, *Hope*, *Journ. of Bot.*, XXVIII, 145; — var. *microloba*, *Christ*, l. c.

Var. Mannii: Fronds resembling *D. filix mas*, *Schott*, much elongated, oblong, 2-pinnatifid; pinnae regularly lobed, lobes simple, only the lowest pinnae 2-pinnatifid, produced on the posterior side.

Var. microloba: Stipe and rachis purpureous, densely clothed with red-black, spreading hairs; lower pinnae $1\frac{1}{2}-2$ c.M. long, $\frac{1}{2}$ c.M. broad; lobes close, ± 6 on each side, 2—4 m.M. broad.

Mindanao.

(56a) **D. balabacensis**, *Christ*, *Philipp. Journ.*, II, 215.

Stipes ± 55 c.M. long, angular, naked like the whole plant, rufo-stramineous or castaneous, as are rachises. Fronds 5-pinnate, deltoid, ± 50 c.M. long, as broad or broader, 5-partite at the base. Pinnae $\pm 8-10$ below the pinnatifid apex, distant, the lowest ± 14 c.M. apart, ± 50 c.M. or more long, ± 25 c.M. broad at the base, deltoid, on stalks ± 5 c.M. long, the lowest posterior pinnulae $\pm 15\frac{1}{2}$ c.M. long, ± 7 c.M. broad, the tertiary segments ovate-oblong, 4—5 c.M. long, ± 2 c.M. broad, obtusely inciso-lobate; higher pinnae sessile, oblong, acuminate, cut down to the costa in the

lowest part, lobed towards the apex, the lobes decurrent, oblong, subacute, \pm 8 m.M. broad, with the sinuses acute, toothed, the teeth decumbent. Texture coriaceous; colour yellowish-green, rather glossy; veins pinnate and 2—3-furcate in the lobes. Sori numerous, minute, medial or submarginal; indusium black, carnosu-coriaceous, soon convolute, persistent.

Balabac, Palawan, Sibuyan.

(58a) **D. dubia**, *Copel.*, Leafl., I, 255.

Rhizome ascending, clothed with lanceolate, castaneous scales. Stipes approximate, 10—20 c.M. long, red-castaneous, clothed towards the frond (like the rachis) with minute hairs and small, distant scales. Fronds lanceolate-deltoid, to 50 c.M. long, to 20 c.M. broad, acuminate, 4-pinnatifid. Pinnae opposite or the higher ones alternate, the lower deltoid, to 12 c.M. long, to 10 c.M. broad. Ultimate divisions lanceolate, cut down nearly to the costa into obtuse, broadly oblanceolate segments. Texture membranaceous; surfaces nearly naked; veins simple. Sori terminal; indusium athyroid, sometimes nephrodioid with a broad or nearly closed sinus, rarely asplenioid. — Judging by the sori this fern would be an Athyrium, but the frond's form and the colour and texture of the indusium mark it as a relative of *Dryopteris rhodolepis*, *C. Chr.* (Copel.).

Luzon.

* (59) **D. setigera**, *O. Ktze.* — Insert as synonyms: *Lastraea flaccida*, *Bedd.*, Ferns S.I., tab. IC; *Cheilanthes setigera*, *Bl.*, (oldest name). — My var. *pallida* seems to be the common form or even the type itself. Omit its synonyms, which are synonyms of *Phegopteris ornata*, *Fée.*

* (61a) **D. Backeri**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XVIII, 8; — var. *aspera*, *v. A. v. R.*, l. c., XXI, 5; *Polypodium asperum*, *Zipp.*, MS. in Herb. Bog.

Stipes and rachises pale-brown, the upper side pubescent with a line of fine, not very short, pale-coloured hairs, the rest provided throughout with numerous squarrose, deciduous, narrow, linear-subulate, pale-brown, finely and deciduously ciliate scales leaving the stipes and rachises rough by their persistent bases. Fronds like those of *D. setigera*, *O. Ktze.*, but the primary and secondary segments looser. Texture firm-herbaceous when dry; tertiary rachises narrowly winged, pubescent above and scaly beneath, as are the costae; upper surface nearly naked, lower hairy on the costulae and veins with whitish

hairs; costulae (main veins) pinnate in the ultimate lobes. Sori numerous, minute, in rows near the costae (central ribs of the tertiary segments), 1—2 on each lobe; indusium minute, membranaceous, whitish, soon hidden among the capsules, visible by $\pm 50 \times$ magnifying power. — Mr. C. A. Backer found this plant in the new vegetation of the small volcanic island Krakatau, which was totally destroyed by an eruption in 1885. It may be a hybrid between *D. setigera*, *O. Ktze* and *multisetia*, *C. Chr.* or *Phegopteris ornata*, *Fée*. It differs: *a*, from *D. setigera* in having the stipes and rachises provided with deciduous, subulate scales with persistent, rough bases; *b*, from *D. multisetia* in having a line of hairs on the upper side of stipes and rachises; *c*, from *Phegopteris ornata* in having the fronds much smaller, with the rachises not winged and the sori minutely indusiate.

Var. aspera: Differs from Mr. Backer's plant in having the rachises not beset with deciduous, flat, subulate scales, but with \pm persistent, subulate, subterete bristles of the same colour, the pinnulae horizontally spreading, the tertiary segments rather small, with the edges much recurved when dry. Indusium somewhat larger, visible by $\pm 15 \times$ magnifying power, but also hidden among the capsulus. Rhizome creeping. — Perhaps a distinct species.

Krakatau; — var. *aspera*: *Java* (*Preanger Regencies*, Zippelius N^o. 268).

§ 2. NEPHRODIUM.

* (65) **D. pteroides**, *O. Ktze*. — Insert as a synonym: *Polypodium pteroides*, *Retz*, (oldest name).

(66a) **D. mindanaensis**, *Christ*, *Philipp. Journ.*, II., 194.

Rhizome creeping or very oblique. Stipes distant, ± 30 c.M. long, plumbeo-rufous, short-puberulous like the whole plant, the base clothed with brown, opaque, ovate, acuminate scales. Fronds ± 70 c.M. long, ± 40 c.M. broad, ovate-acuminate, caudate, not or hardly narrowed at the base. Pinnae ± 28 on each side below the incised apex, spreading, tolerably approximate, sessile, not narrowed at the base, $\pm 2\frac{1}{2}$ c.M. broad, caudate-acuminate, the edge cut to $\frac{1}{2}$ -way down to the costa. Lobes ± 45 —50 on each side, close, pectinate, subfalcate, obtuse but mucronulate, nearly without sinuses between them, $\pm \frac{3}{4}$ c.M. long, $\pm \frac{1}{2}$ c.M. broad. Texture herbaceous; colour bright-green, opaque; veins stramineous and densely white-puberulous, as are the costae, ± 10 —12 on each side, simple, 1—2 lower anastomosing. Sori 5—8 on each side; indusium flaccid, grey, ciliate.

Mindanao.

(69) **D. procurrens**, *O. Ktze.* — Also found in *Luzon* and *Mindanao*

* (72) **D. invisiva**, *O. Ktze.* — Insert as a synonym: *Polypodium invisum*, *Forst.*, (oldest name).

* (74) **D. cucullata**, *Christ*, — Insert as a variety: — var. *mucronata*, *Christ*, *Philipp. Journ.*, II, 195; *Nephrodium mucronatum*, *J.Sm.*, *Journ. of Bot.*, III, 412.

Var. mucronata. Pinnae 2 c.M. or more broad, the lower ones narrowed towards the base, the points of insertion of the pinnae callous; veins ± 12 on each side of the main veins, 6 lower anastomosing. — Intermediate between the type and *D. callosa*, *C. Chr.* — *Luzon*, *Mindanao*.

(78) **D. hirsuta**, *O. Ktze.*; *Nephrodium hirsutum*, *J. Sm.* — Insert: *Bedd.*, *Ferns Br.I.*, tab. XLVI.

* (81) **D. glandulosa**, *O. Ktze.*; *Nephrodium glandulosum*, *Moore.* — Insert: *Bedd.*, *Ferns Br.I.*, I, tab. CXXXII.

* (85) **D. urophylla**, *C. Chr.* — Insert as a variety: var. *pustulosa*, *Christ*, *Philipp. Journ.*, II, 205.

Var. pustulosa: Surfaces hispid, rough-pustulous. — *Luzon*.

* (91) **D. callosa**, *C. Chr.* — Insert as a synonym: *Aspidium callosum*, *Bl.*, (oldest name).

(95) **D. brachyodus**, *O. Ktze.*; *Nephrodium brachyodon*, *Hk.* — Insert: *Bedd.*, *Ferns S.Br.I.*, *Suppl.*, tab. CCCLXXIX: *Polypodium brachyodus*, *Kze.*, (oldest name).

Veins all free or the lowest anastomosing and with sometimes an excurrent veinlet; a single vein generally rises from the costa between each pair of lobes and runs to the margin.

* (94) **D. ferox**, *O. Ktze.* — Insert as a variety: — var. *calvescens*, *Christ*, *Philipp. Journ.*, II, 195.

Var. calvescens: Axillar setae and pustules nearly vanished. — *Mindanao*.

(95) **D. Loheriana**, *C. Chr.* — This species, with all the appearances of *Lastraea*, has the basal veins not always joined. Christ considers it now as a *Lastraea*.

* (97) **D. piloso-spuamata**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XXI, 4.

(99) *D. caudiculata*, *v. A. v. R.*, as construed on page 225, includes 2 distinct forms: *D. basilaris*, *C. Chr.* and *caudiculata*, *v. A. v. R.*

(99a) **D. basilaris**, *C. Chr.*, Ind. Fil.; Christ, in Philipp. Journ., II, 196; *Nephrodium basilaris*, *Pr.*, Epim., 258; *N. philippinense*, *Bk.*, Copel., Polypod. Philipp., 31.

Detailed description. — Rhizome oblique, suberect. Stipes \pm 4—5, subtufted, naked, rufo-stramineous, to \pm 45 c.M. long, provided on each side with 8—10 rudimentary, distant, obtuse auricles, the base clothed with long, lanceolate-subulate, hair-pointed, brown scales. Fronds ovate-acuminate, 60 c.M. long or longer. Pinnae alternate, erecto-patent, numerous, sessile or short-stalked, \pm 14 c.M. long or longer, $1-1\frac{1}{2}$ c.M. broad, cut to $\frac{1}{3}$ -way down to the costa, the base truncate on the upper, semicordate on the lower side, the apex caudate-acuminate; terminal pinna often very elongated. Lobes decumbent, \pm 3 m.M. long, subobtuse. Texture subcoriaceous; surfaces naked like the rachis; colour brownish-green; costae distinct, pale; veins \pm 5—8 on each side, only the lowest anastomosing. Sori minute, medial, \pm 5 on each side of the costulae; indusium grey, persistent. — *Philippines*.

(99b) **D. caudiculata**, *v. A. v. R.*; *D. microloncha*, *Christ*, Philipp. Journ., II, 202; *Nephrodium caudiculatum*, *J. Sm.* (not *Pr.*), Journ. of Bot., III, 411.

Near *D. basilaris*, *C. Chr.* Rhizome thick, oblique, black-brown. Stipes tufted, 2—3 c.M. long, rufo-stramineous, whether or not clothed with some scattered, brown scales. Fronds oblong, 16—24 c.M. long, 4—6 c.M. broad, caudate. Pinnae \pm 12 on each side, \pm 3 c.M. long, \pm $\frac{1}{2}$ c.M. broad, sessile, the base nearly hastate or auriculate on both sides, the anterior auricle the largest, the edge cut hardly $\frac{1}{2}$ -way down to the costa into truncate lobes; lower pinnae gradually reduced into mere, obtuse auricles 5—5 m.M. each way. Surfaces naked; rachis stramineous, sparingly furfuraceous; main veins sparingly pinnate in the lobes, the lowest veins anastomosing irregularly. Sori

numerous, in 3 irregular rows; indusium black-brown, orbicular, persistent. — *Philippines*.

(99c) **D. luzonica**, *Christ*, Philipp. Journ., II, 196; — var. *puberula* & *polyotis*, *Christ*, l. c., 197.

Like *D. basilaris*, *C. Chr.*, but distinguished by the following characters: Stipes slender, flavido-stramineous, naked, or minutely scaly at the base. Pinnae more distant, much elongated to an almost filiform apex, opposite, horizontal, dilated (substipulate) at the base on both sides; 1—2 lowest pairs of pinnae reduced to mere auricles. Lobes blunt, often truncate and denticulate. Texture thin, diaphanous; colour pale-emerald-green; veins often 5 on each side. Indusium thin, caducous.

Var. puberula: Rachis, costae and veins puberulous.

Var. polyotis: Pinnae \pm 16 m.M. broad at the base, short-acuminate; lower pinnae reduced to mere auricles; auricles \pm 20 on each side, reaching to the base of the stipe.

Luzon.

* (102) **D. parasitica**, *O. Ktze.* — Insert as a variety: — var. *falcatula*, *Christ*, Philipp. Journ., II, 197.

Var. falcatula: Lower pinnae opposite, refracted, stipulate at the base on the anterior side; lobes deeply incised, falcate. — *Mindanao*.

(111a) **D. todayensis**, *Christ*, Philipp. Journ., II, 195.

Rhizome suberect, black-brown, nearly naked. Stipes thick, firm, grey-brown, \pm 40 c.M. long, naked or sparingly covered with lanceolate, brown scales. Fronds 70 c.M. or more long, \pm 20 c.M. broad, oblong, acuminate, the rachis grey-brown, puberulous. Pinnae numerous, 40—55 on each side, close, \pm 11 c.M. long, \pm 11 m.M. broad, the apex caudato-acuminate, the base subcallous, cuneato-truncate, nearly unequal-sided, the margin cut to $\frac{1}{2}$ -way down to the costa, the lobes close, \pm 45 on each side, lanceolate, \pm 2 m.M. broad, acute, strongly falcate; lower pinnae rudimentary, suddenly reduced into 8—10 short auricles. Texture herbaceous, flaccid; colour black-green; upper surface puberulous, lower clothed with adpressed hairs on the costae and costulae; veins 7—8 on each side, the lowest anastomosing, the next following terminating in the acute sinuses. Sori minute, medial; indusium small, brown, puberulous,

Mindanao, Negros.

(111*b*) **D. Bordenii**, *Christ*, Philipp. Journ., II, 204.

Rhizome oblique, thick, sparingly clothed with minute, crisped, lanceolate, pale-brown scales. Stipes subfasciculate, firm, to 30 c.M. long, often shorter, yellowish or brownish, puberulous. Fronds ovate, 25—35 c.M. long above the auricles, 15—20 c.M. broad, pinnate, the apex pinnatifid, acuminate. Pinnae lanceolate, ± 15 on each side, horizontal, sessile, ± 10 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad, the base truncate, the edge cut to $\frac{1}{2}$ -way down to the costa, the apex acuminate; higher pinnae adnate, close, lower not-reduced ones sharply deflexed, cuneate at the base; lower pinnae suddenly reduced into ± 10 small, triangular, subacute auricles or minute lobes or glands occupying the stipe to the base. Secondary segments oblique, subfalcate, close, pectinate, ± 3 m.M. long, $\pm 2\frac{1}{2}$ m.M. broad, entire, obtuse. Texture herbaceous; colour dark-green; rachis and surfaces shortly pubescent; veins not very distinct, ± 7 on each side, only the lowest anastomosing. Sori medial, small; indusium small, dark-grey, soon evanished.

Luzon, Palawan.

* (112) **D. truncata**, *O. Ktze.* — Insert as a synonym: *Polypodium truncatum*, *Poir.*, (oldest name).

23. MESOCHLAENA, *R. Brown.*

* (1) **M. polycarpa**, *Bedd.* — Insert as a synonym: *Dryopteris polycarpa*, *Christ*, Philipp. Journ., II, 202.

* (2) **M. larutensis**, *v. A. v. R.*, var. *borneensis*. — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 8.

25. ASPIDIUM, *Swartz.*

§ 1. TECTARIA.

(4) **A. calcareum**, *Pr.* — Insert as a synonym: *Sagenia calcarea*, *J. Sm.*, (oldest name).

* (5) **A. angulatum**, *J. Sm.* — Insert as a synonym: *Polypodium angulatum*, *Willd.*, (oldest name).

* (8) **A. ternatense**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 9; *Tectaria ternatensis*, *v. A. v. R.*, l. c.

§ 2. SAGENIA.

* (12) **A. vastum**, *Bl.* — Insert as a synonym: — *Nephrodium vastum*, *Bk.*, *Hk. Bk.*, *Syn. Fil.*, 296.

* (15) **A. subcaudatum**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 9; *Tectaria subcaudata*, *v. A. v. R.*, l. c.

* (20) **A. semibipinnatum**, *Wall.* — Insert as a synonym: *Sagenia semibipinnata*, *Moore*, *Bedd.*, *Ferns Br.I.*, I, tab. CXXXVII.

* (25) **A. subtriphylum**, *Hk.* — Insert as a synonym: *Polypodium subtriphylum*, *Hk. & Arn.*, (oldest name).

* (31) **A. variolosum**, *Wall.* — Insert: *Bedd.*, *Ferns S.Br.I.*, *Suppl.*, tab. CCCLXV.

* (33) **A. latifolium**, *J. Sm.* — Insert as a synonym: *Polypodium latifolium*, *Forst.*, (oldest name).

* (59) **A. siifolium**, *Mett.* — Insert as a synonym: *Polypodium siifolium*, *Willd.*, (oldest name).

28. ODONTOSORIA, *Presl.*

§ 1. EU-ODONTOSORIA.

* (4) **O. lindsayae**, *v. A. v. R.*, Bull. Dép. Agr. I. N., 1908, XXI, 4.

Stipes and rachises pale-brownish-stramineous, naked. Fronds 4—5-pinnate. Ultimate leaflets cuneate, sessile, simple or 2—5-fid, $\frac{3}{4}$ — $1\frac{1}{4}$ c.M. long, 2—5 m.M. broad at the apex, the incised ones the broadest, the apices rather horizontally or \pm obliquely truncate. Texture subcoriaceous or firm-herbaceous; surfaces naked; veins flabellate, dichotomous. Sori narrow, terminal on the ultimate segments (leaflets or lobes), occupying the whole breadth, uniting the apices of 2 or more veinlets. — In habit this resembles *O. chinensis*, *J. Sm.*, but in the Lindsaya-like sori it approximates *O. retusa*, *J. Sm.*
New Guinea.

29. LINDSAYA, *Dryander.*

§ 1. ODONTOLOMA.

* (1) **L. repens**, *Bedd.* — Insert as synonyms: *Davallia Boryana*, *Pr.*, *Hk. & Grev.*, *lc. Fil.*, tab. CXLIII; *Dicksonia repens*, *Bory*, (oldest name).

§ 2. EULINDSAYA.

* (2) **L. gracilis**, *Bl.* — Insert as a synonym: *L. Lobbiana*, *Hk.*, *Bedd.*, *Ferns Br.I.*, I, tab. XXVIII.

* (18a) **L. cultrata**, *Sw.* — Insert: *Hk. & Grev.*, *lc. Fil.*, tab. CXLIV; *Adiantum cultratum*, *Willd.*, (oldest name).

* (20) **L. scandens**, *Hk.* — Insert: *Bedd.*, *Ferns Br.I.*, II, tab. CCXCVIII.

30. SCHIZOLOMA, *Gaudichaud.*

§ 2. EUSCHIZOLOMA.

(9) **S. Guerinianum**, *Gaud.* — Insert: *Bedd.*, *Ferns Br.I.*, II, tab. CCCXL.

* (11) **S. coriaceum**, *v. A. v. R.* — Insert: *Bull. Dép. Agr. I. N.*, 1908, XVIII, 10.

32. HUMATA, *Cavanilles.*

§ 1. EUHUMATA.

* (5) **H. Gaimardiana**, *J. Sm.* — Insert as a synonym: *Nephrodium Gaimardianum*, *Gaud.*, *Freyc.*, *Voy. Ur. Phys.*, *Bot.*, tab. XII, fig. 1.

* (5) **H. parvula**, *Wall.* — Insert as a synonym: *Acrophorus parvulus*, *Moore*, *Bedd.*, *Ferns Br.I.*, I, tab. XCVII.

34. *DAVALLIA*, *Smith.*

§ 2. COLPOSORIA.

* (5) *D. Griffithiana*, *Hk.* — Insert: *Bedd.*, *Ferns Br.I.*, I, tab. CVI.

(17) *D. trichomanoides*, *Bl.* — Insert. *Bedd.*, *Ferns Br.I.*, II, tab. CLXXVIII.

35. *MICROLEPIA*, *Presl.*

(2) *M. marginalis*, *Bedd.* — Insert as synonyms: *M. urophylla*, *Moore*, *Bedd.*, *Ferns Br.I.*, I, tab. CIII; *Polypodium marginale*, *Thb.*, (oldest name).

* (6) *M. speluncae*, *Moore.* — Insert as a synonym: *Polypodium Speluncae*, *L.*, (oldest name).

(6a) *M. sablanensis*, *Christ*, *Philipp. Journ.*, II, 168.

Fronde large, 5-pinnate, the rachis opaque, fulvo-stramineous, shortly and finely puberulous. Pinnae 35 c.M. and more long, \pm 9 c.M. broad, hardly narrowed at the base, short-stalked, elongate-caudate. Pinnulae approximate, pectinate, numerous (\pm 40 or more pair), horizontal, nearly sessile, acuminate, produced at the base on the upper side. Tertiary segments \pm 20 on each side, close or approximate, separated by narrow sinuses, unequal-sided, the lowest free, addressed to the rachis; upper base truncate, lower cuneate; upper side crenate, lower subentire; lobules \pm 3, minute. Texture herbaceous; colour dark-green; upper surface smooth, lower pilose on the costae and veins; veins distinct, 2—3-furcate in the lobules. Sori minute, 2—3 on the anterior side, in the sinuses of the lobules; indusium inconspicuous, thin. — Very near *M. speluncae*, *Moore.*

Luzon.

* (10) *M. strigosa*, *Pr.* — Insert as a synonym. *Trichomanes strigosum*, *Thb.*, (oldest name).

37. ADIANTUM, *Linné*.

§ 1. EU-ADIANTUM.

* (13) **A. Mettenii**, *Kuhn*. — Also found in *Timor* (Forbes, No. 3755).

* (15a) **A. aculeolatum**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 10.

* (21) **A. suborbiculare**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 11.

38. HYPOLEPIS.

* (3) **H. tenuifolia**, *Bernh.* — Insert as a synonym: *Lonchitis tenuifolia*, *Forst.*, (oldest name).

40. PLAGIOGYRIA, *Mettinius*.

(1a) **P. falcata**, *Copel.*, Philipp. Journ., II, 135, tab. I, B.

Rhizome erect, thick. Stipes tufted, naked, without basal tubercles, 7—24 c.M. long, the fertile fronds on the longest stipes. Barren fronds \pm 20 c.M. long, 6—7 c.M. broad, short-acuminate, cut down nearly or quite to the costa. Pinnae approximate, \pm 5 m.M. broad, acute, straight or subfalcate, sharply serrate, or entire towards the not or hardly dilated, adnate base; lowest pinnae much deflexed, somewhat abbreviate. Texture submembranaceous; surfaces naked; colour black-green; veins conspicuous, mostly simple. Fertile fronds 15—24 c.M. long, $2\frac{1}{2}$ —4 c.M. broad, acuminate, pinnate; pinnae rather distant, falcato-arcuate, 2—3 c.M. long, \pm 2 m.M. broad.

Mindoro.

(3) **P. Christii**, *Copel.* — Also in *Mindoro*.

(7) **P. tuberculata**, *Copel.* — Insert as a variety:— var. *latipinna*, *Copel.*, Philipp. Journ., II, 135.

Var. latipinna: Fertile pinnae broader, 5—4 m.M. broad. — *Mindoro*.

41. **PELLAEA**, *Link.*§ 2. **PTERIDELLA.**

* (4) **P. timorensis**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 11.

45a. **PTERIS**, *Linné.*§ 1. **EUPTERIS.**

* (5) **Pt. orientalis**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 12.

* (6) **Pt. Treubii**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XVIII, 12.

(11a) **Pt. intromissa**, *Christ*, Philipp. Journ., II, 173.

Rhizome short. Stipes fasciculate, slender, rufo- or fusco-stramineous, opaque. Fronds \pm 30 c.M. long, as broad, pinnate, the apex ternate. Pinnæ opposite, medial simple, decurrent, longest \pm 20 c.M. long, linear-lanceolate or linear and caudate, lowest 2—5-partite, decurrent, edges acutely serrate when barren; barren pinnæ or segments \pm 1 $\frac{1}{4}$ c.M. broad, fertile \pm 6 m.M. broad. Texture thinly papyraceous; colour dark-green, opaque, subglaucous; veins slender, not very distinct, oblique, simple or forked. Sori and indusia narrow. — Very near *Pt. cretica*, *L.*

Luzon, Mindanao; China.

* (20) **Pt. Dalhousiae**, *Hk.* — Insert: Bedd., Ferns Br.I., II, tab. CXCI.

* (22a) **Pt. Grevilleana**, *Wall.*, var. *ornata*. — Insert: v. A. v. R., Bull. Dép. Agr. I. N., 1908, XXI, 5.

* (23) **Pt. longipinnula**, *Wall.*, var. *sumatrana*. — Insert: v. A. v. R., Bull. Dép. Agr. I. N., 1908, XXI, 5.

48. LOMARIA, *Willdenow.*

* (1) **L. Patersoni**, *Spr.*, var. *elongata*; *L. elongata*, *Bl.* — Insert: *Bedd., Ferns S.I., tab. XXVIII—XXVIII, A.*

(4) *L. areolaris*, *Harr.* — Omit this, it being referred to *Stenochlaena* further on.

53. DIPLAZIUM, *Swartz.*

§ 1. EUDIPLAZIUM.

* (2b) **D. zeylanicum**, *Moore.* — Insert: *Bedd. Ferns S.I., tab. CCXXVIII.*

* (7) **D. silvaticum**, *Sw.* — A form found in Java (*Depok, v. A. v. R.*) has the sori on the pinnae small and irregular.

(50) **D. Prescottianum**, *Moore.* — Insert: *Bedd., Ferns Br.I., II, tab. CCXLIII; D. silvaticum, Sw., var. Prescottianum, Bedd., Handb. Ind. Ferns, 178.*

* (42a) **D. umbrosum**, *Bedd., Handb. Ind. Ferns, 188; Asplenium umbrosum, J. Sm., Rac., Flor. Btz. I, 228; Athyrium umbrosum, Pr., Christ, Farnkr. d. Erde, 225; C. Chr., Ind. Fil.; A. australe, Bedd., Ferns S.I., tab. CLVIII; Polypodium umbrosum, Ait., (oldest name).*

Caudex erect, to 25 c.M. high, densely covered with the bases of fallen stipes, roots and brown, lanceolate scales. Stipes 30—80 c.M. long, firm, scaly at the base. Fronds often 90—160 c.M. long, 60—120 c.M. broad, ovate-lanceolate to deltoid. Pinnae ovate- to deltoid-lanceolate, 7—20 c.M. broad. Pinnulae close, pinnatifid at the apex, cut down nearly or quite to the costa into unequal-sided, elliptico-rhomboidal segments 5—15 m.M. long, 2—5 m.M. broad. Segments sessile, the apex rounded, the edge toothed to deeply inciso-crenate, the base narrowed or broad and decurrent. Texture herbaceous; surfaces naked; central ribs of the segments pinnate; veins simple or forked. Sori copious, oblong, the lower often double; indusium inflated, entire. — Variable.

Trop. Asia & Australia, New Zealand, Africa.

* (45c) **D. brevisorum**, *J. Sm.*, Journ. of Bot., III, 408, *Asplenium basilare*, *Moore*, Rac., Flor. Btz., I, 229; *A. woodwardioides*, *Bk.*, Hk. Bk., Syn. Fil., 229; *Athyrium woodwardioides*, *Christ.*, Fil. Saras., III, 253; *A. silvaticum*, *Milde*, Bot. Zeit., 1866, 576; *C. Chr.*, Ind. Fil.; *Allantodia sylvatica*, *Bl.*, Enum., 175.

Rhizome erect. Stipes firm, erect, naked, dark-brown, to \pm 70 c.M. or more long. Fronds lanceolate-triangular, 60—100 c.M. long, 50—60 c.M. broad. Lower pinnae largest, to 50 c.M. long, $7\frac{1}{2}$ —10 c.M. broad, pinnate. Pinnulae subpetiolate, 1—2 c.M. broad, narrowed towards the apex, cut \pm $\frac{2}{3}$ -way down or more to the costa into \pm close, linear-oblong, blunt, toothed, subfalcate segments to \pm 1 c.M. long, to \pm $\frac{1}{2}$ c.M. broad. Texture herbaceous; rachis and surfaces naked; main veins pinnate in the lobes; veins simple or forked, \pm 4—5 on a side. Sori short-oblong, close to the costulae, the lower ones often double, close to the costae, not reaching more than $\frac{1}{2}$ -way the edge; indusium ciliate or eroso-fimbriate. — Resembles in habit *D. polypodioides*, *Bl.* The javan form has the dimensions not rarely $2 \times$ as large.

Java, *Cetebes*, *Philippines*.

(45c) **D. muricatum**, *v. A. v. R.*; *Asplenium muricatum*, *Mett.*, Ann. Mus. Bot. L. B., II, 259; *Athyrium umbrosum*, *Pr. var.*, *C. Chr.*, Ind. Fil.

Stipes nodulous or aculeate. Fronds \pm 60 c.M. long, deltoid, \pm 5-pinnatisect. Lowest pinnae distinctly stalked, \pm 20 c.M. long, oblong, acuminate. Pinnulae to \pm 5 c.M. long, $\frac{3}{4}$ c.M. broad, short-stalked, oblong, narrowed towards the blunt apex, the base truncate, not rarely oblique. Lobes oblong, obtuse; upper edge slightly toothed, lower subpinnatifid. Texture firm-membranaceous; colour opaque-green; upper surface naked, lower sparingly scaly; veins forked. Sori near the costulae; indusium curved, pale, membranaceous, entire. — Near *D. umbrosum*, *Bedd.*

Java.

§ 2. ANISOGONIUM.

* (55) **D. proliferum**, *Thouars*; *Callipteris prolifera*, *Bory.* — Insert: *Bedd.*, Ferns Br.I, I, tab. LXX.

Pinnae often proliferous in the axils.

54. ATHYRIUM, Roth.

* (1b) **A. pusillum**, *v. A. v. R.* — Insert: Bull. Dép. Agr. I. N., 1908, XXI, 5, tab. I, a—c.

(15) **A. umbrosum**, *Pr.* — Omit this, it being referred to Diplazium.

(16) **A. silvaticum**, *Milde.* — Omit this, it being referred to Diplazium brevisorum, *J. Sm.*

55. ASPLENIUM, Linné.

§ 1. THAMNOPTERIS.

* (5c) **A. nidus**, *L.* — Insert as synonyms: *A. Phyllitidis*, *Don*, Prod. Flor. Nep., 7; *Thamnopteris Phyllitidis*, *Pr.*, Bedd., Ferns S.I., tab. CXXIII.

§ 3. EU-ASPLENIUM.

* (57) **A. macrophyllum**, *Sw.*, var. minus. — Insert: (? *A. integerrimum*, *Wall.*, Hk. & Grev., Ic. Fil., tab. CXXXVI).

* (69) **A. praemorsum**, *Sw.* — Insert as a synonym: *A. cuneatum*, *Hk. & Grev.*, Ic. Fil., tab. CLXXXIX.

60a. PHEGOPTERIS, Fée.

§ 5. MENISCIUM.

* (41) **Ph. triphylla**, *Mett.*; *Meniscium triphyllum*, *Sw.* — Insert: Hk. & Grev., Ic. Fil.; tab. CXX.

63. CEROPTERIS, Link.

* (1) **C. tartarea**, *Link.* — Insert as a synonym: *Neurogramme tartareum*, *Diels*, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, 264.

66. SYNGRAMMA, J. Smith.

§ 4. Hemigramma. — Omit this as a section, it being a distinct genus.

(10) *S. Zollingeri*, *Diels.* — Omit this, it is *Hemigramma Zollingeri*, *Christ.*

67. HEMIONITIS, Linné.

(1) *H. gymnopteroidea*, *Copel.* — Omit this, it is *Hemigramma Zollingeri*, *Christ.*

67a. HEMIGRAMMA, Christ

Sori linear, running along the veins, in \pm 2—3 longitudinal rows on each side of the costa, approximate or contiguous, united by transversal bars, at length not rarely confluent, sometimes acrostichoid. Paraphyses? No indusium.

Rhizome oblique or erect. Fronds dimorphous, not articulate to the rhizome, simple or irregularly incised or divided. Barren fronds sessile or stalked, clustered, if sessile or subsessile then not rarely in a dense rosette; venation as in *Aspidium*; main veins distinct; areolae irregular, with free veinlets. Fertile fronds long-stalked, contracted. (*)

Malaya.

* **H. Zollingeri**, *Christ*, *Philipp. Journ.*, II, 170; *H. latifolia*, *Copel.*, *Philipp. Journ.*, II, 406, p.p.; *Syngramma Zollingeri*, *Diels*, in *Engl. & Prantl. Nat. Pfl.Fam.*, I⁴, 257; *C. Chr.*, *Ind. Fil.*; *Hemionitis Zollingeri*, *Kurz*, *Rac.*, *Flor. Btz.*, I, 64; *Christ*, *Farnkr. d. Erde*, fig. 147; *H. gymnopteroidea*, *Copel.*, *Polypod. Philipp.*, 67; — var. *major*, *Christ*, *Philipp. Journ.*, l.c.

Rhizome erect or oblique, the scales dense, brown, often crisped, lanceolate or subulate, entire or variously fringed. Fronds dimorphous, simple or irregularly divided. Barren fronds sessile or stalked, clustered or placed in a

(*) Differs from **Syngramma** and **Hemionitis** by the aspidioid venation, the distinct main veins and the areolae provided with included free veinlets. Its natural place is near **Leptochilus**, from which genus it is hardly distinguishable in the forms with confluent sori. The sporangia are densely intermixed with paraphysiform corpuscles, probably the persistent stalks of fallen sporangia.

dense rosette, 7—25 c.M. long, 5—10 c.M. broad, oblanceolate, acute or obtuse, the edge entire or repand to irregularly undulate or incised, the lower part broad or contracted, the base cordate (var. *Teysmannii*), rounded or acute. Texture varying from membranaceous to subcoriaceous; surfaces naked; costa sometimes scaly like the rather short stipe; main veins distinct, not reaching the edge; areolae copious, irregular, with few or many free veinlets. Fertile fronds linear-lanceolate or ligulate, 5—15 c.M. long, 3—12 m.M. broad, narrowed gradually to the apex and a stipe 3—15 c.M. long, the edge subentire, irregularly repand or interrupted. — Exceedingly variable. A form gathered in Borneo by Dr. Hallier (no. 1822), determined by Dr. Christ has the sori decidedly acrostichoid.

Var. major: Stipes of barren fronds to 16 c.M. long, scaly, the scales subulato-setiform, black. Barren fronds varying from simple to pinnate; pinnae ± 5 on each side, connected by a narrow wing. Fertile fronds contracted; pinnae ± 2 on each side, ± 5 c.M. long, ± 5 m.M. broad. Texture herbaceous-membranaceous, nearly diaphanous. — *Leptochilus latifolius*, *C. Chr.* is distinguishable from this by its firmer texture, not diaphanous, its colour, black when dry, its often proliferous fronds.

Malaya; — var. major: *Luzon, Mindanao.*

71. POLYPODIUM, *Linné.*

§ 1. EUPOLYPODIUM.

(60) **P. cornigerum**, *Bk.* — Insert: *Bedd., Ferns S.Br.I., Suppl., tab. CCCLXXXI.*

§ 2. PROSAPTIA.

* (111) **P. contiguum**, *J. Sm.;* *Davallia contigua*, *Spr.* — Insert: *Hk. & Grev., Ic. Fil., tab. CXLI.*

§ 6. PLEOPELTIS.

* (198) **P. hastatum**, *Thb.* — Insert as a synonym: *Pleopeltis malacodon*, *Bedd., Ferns S.Br.I., Suppl., tab. CCCLXXXVII (not Handb. Ind. Ferns, 563).*

79. STENOCHLAENA, *J. Smith.*

§ 1. LOMARIOPSIS.

(4a) **S. areolaris**, *Copel.*, Philipp. Journ., II, 406; *Lomaria areolaris*, *Harr.*, Journ. of Linn. Soc., Bot., XVI, 28; *Blechnum areolare*, *Copel.*, Polypod. Philipp., 90.

Rhizome long, slender, naked, twisted. Stipes of barren fronds \pm 15 c.M. long, naked. Barren fronds ovate, 10—15 c.M. long, 7—10 c.M. broad, pinnate. Pinnae 7—9, lateral nearly sessile, terminal stalked, linear-lanceolate, 5—8 c.M. long, 1—2 c.M. broad, acute, finely serrate, truncate or rounded at the base. Texture coriaceous; surfaces naked; costae and veins slightly chaffy beneath; veins free, simple or forked, united at the base by a series of arches \pm close to the midribs, thus forming on each side a row of long, narrow costal areolae. Fertile fronds on stipes \pm 30 c.M. long, the pinnae narrowly linear, $5\frac{1}{2}$ —5 c.M. long, rather distant. Sori protected by the specialized, recurved, indusium-like margin when young. — Near *S. palustris*, *Bedd.* I saw a plant, probably a form of this, which had the pinnae with 2 glands at the base, 1 on the upper and 1 on the lower edge.

Philippines.

90. ANGIOPTERIS, *Hoffmann.*

(5) **A. evecta**, *Hoffm.* — This variable species is construed as including several forms (considered by de Vriese as distinct species) united by intermediates which are difficult of determination (*).

a. Recurrent veinlets present, reaching at least $\frac{1}{2}$ -way down to the costa.

b. Pinnae rather thick, chartaceous to coriaceous.

c. Largest pinnulae \pm $17\frac{1}{2}$ m.M. or more broad. (3a) *A. evecta*.

(3b) *A. Miqueliana*.

c.c. Largest pinnulae \pm 9— $17\frac{1}{2}$ m.M. broad.

d. Pinnulae pruinose beneath.

(3c) *A. pruinosa*.

d.d. Pinnulae not pruinose beneath.

(*) Determining these forms one should bear in mind that: *a*, the complete fronds are very large; *b*, the collectors commonly gather only some pinnae of a frond; *c*, the gathered pinnae often are those having not the largest pinnulae; *d*, the limit between opaque (impellucid) and pellucid veins or veinlets is not well-marked and it is often uncertain whether a transparent vein or veinlet should be regarded as an opaque or pellucid one; and *e*, the cuneate or rounded bases of the pinnulae are not rarely imperfectly cuneate or rounded, i. e. often \pm intermediate between both.

- e.* Veins beneath pale, opaque. (3d) *A. crassifolia*.
- e.e.* Veins beneath black or fuscous.
- f.* Pinnulae \pm cordate at the base on both sides.
(3e) *A. javanica*.
- f.f.* Pinnulae not cordate at the base; upper base \pm cuneate, lower \pm rounded. (3f) *A. Hartingiana*.
(3g) *A. angustifolia*.
- c.c.c.* Largest pinnulae under 9 m.M. broad. (3h) *A. angustata*.
(3i) *A. caudata*.
- h.b.* Texture rather thin, membranaceous to papyraceous.
- 1.* Pinnulae white-waxy beneath. (3j) *A. hypoleuca*.
- 1.1.* Pinnulae not white-waxy beneath.
- 2.* Largest pinnulae less than 14 c.M. long.
- 3.* Pinnulae \pm equal-sided at the base. (3k) *A. approximata*.
- 3.3.* Pinnulae distinctly unequal-sided at the base.
- 4.* Upper base narrower than the lower.
(3l) *A. aphanosorus*.
- 4.4.* Upper base broader than the lower.
(3m) *A. cuspidata*.
(3n) *A. Teysmanniana*.
- 2.2.* Largest pinnulae more than 14 c.M. long.
- \ddagger . Pinnulae at least 20 c.M. long. (3o) *A. Willinckii*.
- $\ddagger.\ddagger$. Pinnulae at best 20 c.M. long.
- $+$. Veins beneath pale or indistinct.
- $*$. Veins oblique. (3p) *A. Presliana*.
(3q) *A. pallescens*.
- $*.*$. Veins rather subhorizontal. (3r) *A. similis*.
(3s) *A. D'Urvilleana*.
- $+.+$. Veins beneath fuscous or dark-coloured, distinct.
(3t) *A. microsporangia*.
(3u) *A. ankolana*.
- a.a.* Recurrent veinlets wanting or very short, often hardly visible.
- b.* Texture thin, rather membranaceous or herbaceous.
- c.* Veins pellucid. (3v) *A. amboinensis*.
- c'.c'*. Veins not pellucid. (3w) *A. muricata*.
(3x) *A. crassipes*.
- b'.b'*. Texture firm, rather chartaceous or coriaceous. (3z) *A. uncinata*.

(3a) **A. evecata**, *Hoffm.*, de Vr., Mon. Maratt., 16, tab. III—IV, fig. 10.

Pinnulae subpellucid, subpetiolulate, lanceolate-elongate, straight or falcate, acuminate, the base unequally subrotundate or subtruncate, rarely subcuneate; largest pinnulae $\pm 17\frac{1}{2}$ c.M. long, to $\pm 2\frac{1}{4}$ c.M. broad at the centre; upper surface dark-green, lower paler; costa beneath black-brown; veins subhorizontal at the base, \pm oblique towards the apex, pale; recurrent veinlets subpellucid; sori subcontiguous, intramarginal, consisting of ± 10 —13 capsules; indusium membranaceous, fimbriate. — (? *Trop. Asia & Africa*), *Tahiti*.

(3b) **A. Miqueliana**, *de Vr.*, l. c., 26.

Pinnulae petiolate, lanceolate, short-acuminate, the base nearly equally cordate or truncate on both sides or rounded on the upper side; lower and central pinnulae ± 17 c.M. long, ± 2 c.M. broad; colour pale-green; costa and veins beneath sometimes sparingly furfuraceous or piloso-paleaceous; veins pale, opaque (? subtransparent); recurrent veinlets pellucid; sori contiguous, remote from the margin, consisting of 8—10 (after Miquel 4—5) capsules, encircled by a whorl of hairs. — *Java*.

(3c) **A. pruinosa**, *Hæe*, Bot. Zeit., 1846, 417; de Vr., l. c., 26.

Pinnulae short-stalked, lanceolate-oblong, hardly linear, narrowly acuminate, the base unequally subcordate, the upper sometimes subcuneate; lowest pinnulae $\pm 14\frac{1}{4}$ c.M. long, $\pm 1\frac{1}{3}$ c.M. broad; upper surface opaque-olivaceous in dried material, lower white-pruinose; veins black-fuscous; recurrent veinlets whether or not pellucid; sori intramarginal, consisting of ± 8 —17 capsules. — *Java, Philippines, New Guinea*.

(3d) **A. crassifolia**, *de Vr.*, l. c., 17.

Pinnulae subpetiolate, linear-lanceolate, straight or falcate, long-acuminate, the base \pm cuneate on the upper, \pm rotundato-cuneate on the lower side; largest pinnulae ± 14 c.M. long, $\pm 1\frac{1}{2}$ c.M. broad; colour pale, yellowish or yellow-green; costa and veins sometimes minutely paleaceous, rather pale; veins often oblique; recurrent veinlets reaching \pm the costa, opaque, as are the veins; sori close or subcontiguous, intramarginal, consisting of ± 8 —10 capsules; indusium thick. — *Java, Sumatra*.

(3e) **A. javanica**, *Pr.*, Tent. Pterid., Suppl., 20; *A. Dregeana*, *de Vr.*, l. c., 17, tab. III—IV, fig. 8.

Pinnulae petiolulate, linear or linear-oblong, acuminate, the base unequally

cordate or truncate; largest pinnulae $\pm 15-17\frac{1}{2}$ c.M long, $\pm 1\frac{1}{4}-1\frac{1}{3}$ c.M. broad; under surface paler in colour than the upper; costa clothed towards the base beneath with scattered, chaffy, hair-like or filiform, caducous scales; veins close, subhorizontal, generally blackish or fuscous, pellucid; recurrent veinlets often very pellucid; sori intramarginal, rather close, consisting of $\pm 6-12$, mostly 9 capsules. — *Java, Sumatra.*

(3f) **A. Hartingiana, de Vr., l. c., 25.**

Pinnulae petiolulate, linear-lanceolate, acuminate; upper base \pm cuneate, lower \pm rounded; largest pinnulae ± 15 c.M. long, ± 1 c.M. broad; colour black-green above; costa and veins black; recurrent veinlets pellucid; sori slightly remote from the margin, contiguous, consisting of 10—18 capsules. — *Java.*

(3g) **A. angustifolia, Pr., l. c., 21; de Vr, l. c. 18.**

Pinnulae subsessile, narrowly linear, long-acuminate; upper base \pm cuneate, lower \pm rounded; largest pinnulae ± 15 c.M. long, $\pm 1\frac{1}{4}$ c.M. broad; colour above deep-green, beneath pale-green; costa rather deciduously, sparingly and minutely setaceo-paleaceous beneath; veins close, slender, nearly horizontal, fuscous beneath, subtransparent; recurrent veinlets long, pellucid or opaque; sori often contiguous, remote from the margin, consisting of $\pm 5-12$ capsules. — *Luzon, Mindanao, New Guinea.*

(3h) **A. angustata, Miq., de Vr., l. c., 26.**

Pinnulae short-stalked, linear, acute, the base slightly cordate, subinequilateral; central pinnulae 10—12 c.M. long, $\pm 6-9$ m.M. broad; under surface pale; costa beneath fuscous; veins opaque; sori contiguous, \pm marginal, consisting of $\pm 6-8$ capsules. — *Java.*

(3i) **A. caudata, de Vr., l. c., 20.**

Pinnulae approximate, stalked, linear, caudato-acuminate, the base unequally subcordate, the lower base the broadest; largest pinnulae $\pm 16\frac{1}{2}$ c.M. long, $\pm 5-8$ m.M. broad; upper surface dark-green, lower fuscous, rather woolly on the costa and in the lower part; veins beneath fuscous, opaque, rather horizontal; recurrent veinlets long, reaching \pm the costa; sori contiguous, subremote from the margin; consisting of $\pm 9-18$ capsules. — *Philippines; Bismarck Islands.*

(3j) **A. hypoleuca, de Vr., l.c., 21.**

Pinnulae subsessile, lanceolate-oblong, \pm falcate, long-acuminate; upper base oblique, cuneate, lower broader, cuneate or rounded; largest pinnulae to \pm 16 c.M. long, to \pm $4\frac{1}{2}$ c.M. broad at the centre; upper surface yellow-green, lower white-waxy; veins oblique, pellucid; recurrent veinlets not reaching the costa, pellucid. — *Java*.

(3k) **A. approximata**, *de Vr.*, l. c., 25,

Pinnulae short-staked, linear-lanceolate, straight, the base \pm equilateral, rounded or subcordate; largest pinnulae \pm $13\frac{1}{2}$ c.M. long, \pm $1\frac{3}{4}$ c.M. broad at the centre; costa fuscous; veins opaque, subhorizontal; recurrent veinlets reaching nearly to the costa, indistinct, partially pellucid; sori contiguous, distant from the margin, consisting of \pm 9—10 capsules. — *Sumatra*.

(3l) **A. aphanosorus**, *de Vr.*, l. c., 19.

Pinnulae petiolate, lanceolate, subfalcate, rather straight, the base oblique on the upper, broadly cuneate to subtruncate on the lower side; largest pinnulae \pm 11 c.M. long, \pm $1\frac{1}{2}$ c.M. broad; texture pellucid; surfaces very minutely tomentose; costa castaneous; veins subfuscous; recurrent veinlets pellucid; sori minute, distant from the margin, consisting of \pm 10—12 capsules. — *Sumatra*.

(3m) **A. cuspidata**, *de Vr.*, l. c., 18, tab. III—IV, fig. 7.

Pinnulae petiolulate, linear, long-cuspidate; upper base rounded, lower cuneate; largest pinnulae \pm 12 c.M. long, \pm 1 c.M. broad; upper surface dark-green, lower paler; costa beneath woolly-tomentose, fuscous; veins fuscous, opaque; recurrent veinlets pellucid; Sori small, consisting of \pm 7—9 minute capsules. — *Java*.

(3n) **A. Teysmanniana**, *de Vr.*, l. c., 24, tab. I—II.

Pinnulae petiolate, subulate-lanceolate, straight or curved, acute; upper base broadly cuneate or rounded, lower narrower; largest pinnulae \pm 11 c.M. long, \pm $1\frac{3}{4}$ c.M. broad at the base; upper surface dark-green, lower paler, opaque; costa finely fusco-paleaceous beneath; veins distinct, pellucid like the recurrent veinlets; sori distant or approximate, remote from the margin, consisting of \pm 9—13 capsules. — *Java*.

(3o) **A. Willinckii**, *Miq.*, *de Vr.*, l. c., 21.

Pinnulae petiolulate, elongate, very acuminate, the base rounded or sub-

truncate, subinequilateral; barren pinnulae 20—25 c.M. long, \pm 3 c.M. broad, fertile 15—20 c.M. long, \pm 2½ c.M. broad; upper surface dark-green, lower pale-green; costa and veins fuscous; veins and recurrent veinlets \pm oblique, pellucid; sori distant from the margin, consisting of \pm 6—9 capsules. — *Java*.

(3p) **A. Presliana**, *de Vr.*, l. c., 20.

Pinnulae nodoso-petiolate, lanceolate, acute or long-acuminate, the base cuneate on the upper, rounded on the lower side; pinnulae \pm 17½ c.M. long; upper surface dark-green, lower pale, subglaucous, opaque; veins oblique, pellucid. — *Java*.

(3q) **A. pallescens**, *de Vr.*, l. c., 25.

Pinnulae petiolate, lanceolate, long-acuminate, the base unequally subtruncate or rotundato-cuneate; largest pinnulae not rarely coriaceous, \pm 17¾ c.M. long, \pm 1¾ c.M. broad; under surface pale-green, sparingly black-scaly; veins pale, suboblique at the base, oblique towards the apex; recurrent veinlets opaque; sori subremote from the margin, hardly contiguous, consisting of \pm 5—8 capsules. — *Java, Sumatra*.

(3r) **A. similis**, *Pr., de Vr.*, l. c., 17.

Pinnulae pellucid, petiolulate, straight or falcate, long-acuminate, the base unequally rounded; upper base shorter and broader, lower longer and narrower; largest pinnulae \pm 18½ c.M. long, \pm 2 c.M. broad; upper surface green, lower paler; veins and veinlets suboblique or horizontal, subpellucid; recurrent veinlets opaque; sori distant from the margin, not contiguous, consisting of \pm 8—9 capsules. — *Java, Mindanao, New Guinea*.

(3s) **A. D'Urvilleana**, *de Vr.*, l. c., 17, tab. III—IV, fig. 11.

Pinnulae subpellucid, petiolulate, linear, acuminate, the base unequally rounded, broadest on the lower side; upper surface black-green when dry, lower pale-green, fuscous or blackish near the costa; costa fuscous; veins nearly horizontal; recurrent veinlets reaching hardly more than ½-way down to the costa; sori intramarginal, subcontiguous, consisting of \pm 8—10 capsules. — *Philippines; Tahiti*.

(3t) **A. microsporangia**, *de Vr.*, l. c., 18.

Pinnulae subpellucid, petiolulate, linear-lanceolate, long-acuminate, the base

unequally subcuneate to subtruncate, broadest on the lower side; largest pinnulae ± 15 c.M. long, $\pm 1\frac{3}{4}$ c.M. broad; colour fuscous when dry; costa beneath fuscous; veins black or fuscous, sometimes opaque like the recurrent veinlets; sori intramarginal, consisting of $\pm 8-10$ minute capsules. — *Java, Sumatra.*

(3u) **A. ankolana** (*angkolana*) *de Vr.*, l. c., 19, tab. III—IV, fig. 9.

Pinnulae petiolulate, linear-lanceolate, long-acuminate, \pm equal- or unequal-sided at the base, the largest ± 17 c.M. long, $\pm 1\frac{3}{4}$ c.M. broad; costa fuscous or castaneous, sparingly paleaceous beneath; veins fuscous, close, pellucid; recurrent veinlets long, fuscous, pellucid; sori consisting of $\pm 10-12$ capsules. — *Java, Sumatra, ? New Guinea.*

(3v) **A. amboinensis**, *de Vr.*, l. c., 52.

Pinnulae petiolate, lanceolate, straight, suddenly acuminate, the base sub-equilateral, subtruncate or subrotundate; largest pinnulae ± 15 c.M. long, ± 2 c.M. broad, subpellucid; costa beneath fuscous; veins fuscous, pellucid, \pm horizontal; sori intramarginal, hardly contiguous, consisting of $\pm 10-20$ capsules. — *Amboyna.*

(3w) **A. muricata**, *Pr.*, de Vr., l. c., 50.

Pinnulae sessile, pellucid, lanceolate, narrowly acuminate; upper base broadly cuneate, lower narrowly rounded; largest pinnulae ± 11 c.M. long, $\pm 2\frac{1}{2}$ c.M. broad; costa black-fuscous; veins oblique, blackish; recurrent veinlets present, short; sori hardly contiguous, distant from the margin, consisting of $\pm 10-12$ capsules. — *Borneo.*

(3x) **A. crassipes**, *Wall.*, de Vr., l. c., 27; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 238, A.

Pinnulae pellucid, subpetiolate, linear-lanceolate, or lanceolate-oblong, the base unequally cuneate or rounded; largest pinnulae $\pm 16\frac{1}{2}$ c.M. long or longer, ± 3 c.M. broad or broader; colour pale- or yellow-green; veins sub-oblique, opaque, fuscous beneath; sori contiguous, near the edge, consisting of $\pm 10-20$ capsules; indusium fimbriato-laciniate. — *Sumatra, Malacca; North. India, Burmah, South. China.*

(3z) **A. uncinata**, *de Vr.*, l. c., 29, tab. III, fig. 6.

Pinnulae petiolate, lanceolate-oblong, the apex acuminate, rather conspicu-

ously upcurved, the base unequally cuneate, broadest on the upper side: largest pinnulae \pm 11 c.M. long, \pm 2—2½ c.M. broad; under surface minutely punctulate; costa fuscous beneath; veins pellucid; sori contiguous, distant from the edge, consisting of \pm 9—10 capsules. — *Amboyna; New Caledonia.*

INDEX.

INDEX.

A.

Abrodictyum	83
<i>Cumingii</i> , Pr.	99
Aconiopteris	711
Acrophorus	295
<i>affinis</i> , Moore	295
<i>immersus</i> , Moore	295
<i>nodosus</i> , Pr.	295
<i>parvulus</i> , Moore	824
<i>pulcher</i> , Moore	292
<i>stipellatus</i> , Moore	295
Acropterygium	57
Acrosorus	567
<i>exaltatus</i> , Copel.	614
<i>friderici & pauli</i> , Copel. ...	615
<i>Merrilli</i> , Copel.	614
Acrosticheae	122
Acrostichum	730
<i>aculeatum</i> , Rac.	722
<i>angulatum</i> , Bl.	713
<i>antrophyoides</i> , Bk.	744
<i>apiifolium</i> , Hk.	725
<i>appendiculatum</i> , Willd.	724
<i>articulatum</i> , Hk.	725
<i>aureum</i> , L.	730
<i>auritum</i> , Sw.	726
<i>axillare</i> , Cav.	745
<i>Beccarianum</i> , Bk.	714
<i>bicuspe</i> , Hk.	732
<i>bifurcatum</i> , Cav.	710
<i>Blumeanum</i> , Hk.	746, 747

Acrostichum.

<i>borneense</i> , Burck	840
<i>brachystachyum</i> , Rac.	729
<i>callaeifolium</i> , Bl.	714
<i>calomelanos</i> , L.	528
<i>celebicum</i> , Bk.	741
<i>commutatum</i> , Mett. ...	714
<i>conforme</i> , Bl.	714
<i>conforme</i> , Sw.	712
<i>costatum</i> , Wall.	745
<i>costulatum</i> , Ces.	644
<i>Cumingii</i> , Fée	715
<i>decurrens</i> , Desv.	715
<i>dichotomum</i> , L.	803
<i>digitatum</i> , L.	803
<i>diversifolium</i> , Bl.	740
<i>drynarioides</i> , Hk.	731
<i>exsculptum</i> , Bk.	740
<i>flagelliferum</i> , Wall.	739
<i>gracile</i> , Rac.	722
<i>grande</i> , A. Cunn.	708
<i>gorgoneum</i> , Klf. ...	712
<i>Hamiltonianum</i> , Wall.	724
<i>heteroclitum</i> , Pr.	739
<i>heterophyllum</i> , L.	705
<i>hirtum</i> , Sw.	717
<i>Holtrungii</i> , Bk.	737
<i>inaequale</i> , Willd.	730
<i>lanceolatum</i> , Hk.	745
<i>laurifolium</i> , Hk.	719
<i>laurifolium</i> , Thouars	714
<i>Lessonii</i> , Mett.	716

Acrostichum

<i>Linnaeanum</i> , Hk.	755
<i>lomarioides</i> , Christ	746
<i>marginatum</i> , Wall.	712
<i>melanosticticum</i> , Bl.	716
<i>melanosticton</i> , Bl.	716
<i>miniatum</i> , Christ	716
<i>minus</i> , Mett.	744
<i>modestum</i> , Bk.	745
<i>Nieuwenhuisii</i> , Rac.	724
<i>Norrisii</i> , Ces.	714
<i>Norrisii</i> , Hk.	716
<i>nummularifolium</i> , Bl.	685
<i>obovatum</i> , Bl.	685
<i>obtusifolium</i> , Spr.	715
<i>oligodictyon</i> , Bk.	755
<i>paleaceum</i> , Hk. & Grev.	717
<i>pellucido-marginatum</i> , Christ.	712
<i>petiolatum</i> , Sw.	717
<i>platyrhynchos</i> , Ces.	728
<i>platyrhynchos</i> , Hk.	729
<i>praestantissimum</i> , Bory	750
<i>Preslianum</i> , Hk.	740
<i>punctatum</i> , L.	655
<i>quercifolium</i> , Retz	757
<i>Quoyanum</i> , Gaud.	741
<i>repandum</i> , Bl.	741
<i>Requinianum</i> , Gaud.	747
<i>revolutum</i> , Zoll. var. <i>tenellum</i>	728
<i>rigidum</i> , Wall.	751
<i>scandens</i> , Hk.	720
<i>Smithii</i> , Rac.	720
<i>sorbifolium</i> , L.	720
<i>speciosum</i> , Willd.	750
<i>spectabile</i> , Rac.	720
<i>spicatum</i> , L.f.	728
var. <i>brachystachys</i>	729
<i>squamosum</i> , Sw.	717

Acrostichum

<i>stenosemioides</i> , Bk.	725
<i>subrepandum</i> , Hk.	742
<i>taccaefolium</i> , Hk.	757
<i>tartareum</i> , Cav.	528
<i>Teysmannianum</i> , Bk.	727
<i>thalictroides</i> , L.	751
<i>validinervis</i> , Rac.	729
<i>variabile</i> , Hk.	756
var. <i>axillare</i>	745
var. <i>decurrens</i>	756
<i>virens</i> , Wall.	741
<i>viscosum</i> , Sw.	717
<i>Zollingeri</i> , Kze	742
<i>Actinostachys</i>	115
<i>Adenophorus</i>	
<i>bipinnatus</i> , Gaud.	611
<i>Tamarisci</i> , Hk. & Grev.	611
<i>Adiantellum</i>	519
<i>Adiantum</i>	519, 826
<i>aculeolatum</i> , v. A. v. R.	527, 826
<i>aethiopicum</i> , L.	550, 331
<i>affine</i> , Hk.	525
<i>alatum</i> , Copel.	526
<i>amoenum</i> , Hk. & Grev.	554
<i>capillus veneris</i> , L.	535
<i>cardiochlaena</i> , Kze	528
<i>caudatum</i> , L.	524
var. β	524
var. <i>ciliatum</i>	524
<i>celebicum</i> , Christ.	551
<i>ciliatum</i> , Bl.	524
<i>concinnum</i> , H. B. K.	552
<i>cultratum</i> , Willd.	824
<i>cuneatum</i> , L. & F.	352
<i>denticulatum</i> , Burm.	301
<i>diaphanum</i> , Bl.	323
<i>decorum</i> , Moore.	552

Adiantum			
flabellulatum, <i>L.</i>	554		
formosum, <i>R. Br.</i>	528		
gompophyllum, <i>Bk.</i>	522		
hispidulum, <i>Sw.</i>	555		
Hosei, <i>Bk.</i>	525		
lanceum, <i>L.</i>	275		
Lobbianum, <i>Hk.</i>	527		
lunulatum, <i>Burm.</i>	525		
<i>var. Mettenii</i>	526		
lunulatum, <i>Houtt.</i>	170		
macrophyllum, <i>Sw.</i>	521		
Mettenii, <i>Kuhn</i>	526, 826		
mindanaoense, <i>Copel.</i>	527, 528		
neoguineense, <i>Moore</i>	550		
nobile, <i>Fisch.</i>	550		
opacum, <i>Copcl.</i>	550		
orbiculatum, <i>Lam</i>	270		
Parishii, <i>Hk.</i>	520		
pendactylon, <i>L. & F.</i>	525		
peruvianum, <i>Kl.</i>	521, 526		
philippense, <i>L.</i>	525		
platyphyllum, <i>Sw.</i>	521, 526		
polyphyllum, <i>Willd.</i>	528		
pteroides, <i>L.</i>	545		
pulchellum, <i>Bl.</i>	527		
repens, <i>L.f.</i>	288		
rhizophorum, <i>Sw.</i>	524		
Seemanni, <i>Hk.</i>	521		
Spencerianum, <i>Copel.</i>	525		
stenochlamys, <i>Bk.</i>	554		
suborbiculare, <i>v. A. v. R.</i>	551, 826		
tenerum, <i>Sw.</i>	550		
tinctum, <i>Moore</i>	552		
trapeziforme, <i>L.</i>	525		
<i>Wagneri, Mett.</i>	552		
<i>Zollingeri, Mett.</i>	525		
Aglaomorpha	569		
		Aglaomorpha	
		<i>Mejenima, Schott</i>	671
		Alcicornium	
		<i>bifurcatum, Und.</i>	710
		<i>coronarium, Und.</i>	711
		<i>grande, Und.</i>	708
		<i>sumbawense, Und.</i>	709
		<i>Wallichii, Und.</i>	708
		<i>Wilhelminae Reginae,</i>	
		<i>v. A. v. R.</i>	708
		<i>Willinckii, Und.</i>	710
		Aleuritopteris	557
		Allantodia	140
		<i>Brunoniana, Wall.</i>	140
		<i>javanica, Trev.</i>	140
		<i>nitidula, Kze</i>	454
		<i>Solenopteris, Kze</i>	455
		<i>sylvatica, Bl.</i>	829
		Allosurus	
		<i>Zippelii, Miq.</i>	545
		Alsophila	29, 789
		<i>alternans, Hk.</i>	17
		<i>batjanensis, Christ</i>	40
		<i>Brunoniana, Bedd.</i>	41
		<i>Burbidgei, Bk.</i>	55
		<i>calocoma, Christ</i>	789
		<i>caudata, J. Sm.</i>	785
		<i>celebica, Mett.</i>	42
		<i>Christii, v. A. v. R.</i>	42
		<i>Clementis, Copel.</i>	790
		<i>commutata, Mett.</i>	54
		<i>comosa, Christ</i>	42
		<i>comosa, Wall.</i>	55
		<i>contaminans, Wall.</i>	41, 790
		<i>crenulata, Mett.</i>	27
		<i>crinita, Hk.</i>	40
		<i>cyclodonta, Christ</i>	19
		<i>dimorpha, Christ</i>	56

Alsophila

<i>dubia</i> , <i>Bedd.</i>	51
<i>extensa</i> , <i>Spr.</i>	45
<i>fuliginosa</i> , <i>Christ</i>	59
<i>gigantea</i> , <i>Wall.</i>	52
<i>glabra</i> , <i>Hk.</i>	52, 789
<i>glauca</i> , <i>J. Sm.</i>	41, 790
<i>Haenkei</i> , <i>Pr.</i>	45
<i>intermedia</i> , <i>Mett.</i>	45
<i>Junghuhniana</i> , <i>Kze</i>	28
<i>Kingi</i> , <i>Clarke</i>	56
<i>latebrosa</i> , <i>Wall.</i>	58, 789
<i>lepidoclada</i> , <i>Christ</i>	57
<i>lepifera</i> , <i>J. Sm.</i>	59
<i>var. congesta</i>	789
<i>lunulata</i> , <i>R. Br.</i>	43
<i>lurida</i> , <i>Hk.</i>	44, 791
<i>Margarethae</i> , <i>Schroet.</i>	55
<i>melanopus</i> , <i>Hassk.</i>	40
<i>melanorhachis</i> , <i>Copel.</i>	791
<i>mindanensis</i> , <i>Christ</i>	44
<i>modesta</i> , <i>Bk.</i>	57
<i>obscura</i> , <i>Scort.</i>	54
<i>ornata</i> , <i>Scott</i>	790
<i>polyphlebia</i> , <i>Bk.</i>	45
<i>ramispina</i> , <i>Hk.</i>	54
<i>Ridleyi</i> , <i>Bk.</i>	52
<i>sangirensis</i> , <i>Christ</i>	57
<i>saparuensis</i> , <i>v. A. v. R.</i> . 58,	790
<i>scaberula</i> , <i>Christ</i>	55
<i>siktimensis</i> , <i>Clarke & Bk.</i> . .	790
<i>squamulata</i> , <i>Bedd.</i>	54
<i>squamulata</i> , <i>Hk.</i>	55, 789
<i>tenuisecta</i> , <i>Bl.</i>	257
<i>Teysmanniana</i> , <i>Rac.</i>	51
<i>tomentosa</i> , <i>Hk.</i>	59, 45, 791
<i>trichodesma</i> , <i>Scort.</i>	55
<i>tristis</i> , <i>Bl.</i>	44, 257

Alsophila

<i>truncata</i> , <i>Brack.</i>	42
<i>vexans</i> , <i>Ces.</i>	52
<i>Wallacei</i> , <i>Mett.</i>	56
<i>Warburgii</i> , <i>Christ</i>	204
<i>xantholepis</i> , <i>Christ</i>	44
<i>Amphicosmia</i>	27
<i>alternans</i> , <i>Moore</i>	17
<i>riparia</i> , <i>Gardn.</i>	27
<i>Angiopteris</i>	761, 855
<i>amboinensis</i> , <i>de Vr.</i>	859
<i>angkolana</i> , <i>de Vr.</i>	859
<i>angustata</i> , <i>Miq.</i>	856
<i>angustifolia</i> , <i>Pr.</i>	856
<i>ankolana</i> , <i>de Vr.</i>	859
<i>aphanosorus</i> , <i>de Vr.</i>	857
<i>approximata</i> , <i>de Vr.</i>	837
<i>cartilagens</i> , <i>Christ</i>	762
<i>cartilaginea</i> , <i>Christ</i>	762
<i>caudata</i> , <i>de Vr.</i>	836
<i>crassifolia</i> , <i>de Vr.</i>	855
<i>crassipes</i> , <i>Wall.</i>	859
<i>cuspidata</i> , <i>de Vr.</i>	837
<i>Dregeana</i> , <i>de Vr.</i>	855
<i>D'Urvilleana</i> , <i>de Vr.</i>	838
<i>evecta</i> , <i>Hoffm.</i>	762, 855, 855
<i>Hartingeana</i> , <i>de Vr.</i>	836
<i>hypoleuca</i> , <i>de Vr.</i>	856
<i>javanica</i> , <i>Pr.</i>	855
<i>microsporangia</i> , <i>de Vr.</i>	858
<i>Miqueliana</i> , <i>de Vr.</i>	855
<i>muricata</i> , <i>Pr.</i>	859
<i>pallescens</i> , <i>de Vr.</i>	858
<i>Presliana</i> , <i>de Vr.</i>	838
<i>pruinosa</i> , <i>Kze</i>	855
<i>similis</i> , <i>Pr.</i>	858
<i>Smithii</i> , <i>Rac.</i>	762
<i>Teysmanniana</i> , <i>de Vr.</i>	857

- Angiopteris
 uncinata, de Vr. 859
 Willinckii, de Vr. 857
 Anchistea 590
 Anisogonium 595
 alismaefolium, Christ 425
 cordifolium, Bedd. 422
 elegans, Pr. 425
 esculentum, Pr. 425
 lineolatum, Bedd. 425
 Smithianum, Bedd. 425
 Antigramme 477
 Antrophyum 529
 avenium, Bl. 538
 Boryanum, Bl. 535
 Brookei, Hk. 554
 callifolium, Bl. 551, 555
 var. alatum 555
 Clementis, Christ 551
 coriaceum, Bl. 558
 coriaceum, Wall. 554
 costatum, v. A. v. R. 552
 Cumingii, Fée 555
 elongatum, Fée 555
 falcatum, Bl. 555
 Grevillei, Balf. 555
 immersum, Mett. 551
 involutum, Bl. 557
 lanceolatum, Bl. 558
 latifolium, Bl. 555
 marginale, Bl. 551
 niphoboloides, Kze 687
 obtusum, Bl. 551
 ovatum, v. A. v. R. 554
 parvulum, Bl. 552
 var. elongatum 555
 plantagineum, Klf. 555
 pumilum, Klf. 551
 Antrophyum
 reticulatum, Bedd. 553, 554
 reticulatum, Klf. 555
 semicostatum, Bl. 552, 555
 spathulatum, v. A. v. R. .. 556
 strictum, Mett. 552
 subfalcatum, Brack. 554
 Arcypteris 512
 Arthrobotrya 725
 Arthropteris 154
 glabra, Copel. 155
 obliterata, J. Sm. 155
 ramosa, Mett. 155
 Aspidieae 121
 Aspidium 255, 822
 abortivum, Bl. 218
 var. B & C 218
 aculeatum, Schott 166
 var. obtusum 167
 acutum, Schk. 162
 adiantoides, Bl. 262
 adnatum, Bl. 191
 alatellum, Christ 210
 alatum, Wall. 241
 amabile, Bl. 168
 amabile, Rac. 167
 ambiguum, Diels 521
 andaiense, C. Chr. 514
 Angilogense, Christ 810
 angulatum, Christ 245
 angulatum, Hk. 255
 angulatum, J. Sm. 256, 822
 Arbuscula, Willd. 218
 aridum, Don 212
 aristatum, Sw. 169
 var. conifolium 169
 var. Sarasinorum 169
 atratum, Wall. 180

Aspidium

<i>attenuatum</i> , Kze	184
<i>auriculatum</i> , Sw.	
<i>var. obliquum</i>	809
<i>Barberi</i> , C. Chr.	515
<i>batjanense</i> , Christ	166
<i>Beccarianum</i> , Bk.	258
<i>Beccarianum</i> , Diels	515
<i>Beddomei</i> , Prantl	191
<i>Bencitanum</i> , Gaud.	225
<i>biaristatum</i> , Bl.	167
<i>biseriatum</i> , Christ	251
<i>biserratum</i> , Sw.	162
<i>Bolsteri</i> , Copel.	519
<i>Boryanum</i> , Willd.	201
<i>brachiatum</i> , Zoll. & Mor. . .	252
<i>brachyodus</i> , Gris.	220
<i>Brongniartii</i> , Diels	515
<i>Bryanti</i> , Copel.	514
<i>calcaratum</i> , Bl.	185
<i>calcareum</i> , Pr.	256, 822
<i>callosum</i> , Bl.	819
<i>canescens</i> , Christ	208
<i>forma acrostichoides</i> . . .	505
<i>forma gymnogrammoides</i> . . .	507
<i>forma nephrodiiiformis</i> . .	251
<i>carvifolium</i> , Kze	169
<i>caryotideum</i> , Wall.	255
<i>Cesatianum</i> , C. Chr.	258
<i>chaerophyllaefolium</i> , Zipp. . .	194
<i>cicutarium</i> , Christ	248
<i>cicutarium</i> , Sw.	249
<i>coadunatum</i> , Wall.	248
<i>coniifolium</i> , Wall.	169
<i>conioneuron</i> , Fée.	210
<i>Copelandii</i> , C. Chr.	246
<i>costale</i> , Bl.	455
<i>crassifolium</i> , Bl.	182

Aspidium

<i>crenatum</i> , Kuhn	202
<i>cucullatum</i> , Bl.	215
<i>curvifolium</i> , Kze	169
<i>cyatheoides</i> , Klf.	216
<i>Dayi</i> , Bedd.	188
<i>decurrens</i> , Pr.	247
<i>densum</i> , Zipp.	167
<i>deversum</i> , Kze	189
<i>devexum</i> , Kze	811
<i>diaphanum</i> , Zoll. & Mor. . . .	165
<i>difforme</i> , Bl.	520
<i>discretum</i> , Don	167
<i>dissectum</i> , Christ	196
<i>dissectum</i> , Mett.	212
<i>echinatum</i> , Mett.	182
<i>edulentum</i> , Kze	194
<i>erythrosorum</i> , Eat.	815
<i>Everettii</i> , C. Chr.	241
<i>excellens</i> , Bl.	171
<i>exiguum</i> , Kze	190
<i>extensum</i> , Bl.	210
<i>falcatum</i> , Sw.	255
<i>Fauriei</i> , Christ	
<i>var. elatius</i>	429
<i>ferox</i> , Bl.	221
<i>ferrugineum</i> , Diels	516
<i>Filix-mas</i> , Sw.	192
<i>var. fibrillosum</i>	192
<i>var. megaphyllum</i>	204
<i>flaccidum</i> , Bl.	195
<i>floccigerum</i> , Bl.	807
<i>giganteum</i> , Bl.	811
<i>var. minor</i>	811
<i>giganteum</i> , Copel.	811
<i>glanduligerum</i> , Kze	186
<i>glandulosum</i> , Bl.	215
<i>goggilodus</i> , Schk.	212

Aspidium

<i>gongyloides</i> , Schk.	212
<i>gracilescens</i> , Bl.	186
<i>grammitoides</i> , Christ	181
<i>grande</i> , J. Sm.	249
<i>grandifolium</i> , Mett.	245
<i>grandifolium</i> , Pr.	249
<i>Haenkei</i> , Pr.	257
<i>Hallieri</i> , Christ	184
<i>heterocarpon</i> , Bl.	228
<i>heterodon</i> , Copel.	246
<i>heterophyllum</i> , Hk.	171
<i>hirsutum</i> , Kze	214
<i>hirtipes</i> , Bl.	180
<i>hispidulum</i> , Dec.	228
<i>Hookeri</i> , Bk.	251
<i>Hookeri</i> , Wall.	218
<i>Hosei</i> , C. Chr.	247
<i>hymenophylloides</i> , Bl.	294
<i>immersum</i> , Bl.	188
<i>immersum</i> , Hk.	252
<i>intermedium</i> , Bl.	199
<i>intermedium</i> , Christ	202
<i>invisum</i> , Christ	212
<i>irriguum</i> , J. Sm.	245
<i>irregulare</i> , C. Chr.	520
<i>Keckii</i> , Luer.	248
<i>Kehdingianum</i> , Luer.	255
<i>Kingii</i> , Hance	165
<i>Koordersii</i> , Christ	181
<i>Kuhnii</i> , C. Chr.	520
<i>Künstleri</i> , Bedd.	256
<i>labrusca</i> , Christ	517
<i>laeve</i> , Kze	220
<i>lamoense</i> , Copel.	517
<i>latebrosum</i> , Kze	221
<i>latifolium</i> , J. Sm.	249, 825
<i>latipinna</i> , Hance	217

Aspidium

<i>Leuzeanum</i> , Kze	172
<i>ligulatum</i> , Kze	184
<i>lineatum</i> , Bl.	209
<i>lobatum</i> , Sw.	167
<i>Lobbii</i> , Hk.	244
<i>Loherianum</i> , Christ	221
<i>longipes</i> , Bl.	211
<i>Luerssenii</i> , Christ	190
<i>macrocarpon</i> , Bl.	451
<i>macrodon</i> , Keys.	516
<i>malayense</i> , Christ	249
<i>mamillosum</i> , C. Chr.	247
<i>marginatum</i> , Wall.	814
<i>megalocarpum</i> , C. Chr.	172
<i>megaphyllum</i> , Christ	204
<i>megaphyllum</i> , Mett.	218
<i>melanocaulon</i> , Bl.	245
<i>melanorachis</i> , C. Chr.	250
<i>membranaceum</i> , Hk.	811
<i>membranifolium</i> , Kze	175
<i>meniscocarpon</i> , Mett.	251
<i>menyanthidis</i> , Pr.	244
<i>moluccense</i> , Bl.	166
<i>Motleyanum</i> , Christ	182
<i>moulmeinense</i> , Christ	217
<i>mucronifolium</i> , Bl.	166
<i>multicaudatum</i> , Wall.	250
<i>multilineatum</i> , Christ ...	215, 217
<i>musifolium</i> , Bl.	807
<i>nebulosum</i> , C. Chr.	246
<i>nephrodioides</i> , Hk.	251
<i>nigripes</i> , Bl.	455
<i>nudum</i> , Diels	250
<i>obliquum</i> , Don	809
<i>obtusilobum</i> , Prantl	203
<i>obtusum</i> , Kze	166
<i>ochtodes</i> , Kze	215

Aspidium

<i>oligophyllum</i> , <i>Rst.</i>	257
<i>ornatum</i> , <i>Christ</i>	495
<i>otaria</i> , <i>Kze</i>	210
<i>pachyphyllum</i> , <i>Kze</i>	259, 252
<i>paleaceum</i> , <i>Rac.</i>	191
<i>palmatum</i> , <i>Mett.</i>	247
<i>parallelogrammum</i> , <i>Kze</i>	192
<i>parasiticum</i> , <i>Sw.</i>	224
<i>patens</i> , <i>Sw.</i>	189
<i>pectinatum</i> , <i>Christ</i>	185
<i>pennigerum</i> , <i>Bl.</i>	218
<i>pentaphyllum</i> , <i>v. A. v. R.</i>	519
<i>perakense</i> , <i>Bedd.</i>	226
<i>persoriferum</i> , <i>Copel.</i>	251
<i>phaeostigma</i> , <i>Ces.</i>	181
<i>platanifolium</i> , <i>Mett.</i>	255
<i>platyphyllum</i> , <i>Pr.</i>	258
<i>platypus</i> , <i>Kze</i>	810
<i>polycarpum</i> , <i>Bl.</i>	252
<i>polymorphum</i> , <i>Wall.</i>	259, 243
<i>Preslianum</i> , <i>Mett.</i>	164
<i>procurrens</i> , <i>Mett.</i>	211
<i>profereoides</i> , <i>Christ</i>	812
<i>productum</i> , <i>Klf.</i>	250
<i>psilopodium</i> , <i>C. Chr.</i>	245
<i>pteroides</i> , <i>Sw.</i>	209
<i>pteropodium</i> , <i>Diels</i>	242
<i>pteropus</i> , <i>Kze</i>	247
<i>pulvinuliferum</i> , <i>Rac.</i>	200
<i>purpurascens</i> , <i>Bl.</i>	201
<i>repandum</i> , <i>Willd.</i>	258
<i>Riedleanum</i> , <i>Gaud.</i>	250
<i>rufescens</i> , <i>Bl.</i>	495
<i>sagenioides</i> <i>Mett.</i>	191
<i>sagenioides</i> , <i>Rac.</i>	197
<i>sagittae-folium</i> , <i>Bl.</i>	222
<i>saxicola</i> , <i>Bl.</i>	515

Aspidium

<i>semibipinnatum</i> , <i>Wall.</i>	244, 825
<i>semicordatum</i> , <i>Sw.</i>	165
<i>setigerum</i> , <i>Kuhn</i>	202
<i>siifolium</i> , <i>Mett.</i>	251, 825
<i>var. pachyphyllum</i>	252
<i>simplicifolium</i> , <i>Hk.</i>	500
<i>singaporianum</i> , <i>Wall.</i>	255
<i>sparsum</i> , <i>Spr.</i>	197
<i>splendens</i> , <i>Christ</i>	195
<i>stenophyllum</i> , <i>Diels</i>	247
<i>stipellatum</i> , <i>Bl.</i>	227
<i>subcaudatum</i> , <i>v. A. v. R.</i>	242
<i>subdecurrens</i> , <i>C. Chr.</i>	514
<i>subdigitatum</i> , <i>Bl.</i>	485
<i>subtriphyllum</i> , <i>Hk.</i>	246, 248, 825
<i>sumatranum</i> , <i>C. Chr.</i>	518
<i>syrmaticum</i> , <i>Willd.</i>	195
<i>tenuisectum</i> , <i>Bl.</i>	257
<i>ternatense</i> , <i>v. A. v. R.</i>	258, 825
<i>ternatum</i> , <i>Diels</i>	242
<i>Teysmannianum</i> , <i>Hk.</i>	251
<i>tricuspe</i> , <i>Bedd.</i>	255
<i>triste</i> , <i>Bl.</i>	257
<i>truncatum</i> , <i>Gaud.</i>	227
<i>unitum</i> , <i>Mett.</i>	212
<i>unitum</i> , <i>Sieb.</i>	215
<i>urophyllum</i> , <i>Christ.</i>	216
<i>variolosum</i> , <i>Wall.</i>	248, 825
<i>varium</i> , <i>Sw.</i>	168
<i>vastum</i> , <i>Bl.</i>	241, 247, 518, 823
<i>var. heterosorum.</i>	518
<i>vestitum</i> , <i>Sw.</i>	167
<i>vile</i> , <i>Kze</i>	199
<i>vitis</i> , <i>C. Chr.</i>	516
<i>vulcanicum</i> , <i>Bl.</i>	166
<i>Wallichii</i> , <i>Hk.</i>	155
<i>Warburgii</i> , <i>Kuhn & Christ</i>	180

Aspidium

<i>Warburgii</i> , var. <i>debile</i>	501
<i>Weinlandii</i> , <i>Christ</i>	244
<i>Whitfordi</i> , <i>Copel.</i>	519
<i>Wigmanii</i> , <i>Rac.</i>	192
<i>Zippelianum</i> , <i>C. Chr.</i>	516
<i>Zollingerianum</i> , <i>Bedd.</i>	248, 252
Aspidopodium	568
Aspidotis	555
Asplenidictyum	457
Asplenieae	122
Asplenium	456, 850
<i>acrostichoides</i> , <i>Sw.</i>	451
<i>acuminatum</i> , <i>Wall.</i>	409
<i>acutiusculum</i> , <i>Bl.</i>	463
<i>adiantoides</i> , <i>C. Chr.</i>	465, 464, 465
<i>aequibasale</i> , <i>Bk.</i>	404
<i>affine</i> , <i>Sw.</i>	472
<i>alismaefolium</i> , <i>Hk.</i>	425
<i>alternifolium</i> , <i>Mett.</i>	405
<i>amboinense</i> , <i>Willd.</i>	441
<i>amoenum</i> , <i>Pr.</i>	466
<i>anisodontum</i> , <i>Pr.</i>	463
<i>amplissimum</i> , <i>Bk.</i>	419
<i>apoense</i> , <i>Copel.</i>	444
<i>approximatum</i> , <i>Bl.</i>	462
<i>arayatense</i> , <i>Christ</i>	460
<i>asperum</i> , <i>Hk.</i>	420
<i>arifolium</i> , <i>Burm.</i>	547
<i>australasicum</i> , <i>Hk.</i>	459
<i>Bakeri</i> , <i>C. Chr.</i>	475
<i>bantamense</i> , <i>Bk.</i>	405
<i>barisanicum</i> , <i>Bk.</i>	404
<i>basilare</i> , <i>Moore</i>	829
<i>batuense</i> , v. <i>A. v. R.</i>	443
<i>Beccarianum</i> , <i>Ces.</i>	473
<i>Belangeri</i> , <i>Bory</i>	457

Asplenium

<i>Belangeri</i> , <i>Kze</i>	474
<i>biseriale</i> , <i>Bk.</i>	405
<i>brachypodium</i> , <i>Bk.</i>	413
<i>Brackenridgei</i> , <i>Bk.</i>	415
<i>borneense</i> , <i>Hk.</i>	457
<i>bulbiferum</i> , <i>Forst.</i>	471
<i>bullatum</i> , <i>Wall.</i>	471
<i>camarinum</i> , <i>Bk.</i>	402
<i>canaliculatum</i> , <i>Bl.</i>	465
<i>caulatum</i> , <i>Bedd.</i>	464
<i>caudatum</i> , <i>Forst.</i>	460, 469
var. <i>horridum</i>	468
<i>ceramicum</i> , <i>Miq.</i>	421
<i>Cesatianum</i> , <i>Bk.</i>	465
<i>cheilosorum</i> , <i>Kze</i>	466
<i>chlorophyllum</i> , <i>Bk.</i>	401
<i>comosum</i> , <i>Christ</i>	446
<i>colubrinum</i> , <i>Christ</i>	458
<i>concolor</i> , <i>Hk.</i>	442
<i>confertum</i> , <i>Bk.</i>	404
<i>confluens</i> , <i>Kze</i>	428
<i>contiguum</i> , <i>Bedd.</i>	460
<i>contiguum</i> , <i>Klf.</i>	461
<i>cordifolium</i> , <i>Mett.</i>	422
<i>crenato-serratum</i> , <i>Bl.</i>	407
<i>crinicaule</i> , <i>Hance</i>	463
<i>crinitum</i> , <i>Bk.</i>	414
<i>cultratum</i> , <i>Hk.</i>	407
<i>cuneatum</i> , <i>Hk. & Grev.</i>	850
<i>cuneatum</i> , <i>Lam.</i>	470
<i>curtisorum</i> , <i>Christ</i>	459
<i>cyatheifolium</i> , <i>Rich.</i>	416
<i>cymbifolium</i> , <i>Christ.</i>	440
<i>decorum</i> , <i>Kze</i>	474
<i>decussatum</i> , <i>Sw.</i>	424
<i>deltoideum</i> , <i>Pr.</i>	418
<i>densum</i> , <i>Brack.</i>	452

Asplenium

<i>denticulatum</i> , Bl.	467
<i>dichotomum</i> , Hk.	474
<i>dimidiatum</i> , Sw.	462
<i>diversifolium</i> , Bl.	454
<i>forma concolor</i>	442
<i>forma indivisa</i>	442
<i>Doreyi</i> , Kze	458
<i>echinosporum</i> , Rac.	445
<i>Elmeri</i> , Christ	469
<i>elongatum</i> , Sw.	458
<i>epiphyticum</i> , Copel.	441
<i>erectum</i> , Bl.	457
<i>erectum</i> , Bory	458
<i>esculentum</i> , Pr.	425
<i>exiguum</i> , Bedd.	455
<i>falcatum</i> , Bedd.	465
<i>falcatum</i> , Lam.	464
<i>fejeense</i> , Brack.	441
<i>filipes</i> , Copel.	467
<i>Filix Femina</i> , Bernh. var.	454
<i>Finlaysonianum</i> , Wall.	476
<i>Forbesii</i> , Bk.	422
<i>fraxinifolium</i> , Wall.	405
<i>fuliginosum</i> , Hk.	456
<i>furcatum</i> , Thb.	470
<i>Gardneri</i> , Bk.	464
<i>gedeanum</i> , Rac.	429
<i>glochidiatum</i> , Rac.	444, 445
<i>grammitoides</i> , Hk.	410
<i>griseum</i> , Copel.	459
<i>gymnogrammoides</i> , Kl.	454
<i>Hagenii</i> , v. A. v. R.	468
<i>Hancei</i> , Bk.	465
<i>heterocarpum</i> , Wall.	466
<i>heterodon</i> , Mett.	456
<i>heteropteron</i> , Mett.	425
<i>hirtum</i> , Klf.	462

Asplenium

<i>horizontale</i> , Bk.	456
<i>horridum</i> , Klf.	461, 468
<i>Huttoni</i> , Bk.	421
<i>impersum</i> , Christ	464
<i>insigne</i> , Copel.	459
<i>insititium</i> , Brack.	461
<i>integerrimum</i> , Wall.	850
<i>integrifolium</i> , Mett.	422
<i>japonicum</i> , Thb.	408
<i>javanicum</i> , Bl.	140
<i>Labillardieri</i> , Kze.	454
<i>laciniatum</i> , Don.	468
<i>laserpitiifolium</i> , Bedd.	471
<i>laserpitiifolium</i> , Lam.	472
<i>lasiopteris</i> , Mett.	411
<i>latifolium</i> , Don	416
<i>Lauterbachii</i> , Christ	475
<i>laxivenum</i> , Copel.	461
<i>lepturus</i> , J. Sm.	461
<i>lineolatum</i> , Mett.	425
<i>Linza</i> , Ces.	479
<i>Lobbianum</i> , Hk.	408
<i>Loherianum</i> , Christ	452
<i>lomariaceum</i> , Christ	401
<i>longissimum</i> , Bl.	457
<i>lugubre</i> , Hk.	456
<i>lunulatum</i> , Sw.	458
<i>macrocarpum</i> , Bl.	451
<i>macrophyllum</i> , Sw.	464, 465, 466, 850
<i>Mactierii</i> , Bedd.	445
<i>manilense</i> , Spr.	417
<i>Meyenianum</i> , Mett.	418
<i>Meyenianum</i> , Pr.	459
<i>microxiphion</i> , Bk.	442
<i>militare</i> , Copel.	462
<i>minus</i> , Bl.	451

Asplenium

<i>monotis</i> , Christ	457
<i>multijugum</i> , Wall.	455
<i>multilineatum</i> , Hk.	453
<i>muricatum</i> , Mett.	829
<i>musaeifolium</i> , Mett.	459
<i>myriophyllum</i> , Muell.	475
<i>natunae</i> , Bk.	444
<i>nidus</i> , L.	459, 850
<i>var. pachyphyllum</i>	459
<i>var. Simonsianum</i>	458
<i>nigrescens</i> , Bl.	461
<i>nigripes</i> , Bl.	455
<i>nitidulum</i> , Moore	454
<i>nitidum</i> , Sw.	471
<i>normale</i> , Don	455
<i>normale</i> Mett. ..	451
<i>novo-guineense</i> , Rst.	476
<i>obscurum</i> , Bl.	467
<i>obtusilobum</i> , Hk.	474
<i>oceanicum</i> , C. Chr.	474
<i>oxyphyllum</i> , Hk.	455
<i>pachyphyllum</i> , Kze	459
<i>paradoxum</i> , Bl.	465
<i>pallidum</i> , Bl.	405
<i>pellucidum</i> , Lam. ...	454, 459, 462
<i>persicifolium</i> , J. Sm.	460
<i>Petersenii</i> , Kze	411
<i>Phyllitidis</i> , Don	459, 850
<i>plicatum</i> , Zipp.	459
<i>polypodioides</i> , Mett.	419, 420
<i>porphyrorachis</i> , Bk.	401
<i>porrectum</i> , Wall.	407
<i>praemorsum</i> , Sw.	470, 850
<i>Prescottianum</i> , Wall.	415
<i>prionurus</i> , J. Sm.	459
<i>proliferum</i> , Lam.	424
<i>pusillum</i> , Bl.	428

Asplenium

<i>remotum</i> , Moore	455
<i>resectum</i> , Sm.	466
<i>salignum</i> , Bl.	454
<i>salignum</i> , Hk.	454
<i>Sancti-Christophori</i> , Christ ..	465
<i>Sarasinorum</i> , Copel.	452
<i>scandens</i> , J. Sm.	475
<i>Schoggersii</i> , v. A. v. R. ...	460
<i>scolopendrioides</i> , J. Sm. ...	444, 446
<i>scolopendrioides</i> , Muell.	442
<i>scolopendropsis</i> , Muell.	477
<i>Scortechinii</i> , Bedd.	445
<i>Selenopteris</i> , Mett.	455
<i>serraeforme</i> , Mett.	467
<i>serrulatum</i> , Pr.	417
<i>setisectum</i> , Bl.	472
<i>shepherdiioides</i> , Bk.	408
<i>Simonsianum</i> , Hk.	458
<i>simplex</i> , Bl.	442
<i>simplex</i> , Mett.	459
<i>Smathianum</i> , Bk.	424
<i>sorzogonense</i> , Pr.	412
<i>var. majus</i> ...	414
<i>spathulatum</i> , Bk.	442
<i>spathulinum</i> , J. Sm.	472
<i>speciosum</i> , Mett.	415
<i>spinulosum</i> , Mett.	424, 425
<i>squamulatum</i> , Bl.	445
<i>Stantoni</i> , Copel.	452
<i>Steerei</i> , Harr.	459
<i>stereophyllum</i> , Kze	469
<i>subaquatile</i> , Ces.	455
<i>subavenium</i> , Hk.	455
<i>subnormale</i> , Copel.	451
<i>subserratum</i> , Bl.	400
<i>subserratum</i> , Ces.	480
<i>sulcatum</i> , Mett.	472

Asplenium

<i>sumatranum</i> , Hk.	455
<i>sundense</i> , Bl.	444
<i>sylvaticum</i> , Pr.	402
<i>taeniosum</i> , Kze	442
<i>tenerum</i> , Forst.	458
<i>var. Belangeri</i>	474
<i>tenuifolium</i> , Don	469
<i>thelypteroides</i> , Mchx	451
<i>tomentosum</i> , Hk.	414
<i>tomentosum</i> , Mett.	411
<i>Toppingianum</i> , Copel.	428
<i>Toppingii</i> , Copel.	429
<i>trichomanes</i> , L.	452
<i>truncatilobum</i> , Fée	460
<i>umbrosum</i> , J. Sm.	828
<i>unilaterale</i> , Lam.	466, 467
<i>vacillans</i> , Kze	405
<i>vestitum</i> , Hk.	417
<i>vittaeforme</i> , Cav.	444
<i>vulcanicum</i> , Bl.	456
<i>Warburgianum</i> , Christ	467
<i>Wernerii</i> , Rst.	445
<i>Wightianum</i> , Ces.	463
<i>Wightianum</i> , Wall.	456
<i>woodwardioides</i> , Bk.	829
<i>xiphophyllum</i> , Bk.	406
<i>zamiaefolium</i> , Pr.	464
<i>zamiaefolium</i> , Willd.	462
<i>zamiioides</i> , Hk.	465
<i>zeylanicum</i> , Ces.	401
<i>zeylanicum</i> , Hk.	400

Atactosia 568

Athyrium 426, 850

<i>acrostichoides</i> , Diels	451
<i>allantodioides</i> , Bedd.	451
<i>anisopteron</i> , Christ	429
<i>anisopteron</i> , Christ	429

Athyrium

<i>aristulatum</i> , Copel.	452
<i>australe</i> , Bedd.	828
<i>benguetense</i> , Christ	450
<i>Copelandi</i> , Christ	450
<i>drepanopteron</i> , Moore	455
<i>drepanopteron</i> , R. Br.	455
<i>gedeanum</i> , Christ.	429
<i>gymnogrammoides</i> , Bedd.	454
<i>hyalostegium</i> , Copel.	451
<i>Loheri</i> , Christ	452
<i>macrocarpum</i> , Bedd.	451
<i>nanum</i> , Christ	428
<i>nigripes</i> , Bedd.	455
<i>nigripes</i> , Moore	455
<i>nitidulum</i> , Milde	454
<i>oxyphyllum</i> , Moore	455
<i>pectinatum</i> , Bedd.	455
<i>philippinense</i> , Christ	452
<i>pusillum</i> , v. A. v. R.	428, 850
<i>Sarasinorum</i> , Christ	452
<i>selenopteris</i> , Moore	455
<i>silvaticum</i> , Milde	829
<i>solenopteris</i> , Moore	455
<i>thelypteroides</i> , Desv.	451
<i>Toppingianum</i> , Copel.	428
<i>umbrosum</i> , Pr.	828
<i>var. muricatum</i>	829
<i>woodwardioides</i> , Christ	829
<i>Austrogramme</i>	559

B.

Balantium

<i>Blumei</i> , Kze	47
<i>Bipinnatifidae</i> (Pteris)	552
<i>Blechnae</i>	122
<i>Blechnum</i>	585

Blechnum

<i>areolare</i> , Copel.	855
<i>Blumei</i> , Moore	587
<i>capense</i> , Schl.	581
<i>cartilagineum</i> , Sw.	586
<i>colubrinum</i> , Blanco	152
<i>egenolfooides</i> , C. Chr.	580
<i>egregium</i> , Copel.	588
<i>elongatum</i> , Gaud.	586
<i>Finlaysonianum</i> , Wall.	587
<i>Fraseri</i> , Luerss.	582
<i>lanceola</i> , Sw.	586
<i>nitidum</i> , Pr.	586
<i>var. contracta</i>	586
<i>orientale</i> , L.	587
<i>Patersoni</i> , Mett.	579
<i>var. majus</i>	579
<i>punctulatum</i> , Sw.	580
<i>radicans</i> , L.	591
<i>serrulatum</i> , Rich.	588
<i>stenopterum</i> , Hance	541
<i>striatum</i> , R. Br.	588
<i>Treubii</i> , v. A. v. R.	586
<i>vestitum</i> , Kuhn	581
<i>vulcanicum</i> , Kuhn	579
Botrychium	778
<i>daucifolium</i> , Wall.	778
<i>lanuginosum</i> , Wall.	778
<i>subcarnosum</i> , Wall.	778
<i>virginianum</i> , Sw.	
<i>var. lanuginosum</i>	778
<i>virginicum</i> , Willd.	
<i>var. lanuginosum</i>	778

C.

Callipteris

<i>alismaefolia</i> , J. Sm.	425
<i>cordifolia</i> , Copel.	422

Callipteris

<i>elegans</i> , J. Sm.	425
<i>esculenta</i> , J. Sm.	425
<i>pariens</i> , Copel.	424
<i>prolifera</i> , Bory	424, 829
<i>Smithiana</i> , Bedd.	425
<i>sylvatica</i> , Bory	402
Callogramme	539
<i>Callymodon cucullatus</i> , Pr. ...	589
Campyloneuron	568
Campteria	552
<i>anamallayensis</i> , Bedd.	571
<i>biaurita</i> , Hk.	571
<i>heterophlebia</i> , Moore	572
<i>Junghuhnii</i> , Reinw.	574
<i>Kleiniana</i> , Pr.	571
<i>Wallichiana</i> , Moore	572
Camptosorus	477
Cardiomanes	85
Cephalomanes	84
<i>asplenioides</i> , Pr.	101
<i>auriculatum</i> , V. D. B.	96
<i>javanicum</i> , V. D. B.	800
<i>rhomboideum</i> , V. D. B.	800
<i>Zollingeri</i> , V. D. B.	800
Ceratopteris	751
<i>thalictroides</i> , Brong.	751
Ceropteris	527, 850
<i>calomelanos</i> , Und.	528
<i>chrysosora</i> , v. A. v. R. ...	529
<i>tartarea</i> , Link	528, 850
Ceterach	
<i>pedunculata</i> , Hk. & Grev. ..	675
Cheilanthes	356
<i>alpina</i> , Bl.	355
<i>argentea</i> , Kze	337
<i>var. chrysophylla</i>	338
<i>Belangeri</i> , C. Chr.	359

Cheilanthes			
Boltoni, <i>Copel.</i>	557		
<i>chrysophylla</i> , <i>Hk.</i>	558		
<i>distans</i> , <i>Mett.</i>	484		
<i>farinosa</i> , <i>Klf.</i>	558		
<i>fragilis</i> , <i>Hk.</i>	540		
<i>gigantea</i> , <i>Ces.</i>	494		
<i>hirsuta</i> , <i>Mett.</i>	485		
<i>hirta</i> , <i>Sw.</i>	540		
<i>hispidula</i> , <i>Kze.</i>	559		
<i>multifida</i> , <i>Sw.</i>	559		
<i>pallida</i> , <i>Bl.</i>	556		
<i>parviloba</i> , <i>Sw.</i>	540		
<i>pteroides</i> , <i>Sv.</i>	545		
<i>resinifera</i> , <i>Bl.</i>	142		
<i>rufa</i> , <i>Don</i>	558		
<i>setigera</i> , <i>Bl.</i>	817		
<i>tenuifolia</i> , <i>Sw.</i>	559		
<i>varians</i> , <i>Hk.</i>	559		
Cheiroglossa	772		
Cheiropleuria	752		
<i>bicuspis</i> , <i>Pr.</i>	752		
Christensenia	767		
<i>aesculifolia</i> , <i>Max.</i>	768		
<i>Cumingiana</i> , <i>Christ</i>	768		
Christopteris	704		
<i>sagitta</i> , <i>Copel.</i>	704		
Chrysodium	755		
<i>aureum</i> , <i>Mett.</i>	750		
<i>praestantissimum</i> , <i>Mett.</i>	750		
Chrysopteris	569		
Cibotium	47, 792		
<i>Assamicum</i> , <i>Hk.</i>	795		
<i>Baranetz</i> , <i>J. Sm.</i>	795		
<i>barometz</i> , <i>J. Sm.</i>	48, 792		
<i>Cumingii</i> , <i>Kze.</i>	795		
<i>sumatranum</i> , <i>Christ</i>	795		
Cincinalis	484		
Colposoria	296		
Compositae (Polypodium)	567		
Compositae (Pteris)	552		
Coniogramme	527		
<i>fraxinea</i> , <i>Diels</i>	527		
<i>Craspedoneuron</i>			
<i>album</i> , <i>V. D. B.</i>	95		
<i>Braunii</i> , <i>V. D. B.</i>	95		
<i>pallidum</i> , <i>V. D. B.</i>	95		
<i>Crepidomanes</i>			
<i>humile</i> , <i>V. D. B.</i>	94		
Crypsinus	568		
Cryptogramme			
<i>aurata</i> , <i>Prantl</i>	550		
<i>japonica</i> , <i>Prantl</i>	551		
<i>Ctenopteris</i>			
<i>repandula</i> , <i>Kze.</i>	599		
<i>solida</i> , <i>Kze.</i>	606		
Cuneatae (Adiantum)	519		
Cyathea	15, 785		
<i>adenochlamys</i> , <i>Christ</i>	785		
<i>alternans</i> , <i>Pr.</i>	17, 784		
<i>aneitensis</i> , <i>Hk.</i>	16		
<i>arachnoidea</i> , <i>Hk.</i>	26		
<i>arborea</i> , <i>Sm.</i>			
<i>var. pallida</i>	21		
<i>assimilis</i> , <i>Hk.</i>	20		
<i>Beccariana</i> , <i>Ces.</i>	20		
<i>Brunonis</i> , <i>Wall.</i>	15, 785		
<i>callosa</i> , <i>Christ</i>	787		
<i>capensis</i> , <i>Sm.</i>	27		
<i>caudata</i> , <i>Copel.</i>	785		
<i>celebica</i> , <i>Bl.</i>	42		
<i>celebica</i> , <i>v. A. v. R.</i>	26, 788		
<i>Christii</i> , <i>Copel.</i>	785		
<i>crenulata</i> , <i>Bl.</i>	21		
<i>cyclodonta</i> , <i>v. A. v. R.</i>	19, 784		
<i>deparoides</i> , <i>Ces.</i>	626		

Cyathea		Cyathea	9
<i>dulitensis</i> , <i>Bk.</i>	16	Cyclopeltis	163, 808
<i>excelsa</i> , <i>Kze.</i>	21	<i>Presliana</i> , <i>J. Sm.</i>	164, 808
<i>ferruginea</i> , <i>Christ</i>	784	<i>semicordata</i> , <i>J. Sm.</i>	165, 808
<i>fusca</i> , <i>Bk.</i>	19	Cyclophorus	678
<i>Havilandii</i> , <i>Bk.</i>	22	<i>abbreviatus</i> , <i>C. Chr.</i>	692
<i>hymenodes</i> , <i>Mett.</i>	24	<i>acrostichoides</i> , <i>Pr.</i>	685
<i>inquinans</i> , <i>Christ</i>	23	<i>adnascens</i> , <i>Desv.</i>	681
<i>integra</i> , <i>J. Sm.</i>	20	<i>albicans</i> , <i>Pr.</i>	687, 688
<i>javanica</i> , <i>Bl.</i>	23	<i>angustatus</i> , <i>Desv.</i>	691
<i>Korthalsii</i> , <i>Mett.</i>	21	<i>anetioides</i> , <i>C. Chr.</i>	682
<i>leucophaës</i> , <i>Hassk.</i>	24	<i>argyrolepis</i> , <i>Christ</i>	694
<i>leucotricha</i> , <i>Christ</i>	18	<i>asterosorus</i> , <i>C. Chr.</i>	694
<i>Lobbiana</i> , <i>Hk.</i>	17	<i>Beddomeanus</i> , <i>C. Chr.</i>	695
<i>Loheri</i> , <i>Christ</i>	787	<i>Christii</i> , <i>C. Chr.</i>	695
<i>Macgregorii</i> , <i>Muell.</i>	17	<i>confluens</i> , <i>C. Chr.</i>	690
<i>moluccana</i> , <i>R. Br.</i>	15, 785	<i>distichocarpus</i> , <i>C. Chr.</i>	695, 694
<i>Muelleri</i> , <i>Bk.</i>	25	<i>flocciger</i> , <i>Pr.</i>	687
<i>negrosiana</i> , <i>Christ</i>	786	<i>heteractis</i> , <i>C. Chr.</i>	684
<i>oinops</i> , <i>Hassk.</i>	25	<i>lanuginosus</i> , <i>C. Chr.</i>	685
<i>orientalis</i> , <i>Moore</i>	21, 785	<i>Lauterbachii</i> , <i>C. Chr.</i>	690
<i>petiolata</i> , <i>J. Sm.</i>	20	<i>macropodus</i> , <i>C. Chr.</i>	685
<i>philippinensis</i> , <i>Bk.</i>	16, 785	<i>mollis</i> , <i>Pr.</i>	688
<i>polypoda</i> , <i>Bk.</i>	18	<i>nummularifolius</i> , <i>C. Chr.</i>	685
<i>rufopannosa</i> , <i>Christ</i>	784	<i>var. obovatus</i>	685
<i>runensis</i> , <i>v. A. v. R.</i>	22, 786	<i>obovatus</i> , <i>v. A. v. R.</i>	685
<i>saccata</i> , <i>Christ</i>	19	<i>pachydermus</i> , <i>C. Chr.</i>	680
<i>sarawakensis</i> , <i>Hk.</i>	17	<i>pannosus</i> , <i>C. Chr.</i>	685
<i>schizochlamys</i> , <i>Bk.</i>	25	<i>penangianus</i> , <i>C. Chr.</i>	686
<i>sinops</i> , <i>Rac.</i>	25	<i>porosus</i> , <i>Pr.</i>	687, 688
<i>spinulosa</i> , <i>Wall.</i>	25, 786	<i>princeps</i> , <i>C. Chr.</i>	684
<i>strigosa</i> , <i>Christ</i>	24	<i>rasamalae</i> , <i>C. Chr.</i>	686
<i>suluensis</i> , <i>Bk.</i>	18	<i>rupestris</i> , <i>C. Chr.</i>	680
<i>sumatrana</i> , <i>Bk.</i>	23	<i>samarensis</i> , <i>C. Chr.</i>	691
<i>tripinnata</i> , <i>Copcl.</i>	788	<i>Sarasinorum</i> , <i>C. Chr.</i>	695
<i>Wernerii</i> , <i>Rst.</i>	786	<i>sphaerostichus</i> , <i>C. Chr.</i>	682
<i>Zollingeriana</i> , <i>Mett.</i>	20, 785	<i>splendens</i> , <i>C. Chr.</i>	696
Cyatheaaceae	9	<i>sticticus</i> , <i>C. Chr.</i>	687

Cyclophorus

stigmatosus, <i>Desv.</i>	693
varius, <i>Gaud.</i>	682
Warburgii, <i>C. Chr.</i>	692
Cyclosorus	175
Cyrtomium	255
<i>caryotideum</i> , <i>Pr.</i>	255
<i>falcatum</i> , <i>Pr.</i>	255
Cystodium	282
<i>sorbifolium</i> , <i>J. Sm.</i>	282
Cystopteris	257
<i>tenuisecta</i> , <i>Mett.</i>	257
<i>tristis</i> , <i>Mett.</i>	44, 257

D.

Darea	457
Davallia	296, 825
<i>achilleaefolia</i> , <i>Wall.</i>	722
<i>affinis</i> , <i>Hk.</i>	295
<i>alpina</i> , <i>Bl.</i>	290
<i>amboynensis</i> , <i>Hk.</i>	515
<i>angustata</i> , <i>Wall.</i>	286
<i>asperrima</i> , <i>Ces.</i>	282
<i>Beccariana</i> , <i>Ces.</i>	501
<i>Belangeri</i> , <i>Bory</i>	291
<i>biflora</i> , <i>Klf.</i>	259
<i>bipinnatifida</i> , <i>Bk.</i>	290
<i>bipinnatifida</i> , <i>Bl.</i>	291
<i>Blumeana</i> , <i>Ces.</i>	265
<i>Blumeana</i> , <i>Hk.</i>	265
<i>borneensis</i> , <i>J. Sm.</i>	198
<i>Boryana</i> , <i>Pr.</i>	824
<i>botrychioides</i> , <i>Bk.</i>	290
<i>brevipes</i> , <i>Copel.</i>	506
<i>bullata</i> , <i>Wall.</i>	505
<i>caudata</i> , <i>Cav.</i>	505
<i>celebica</i> , <i>Christ.</i>	145
<i>chaerophylla</i> , <i>Ces.</i>	291

Davallia

<i>chaerophylla</i> , <i>Wall.</i>	292
<i>cicutarioides</i> , <i>Bk.</i>	292
<i>ciliata</i> , <i>Hk.</i>	299
<i>contigua</i> , <i>Spr.</i>	616, 852
<i>corniculata</i> , <i>Ces.</i>	501
<i>corniculata</i> , <i>Moore</i>	502
<i>Cumingii</i> , <i>Hk.</i>	289
<i>decipiens</i> , <i>Ces.</i>	259
<i>decurrens</i> , <i>Hk.</i>	502
<i>delicatula</i> , <i>Christ</i>	262
<i>Denhami</i> , <i>Hk.</i>	516
<i>denticulata</i> , <i>Mett.</i>	501, 502
<i>var. elata</i>	502
<i>deparoides</i> , <i>Ces.</i>	626
<i>dissecta</i> , <i>Moore</i>	506
<i>divaricata</i> , <i>Bl.</i>	505, 506
<i>elata</i> , <i>Spr.</i>	502
<i>elegans</i> , <i>Hedw.</i>	505
<i>elegans</i> , <i>Sw.</i>	501
<i>var. pulchra</i>	502
<i>embolostegia</i> , <i>Copel.</i>	505
<i>Emersoni</i> , <i>Hk. & Grev.</i>	615
<i>epiphylla</i> , <i>Bl.</i>	502
<i>exaltata</i> , <i>Copel.</i>	614
<i>falcinella</i> , <i>Pr.</i>	292
<i>fejeensis</i> , <i>Hk.</i>	507
<i>fijiensis</i> , <i>Hk.</i>	504, 507
<i>flaccida</i> , <i>R. Br.</i>	510
<i>flagellifera</i> , <i>Wall.</i>	515
<i>Friderici et Pauli</i> , <i>Christ</i> ...	615
<i>fructuosa</i> , <i>Christ</i>	504
<i>Gaimardiana</i> , <i>Pr.</i>	286
<i>gracilis</i> , <i>Bl.</i>	515
<i>Griffithiana</i> , <i>Hk.</i>	500, 825
<i>hemiptera</i> , <i>Bory</i>	261
<i>hirsuta</i> , <i>v. A. v. R.</i> ...	294, 299
<i>hirta</i> , <i>Klf.</i>	515

Davallia

<i>Hosei</i> , Bk.	295
<i>hymenophylloides</i> , Bk.	264
<i>hymenophylloides</i> , Kuhn.	295
<i>immersa</i> , Wall.	292
<i>inaequalis</i> , Kze	
<i>var. minor</i>	282
<i>intermarginalis</i> , Bl.	287
<i>intramarginalis</i> , Ces.	515
<i>Kingii</i> , Bk.	294
<i>Korthalsii</i> , Kuhn	500
<i>Kunzeana</i> , Hk	274
<i>Kurzii</i> , Clarke	512
<i>lanceolata</i> , Bk.	475
<i>Lapeyrousii</i> , Hk.	264
<i>lepida</i> , Pr.	289
<i>Lindleyi</i> , Hk.	505
<i>Lobbiana</i> , Moore	500
<i>lonchitidea</i> , Wall.	511
<i>longipinnula</i> , Ces.	514
<i>longula</i> , Kze	289
<i>Lorrainii</i> , Hance	505
<i>Luzonica</i> , Hk.	515
<i>manilensis</i> , Goldm.	509
<i>marginalis</i> , Bk.	509
<i>moluccana</i> , Bl.	282
<i>moluccana</i> , Rb.	515
<i>Mooreana</i> , Masters	501
<i>nephrodioides</i> , Bk.	295
<i>nodosa</i> , Hk.	295
<i>Novae Guineae</i> , Rst.	285
<i>oligophlebia</i> , Bk.	514
<i>ornata</i> , Wall.	505
<i>pallida</i> , Mett.	501
<i>parallela</i> , Wall.	287
<i>parvula</i> , Wall.	287
<i>pectinata</i> , Bl.	616
<i>pectinata</i> , Sm.	288

Davallia

<i>pedata</i> , Sm.	288
<i>pentaphylla</i> , Bl.	297
<i>philippinensis</i> , Harr.	511
<i>pinnata</i> , Cav.	514
<i>pinnatifida</i> , Bk.	288
<i>platyphylla</i> , Don	511
<i>polyantha</i> , Hk.	505
<i>polypodioides</i> , Don	
<i>var. hispida</i>	515
<i>var. pubescens</i>	510
<i>var. rhomboidea</i>	515
<i>var. subglabra</i>	510
<i>proxima</i> , Bl.	512
<i>pulchella</i> , Hk.	262
<i>pulchra</i> , Don	292
<i>pusilla</i> , Mett.	289
<i>pyxidata</i> , Cav.	505
<i>repens</i> , Desv.	261
<i>retusa</i> , Cav.	259
<i>rhomboidea</i> , Wall.	515
<i>Sarasinorum</i> , Christ	509
<i>Schlechteri</i> , C. Chr.	614
<i>serraeformis</i> , Wall.	615
<i>sessilifolia</i> , Bl.	287
<i>solida</i> , Sw.	505, 506
<i>var. fijiensis</i>	507
<i>Speluncae</i> , Bk.	146, 510
<i>var. exserta</i>	511
<i>var. trichosticha</i>	510
<i>stenoloba</i> , Bk	501
<i>stenomera</i> , Kze	504
<i>strigosa</i> , (Sw.) Kze	512
<i>var. rhomboidea</i>	515
<i>tenuifolia</i> , Sw.	259
<i>trapeziformis</i> , Rb.	515
<i>trichomanoides</i> , Bl.	504, 825
<i>trichosticha</i> , Hk.	510

- Davallia**
triphylla, *Hk.* 297
triquetra, *Bk.* 263
tristis, *Rac.* 257
Veitchii, *Bk.* 506
vestita, *Bl.* 291
villosa, *Wall.* 509
viscidula, *Mett.* 294
Wagneriana, *Copel.* 504
- Davallieae** 121
- Davallodes** 297
grammatorum, *Copel.* . 299, 500
gymnocarpum, *Copel.* . 299, 500
hirsutum, *Copel.* 299, 500
- Dennstaedtia** 140, 804
ampla, *Bedd.* 142
cuneata, *Moore* 144, 805
deltoidea, *Moore* 145
dennstaedtioides, *Copel.* 804
Elmeri, *Copel.* 805
erythrorachis, *Diels* 142, 804
flaccida, *Bernh.* 145, 805
glabrata, *C. Chr.* 145
gomphophylla, *C. Chr.* 146
Hooveri, *Christ* 805
javanica, *Christ* 145
Kingi, *Bedd.* 145
Merrilli, *Copel.* 804
moluccana, *Moore* 146, 147
remota, *Diels* 144
resinifera, *Mett.* 142
rhombofolia, *C. Chr.* 141
samoensis, *Moore* 145, 804
scabra, *Moore* 145
scandens, *Moore* 146, 147
Smithii, *Moore* 145
sumatrana, *v. A. v. R.* . 146, 806
Williamsi, *Copel.* 804
- Diacalpe** 139
aspidioides, *Bl.* 139
pseudo-caenopteris, *Kze* 139
- Diblemma**
samarensis, *J. Sm.* 651
- Dicksonia** 47, 791
ampla, *Bedd.* 142
Barometz, *Link* 48
Blumei, *Moore* 47, 791
chrysotricha, *Moore* 47
Copelandi, *Christ* 791
cuneata, *Hk.* 144
delicata, *Muell.* 145
deltoidea, *Hk.* 145
erythrorachis, *Christ* 142
flaccida, *Sw.* 145
glabrata, *Ces.* 145
gomphophylla, *Bk.* 146
grandis, *Rst.* 792
javanica, *Bl.* 145
Kingi, *Bedd.* 145
moluccana, *Bl.* 146
nephrolepioides, *Christ* 156
remota, *Christ* 144
repens, *Bory* 824
rhombofolia, *Bk.* 141
samoensis, *Bk.* 145
scabra, *Wall.* 145
scandens, *Bl.* 146
Smithii, *Hk.* 145
sorbifolia, *Sm.* 282
Zippeliana, *Kze* 146
- Dicksonieae** 9
- Dicranopteris** 56
dolosa, *Copel.* 62
- Dictyogramme** 540
- Dictyopteris** 511
ambigua, *v. A. v. R.* 521

Dictyopteris

<i>andaiensis</i> , v. A. v. R.	514
<i>Barberi</i> , Bedd.	515
<i>Beccariana</i> , v. A. v. R.	515
<i>Bolsteri</i> , v. A. v. R.	519
<i>Bryanti</i> , v. A. v. R.	514
<i>difformis</i> , Moore	520
<i>ferruginea</i> , v. A. v. R.	516
<i>Hancockii</i> , v. A. v. R.	518
<i>heterosora</i> , Bedd.	518
<i>irregularis</i> , Pr.	520
<i>labrusca</i> , v. A. v. R.	517
<i>lamoensis</i> , v. A. v. R.	517
<i>macrodongta</i> , Pr.	516
<i>pentaphylla</i> , v. A. v. R.	519
<i>petrophylla</i> , Moore	515
<i>polycarpa</i> , Bedd.	520
<i>pteroides</i> , Pr.	515
<i>saxicola</i> , v. A. v. R.	515
<i>subdecurrens</i> , v. A. v. R. .	514
<i>vitis</i> , v. A. v. R.	516
<i>Whitfordi</i> , v. A. v. R.	519
<i>Didymochlaena</i>	170
<i>lunulata</i> , Desv.	170
<i>polycarpa</i> , Bk.	252
<i>sinuosa</i> , Desv.	170
<i>truncatula</i> , J. Sm.	170
<i>Didymoglossum</i>	
<i>capillatum</i> , V. D. B.	95
<i>Filicula</i> , Desv.	95
<i>late-alatum</i> , V. D. B.	95
<i>laxum</i> , V. D. B.	94
<i>serrulatum</i> , Pr.	798
<i>Digrammaria</i>	
<i>ambigua</i> , Pr.	521
<i>robusta</i> , Fée	424
<i>Diplaziopsis</i>	159, 805
<i>javanica</i> , C. Chr.	140

<i>Diplazium</i>	595, 828
<i>accedens</i> , Bl.	424
<i>acrotis</i> , Christ	405
<i>acuminatum</i> , Bl.	409
<i>acuminatum</i> , Pr.	402
<i>aequibasale</i> , C. Chr.	404
<i>alismacfolium</i> , Pr.	545
<i>alismiifolium</i> , v. A. v. R. ..	425
<i>alternifolium</i> , Bl.	405
<i>amplissimum</i> , Diels	419
<i>angustatum</i> , Amann	425
<i>asperum</i> , Bl.	420
<i>atratum</i> , Christ	418
<i>bantamense</i> , Bl.	405
<i>barisanicum</i> , C. Chr.	404
<i>biseriale</i> , C. Chr.	405
<i>Bolsteri</i> , Copel.	415
<i>brachypodium</i> , Copel.	415
<i>brachysoroides</i> , Copel.	412
<i>Brackenridgei</i> , Diels	415
<i>brevisorum</i> , J. Sm.	829
<i>bulbiferum</i> , Brack.	415
<i>Burchardi</i> , Rst.	411
<i>camarinum</i> , Copel.	402
<i>ceramicum</i> , C. Chr.	421
<i>Ceylanicum</i> , Christ	400
<i>chlorophyllum</i> , Bedd.	401
<i>Christii</i> , C. Chr.	409
<i>confertum</i> , C. Chr.	404
<i>cordifolium</i> , Bl.	422
<i>crenato-serratum</i> , Moore ...	407
<i>crinitum</i> , C. Chr.	414
<i>cultratum</i> , Pr.	407
<i>Cumingii</i> , C. Chr.	425
<i>cyatheifolium</i> , Pr.	416
<i>davaoense</i> , Copel.	416
<i>decussatum</i> , Bedd.	408
<i>decussatum</i> , Christ	424

Diplazium

<i>deltoideum</i> , Pr.	418
<i>dilatatum</i> , Bl.	416
<i>dolichosorum</i> , Copel.	417
<i>ebeneum</i> , J. Sm.	420
<i>elegans</i> , Hk.	425
<i>esculentum</i> , Sw.	425
<i>Forbesii</i> , C. Chr.	422
<i>fraxineum</i> , Don	527
<i>fraxinifolium</i> , Bedd.	405
<i>fraxinifolium</i> , Pr.	425
<i>fructuosum</i> , Copel.	420
<i>grammitoides</i> , Pr.	409, 410, 415
<i>Hosei</i> , Christ	407
<i>Huttoni</i> , C. Chr.	421
<i>inconspicuum</i> , Christ	401
<i>japonicum</i> , Bedd.	408
<i>larutense</i> , Bedd.	406
<i>Lasiopteris</i> , Kze	411
<i>latifolium</i> , Moore	416
<i>lineolatum</i> , Bl.	425
<i>Lobbianum</i> , Moore	408
<i>manilense</i> , C. Chr.	417
<i>maximum</i> , C. Chr.	416
<i>Merrilli</i> , Copel.	400
<i>Meyenianum</i> , Pr.	418
<i>muricatum</i> , v. A. v. R.	829
<i>oligosorum</i> , Copel.	419
<i>palauanense</i> , Copel.	406
<i>pallidum</i> , Moore	405
<i>pariens</i> , C. Chr.	424
<i>Petersenii</i> , Christ	411
<i>petiolare</i> , Pr.	409
<i>polypodioides</i> , Bedd.	420
<i>polypodioides</i> , Bl.	419, 420, 456
<i>porphyrorachis</i> , Diels	401
<i>porrectum</i> , Pr.	407
<i>Prescottianum</i> , Moore	415, 828

Diplazium

<i>prionophyllum</i> , Kze	420
<i>proliferum</i> , Thouars	424, 829
<i>shepherdoides</i> , C. Chr.	408
<i>silvaticum</i> , Sw.	402, 828
<i>var. Prescottianum</i>	828
<i>Smithianum</i> , Diels	425
<i>sorzogonense</i> , Pr.	412, 415
<i>speciosum</i> , Bl.	415
<i>spinulosum</i> , Bl.	425
<i>subserratum</i> , Moore	400
<i>tabacinum</i> , Copel.	404
<i>tomentosum</i> , Bl.	414
<i>umbrosum</i> , Bedd.	828
<i>vacillans</i> , C. Chr.	405
<i>vestitum</i> , Pr.	417
<i>Weinlandii</i> , Christ	402
<i>Whitfordi</i> , Copel.	415
<i>Williamsi</i> , Copel.	409
<i>Woodii</i> , Copel.	410
<i>xiphophyllum</i> , C. Chr.	406
<i>zeylanicum</i> , Moore	400, 828
<i>Diplopterygium</i>	57
<i>Dipteris</i>	521
<i>bifurcata</i> , Christ	522
<i>conjugata</i> , Reinw.	525
<i>Dipteris</i> , Christ	525
<i>Horsfieldii</i> , Bedd.	525
<i>Lobbiana</i> , Moore	522
<i>Nieuwenhuisii</i> , Christ	522
<i>quinquefurcata</i> , Christ	525
<i>Disphenia</i>	
<i>orientalis</i> , Kze	785
<i>Doodia</i>	589
<i>dives</i> , Kze	589, 590
<i>media</i> , R. Br.	590
<i>Doryopteridastrum</i>	546
<i>Doryopteris</i>	546

Doryopteris

concolor, <i>Kuhn</i>	547
ludens, <i>J. Sm.</i>	547
Drymoglossum	702
carnosum, <i>J. Sm.</i>	702
carnosum, <i>Rac.</i>	705
confertum, <i>Copel.</i>	558
heterophyllum, <i>C. Chr.</i> .	681, 705
novo-guineae, <i>Christ</i>	705
piloselloides, <i>Pr.</i>	705
rigidum, <i>Hk.</i>	705
<i>var. longistipitatum</i>	644
Drynaria	696
<i>alata</i> , <i>Brack.</i>	658
cornucopia, <i>v. A. v. R.</i> ...	700
coronans, <i>J. Sm.</i>	670
decurrens, <i>Brack.</i>	665
descensa, <i>Copel.</i>	698
<i>Heracleum</i> , <i>Moore</i>	671
involuta, <i>v. A. v. R.</i>	700
<i>Linnaei</i> , <i>Bedd.</i>	699
<i>lomarioides</i> , <i>J. Sm.</i>	625
nectarifera, <i>Diels</i>	700
pleuridioides, <i>Pr.</i>	697, 698
propinqua, <i>J. Sm.</i>	698
quercifolia, <i>J. Sm.</i> .	698, 699, 701
revoluta, <i>J. Sm.</i>	658
rigidula, <i>Bedd.</i>	699
sparsisora, <i>Moore</i>	699
splendens, <i>Bedd.</i>	701
<i>tenuiloris</i> , <i>J. Sm.</i>	651
Drynariopsis	569
Dryopteris	174, 815
abortiva, <i>O. Ktze</i>	217
aciculata, <i>C. Chr.</i>	200
acromanes, <i>Christ</i>	505
adenophora, <i>C. Chr.</i>	214
adnata, <i>v. A. v. R.</i>	191

Dryopteris

<i>africana</i> , <i>C. Chr.</i>	498
alatella, <i>C. Chr.</i>	210
amboinensis, <i>O. Ktze</i>	219
aortisora, <i>C. Chr.</i>	501
appendiculata, <i>C. Chr.</i>	215
arbuscula, <i>O. Ktze</i>	218
arfakiana, <i>C. Chr.</i>	502
arida, <i>O. Ktze</i>	212
asperula, <i>C. Chr.</i>	495
attenuata, <i>O. Ktze</i> .	184, 185, 815
Backeri, <i>v. A. v. R.</i>	817
Bakeri, <i>Copel.</i>	208
balabacensis, <i>Christ</i>	816
basilaris, <i>C. Chr.</i>	820
Beccariana, <i>C. Chr.</i>	185
Beddomei, <i>O. Ktze</i>	191
Benoitiana, <i>v. A. v. R.</i>	225
Blumei, <i>v. A. v. R.</i>	231
Bordenii, <i>Christ</i>	822
borneensis, <i>O. Ktze</i>	198
Boryana, <i>C. Chr.</i>	201
brachyodus, <i>O. Ktze</i> ...	220, 819
brevipinna, <i>C. Chr.</i>	510
brunneo-villosa, <i>C. Chr.</i> ...	494
calcarata, <i>O. Ktze</i>	185, 228
callosa, <i>C. Chr.</i>	220, 819
cana, <i>O. Ktze</i>	489
canescens, <i>C. Chr.</i> .	251, 505, 507
<i>var. Bakeri</i>	208
<i>var. heterophylla</i>	171
<i>var. simplicifolia</i>	500
caudiculata, <i>v. A. v. R.</i> ...	225
<i>ceramica</i> , <i>v. A. v. R.</i>	506
<i>Cesatiana</i> , <i>C. Chr.</i>	509
chaerophyllifolia, <i>C. Chr.</i> ..	194
chamaeotaria, <i>Christ</i>	505

Dryopteris

<i>cochleata</i> , C. Chr.	195
<i>conterminoides</i> , C. Chr. . . .	190
<i>Copelandi</i> , Christ	492
<i>crassifolia</i> , O. Ktze	182
<i>Creaghii</i> , C. Chr.	185
<i>crenata</i> , O. Ktze	202
<i>crinipes</i> , O. Ktze	229
<i>cucullata</i> , Christ	212, 215, 819
<i>cuspidata</i> , Christ	511
<i>cyatheoides</i> , O. Ktze	216
<i>Dayi</i> , C. Chr.	188
<i>debilis</i> , C. Chr.	501
<i>didymosora</i> , C. Chr.	225
<i>dissecta</i> , O. Ktze	196
<i>diversifolia</i> , v. A. v. R.	189, 815
<i>diversiloba</i> , Christ	505
<i>dubia</i> , Copel.	817
<i>echinata</i> , O. Ktze	182
<i>edulenta</i> , O. Ktze	194
<i>eminens</i> , C. Chr.	214
<i>erubescens</i> , C. Chr.	489
<i>erythrosora</i> , O. Ktze	195, 815
<i>exigua</i> , O. Ktze	190, 814
<i>extensa</i> , O. Ktze	210
<i>ferox</i> , O. Ktze	221, 819
<i>filix mas</i> , Schott	181, 192, 814
<i>var. adnata</i>	191
<i>firmula</i> , C. Chr.	501
<i>flaccida</i> , O. Ktze	195, 198, 815
<i>Foxii</i> , Christ	814
<i>glandulosa</i> , O. Ktze	215, 819
<i>glaucostipes</i> , C. Chr.	226
<i>gongyloides</i> , O. Ktze	212
<i>gracilescens</i> , O. Ktze	186
<i>grammitoides</i> , C. Chr.	181
<i>granulosa</i> , C. Chr.	505
<i>gymnopoda</i> , C. Chr.	182

Dryopteris

<i>Haenkeana</i> , O. Ktze	214
<i>Hallieri</i> , C. Chr.	184
<i>Hasseltii</i> , C. Chr.	494
<i>heleopteroides</i> , Christ	814
<i>heterocarpa</i> , O. Ktze	228
<i>hirsuta</i> , O. Ktze	214, 819
<i>hirtipes</i> , O. Ktze	180
<i>hispidula</i> , O. Ktze	228
<i>holophylla</i> , C. Chr.	500
<i>Hosei</i> , C. Chr.	510
<i>immersa</i> , O. Ktze	188
<i>var. ligulata</i>	815
<i>imponens</i> , C. Chr.	506
<i>indica</i> , v. A. v. R.	224
<i>intermedia</i> , O. Ktze	199, 816
<i>invisa</i> , O. Ktze	212, 819
<i>Keraudreniana</i> , C. Chr.	496
<i>Kingii</i> , C. Chr.	497
<i>Koordersii</i> , C. Chr.	181
<i>labuanensis</i> , C. Chr.	500
<i>laevis</i> , C. Chr.	220
<i>larutensis</i> , C. Chr.	252
<i>laserpitiifolia</i> , C. Chr.	494
<i>latebrosa</i> , C. Chr.	221
<i>latipinna</i> , O. Ktze	217
<i>ligulata</i> , O. Ktze	184, 815
<i>lineata</i> , C. Chr.	209
<i>Loheriana</i> , C. Chr.	221, 820
<i>longipes</i> , O. Ktze	211
<i>loxoscapoides</i> , C. Chr.	495
<i>Luerssenii</i> , C. Chr.	190
<i>luzonica</i> , Christ	821
<i>marginata</i> , Christ	814
<i>megaphylla</i> , C. Chr.	218, 219
<i>Merrillii</i> , Christ	505
<i>Metteniana</i> , Hier.	815
<i>microchlamys</i> , v. A. v. R.	226

Dryopteris

<i>microloncha</i> , Christ	820
<i>Millettii</i> , C. Chr.	198
<i>mindanaensis</i> , Christ	818
<i>monodonta</i> , C. Chr.	187
<i>Molleyana</i> , C. Chr.	182
<i>moulmeinensis</i> , C. Chr. ...	217
<i>multilineata</i> , O. Ktze	219
<i>multisetata</i> , C. Chr.	205
<i>nephrodioides</i> , O. Ktze.	251
<i>obscura</i> , O. Ktze	492
<i>obtusiloba</i> , O. Ktze	205
<i>ochtodes</i> , C. Chr.	215
<i>oosora</i> , C. Chr.	250
<i>opaca</i> , C. Chr.	498
<i>orbicularis</i> , C. Chr.	251
<i>orientalis</i> , C. Chr.	185
<i>var. Webbiana</i>	815
<i>ornata</i> , C. Chr.	495
<i>otaria</i> , O. Ktze	209
<i>oxyodus</i> , C. Chr.	491
<i>padangensis</i> , C. Chr.	198
<i>papuana</i> , C. Chr.	196
<i>parasitica</i> , O. Ktze . 211, 217, 224,	250, 821
<i>var. amboinensis</i>	219
<i>var. latipinna</i>	217
<i>patens</i> , O. Ktze	189, 814
<i>pectinata</i> , v. A. v. R. . 185, 813	
<i>Peekeli</i> , v. A. v. R. ... 188, 813	
<i>penangiana</i> , C. Chr.	229
<i>pennigera</i> , C. Chr.	504
<i>perakensis</i> , C. Chr.	226
<i>peranemiformis</i> , C. Chr.	
<i>var. obtusiloba</i>	205
<i>phaeostigma</i> , C. Chr.	181
<i>phegopteroidea</i> , C. Chr.	195
<i>philippina</i> , C. Chr.	190

FILICES.

Dryopteris

<i>pilosiuscula</i> , C. Chr. ... 185, 215	
<i>piloso-squamata</i> , v. A. v. R. . 222	
.....	820
<i>polycarpa</i> , Christ	822
<i>polytricha</i> v. A. v. R.	187
<i>Preslii</i> , O. Ktze	194
<i>procurrens</i> , O. Ktze ... 211, 819	
<i>producta</i> , C. Chr.	250
<i>prolifera</i> , C. Chr.	504
<i>pseudoreptans</i> , C. Chr.	225
<i>pteroides</i> , O. Ktze	209, 818
<i>pulvinulifera</i> , O. Ktze	200
<i>punctata</i> , C. Chr. 535, 495	
<i>purpurascens</i> , v. A. v. R. .. 201	
<i>quadriaurita</i> , Christ	489
<i>Raciborskii</i> , v. A. v. R. . 197, 816	
<i>Ramosii</i> , Christ	502
<i>recedens</i> , O. Ktze	198, 816
<i>rhodolepis</i> , C. Chr. . 199, 202, 817	
<i>var. sawarakensis</i>	200
<i>Riedleana</i> , v. A. v. R.	250
<i>rizalensis</i> , Christ	815
<i>rubida</i> , C. Chr.	502
<i>rufescens</i> , C. Chr.	495
<i>sagenioides</i> , O. Ktze	191
<i>var. obscura</i>	492
<i>sagittifolia</i> , O. Ktze	222
<i>sakayensis</i> , C. Chr.	224
<i>salicifolia</i> , C. Chr.	511
<i>sawarakensis</i> , v. A. v. R. .. 200	
<i>setigera</i> , O. Ktze .. 57, 202, 817	
<i>setosa</i> , C. Chr.	492
<i>setosa</i> , O. Ktze	198
<i>simillima</i> , C. Chr.	225
<i>simplicifolia</i> , Christ	500
<i>singalanensis</i> , C. Chr.	192
<i>Smithii</i> , v. A. v. R.	490

Euhypolepis	535
Eulindsaya	260
Eulygodium	109
Eumonogramma	551
Eunephrolepis	156
Eunotholaena	485
Eu-odontosoria	258
Eu-ophioglossum	772
Eu-osmunda	755
Euphegopteris	486
Euplatycerium	707
Eupodium	765
Eupolybotrya	725
Eupolypodium	567
Eupteris	552
Eusaccoloma	282
Euschizaea	115
Euschizoloma	276
<i>Euscolopendrium</i>	477
Eutrichomanes	85
Euvittaria	555
Euwoodwardia	590

F.

Féea	85
Furcatae (Polypodium)	567

G.

Gleichenia	55, 794
amboinensis, v. A. v. R.	62, 796
arachnoidea, Mett.	58
bifurcata, Bl.	59
candida, Rst.	796
circinata, Sw.	56, 794
crassifolia, Copel.	795
dicarpa, R. Br.	794
var. <i>a'pina</i>	56

Gleichenia	
<i>dichotoma</i> , Hk.	59
dolosa, C. Chr.	62
excelsa, J. Sm.	795
ferruginea, Bl.	59
flabellata, R. Br.	60
flagellaris, Spr.	59
gigantea, Wall.	795
glauca, Hk.	58, 794
var. <i>arachnoidea</i>	58
Hallieri, Christ	61
Hermanni, R. Br.	59
hecistophylla, A. Cunn.	56
hirta, Bl.	60
hispida, Mett.	61, 796
Koordersii, Christ	60
laevigata, Hk.	59, 769
laevissima, Christ	795
linearis, Clarke	59, 795
Loheri, Christ	796
longissima, Bl.	58, 794, 795
var. <i>arachnoides</i>	58
var. <i>rigida</i>	58
microphylla, R. Br.	56
Norrisii, Mett.	58
oceanica, Kuhn	60
rigida, J. Sm.	795
semivestita, Lab.	56, 794
vestita, Bl.	61, 796
vulcanica, B.	56
Warburgii, Christ	58, 795
Gleicheniaceae	55
<i>Gnoophora</i>	
<i>glauca</i> , Bl.	41
<i>lurida</i> , Bl.	791
<i>tomentosa</i> , Bl.	791
Goniophlebium	568
<i>argutum</i> , J. Sm.	621

Goniophlebium

<i>cuspidatum</i> , Pr.	619
<i>Korthalsii</i> , Bedd.	618
<i>molle</i> , Bedd.	621
<i>Prainii</i> , Bedd.	617
<i>Reinwardtii</i> , de Vr.	622
<i>serratifolium</i> , Brack.	622
<i>subauriculatum</i> , Pr.	622
<i>verrucosum</i> , J. Sm.	619
<i>Goniopteris</i>	486
<i>lineata</i> , Bedd.	217
<i>multilineata</i> , Bedd.	217
<i>penangiana</i> , Bedd.	229
<i>rubida</i> , J. Sm.	502
<i>Gonocormus</i>	85
<i>diffusus</i> , V. D. B.	90
<i>minutus</i> , V. D. B.	90
<i>palmaris</i> , V. D. B.	90
<i>Teysmannii</i> , V. D. B.	91
<i>Grammitideae</i>	122
<i>Grammitis</i>	567
<i>adpersa</i> , Bl.	578
<i>australis</i> , R. Br.	578
<i>blechnoides</i> , Grev.	605
<i>Billardieri</i> , Willd.	578
<i>bisulcata</i> , Hk.	576
<i>clavifer</i> , Hk.	608
<i>congener</i> , Bl.	582
<i>cucullata</i> , Bl.	589
<i>decurrens</i> , Wall.	677
<i>denticulata</i> , Bl.	590
<i>fasciata</i> , Bl.	579
<i>fasciculata</i> , Bl.	587
<i>heterocarpa</i> , Bl.	676
<i>hirta</i> , Bl.	586
<i>longifolia</i> , Bl.	658
<i>macrophylla</i> , Bl.	674
<i>obscura</i> , Bl.	586

Grammitis

<i>pilosiuscula</i> , Bl.	580
<i>pubinervis</i> , Bl.	586
<i>pusilla</i> , Bl.	
<i>var. alpestris</i>	582
<i>var. lasiosora</i>	580
<i>Reinwardtii</i> , Bl.	583
<i>sessilifolia</i> , J. Sm.	577
<i>setosa</i> , Bl.	585
<i>subpinnatifida</i> , Bl.	590
<i>Totta</i> , Pr.	497
<i>Gymnogramme</i>	
<i>acuminata</i> , Bk.	673
<i>alismaefolia</i> , Hk.	543
<i>ambigua</i> , Hk.	521
<i>appendiculata</i> , Bl.	213
<i>aspidioides</i> , Hk.	508
<i>avenia</i> , Bk.	538
<i>Boerlageana</i> , Christ	528
<i>borneensis</i> , Hk.	541
<i>calomelanos</i> , Klf.	528
<i>var. chrysophylla</i>	528
<i>campyloneuroides</i> , Bk.	674
<i>canescens</i> , Bl.	507
<i>cartilaginea</i> , Bk.	542
<i>chrysophylla</i> , Klf.	528
<i>chrysosora</i> , Bk.	529
<i>Dayi</i> , Bedd.	541
<i>decurrens</i> , Hk.	677
<i>edulis</i> , Ces.	425
<i>elliptica</i> , Bk.	677
<i>Féei</i> , Hk.	675
<i>fraxinea</i> , Bedd.	527
<i>grandis</i> , Rac.	537
<i>Hamiltoniana</i> , Hk.	675
<i>heterocarpa</i> , Hk.	676
<i>iridifolia</i> , Christ	539
<i>involuta</i> , Hk.	537

- Gymnogramme*
- javanica*, Bl. 527
- lanceolata*, Hk. 538
- Lobbiana*, Hk. 541
- Lowii*, Hk. & Grev. 497
- macrophylla*, Hk. 674
- Maingayi*, Bk. 677
- membranacea*, Hk. 675
- obtusata*, Bl. 498
- obtusifolia*, Hk. 541
- ochracea*, Pr. 528
- opaca*, Spr. 498
- pinnata*, Hk. 546
- pteridiformis*, Ces. 484
- quinata*, Hk. 545
- regularis*, Bk. 675
- serrulata*, Bl. 527
- spuria*, Bk. 674
- Stegnogramme*, Bl. 508
- tartarea*, Desv. 528
- var. ochracea* 528
- Totta*, Schl. 497
- valleculata*, Bk. 540
- vittaeformis*, Mett. 545
- Wallichii*, Hk. 545
- Gymnogrammeae* 122
- Gymnopteris* 755
- aurita*, Keys. 726
- axillaris*, Pr. 745
- contaminans*, Bedd. 741
- costata*, Bedd. 745
- decurrens*, Hk. 736
- decurrens*, Moore 756
- Feei*, Moore 745
- flagellifera*, Bedd. 759
- Hollrungii*, Kuhn 757
- inconstans*, Copel. 758
- lanceolata*, Bedd. 745
- Gymnopteris*
- latifolia*, Pr. 757
- Linnaeana*, Christ 755
- lomarioides*, Christ 746
- minor*, Hk. 744
- minus*, Bedd. 744
- Naumannii*, Diels 742
- normale*, J. Sm. 745
- platyrhynchos*, J. Sm. 729
- Prestiana*, J. Sm. 740
- quercifolia*, Bernh. 757
- repanda*, Christ 741
- subquinquefida*, Pr. 757
- subrepanda*, J. Sm. 742
- taccaefolia*, J. Sm. 757
- trilobata*, J. Sm. 737
- variabilis*, Bedd. 756
- var. axillaris* 745
- var. lanceolata* 745
- Vespertilio*, Hk. 752
- virens*, Keys. 741
- Gymnosphaera*
- glabra*, Bl. 789
- squamulata*, Bl. 789
- squamulata*, J. Sm. 55
- Gymnotheca* 765
- Gyrosorium*
- samarense*, Pr. 691
- H.**
- Habrodictyon* 85
- Cumingii*, V. D. B. 99
- Helmintostachys* 777
- dulcis*, Klf. 777
- zeylanica*, Hk. 777
- Hemidictyum* 457
- Finlaysonianum*, Moore 476

- Hemigramma 851
 latifolia, Copel. 757, 851
 Zollingeri, Diels 851
 Hemionitis 546, 851
 arifolia, Moore 547
 cordata, Ilk. & Grev. 547
 cordifolia, Rb. 547
 coriacea, Don 554
 elongata, Brack. 546
 esculenta, Retz 425
 gymnopteroidea, Copel. 851
 Hosei, Bk. 545
 immersa, Bory 551
 lanceolata, Hk. 545
 opaca, Don 498
 palmata, L. 547
 plantaginea, Cav. 555
 prolifera, Retz 504
 reticulata, Forst. 555
 Zollingeri, Kurz 851
 Hemiplebium 83
 Hemipteris 575
 Wernerii, Rst. 576
 Hemitelia 27, 788
 alternans, Hk. 17
 capensis, R. Br. 27, 788
 crenulata, Mett. 27
 Junghuhniana, Mett. 28
 latebrosa, Mett. 58
 sumatrana, v. A. v. R. . 28, 788
 truncata, Christ 42
 Heteronevron
 Naumannii, Kuhn 742
 Prestianum, Fée 740
 sculpturatum, Fée 745
 Heteropterygium 57
 Hewardia 519
 Histiopteris 549
 incisa, J. Sm. 549
 montana, Copel. 549
 Holopterygium 57
 Humata 284, 824
 alpina, Moore 289, 290, 291
 angustata, J. Sm. 286
 botrychioides, Brack. 290
 cicutarioides, Diels 292
 Cumingii, Brack. 289
 falcinella, Copel. 292
 Gaimardiana, J. Sm. ... 286, 824
 heterophylla, Desv. 286
 var. angustata 286
 Hosei, v. A. v. R. 295
 hymenophylloides, Copel. ... 295
 immersa, Mett. 292
 intermedia, C. Chr. 288
 Kingii, Diels 294
 lepida, Moore 289
 neoguineensis, C. Chr. 290
 nephrodioides, v. A. v. R. . 295
 parallela, Brack. 286
 parvula, Mett. 287, 824
 pectinata, Desv. 288
 pedata, J. Sm. 288
 pinnatifida, Bedd. 288
 polypodioides, Brack. 287
 pulchra, Diels 292
 pusilla, Carr. 289
 repens, Diels . 288, 289, 290, 291
 var. alpina 290
 sessilifolia, Mett. 287
 vestita, Moore 291
 viscidula, v. A. v. R. 294
 Hydroglossum 109
 Hymenolepis 727
 brachystachys, J. Sm. 729

Hymenolepis

<i>callifolia</i> , <i>Christ</i>	729
<i>platyrhynchos</i> , <i>Kze</i>	729
<i>revoluta</i> , <i>Bl.</i>	728
<i>rigidissima</i> , <i>Christ</i>	728
<i>spicata</i> , <i>Pr.</i>	728, 729
<i>var. brachystachys</i>	729
Hymenophyllaceae	65
Hymenophyllum	65, 797
<i>aculeatum</i> , <i>Bedd.</i>	81, 82
<i>aculeatum</i> , <i>Rac.</i>	81, 82, 798
<i>affine</i> , <i>Rac.</i>	78
<i>atrovirens</i> , <i>Colen.</i>	69
<i>australe</i> , <i>Willd.</i>	69, 797
<i>badium</i> , <i>Hk. & Grev.</i>	70
<i>bismarckianum</i> , <i>Christ</i>	82
<i>blandum</i> , <i>Rac.</i>	78
<i>Blumeanum</i> , <i>Spr.</i>	71, 797
<i>borneense</i> , <i>Hk.</i>	74
<i>brachyglossum</i> , <i>A. Br.</i>	82
<i>campanulatum</i> , <i>Christ</i>	798
<i>crispatum</i> , <i>Wall.</i>	797
<i>demissum</i> , <i>Sw.</i>	69, 797
<i>denticulatum</i> , <i>Sw.</i>	81, 82
<i>dilatatum</i> , <i>Bl.</i>	75
<i>dilatatum</i> , <i>Sw.</i>	68, 72
<i>discosum</i> , <i>Christ</i>	67
<i>edulentum</i> , <i>C. Chr.</i>	76
<i>erosum</i> , <i>Bl.</i>	69
<i>eximium</i> , <i>Kze</i>	68
<i>fimbriatum</i> , <i>J. Sm.</i>	69
<i>flabellatum</i> , <i>Lab.</i>	68
<i>formosum</i> , <i>Brack.</i>	72
<i>fraternum</i> , <i>Harr.</i>	80
<i>fuscum</i> , <i>V. D. B.</i>	75
<i>halconense</i> , <i>Copel.</i>	797
<i>holochilum</i> , <i>C. Chr.</i>	78
<i>Junghuhnii</i> , <i>V. D. B.</i>	75

Hymenophyllum

<i>integrum</i> , <i>V. D. B.</i>	71
<i>javanicum</i> , <i>Spr.</i>	69
<i>var. badium</i>	70
<i>klabatense</i> , <i>Christ</i>	79
<i>Kurzii</i> , <i>Praatl</i>	77, 797
<i>Lobbii</i> , <i>Moore</i>	77
<i>Merrillii</i> , <i>Christ</i>	798
<i>micranthum</i> , <i>V. D. B.</i>	69
<i>modestum</i> , <i>V. D. B.</i>	68
<i>multifidum</i> , <i>Sw.</i>	79, 798
<i>Neesii</i> , <i>Hk.</i>	81
<i>nitens</i> , <i>R. Br.</i>	68
<i>obtusum</i> , <i>Hk. & Arn.</i>	74
<i>ooides</i> , <i>Muell. & Bk.</i>	67
<i>pachydermicum</i> , <i>Ces.</i>	75
<i>paniculiflorum</i> , <i>Pr.</i>	70
<i>pedicularifolium</i> , <i>Ces.</i>	74
<i>physocarpum</i> , <i>Christ</i>	80
<i>polyanthos</i> , <i>Sw.</i>	71
<i>var. dentatum</i>	79
<i>praetervisum</i> , <i>Christ</i>	79
<i>productum</i> , <i>Kze</i>	79
<i>pycnocarpum</i> , <i>V. D. B.</i>	71, 797
<i>rarum</i> , <i>R. Br.</i>	72
<i>Reinwardtii</i> , <i>V. D. B.</i>	80
<i>ringens</i> , <i>Christ</i>	70
<i>sabinifolium</i> , <i>Bk.</i>	81
<i>salakense</i> , <i>Rac.</i>	71
<i>semibivalve</i> , <i>Hk. & Grev.</i>	72
<i>serrulatum</i> , <i>C. Chr.</i>	97, 798
<i>Smithii</i> , <i>Hk.</i>	78, 79
<i>Steerei</i> , <i>C. Chr.</i>	80
<i>subdemissum</i> , <i>Christ</i>	797
<i>subflabellatum</i> , <i>Ces.</i>	76
<i>tusmanicum</i> , <i>V. D. B.</i>	69
<i>thuidium</i> , <i>Harr.</i>	71
<i>Treubii</i> , <i>Rac.</i>	72

Hymenophyllum

- tunbridgense, Sm. 78, 797
 violaceum, Meyen 79
 Zollingerianum, Kze 73

Hypodematium

- phegopteroideum, Kuhn 195

Hypolepis 554, 826

- alpina, Hk. 555
 pallida, Hk. 556
 punctata, Bedd. 555
 punctata, Mett. 495
 resinifera, Hk. 142
 rugulosa, J. Sm. 495
 tenuifolia, Bernh. 556, 826

I.

Integrifoliae (Polypodium) 567

Isoloma 276

- lanuginosum, J. Sm. 157

K.

Kaulfussia 767

- aesculifolia, Bl. 768

- assamica, Griff. 768

L.

Lacostea 85

Lastraea 175

- amabilis Moore 168

- aristata, Moore 169

- attenuata, J. Sm. 815

- Blumei, Moore 202

- borneensis, Hk. 198

- Boryana, Moore 201

- calcarata, Moore 185

- var. sericea 186

- cochleata, Moore 193

- coniifolia, Moore 169

Lastraea

- crassifolia, Moore 182

- crenata, Bedd. 202

- Dayi, Bedd. 188

- dissecta, Carr. 196

- divisa, Moore 201

- elongata, Pr. 814

- eriocarpa, Pr. 202

- exigua, J. Sm. 814

- falciloba, Hk. 183

- Filix-mas, Pr. 192

- var. elongata 814

- flaccida, Bedd. 817

- flaccida, Moore 815

- fuscipes, Moore 191

- gracilescens, Bedd. 191

- gracilescens Moore 186

- hirtipes, Moore 180

- immersa, Moore 188

- ligulata, J. Sm. 815

- melanopus, Bedd. 191

- membranifolia, Bedd. 196

- nephrodioides, Bedd. 182

- obtusiloba, Bedd. 205

- odontoloma, Moore 192

- padangensis, Bedd. 198

- platypus, Moore 810

- Presliana, J. Sm. 808

- pulvinulifera, Bedd. 200

- recedens, J. Sm. 198

- sericea, Scott 186

- singalanensis, Bedd. 192

- sparsa, Moore 197

- spectabilis, J. Sm. 195

- splendens, Bedd. 195

- syrmatica, Moore 195

- tenericaulis, Moore 202

- unidentata, Bedd. 187

- Lastraea*
viscosa, J. Sm. 186
- Lecanopteris* 568
carnosa, Bl. 625
Curtisii, Bk. 627
deparoides, Bk. 626
incurvata, Bk. 627
Macleayi, Bk. 627
Nieuwenhuisii, Christ 626
pumila, Bl. 625
- Lepisorus* 569
- Leptochilus* 755
antrophyoides, C. Chr. 744
axillaris, Klf. 745
celebicus, C. Chr. 741
cuspidatus, C. Chr. 741
decurrens, Bl. 756
diversifolius, C. Chr. 740
exculptus, C. Chr. 740
exsculptus, C. Chr. 740
Harlandii, C. Chr. 756
heteroclitus, C. Chr. 759
var. inconstans 758
var. Linnaeanus 755
Hollrungii, C. Chr. 757
hydrophyllus, Copel. 744
inconstans, C. Chr. 758
latifolius, C. Chr. 757
lanceolatus, Fée 745
Linnaeanus, Fée 735
lomarioides, Bl. 746, 747
minor, Fée 744
modestus, C. Chr. 745
Naumanni, C. Chr. 742
normalis, Copel. 744, 745
oligodictyus, C. Chr. 755
perakensis, C. Chr. 747
Preslianus, C. Chr. 740
- Leptochilus*
prolifer, C. Chr. 758
Raapii, v. A. v. R. 746
rizalianus, Christ 744
scalpturatus, C. Chr. 745
stolonifer, Christ 759
trifidus, v. A. v. R. 756
virens, C. Chr. 741
zeylanicus, C. Chr. 757
Zollingeri, Fée 742
- Leptoclonium* 66
acanthoides, V. D. B. 81
aculeatum, V. D. B. 82
affine, V. D. B. 78
Braunii, V. D. B. 82
denticulatum, V. D. B. 81
edulentum, V. D. B. 76
holochilum, V. D. B. 78
Neesii, V. D. B. 81
violaceum, V. D. B. 79
- Leptogramma(e)* 486
opaca, Bedd. 498
Totta, J. Sm. 497
- Leptolepia* 285
novae guineae, v. A. v. R. 285
- Leptomanes* 84
- Leptopteris* 756
alpina, C. Chr. 757
Wilkesiana, Christ 757
- Leucostegia* 284
hirsuta, J. Sm. 299
hymenophylloides, Bl. 294
immersa, Pr. 295
pulchra, J. Sm. 292
- Lindsaya* 260, 824
acutifolia, Desv. 157
adiantoides, J. Sm. 268
adiantoides, Kuhn 262

Lindsaya

<i>ambigens</i> , <i>Ces.</i>	275
<i>apiculata</i> , <i>Kze</i>	269
<i>apoensis</i> , <i>Copel</i>	265
<i>azurea</i> , <i>Christ</i>	275
<i>blanda</i> , <i>Mett.</i>	262
<i>Blumeana</i> , <i>Kuhn</i>	265
<i>borneensis</i> , <i>Hk.</i>	271
<i>capillacea</i> , <i>Christ</i>	264
<i>caudata</i> , <i>Hk.</i>	275
<i>chinensis</i> , <i>Mett.</i>	259
<i>concinna</i> , <i>J. Sm.</i>	262, 267
<i>Copelandi</i> , <i>C. Chr.</i>	267
<i>cordata</i> , <i>Gaud.</i>	279
<i>crassipes</i> , <i>Rst.</i>	269
<i>crispa</i> , <i>Bk.</i>	268
<i>cultrata</i> , <i>Sw.</i>	269, 824
<i>cuneifolia</i> , <i>Pr.</i>	259
<i>cyathicola</i> , <i>Copel.</i>	265
<i>davallioides</i> , <i>Bl.</i>	274, 275
<i>decomposita</i> , <i>Willd.</i> ...	274, 275
<i>delicatula</i> , <i>Christ</i>	262
<i>divergens</i> , <i>Hk. & Grev.</i>	278
<i>ensifolia</i> , <i>Sw.</i>	280
<i>falcata</i> , <i>Dry.</i>	268
<i>flabellulata</i> , <i>Dry.</i>	270
<i>gomphophylla</i> , <i>Bk.</i>	270
<i>gracilis</i> , <i>Bl.</i>	261, 267, 824
<i>gracillima</i> , <i>Copel.</i>	267
<i>Gueriniana</i> , <i>Gaud.</i>	280
<i>Haviceii</i> , <i>Copel.</i>	265
<i>heterophylla</i> , <i>Dry.</i>	281
<i>Hosei</i> , <i>C. Chr.</i>	280
<i>humilis</i> , <i>Kuhn</i>	268
<i>hymenophylloides</i> , <i>Bl.</i>	264
<i>impressa</i> , <i>Christ</i>	272
<i>indurata</i> , <i>Bk.</i>	277
<i>Jamesonioides</i> , <i>Bk.</i>	276

Lindsaya

<i>lancea</i> , <i>Bedd.</i>	275
<i>lanuginosa</i> , <i>Hk.</i>	157
<i>Lapeyroussii</i> , <i>Bk.</i>	264
<i>Leprieurii</i> , <i>Hk.</i>	268
<i>lobata</i> , <i>Poir.</i>	274
<i>Lobbiana</i> , <i>Hk.</i>	267, 824
<i>Loheriana</i> , <i>Christ</i>	267
<i>longissima</i> , <i>Christ</i>	272
<i>Merrilli</i> , <i>Copel.</i>	264
<i>montana</i> , <i>Copel.</i>	267
<i>natunae</i> , <i>Bk.</i>	272
<i>nitens</i> , <i>Bl.</i>	274
<i>oblongifolia</i> , <i>Reinw.</i>	269
<i>orbiculata</i> , <i>Mett.</i> ..	267, 270, 281
<i>ovata</i> , <i>J. Sm.</i>	277
<i>pectinata</i> , <i>Bl.</i> ..	261, 264, 269, 270
<i>pellaeiformis</i> , <i>Christ</i>	278
<i>polymorpha</i> , <i>Wall.</i>	270
<i>propinqua</i> , <i>Hk.</i>	274
<i>pulchella</i> , <i>Mett.</i>	262
<i>recurvata</i> , <i>Wall.</i>	274
<i>repens</i> , <i>Bedd.</i>	261, 824
<i>repens</i> , <i>Kze</i>	296, 270
<i>retusa</i> , <i>Mett</i>	259
<i>rigida</i> , <i>J. Sm.</i>	271
<i>Sarasinorum</i> , <i>Christ</i>	275
<i>scandens</i> , <i>Hk.</i>	270, 824
<i>striata</i> , <i>Bl.</i>	270
<i>tenera</i> , <i>Dry.</i>	270
<i>tenuifolia</i> , <i>Bl.</i>	265
<i>tenuifolia</i> , <i>Mett.</i>	259
<i>trapeziformis</i> , <i>Dry.</i>	268, 275
<i>tricrenata</i> , <i>Bk.</i>	271
<i>trilobata</i> , <i>Bk.</i>	280
<i>triquetra</i> , <i>Bk.</i>	265
<i>Walkeræ</i> , <i>Hk.</i>	278

- Lindsaya
 Weneri, *Rst.* 274
- Lindsayopsis 156
- Litobrochia 552
aurita, *Pr.* 549
incisa, *Pr.* 549
ludens, *Bedd.* 547
marginata, *Pr.* 575
microdictyon, *Fée* 574
pedata, *Bedd.* 547
tripartita, *Pr.* 575
- Lomagamma(e) 755
articulata, *Copel.* 725
lomarioides, *J. Sm.* 746
perakensis, *Bedd.* 747
pteroides, *J. Sm.* 746
- Lomaria 578, 828
aculeata, *Bl.* 722
adnata, *Bl.* 542
areolaris, *Harr.* 855
articulata, *Muell.* 545
auriculata, *Bl.* 587
capensis, *Willd.* 581
concinna, *Bk.* 541
egenolfioides, *Bl.* 580
elongata, *Bl.* 579, 828
 var. Cumingiana 579
euphlebia, *Kze* 545
Fraseri, *A. Cunn.* 581
glauca, *Bl.* 542
gracilis, *Bl.* 722
Patersoni, *Spr.* 579, 828
procera, *Spr.* 581
punctulata, *Kze.* 580
pycnophylla, *Kze* 545
speciosa, *Bl.* 751
spectabilis, *Kze* 720
stenoptera, *Bk.* 541
- Lomaria
vestita, *Bl.* 581
vulcanica, *Bl.* 579
- Lomaria* (*Blechnum* §) 578
- Lomariopsis 718
palustris, *Kuhn* 720
sorbifolia *Fée* 720
- Lonchitis
tenuifolia, *Forst.* 826
- Lophidium 115
- Lorinseria 590
- Loxogramme 556
avenia, *Pr.* 558
Blumeana, *Pr.* 558
conferta, *Copel.* 558
coriacea, *Pr.* 558
grandis, *Copel.* 557
involuta, *Pr.* 557
iridifolia, *Copel.* 559
lanceolata, *Pr.* 558
parallela, *Copel.* 557
- Loxoscapha 457
- Luerssenia 255
Kehdingiana, *Kuhn* 255
- Lygodium 109, 801
altum, *v. A. v. R.* 114
basilanicum, *Christ* 802
circinatum, *Sw.* .. 111, 112, 802
 var. trifurcatum 112
dichotomum, *Sw.* 111, 112
digitatum, *Pr.* 112
flexuosum, *Sw.* ... 111, 114, 802
 var. alta 114
japonicum, *Sw.* 114, 802
Matthewi, *Copel.* 805
Mearnsii, *Copel.* 802
Merrilli, *Copel.* 805
microphyllum, *R. Br.* 115

Lygodium

- pinnatifidum*, Prantl 115
pinnatifidum, Sw. 115, 114
polystachyum, Wall. 115
salicifolium, Pr. 115
scandens, Sw. 115, 802
semihastatum, Desv. 111
Teysmannii, v. A. v. R. . 111, 801
trifurcatum, Bk. 112, 802

M.

- Marattia** 765
alata, Sm. 766
callioidus, de Vr. 765
fraxinea, Sm. 765
pellucida, Pr. 765
sambucifolia, Christ 764
sambucina, Bl. 764
silvatica, Bl. 765
Smithii, Mett. 766
ternatea, de Vr. 766
vestita, Christ 764
Wernerii, Rst. 767
- Marattiaceae** 761
- Marginaria** 568
nummularia, Pr. 654
- Matonia** 51
pectinata, R. Br. 51
sarmentosa, Bk. 51
- Matoniaceae** 51
- Meniscium** 486
Beccarianum, Ces. 509
cuspidatum, Bl. 511
Hosei, Bk. 510
proliferum, Sw. 504, 510
salicifolium, Bk. 510
stenophyllum, Bk. 510
triphyllum, Sw. 509, 850
- Mertensia** 56
crassifolia, Pr. 795
laevigata, Willd. 796
- Mesocarpus** 763
- Mesochlaena** 251, 822
javanica, R. Br. 252
larutensis, v. A. v. R. . 252, 822
polycarpa, Bedd. . 185, 252, 822
rar. microchlamys 226
- Mesosorus** 765
- Microgonium**
bimarginatum, V. D. B. ... 86
Motleyi, V. D. B. 85
sublimbatum, V. D. B. 86
- Microlepia** 507, 825
ciliata, Copel. 299
dennstaedtioides, Copel. 804
exserta, Mett. 511
firmula, C. Chr. 508
hirsuta, Pr. 299
hirta, Pr. 515
Kurzii, Bedd. 512
manilensis, Pr. 509
marginalis, Bedd. 509, 825
marginata, C. Chr. 509
mohuccana, Bedd. 285
philippinensis, Copel. 511
pinnata, J. Sm. 515
var. gracilis 515
var. Luzonica 515
platyphylla, J. Sm. 511, 512
polypodioides, Bedd. 515
proxima, Pr. 512
rhomboidea, Pr. 515
sablanensis, Christ 825
scaberula, Mett. 509
scabra, J. Sm. 509
speluncae, Moore . 510, 515, 825

Microlepia

<i>strigosa</i> , Pr.	312, 515, 825
<i>trapeziformis</i> , Kuhn	515
<i>trichosticha</i> , J. Sm.	510
<i>urophylla</i> , Moore	825
Monachosorum	485
<i>subdigitatum</i> , Kuhn	485
Monogramma	551
<i>dareicarpa</i> , Hk.	551
<i>intermedia</i> , Copel.	552
<i>interrupta</i> , Bk.	552
<i>Junghuhnii</i> , Hk.	552
<i>Loheriana</i> , v. A. v. R.	555
<i>paradoxa</i> , Bedd.	552
<i>trichoidea</i> , J. Sm.	552
Myrmecophila	568

N.

Neottopteris	456
Nephrodium	175
<i>abortivum</i> , J. Sm.	217
<i>abruptum</i> , Bedd.	218
<i>abruptum</i> , J. Sm.	227
<i>aciculatum</i> , Bk.	200
<i>acrostichoides</i> , J. Sm.	505
<i>alatellum</i> , Christ	210
<i>albo-punctatum</i> , Desv.	185
<i>amboinense</i> , Pr.	219
<i>angustifolium</i> , Pr.	228
<i>aortisorum</i> , Copel.	501
<i>appendiculatum</i> , Rac.	215
<i>Arbuscula</i> , Desv.	218
<i>aridum</i> , J. Sm.	212
<i>aristatum</i> , Hk.	209
<i>asperulum</i> , Copel.	495
<i>attenuatum</i> , Fée	184
<i>auriculatum</i> , Diels	490
<i>Bakeri</i> , Harr.	208

Nephrodium.

<i>basilare</i> , Pr.	820
<i>Beccarianum</i> , Ces.	185
<i>Beddomei</i> , Bk.	191
<i>Benoitianum</i> , Pr.	225
<i>Blumei</i> , Hk.	202
<i>Blumei</i> , J. Sm.	251
<i>Boryanum</i> , Bk.	201
<i>borneense</i> , Hk.	198
<i>brachyodon</i> , Hk.	220, 819
<i>calcaratum</i> , Hk.	186
<i>callosum</i> , Keys.	220
<i>canescens</i> , Christ	251, 507
<i>caudiculatum</i> , J. Sm.	820
<i>conjugatum</i> , T. & B.	196
<i>costatum</i> , Bedd.	229
<i>crassifolium</i> , Hk.	182
<i>Creaghii</i> , Bk.	185
<i>crenatum</i> , Bk.	202
<i>crinipes</i> , Hk.	229
<i>cucullatum</i> , Bk.	215
<i>cuspidatum</i> , Pr.	741
<i>cyatheoides</i> , Pr.	216
<i>Dayi</i> , Bedd.	188
<i>debile</i> , Bk.	225
<i>debile</i> , Diels	501
<i>decurrens</i> , Bk.	247
<i>didymosorum</i> , Par.	225
<i>dissectum</i> , Desv.	196
<i>dissitifolium</i> , Bk.	196
<i>diversilobum</i> , Pr.	505
<i>divisum</i> , Hk.	201
<i>echinatum</i> , Bk.	182
<i>edulentum</i> , Bk.	194
<i>elongatum</i> , Hk. & Grev.	814
<i>eminens</i> , Bk.	214
<i>erubescens</i> , Diels	489
<i>erythrosorum</i> , Hk.	195

Nephrodium

<i>eusorum</i> , Bedd.	227
<i>excellens</i> , Hk.	171
<i>exiguum</i> , Hk.	190
<i>extensum</i> , Moore	210
<i>var. minor</i>	219
<i>Everettii</i> , Bk.	241
<i>falcilobum</i> , Hk.	185
<i>Fauriei</i> , Christ	
<i>var. elatius</i>	429
<i>ferox</i> , Moore	221
<i>Filix-mas</i> , Rich.	192
<i>var. marginata</i>	814
<i>flaccidum</i> , Hk.	193
<i>Foxii</i> , Copel.	814
<i>Gaimardianum</i> , Gaud.	824
<i>giganteum</i> , Bk.	175, 811
<i>glandulosum</i> , J. Sm.	209
<i>glandulosum</i> , Moore ...	215, 819
<i>glaucostipes</i> , Bedd.	226
<i>gongyloides</i> , Schott	212
<i>gracilescens</i> , Hk.	186
<i>grammitoides</i> , Copel.	181
<i>grande</i> , Bk.	249
<i>granulosum</i> , J. Sm.	505
<i>gymnopodium</i> , Bk.	182
<i>Haenkeanum</i> , Pr.	214
<i>heterocarpon</i> , Moore	228
<i>heterophyllum</i> , Hk.	171
<i>hirsutum</i> , J. Sm.	214, 819
<i>hirtipes</i> , Hk.	180
<i>hirtum</i> , Copel.	492
<i>hispidulum</i> , Bk.	228
<i>Hosei</i> , Bk.	247
<i>immersum</i> , Hk.	188
<i>intermedium</i> , Bk.	202
<i>invisum</i> , Carr.	212
<i>irriguum</i> , Bk.	245

Nephrodium

<i>javanicum</i> , Hk.	232
<i>Keckii</i> , Bk.	248
<i>Keraudrenianum</i> , Diels	496
<i>Labrusca</i> , Rac.	517
<i>larutense</i> , Bedd.	252
<i>latifolium</i> , Bk.	249
<i>latipinna</i> , Hk.	217
<i>Leuzeanum</i> , Hk.	172
<i>ligulatum</i> , Hk.	184
<i>lineatum</i> , Bedd.	216
<i>lineatum</i> , Pr.	209
<i>Lobbii</i> , Bk.	244
<i>longipes</i> , Moore	211
<i>Loherianum</i> , Copel.	221
<i>Luerssenii</i> , Harr.	190
<i>Mannii</i> , Hope	816
<i>megaphyllum</i> , Bk.	204
<i>melanocaulon</i> , Bk.	245
<i>melanopus</i> , Hk.	191
<i>melanorachis</i> , Bk.	250
<i>membranifolium</i> , Pr.	175
<i>Menyanthidis</i> , Bk.	244
<i>microchlamys</i> , Bk.	226
<i>molle</i> , R. Br.	224
<i>var. amboinense</i>	219
<i>var. major</i>	227
<i>Motleyanum</i> , Hk.	182
<i>moulmeinense</i> , Bedd.	217
<i>mucronatum</i> , J. Sm.	819
<i>multicaudatum</i> , Clarke	250
<i>multijugum</i> , Bk.	215
<i>multilineatum</i> , Bedd.	219
<i>multilineatum</i> , Pr.	214
<i>multisetum</i> , Bk.	205
<i>nebulosum</i> , Bk.	246
<i>nudum</i> , Bk.	250
<i>obliteratum</i> , R. Br.	155

Nephrodium

<i>obscurum</i> , Diels	492
<i>obscurum</i> , Moore	212
<i>obtusatum</i> , Diels	498
<i>obtusilobum</i> , Bk.	205
<i>ochtodes</i> , Hk.	215
<i>odoratum</i> , Bk.	202
<i>oligodictyon</i> , Bk.	172
<i>oosorum</i> , Bk.	250
<i>ornatum</i> , Christ	495
<i>Otaria</i> , Bk.	209
<i>oxyodon</i> (<i>oxylon</i>), Copel.	491
<i>pachyphyllum</i> , Bk.	252
<i>parasiticum</i> , Desv.	224
<i>patens</i> , Desv.	189
<i>pectinatum</i> , Hier.	185
<i>pennigerum</i> , Bedd.	224
<i>pennigerum</i> , Moore	217, 218
<i>pennigerum</i> , Pr.	504
<i>perakense</i> , Bk.	226
<i>philippinense</i> , Bk.	820
<i>pilosiusculum</i> , Zipp.	215
<i>platypus</i> , Hk.	168
<i>polymorphum</i> , Bk.	245
<i>polytrichum</i> , Bk.	187
<i>polytrichum</i> , Schrad.	189
<i>Preslii</i> , Bk.	194
<i>procurrens</i> , Bk.	211
<i>proliferum</i> , Keys.	504
<i>prolificum</i> , Diels	195
<i>propinquum</i> , R. Br.	212
<i>pteroides</i> , J. Sm.	209
<i>pteropodium</i> , Bk.	242
<i>pulvinuliferum</i> , Bk.	200
<i>punctatum</i> , Diels	495
<i>punctatum</i> , Par.	210
<i>purpurascens</i> , Hk.	201
<i>recedens</i> , Hk.	198

Nephrodium

<i>rhodolepis</i> , Clarke	202
<i>Riedlianum</i> , Moore	250
<i>rubidum</i> , Copel.	502
<i>rufescens</i> , Diels	495
<i>rugulosum</i> , Copel.	495
<i>sagenioides</i> , Bk.	191
<i>sagittaeifolium</i> , Moore	222
<i>sakayense</i> , Zeill.	224
<i>salicifolium</i> , Diels	511
<i>sarawakense</i> , Bk.	200
<i>semibipinnatum</i> , Bk.	244
<i>seligerum</i> , Bk.	202
<i>setosum</i> , Bk.	198
<i>süifolium</i> , Bk.	251
<i>simplicifolium</i> , J. Sm.	500
<i>simulans</i> , Bk.	190, 225
<i>singalanense</i> , Bk.	192
<i>singaporianum</i> , Bk.	255
<i>sparsum</i> , Don	197
<i>spectabile</i> , Hk.	195
<i>Spenceri</i> , Copel.	508
<i>splendens</i> , Hk.	195
<i>Stegnogramme</i> , Diels	508
<i>stenophyllum</i> , Bk.	247
<i>stipellatum</i> , Moore	227
<i>subdigitatum</i> , Bk.	245
<i>subdimorphum</i> , Christ	251
<i>subtriphyllum</i> , Bk.	246
<i>syrmaticum</i> , Desv.	195
<i>tectum</i> , Bedd.	225
<i>tenericaule</i> , Hk.	202, 495
<i>terminans</i> , J. Sm.	209
<i>ternatum</i> , Bk.	242
<i>Totta</i> , Diels	498
<i>triphyllum</i> , Diels	509
<i>truncatum</i> , Pr.	227
<i>tuberculatum</i> , Ces.	491

Nephrodium

<i>unitum</i> , Bory	215
<i>unitum</i> , R. Br.	212
<i>wrophyllum</i> , Keys.	216, 511
<i>variolosum</i> , Bk.	248
<i>vastum</i> , Bk.	825
<i>vile</i> , Bk.	199
<i>viscosum</i> , Bk.	186
<i>Vitis</i> , Rac.	516
<i>vulcanicum</i> , Bk.	185
<i>Webbianum</i> , Hk.	185, 815
<i>Wightii</i> , Clarke	245
<i>Wigmanii</i> , Rac.	192
<i>Zollingerianum</i> , Bk.	255
<i>Nephrolepis</i>	155, 807
<i>acuta</i> , Pr.	162
<i>var. rufescens</i>	160
<i>acutifolia</i> , Christ	157
<i>acuminata</i> , Kuhn	159
<i>acuminata</i> , Pr.	162
<i>barbata</i> , Copel.	161
<i>biserrata</i> , Schott	162
<i>celebica</i> , Zoll.	160
<i>cordifolia</i> , Pr.	160
<i>var. Lauterbachii</i>	160
<i>davalliae</i> , v. A. v. R.	807
<i>davalloides</i> , Kze	159
<i>dicksonioides</i> , Christ	156
<i>Duffi</i> , Moore	159
<i>exaltata</i> , Schott	161, 808
<i>floccigera</i> , Moore	159, 807
<i>glabra</i> , Copel	808
<i>hirsutula</i> , Pr.	160, 808
<i>Lauterbachii</i> , Christ	160
<i>lindsayae</i> , Christ	157
<i>obliterata</i> , J. Sm.	155
<i>punctulata</i> , Pr.	162
<i>radicans</i> , Kuhn	161

Nephrolepis

<i>ramosa</i> , Moore	155
<i>rufescens</i> , Wawra	160
<i>tuberosa</i> , Pr.	160
<i>volubilis</i> , J. Sm.	161
<i>Neurogramme</i>	528
<i>alismaefolia</i> , Christ	545
<i>calomelanos</i> , Diels	528
<i>cartilagidens</i> , Christ	542
<i>fraxinea</i> , Christ	527
<i>quinata</i> , Christ	543
<i>tartareum</i> , Diels	850
<i>Niphobolus</i>	678
<i>abbreviatus</i> , Zoll.	692
<i>acrostichoides</i> , Rich.	685
<i>adnascens</i> , Klf.	681
<i>albicans</i> , Bl.	688
<i>anetioides</i> , Christ	682
<i>angustatus</i> , Pr.	691
<i>Beddomeanus</i> , Gies.	695
<i>Blumeanus</i> , Kze	688
<i>carnosus</i> , Bl	681
<i>caudatus</i> , Klf.	681
<i>Christii</i> , Gies.	695
<i>confluens</i> , Spr.	690
<i>costatus</i> , Pr	695
<i>distichocarpus</i> , Gies.	695
<i>elongatus</i> , Bl. ..	681
<i>fissus</i> , Bedd.	687, 688
<i>fissus</i> , Bl.	685
<i>flocciger</i> , Bl.	687
<i>glaber</i> , Klf.	688
<i>Heteractis</i> , J. Sm.	684
<i>lanuginosus</i> , Gies.	685
<i>Lauterbachii</i> , Christ	690
<i>lingua</i> , Bedd.	684
<i>Lingua</i> , Spr.	684, 692, 695
<i>Mannii</i> , Gies.	687

- Niphobolus*
mollis, Kze 688
nummulariaefolius, J. Sm. ... 685
nummularifolius, Bedd. 705
obovatus, Kze 685
pannosus, Bedd. 684
penangianus, Hk. 686
porosus, Pr. 687
princeps, Gies. 684
puberulus, Bl. 685
Rasamalae, Gies. 686
rupestris, Spr. 680
samarensis, Fée 691
Sarasinorum, Gies. 693
sphaerocephalus, Hk. & Grev. 691
sphaerostichus, Copel. 682
splendens, J. Sm. 696
sticticus, Kze 688
stigosus, Moore 695
varius, Klf. 682
venosus, Bl. 695
Warburgii, Gies. 692
Niphopsis 678
Notho(ch)laena 485
carnosa, Wall. 702
densa, J. Sm. 485
distans, R. Br. 484
hirsuta, Desv. 485
pteridiformis, Bk. 484
- O.**
- Odontoloma* 260
Boryanum, J. Sm. 261
Odontosoria 258, 825
biflora, C. Chr. 259
chinensis, J. Sm. 259, 825
lindsayae, v. A. v. R. 825
- Odontosoria*
retusa, J. Sm. 259, 825
Oleandra 151, 806
ciliata, Kl. 152
colubrina, Copel. 152, 806
var. membranacea 806
var. nitida 806
Cumingii, J. Sm. 153
cuspidata, Bk. 153, 806
hirtella, Miq. 152
mollis, Pr. 152
musifolia, Pr. 153, 807
neriifolia, Pr. 152, 155
neriiformis, Bedd. 155
neriiformis, Cav. 152
var. brachypus 152
ornata, Christ 154
pilosa, Hk. 152
Sibaldii, Grev. 155
tricholepis, Kze 154
Wallichii, Pr. 153
Wernerii, Rst. 806
Whitmeei, Bk. 154
Olfersia 725
aculeata, Bl. 722
gracilis, Bl. 722
Oligosoriae (*Adiantum*) 519
Onychium 550
auratum, Klf. 550
japonicum, Kze 551
lucidum, Spr. 551
siliculosum, C. Chr. 550
tenue, Christ 550
Ophioderma 772
Ophioglossaceae 771
Ophioglossum 771
acuminatum, Houtt. 159
brevipes, Bedd. 774

Ophioglossum

<i>circinatum</i> , <i>Burm.</i>	802
<i>fibrosum</i> , <i>Schum.</i>	774
<i>flexuosum</i> , <i>L.</i>	802
<i>gramineum</i> , <i>Willd.</i>	773
<i>inconspicuum</i> , <i>v. A. v. R.</i>	775
<i>intermedium</i> , <i>Hk.</i>	776
<i>japonicum</i> , <i>Thb.</i>	802
<i>macrorrhizum</i> , <i>Kze</i>	774
<i>moluccanum</i> , <i>Schl.</i>	775
<i>forma inconspicua</i>	774
<i>forma lanceolata</i>	775
<i>forma pumila</i>	774
<i>nudicaule</i> , <i>L.f.</i>	775
<i>obovatum</i> , <i>Miq.</i>	775
<i>parvifolium</i> , <i>Grev. & Hk.</i>	775
<i>petiolatum</i> , <i>Hk.</i>	775
<i>pedunculatum</i> , <i>Desv.</i>	775
<i>pendulum</i> , <i>L.</i>	777
<i>pumilum</i> , <i>v. A. v. R.</i>	774
<i>reticulatum</i> , <i>L.</i>	776
<i>scandens</i> , <i>L.</i>	802
<i>simplex</i> , <i>Ridl.</i>	775
<i>timorense</i> , <i>Miq.</i>	775
<i>Zeylanicum</i> , <i>Houtt.</i>	757
<i>Osmunda</i>	755
<i>Banksiaefolia</i> , <i>Kuhn</i>	756
<i>blechnoides</i> , <i>Zoll.</i>	756
<i>bromeliaefolia</i> , <i>Copel.</i>	756
<i>capensis</i> , <i>L.</i>	581
<i>coronaria</i> , <i>Kön.</i>	711
<i>javanica</i> , <i>Bl.</i>	756
<i>Prestiana</i> , <i>J. Sm.</i>	756
<i>zeylanica</i> , <i>L.</i>	777
<i>Osmundaceae</i>	755
<i>Osmundastrum</i>	755
<i>Oxygonium</i>	
<i>alismaefolium</i> , <i>J. Sm.</i>	425

P.

<i>Paesia</i> (<i>Pteris</i> §)	577
<i>Paesia</i>	576
<i>radula</i> , <i>C. Chr.</i>	576
<i>rugulosa</i> , <i>Kuhn</i>	577
<i>Parasaccoloma</i>	282
<i>Parkeria</i>	
<i>pteridioides</i> , <i>Hk.</i>	751
<i>Parkeriaceae</i>	751
<i>Partitae</i> (<i>Pteris</i>)	552
<i>Patania</i>	140
<i>Pedatae</i> (<i>Adiantum</i>)	519
<i>Pellaea</i>	544, 827
<i>concolor</i> , <i>Bk.</i>	547
<i>Doniana</i> , <i>Hk.</i>	546
<i>falcata</i> , <i>Fée</i>	545
<i>geraniaefolia</i> , <i>Fée</i>	547
<i>ludens</i> , <i>Prantl</i>	547
<i>pteroides</i> , <i>Prantl</i>	545
<i>timorensis</i> , <i>v. A. v. R.</i>	546, 827
<i>Zippelii</i> , <i>Bk.</i>	545
<i>Phegopterideae</i>	122
<i>Phegopteris</i>	486, 850
<i>acromanes</i> , <i>v. A. v. R.</i>	505
<i>aortisora</i> , <i>v. A. v. R.</i>	501
<i>appendiculata</i> , <i>v. A. v. R.</i>	489
<i>arfakiana</i> , <i>v. A. v. R.</i>	502
<i>asperula</i> , <i>v. A. v. R.</i>	495
<i>auriculata</i> , <i>J. Sm.</i>	490
<i>barbata</i> , <i>Mett.</i>	221
<i>Barberi</i> , <i>Mett.</i>	515
<i>borneensis</i> , <i>v. A. v. R.</i>	500
<i>Beccariana</i> , <i>v. A. v. R.</i>	509
<i>Brongniartii</i> , <i>Mett.</i>	515
<i>canescens</i> , <i>Mett.</i>	507
<i>ceramica</i> , <i>v. A. v. R.</i>	506
<i>chamaeotaria</i> , <i>v. A. v. R.</i>	505

Phegopteris

<i>cheilanthoides</i> , v. A. v. R.	494
<i>Copelandi</i> , v. A. v. R.	492
<i>cuspidata</i> , Mett.	511
<i>debilis</i> , Mett.	501
<i>difformis</i> , Mett.	520
<i>distans</i> , Mett.	496
<i>diversiloba</i> , v. A. v. R.	505
<i>erubescens</i> , J. Sm.	489
<i>ferruginea</i> , Mett.	516
<i>firmula</i> , v. A. v. R.	501
<i>granulosa</i> , v. A. v. R.	505
<i>Hasseltii</i> , Mett.	494
<i>hirta</i> , Christ	492
<i>holophylla</i> , v. A. v. R.	500
<i>Hosei</i> , v. A. v. R.	510
<i>imponens</i> , v. A. v. R.	506
<i>irregularis</i> , Mett.	520
<i>Keraudreniana</i> , Mann	496
<i>Kingi</i> , Bedd.	497
<i>laserpitifolia</i> , Bedd.	494
<i>loxoscapoides</i> , v. A. v. R.	495
<i>luxurians</i> , Mett.	504
<i>macrodonta</i> , Mett.	516
<i>Merrillii</i> , v. A. v. R.	505
<i>nervosa</i> , Fée	165
<i>obscura</i> , Fée	492
<i>obtusata</i> , Christ	498
<i>opaca</i> , Mett.	498
<i>ornata</i> , Fée	494
<i>oxyodus</i> , v. A. v. R.	491
<i>pennigera</i> , v. A. v. R.	504
<i>petrophya</i> , Mett.	515
<i>philippinensis</i> , Mett., var. 1.	521
<i>prolifera</i> , Kuhn	504, 510
<i>punctata</i> , Mett.	555, 495
<i>quadriaurita</i> , v. A. v. R.	489
<i>Ramosii</i> , v. A. v. R.	502

Phegopteris

<i>rubida</i> , v. A. v. R.	502
<i>rufescens</i> , Mett.	495
<i>salicifolia</i> , Mett.	511
<i>simplicifolia</i> , v. A. v. R.	500
<i>Smithii</i> , v. A. v. R.	490
<i>Spenceri</i> , v. A. v. R.	508
<i>stegnogramme</i> , Mett.	508
<i>stenophylla</i> , v. A. v. R.	510
<i>subdigitata</i> , Bedd.	485
<i>subconnexa</i> , Christ	496
<i>subdecurrens</i> , Luer.	514
<i>subobscura</i> , Christ	491
<i>totta</i> , Mett.	498
<i>triphylla</i> , Mett.	509, 850
<i>tuberculata</i> , v. A. v. R.	491
<i>xiphioides</i> , v. A. v. R.	501
<i>Phlebodium</i>	568
<i>Photinopteris</i>	751
<i>drynarioides</i> , Bedd.	751
<i>simplex</i> , J. Sm.	731
<i>speciosa</i> , Bl.	751
<i>Phyllitis</i>	476
<i>intermedia</i> , v. A. v. R.	478
<i>mambare</i> , v. A. v. R.	478
<i>schizocarpa</i> , v. A. v. R.	479
<i>scolopendropsis</i> , v. A. v. R.	477
.	478
<i>Phyllotrichium</i>	778
<i>Phymatodes</i>	569
<i>myriocarpa</i> , Pr.	641
<i>Phymatopsis</i>	569
<i>Physapteris</i>	536
<i>Pinnatae</i> (Polypodium)	567
<i>Pinnatae</i> (Pteris)	552
<i>Pinnatifidae</i> (Polypodium)	567
<i>Pinnatifidae</i> (Pteris)	552
<i>Plagiogyria</i>	540, 826

- Plagiogyria
 adnata, *Bedd.* 342
 Christii, *Copel.* 342, 826
 euphlebia, *Mett.* 343
 falcata, *Copel.* 826
 glauca, *Mett.* 342
 pycnophylla, *Mett.* 343
 stenoptera, *Diels* 341
 tuberculata, *Copel.* 343, 826
 Platycerium 707
 alcicorne, *Desv.* 710
 biforme, *Bl.* 711
 bifurcatum, *C. Chr.* 710
 coronarium, *Desv.* 711
 grande, *J. Sm.* 708, 709
 sumbawense, *Christ* 709
 Wallichii, *Hk.* 708
 Wandae, *Rac.* 709
 Wilhelminae Reginae,
 v. A. v. R. . . . 708
 Willinckii, *Moore* 710
 Platyloma 344
 falcatum, *J. Sm.* 345
 Platytaenia 747
 Requiniana, *Kuhn* 747
 Platyzoma 55
 Plenasium 755
 Pleocnemia 170, 810
 aristata, *Bedd.* 210
 devexa, *v. A. v. R.* 811
 excellens, *v. A. v. R.* 171
 gigantea, *Pr.* 173, 811
 heterophylla, *v. A. v. R.* 171, 810
 Leuzeana, *Pr.* 172, 810
 megalocarpa, *Bedd.* 172
 membranacea, *Bedd.* 811
 membranifolia, *Bedd.* 173
 profereoides, *v. A. v. R.* .. 812
 Pleocnemia
 Trimeni, *Bedd.* 811
 Pleopeltis 368
 accedens, *Moore* 632
 angustata, *Pr.* 691
 dilatata, *Bedd.* 663
 ebenipes, *Bedd.* 659
 Griffithiana, *Moore* 645
 hastata, *Moore* 660
 hemionitidea, *Moore* 648
 incurvata, *Moore* 663
 insignis, *Bedd.* 663
 irioides, *Moore* 655
 laciniata, *Bedd.* 664
 linearis, *Moore* 637
 longifolia, *Bedd.* 638
 longissima, *Bedd.* 663
 longissima, *Moore* 666
 malacodon, *Bedd.* 832
 membranacea, *Moore* 649
 musaeifolia, *Moore* 654
 nigrescens, *Carr.* 665
 normalis, *Moore* 641
 nuda, *Hk.* 637
 ovata, *Moore* 653
 oxyloba, *Bedd.* 660
 palmata, *Moore* 669
 Parishii, *Bedd.* 698
 peltata, *Scort.* 632
 Phymatodes, *Moore* 664
 platyphylla, *Bedd.* 647
 pteropus, *Moore* 661
 punctata, *Bedd.* 653
 rhynchophylla, *Moore* 645
 rostrata, *Bedd.* 636
 rupestris, *Moore* 651
 Scortechinii, *Bedd.* 652
 sinuosa, *Bedd.* 625

- Pleopeltis*
- stenophylla*, Moore 638
- superficialis*, Bedd. 631
- var. latifrons* 632
- tridactyla*, Moore 661
- Wightiana*, Bedd. 637
- Zippelii*, Moore 649
- zosteræformis*, Bedd. 640
- Pleuridium* 569
- Pleurogramme* 551
- interrupta*, Christ 552
- Loheriana*, Christ 555
- paradoxa*, Fée 552
- pusilla*, Christ 553, 560
- Poecilopteris* 753
- costata*, Moore 745
- diversifolia*, Pr. 740
- flagellifera*, J. Sm. 759
- Presliana*, Moore 740
- prolifera*, Pr. 738
- repanda*, Bedd. 741
- repanda*, Pr. 741
- semicordata*, Moore 741
- subrepanda*, Pr. 742
- Polybotrya* 722
- apiifolia*, J. Sm. 725
- appendiculata*, J. Sm. . 505, 724
- articulata*, J. Sm. 725
- aurita*, Bl. 726
- Hamiltoniana*, Pr. 724
- latifolia*, Meyen 757
- marginata*, Bl. 724
- Nieuwenhuis(en)ii*, Rac. . . . 724
- prolifera*, Bory 738
- stenosemioides*, Copel. 725
- Polypodiaceae* 121
- Polypodieae* 122
- Polypodium* 568, 832
- Polypodium*
- abbreviatum*, Mett. 692
- accedens*, Bl. 653
- acrostichoides*, Forst. 685
- aculeatum*, L. 809
- adnascens*, Sw. 681
- adpersum*, Bl. 578
- affine*, Bl. 659
- alatellum*, Christ 582
- alatum*, Hk. 658
- albicans*, Mett. 688
- albidosquamatum*, Bl. 668
- albulum*, Christ 651
- alternans*, Wall. 784
- alternidens*, Ces. 589
- andaiense*, Bk. 514
- angulatum*, Willd. 822
- angustatum*, Bl. 669
- angustatum*, Sw. 691
- Annabellæ*, Forb. 659
- anomalum*, Christ 661
- antrophyoides*, v. A. v. R. . 652
- aortisorum*, Harr. 501
- appendiculatum*, Wall. 489
- var. squamaestipes* 490
- arfakianum*, Bk. 502
- argutum*, Wall. 621
- aristatum*, Forst. 810
- asperulum*, J. Sm. 495
- asperum*, Zipp. 817
- asterosorum*, Bk. 694
- auriculatum*, Wall. 490
- australe*, Mett. 578
- avenium*, Mett. 558
- Bakeri*, Luerss. 652
- barathrophyllum*, Bk. 605
- barbatum*, Hk. 221
- Barberi*, Hk. 515

Polypodium

<i>barisanicum</i> , v. A. v. R.	627
<i>Barometz</i> , L.	792
<i>Beccarianum</i> , Ces.	515
<i>Beccarii</i> , v. A. v. R.	635
<i>Beddomei</i> , Bk.	621
<i>benguetense</i> , Copel.	622
<i>bicuspe</i> , Bl.	752
<i>bifurcatum</i> , Bk.	522
<i>Billardieri</i> , C. Chr.	578
<i>bipinnatifidum</i> , Bk.	611
<i>bisulcatum</i> , Hk.	576
<i>blechnoides</i> , Hk.	605
<i>Blumeanum</i> , C. Chr.	538
<i>Blumei</i> , Mett.	558
<i>Bolsteri</i> , Copel.	675
<i>borneense</i> , Hk.	500
<i>brachyodus</i> , Kze	819
<i>brachyphlebium</i> , Bk.	602
<i>brevifrons</i> , Scort.	600
<i>Brongniartii</i> , Bory	513
<i>Burbidgei</i> , Bk.	590
<i>caespitosum</i> , Mett.	582
<i>calophlebium</i> , Copel.	676
<i>campyloneuroides</i> , Bl.	648
<i>canescens</i> , Bl.	507
<i>capense</i> , L.f.	788
<i>carnosum</i> , Christ	625
<i>carnosum</i> , Mett.	681
<i>caudatum</i> , Mett.	681
<i>caudiforme</i> , Bl.	676
<i>caudiforme</i> , Christ	676
<i>celebicum</i> , Bl.	605, 607
<i>Cesatianum</i> , Bk.	605
<i>cheilanthoides</i> , Bk.	494
<i>Christii</i> , Copel.	581
<i>clavifer(um)</i> , Hk.	608
<i>commutatum</i> , Bl.	659

Polypodium

<i>confluens</i> , Hk.	680
<i>confluens</i> , R. Br.	690
<i>congener</i> , Pr.	582
<i>congenerum</i> , Pr.	582
<i>conjugatum</i> , Bk.	670
<i>conjunctisorum</i> , Bk.	596
<i>contiguum</i> , Brack.	605
<i>contiguum</i> , Hk.	658
<i>contiguum</i> , J. Sm.	616, 852
<i>contiguum</i> , Wall.	657
<i>Copelandi</i> , C. Chr.	558
<i>cordifolium</i> , L.	160
<i>cornigerum</i> , Bk.	596, 852
<i>coronans</i> , Wall.	670
<i>costulatum</i> , Bk.	644
<i>craterisorum</i> , Harr.	607
<i>crenatum</i> , Forsk.	202
<i>crenulatum</i> , Kze	669
<i>cryptocarpum</i> , v. A. v. R.	616
<i>cryptosorum</i> , C. Chr.	601
<i>cucullatum</i> , Nees & Bl.	589
<i>Cumingianum</i> , Hk.	516
<i>curtidens</i> , Christ	652
<i>Curtisii</i> , Bk.	604
<i>cuspidatum</i> , Bl.	618
<i>cyclobasis</i> , Bk.	659
<i>damunense</i> , Rst.	654
<i>davalliaceum</i> , Muell. & Bk.	604
<i>davalloides</i> , Mett.	485
<i>debile</i> , Bk.	501
<i>decipiens</i> , Mett.	601
<i>decorum</i> , Brack.	608
<i>decrescens</i> , Christ	605
<i>denticulatum</i> , Pr.	590
<i>deparoides</i> , Christ	626
<i>difforme</i> , Bl.	520
<i>dilatatum</i> , Wall.	665

Polypodium

<i>dilatatum</i> , var. <i>grandidentatum</i>	661
<i>diminutum</i> , <i>Bk.</i>	577
<i>diplosorum</i> , <i>Christ</i>	586
var. <i>setosum</i>	585
<i>Dipteris</i> , <i>Bl.</i>	525
<i>distans</i> , <i>Don</i>	496
<i>distichocarpum</i> , <i>Mett.</i>	695
<i>diversifolium</i> , <i>Sw.</i>	699
<i>dolichosorum</i> , <i>Christ</i>	646
<i>dolichosorum</i> , <i>Copel.</i>	585
<i>dolichopterum</i> , <i>Copel.</i>	662
<i>drepanopteron</i> , <i>Kze</i>	455
<i>drynarioides</i> , <i>Christ</i>	751
<i>dulitense</i> , <i>Bk.</i>	652
<i>duriusculum</i> , <i>Christ</i>	604
<i>ebenipes</i> , <i>Hk.</i>	659
<i>ellipticum</i> , <i>Thb.</i>	677
<i>Elmeri</i> , <i>Copel.</i>	675
<i>erubescens</i> , <i>Wall.</i>	489
<i>erythrotrichum</i> , <i>Copel.</i>	599
<i>euryphyllum</i> , <i>C. Chr.</i>	665
<i>exaltatum</i> , <i>Copel.</i>	614
<i>exaltatum</i> , <i>L.</i>	808
<i>extensum</i> , <i>Forst.</i>	45
<i>evectum</i> , <i>Forst.</i>	762
<i>fasciatum</i> , <i>Pr.</i>	579
<i>Feei</i> , <i>Mett.</i>	675
<i>ferrugineum</i> , <i>Bk.</i>	516
<i>Filix Mas</i> , <i>L.</i>	814
<i>firmulum</i> , <i>Bk.</i>	501
<i>fissum</i> , <i>Bk.</i>	687
<i>flabellivenium</i> , <i>Bk.</i>	574, 650
<i>floccigerum</i> , <i>Mett.</i>	687
<i>Forbesii</i> , v. <i>A. v. R.</i>	657
<i>Friderici et Pauli</i> , <i>Christ</i>	575
<i>Friderici et Pauli</i> , <i>Christ</i>	615
<i>furcatum</i> (<i>fuscatum</i>), <i>Bl.</i>	601

Polypodium

<i>Gaudichaudii</i> , <i>Bory</i>	699
<i>gedeanum</i> , <i>Rac.</i>	611
<i>glabrum</i> , <i>Mett.</i>	690
<i>glandulosum</i> , <i>Hk.</i>	609
<i>glaucopruinatum</i> , <i>C. Chr.</i>	660
<i>glaucum</i> , <i>Kze</i>	660
<i>glaucum</i> , <i>Thb.</i>	794
<i>glossipes</i> , <i>Bk.</i>	642
<i>gracillimum</i> , <i>Copel.</i>	589
<i>grandidens</i> , <i>Kze</i>	619
<i>grandidentatum</i> , <i>Bk.</i>	661
<i>grandifolium</i> , <i>Wall.</i>	649
<i>granulosum</i> , <i>Pr.</i>	505
<i>Griffithianum</i> , <i>Hk.</i>	645
<i>halconense</i> , <i>Copel.</i>	610
<i>Hamiltonianum</i> , <i>Diels</i>	675
<i>hammatisorum</i> , <i>Harr.</i>	634
<i>Hancockii</i> , <i>Bk.</i>	518
<i>harpophyllum</i> , <i>Zenk.</i>	610
<i>Hasseltii</i> , <i>Bl.</i>	494
<i>hastatum</i> , <i>Thb.</i>	660, 852
<i>Havilandii</i> , <i>Bk.</i>	584
<i>Hellwigii</i> , <i>Diels</i>	645
<i>hemiontideum</i> , <i>Wall.</i>	648
<i>heracleum</i> , <i>Kze</i>	671
<i>Heteractis</i> , <i>Mett.</i>	684
<i>heterocarpum</i> , <i>Bl.</i>	655
<i>heterocarpum</i> , <i>Mett.</i>	647, 676
<i>heterolobum</i> , <i>C. Chr.</i>	661
<i>heterosorum</i> , <i>Bk.</i>	518
<i>hirsutulium</i> , <i>Forst.</i>	808
<i>hirtellum</i> , <i>Bl.</i>	582
var. <i>adpersum</i>	578
var. <i>lasiosorum</i>	580
<i>hirtum</i> , <i>Pr.</i>	492
<i>holophyllum</i> , <i>Bk.</i>	500
<i>holophyllum</i> , <i>Bk.</i>	650

Polypodium

<i>Hookeri</i> , Brack.	584
<i>Horsfieldii</i> , R. Br.	525
<i>Hosei</i> , C. Chr.	674
<i>imbricatum</i> , Karst.	624
<i>imponens</i> , Ces.	506
<i>inarticulatum</i> , Copel.	608
<i>inconspicuum</i> , Bl.	596
<i>incurvatum</i> , Bl.	665
<i>induratum</i> , Bk.	657
<i>insigne</i> , Bl.	665
<i>integrius</i> , Copel.	618
<i>interruptum</i> , C. Chr.	673
<i>intromissum</i> , Christ	585
<i>invisum</i> , Forst.	819
<i>involutum</i> , Mett.	537
<i>iridifolium</i> , Diels	559
<i>irioides</i> , Poir.	653
<i>irregulare</i> , Pr.	520
<i>Jagorianum</i> , Mett.	585
<i>Keraudrenianum</i> , Gaud.	496
<i>hasyanum</i> , Hk.	605
<i>knutsfordianum</i> , Bk.	584
<i>Koenigii</i> , Bl.	681
<i>Koningsbergeri</i> , v. A. v. R.	619
<i>Koordersii</i> , Christ	581
<i>Korthalsii</i> , Mett.	618
<i>Labrusca</i> , Hk.	517
<i>laciniatum</i> , Bl.	664
<i>lagopodioides</i> , Christ	640, 651
<i>lagunense</i> , Christ	667
<i>lampongense</i> , Zoll.	622
<i>lancifolium</i> , v. A. v. R.	598
<i>laserpitiiifolium</i> , Scort.	494
<i>lasiosorum</i> , Hk.	580
<i>latifolium</i> , Forst.	825
<i>lecanopteris</i> , Mett.	625
<i>leucophorum</i> , Bk.	650

Polypodium

<i>Leuzeanum</i> , Gaud.	810
<i>Léysii</i> , Bk.	607
<i>lineare</i> , Burm.	59
<i>lineare</i> , Thb.	657, 641
<i>lineatum</i> , Colebr.	229
<i>linguaeforme</i> , Mett.	642
<i>Linn(a)ei</i> , Bory	699
<i>lividum</i> , Mett.	602
<i>lobatum</i> , Huds.	810
<i>Lobbianum</i> , Hk.	522
<i>Lobbianum</i> , Hk.	608
<i>locellatum</i> , Bk.	577
<i>Loherianum</i> , Christ	590
<i>lomarioides</i> , Kze	625
<i>longifolium</i> , Mett.	658
<i>longifrons</i> , Wall.	641
<i>loriforme</i> , Hk.	657
<i>longissimum</i> , Bl.	666
<i>loxogramme</i> , Mett.	538
<i>loxoscaphoides</i> , Bk.	495
<i>ludens</i> , Bk.	591
<i>luzonicum</i> , Copel	651
<i>lycopodioides</i> , L.	640
<i>Macleayi</i> , v. A. v. R.	626
<i>macrochasmum</i> , Bk.	664
<i>macrodon</i> , Reinw.	516
<i>macrophyllum</i> , Bk.	674
<i>macropodium</i> , Bk.	683
<i>macrum</i> , Copel.	609
<i>Maingayi</i> , Diels	677
<i>malaccanum</i> , Bk.	602
<i>malaicum</i> , v. A. v. R.	577
<i>marginale</i> , Thb.	825
<i>Maxwellii</i> , Bk.	576
<i>mediale</i> , Bk. <i>var. parasiticum</i>	581
<i>megalocarpum</i> , Hk.	172

Polypodium

<i>membranaceum</i> , Don.	649
<i>mengtzeense</i> , Copel.	622
<i>Merrillii</i> , Copel.	579
<i>Merritti</i> , Copel.	597
<i>Meyenianum</i> , Hk.	671, 700
<i>millefolium</i> , Bl.	612
<i>millisorum</i> , Bk.	655
<i>mindanaense</i> , Christ	654
<i>minimum</i> , Bk.	579
<i>minutum</i> , Bk.	598
<i>minutum</i> , Bl.	597
<i>mirabile</i> , C. Chr.	624
<i>molle</i> , Mett.	688
<i>mollicomum</i> , Nees & Bl.	601
<i>molliculum</i> , Copel.	620
<i>mollipilum</i> , Bk.	584
<i>monstrosum</i> , Copel.	655
<i>Moseleyi</i> , Bk.	669
<i>multicaudatum</i> , Copel.	605
<i>multilineatum</i> , Wall.	217
<i>musgravianum</i> , Bk.	596
<i>musifolium</i> , Bl.	654
<i>myriocarpum</i> , Mett.	641
<i>myrmecophilum</i> , Christ	624
<i>naviculare</i> , v. A. v. R.	627
<i>nectariferum</i> , Bk.	700
<i>neglectum</i> , Bl.	655
<i>Nieuwenhuisii</i> , v. A. v. R.	626
<i>nigrescens</i> , Bl.	665
<i>nitens</i> , Bk.	696
<i>normale</i> , Don	641
<i>var. sumatranum</i>	651
<i>nummulariaefolium</i> , Mett.	685
<i>nummularium</i> , Mett.	654
<i>nutans</i> , Bl.	606
<i>nutans</i> , J. Sm.	608
<i>obovatum</i> , Mett.	685

Polypodium

<i>oblanceolatum</i> , Bk.	575
<i>obliquatum</i> , Bl.	600
<i>obliquatum</i> , Copel.	605
<i>obliquatum</i> , Hk.	599
<i>obscurum</i> , Hk.	492
<i>obscurum</i> , Mett.	586
<i>obtusifolium</i> , C. Chr.	606
<i>oleandroides</i> , Bk.	575
<i>oodes</i> , Kze	656, 652
<i>ornatissimum</i> , Rst.	575
<i>ornatum</i> , Wall.	494
<i>ovatum</i> , Wall.	655
<i>oxyodon</i> , Bk.	491
<i>oxyphyllum</i> , Kze	649
<i>pachydermum</i> , Bk.	680
<i>padangense</i> , Bk.	586
<i>pallens</i> , Bl.	620
<i>palmatum</i> , Bl.	669
<i>paludosum</i> , Bl.	496
<i>palustre</i> , Burm.	720
<i>pannosum</i> , Mett.	683
<i>papillosum</i> , Bl.	600
<i>papillosum</i> , Ces.	605
<i>papuanum</i> , Bk.	659
<i>parallelum</i> , C. Chr.	557
<i>parasiticum</i> , L.	224
<i>parasiticum</i> , Mett.	581
<i>parvulum</i> , Bedd.	598
<i>parvulum</i> , Bl.	596
<i>patelliferum</i> , Burck	625
<i>patens</i> , Sw.	814
<i>paucijugum</i> , v. A. v. R.	658
<i>paucisorum</i> , Copel.	577
<i>pectinatum</i> , Forsk.	185
<i>pediculatum</i> , Bk.	608
<i>pedunculatum</i> , Mett.	675
<i>peltatum</i> , v. A. v. R.	652

Polypodium

<i>penangianum</i> , Hk.	229, 686
<i>pennigerum</i> , Forst.	504
<i>pentaphyllum</i> , Bk.	662
<i>persicifolium</i> , Desv.	618
<i>petrophyum</i> , Bl.	515
<i>phanerophlebium</i> , Copel. . . .	665
<i>phlebiscopum</i> , Bk.	648
<i>phlebodioides</i> , Copel.	618
<i>phyllomanes</i> , Christ	
<i>var. ovatum</i>	655
<i>phymatodes</i> , L.	664
<i>pilosiusculum</i> , Pr.	580
<i>platyphyllum</i> , Sw.	647
<i>pleiosoroides</i> , Copel.	576
<i>pleuridioides</i> , Mett.	697
<i>pleurogrammoides</i> , Rst.	585
<i>polycarpum</i> , Bk.	520
<i>polymorphum</i> , v. A. v. R. . . .	615
<i>porosum</i> , Wall.	687
<i>Prainii</i> , C. Chr.	617
<i>princeps</i> , Mett.	684
<i>productum</i> , Christ	658
<i>proliferum</i> , Rb.	504
<i>propinquum</i> , Wall.	698
<i>proteus</i> , Copel.	670
<i>pseudoarticulatum</i> , Copel. . . .	601
<i>pseudoconnatum</i> , Copel.	621
<i>pteroides</i> , Pr.	515
<i>pteropus</i> , Bl.	661
<i>pubinerve</i> , Christ	586
<i>punctatum</i> , Sw.	655
<i>punctatum</i> , Thb.	495
<i>pusillum</i> , Christ	582
<i>pyrolifolium</i> , Goldm.	654
<i>quercifolium</i> , L.	698
<i>quinquefidum</i> , Bk.	660
<i>quinquefurcatum</i> , Bk.	525

Polypodium

<i>Raapii</i> , v. A. v. R.	640
<i>Raciborskii</i> , C. Chr.	557
<i>radicans</i> , Burm.	161
<i>Rasamalae</i> , Rac.	686
<i>recedens</i> , J. Sm.	816
<i>regulare</i> , Mett.	675
<i>Reinwardtii</i> , Kze	622
<i>Reinwardtii</i> , Pr.	585
<i>var. obscurum</i>	586
<i>repandulum</i> , Mett.	599
<i>var. malayanum</i>	600
<i>revolutum</i> , C. Chr.	658
<i>rhynchophyllum</i> , Hk.	645
<i>rigidulum</i> , Sw.	699
<i>rivulare</i> , Copel.	662
<i>rostratum</i> , Hk.	656
<i>rubidum</i> , Hk.	502
<i>rudimentum</i> , Copel.	656
<i>rufescens</i> , Bl.	495
<i>rugulosum</i> , Lab.	495
<i>rupestre</i> , Bl.	650, 651
<i>rupestre</i> , R. Br.	680
<i>sablanianum</i> , Christ	642
<i>sagitta</i> , Christ	704
<i>samarense</i> , Mett.	691
<i>Sarasinorum</i> , v. A. v. R.	615
<i>sarawakense</i> , Bk.	651
<i>sarcopus</i> , de Vr. & Teyssm. . . .	624
<i>Sauvinieri</i> , Bk.	668
<i>saxatile</i> , Mett.	650
<i>scabristipes</i> , Bk.	585
<i>Schefferi</i> , v. A. v. R.	609
<i>Schenkii</i> , Harr.	600
<i>Schlechteri</i> , v. A. v. R.	614
<i>Schneideri</i> , Christ	666
<i>Schumannianum</i> , Diels	654
<i>scolopendrinum</i> , C. Chr.	557

Polypodium

<i>Scortechinii</i> , Bk.	242, 652
<i>sculpturatum</i> , Bk.	668
<i>selliguea</i> , Mett.	675
<i>semicordatum</i> , Sw.	808
<i>serraeforme</i> , J. Sm.	615
<i>serrato-dentatum</i> , v. A. v. R.	597
<i>sessilifolium</i> , Hk.	577
<i>setigerum</i> , Bl.	587
<i>setigerum</i> , Hk. & Arn.	584
<i>setosum</i> , Pr.	585
<i>sibarongae</i> , Christ	606
<i>sifolium</i> , Willd.	825
<i>simplicifolium</i> , Hk.	500
<i>sinuosum</i> , Wall.	625
<i>solidum</i> , Mett.	606
<i>soridens</i> , Hk.	655
<i>sparsisorum</i> , Desv.	699
<i>sparsum</i> , Ham.	815
<i>speciosum</i> , Christ	751
<i>Speluncae</i> , L.	825
<i>sphaerostichum</i> , Mett.	682
<i>splendens</i> , Hk.	696, 701
<i>spurium</i> , Mett.	674
<i>Stanleyanum</i> , Bk.	577
<i>stenobasis</i> , Bk.	607
<i>stenophyllum</i> , Bl.	658
<i>stenopteris</i> , Bk.	655
<i>sticticum</i> , Mett.	687
<i>stigmatosum</i> , Sw.	695
<i>streptophyllum</i> , Bk.	595
<i>subaquatile</i> , Christ	668
<i>subarboreum</i> , Bk.	204
<i>subauriculatum</i> , Bl.	622
<i>var. pallens</i>	620
<i>subdichotomum</i> , Rac.	588
<i>subdigitatum</i> , Bl.	485
<i>subdrynariaceum</i> , Christ	654

Polypodium

<i>subecostatum</i> , Hk.	636
<i>subevenosum</i> , Bk.	582
<i>var. sessilifolia</i>	577
<i>subfalcatum</i> , Bl.	599
<i>var. glabrum</i>	599
<i>var. semiintegrum</i>	598
<i>subfasciatum</i> , Rst.	579
<i>subgeminatum</i> , Christ	659
<i>subirideum</i> , Christ	654
<i>subminutum</i> , v. A. v. R. ..	598
<i>subobliquatum</i> , Christ	606
<i>suboppositum</i> , Christ	652
<i>subpinnatifidum</i> , Bl.	590
<i>subpleiosorum</i> , Rac.	587
<i>subrepandulum</i> , Christ	598
<i>subrostratum</i> , C. Chr.	656
<i>subsecundo-dissectum</i> , Zoll. .	598
<i>subselligueum</i> , Bk.	578
<i>subserratum</i> , Hk.	401
<i>subsparsum</i> , Bk.	650
<i>subtriangulare</i> , v. A. v. R. .	595
<i>subtriphyllosum</i> , Hk. & Arn. .	825
<i>sucklingianum</i> , Bk.	580
<i>sumatranum</i> , Bk.	587
<i>sumatranum</i> , Bk.	658
<i>sundense</i> , C. Chr.	658
<i>superficiale</i> , Bl.	651, 652
<i>taeniopsis</i> , Christ	651
<i>tamariscinum</i> , Klf.	611
<i>taxodioides</i> , Bk.	612
<i>tenuilore</i> , Kze	651
<i>tenuisectum</i> , Bl.	612
<i>tomentellum</i> , C. Chr.	620
<i>Toppingii</i> , v. A. v. R.	616
<i>tortile</i> , v. A. v. R.	614
<i>torulosum</i> , Bk.	652
<i>tottum</i> , Willd.	497

Prosaptia

<i>contigua</i> , Bedd.	616
<i>cryptocarpa</i> , Copel.	616
<i>Emersoni</i> , Pr.	615
<i>polymorpha</i> , Copel.	615
<i>serraeformis</i> , Christ	615
<i>Schlechteri</i> , Christ	614
<i>toppingii</i> , Copel.	616
<i>Protogleichenia</i>	56
<i>Psomiocarpa</i>	725
<i>Pterideae</i>	121
<i>Pteridella</i>	544
<i>Pteridium</i>	577
<i>aquilinum</i> , Kuhn	577
<i>Pteris</i>	551, 827
<i>amoena</i> , Bl.	567
<i>aquilina</i> , L.	577
<i>argentea</i> , Gmel.	537
<i>argyraea</i> , Moore	566
<i>asperula</i> , J. Sm.	566
<i>aurita</i> , Bl.	549
<i>Beccariana</i> , C. Chr.	564
<i>biaurita</i> , L.	571, 572, 574
<i>var. quadriaurita</i>	565
<i>var. reducta</i>	555
<i>blechnoides</i> , Willd.	565
<i>Blumeana</i> , Ag.	566
<i>var. asperula</i>	566
<i>var. pluricaudata</i>	564
<i>caesia</i> , Copel.	567
<i>chrysocarpa</i> , Hk. & Grev. ..	550
<i>chryso sperma</i> , Hk. & Grev. .	550
<i>concinna</i> , Bk.	564
<i>concolor</i> , L. & F.	547
<i>crenata</i> , Sw.	560
<i>cretica</i> , L.	558
<i>var. heteromorpha</i>	562
<i>var. melanocaulon</i>	558

Pteris

<i>cretica</i> , <i>var. stenophylla</i>	557
<i>Dalhousiae</i> , Hk.	565, 827
<i>Dalhousiae</i> , Rac.	554
<i>decussata</i> , J. Sm.	568
<i>dimidiata</i> , Willd.	562
<i>dispar</i> , Kze	562
<i>distanis</i> , J. Sm.	562
<i>divaricatissima</i> , Bk.	577
<i>diversifolia</i> , Sw.	555
<i>ensififormis</i> , Burm.	560
<i>esculenta</i> , Forst.	577
<i>excelsa</i> , Gaud.	568
<i>falcata</i> , R. Br.	545
<i>farinosa</i> , Forsk.	558
<i>finisterrae</i> , Rst.	575
<i>furcans</i> , Bk.	567
<i>geminata</i> , Wall.	571
<i>geraniaefolia</i> , Raddi	547
<i>Grevilleana</i> , Wall.	564, 567,
	827
<i>Griffithiana</i> , Hk.	564
<i>heteromorpha</i> , Fée	562
<i>heterophlebia</i> , Kze	572
<i>hirsuta</i> , Poir.	485
<i>inaequalis</i> , Bk.	562
<i>incisa</i> , Thb.	549
<i>Indica</i> , Christ	557
<i>intromissa</i> , Christ	827
<i>Junghuhnii</i> , Bk.	574
<i>Kleiniana</i> , Christ	571
<i>laevigata</i> , Zipp.	554
<i>lanuginosa</i> , Spr.	577
<i>ligulata</i> , Gaud.	570
<i>lineata</i> , L.	561
<i>longifolia</i> , L.	555, 556
<i>longipes</i> , Don	569
<i>longipinnula</i> , Ces.	564

Pteris

<i>longipinnula</i> , Wall.	565, 827
<i>ludens</i> , Wall.	547
<i>marginata</i> , Bory	575
<i>maxima</i> , Bk.	571
<i>melanocaulon</i> , Fée	558
<i>mertensioides</i> , Willd.	575
<i>microdictyon</i> , Hk.	574
<i>Milneana</i> , Bk.	574
<i>mixta</i> , Christ	570
<i>moluccana</i> , Bl.	557
<i>multifida</i> , Poir.	561
<i>nemoralis</i> , Willd.	566
<i>normalis</i> , Don	566
<i>var. B</i>	566
<i>opaca</i> , J. Sm.	556
<i>orientalis</i> , v. A. v. R.	555, 827
<i>papuana</i> , Ces.	558
<i>parviloba</i> , Christ	567
<i>patens</i> , Hk.	568
<i>pellucens</i> , Ag.	569
<i>pellucida</i> , Pr.	557
<i>var. stenophylla</i>	557
<i>philippinensis</i> , Fée	570
<i>platysora</i> , Bk.	556
<i>pluricaudata</i> , Copel.	564
<i>quadriaurita</i> , Retz	565, 567
.	571, 574, 489
<i>var. biaurita</i>	571
<i>var. digitata</i>	564
<i>var. Lauterbachii</i>	570
<i>var. longipinnula</i>	565
<i>radicans</i> , Christ	569
<i>Radula</i> , Bk.	577
<i>rangiferina</i> , Pr.	554
<i>reducta</i> , Bk.	555
<i>rugulosa</i> , Lab.	577
<i>scolopendrina</i> , Bory	561

Pteris

<i>semipinnata</i> , L.	562, 567
<i>var. dispar</i>	562
<i>siliculosa</i> , Desv.	550
<i>serrulata</i> , L.f.	561
<i>stenophylla</i> , Wall.	557
<i>sumatrana</i> , Bk.	561
<i>torricelliana</i> , Christ	565
<i>trachyphylla</i> , Kze	566
<i>Treacheriana</i> , Bk.	558
<i>Treubii</i> , v. A. v. R.	556, 827
<i>tripartita</i> , Sw.	575
<i>var. Junghuhnii</i>	574
<i>var. Milneana</i>	574
<i>umbraculifera</i> , Mett.	565
<i>varians</i> , Wall.	559
<i>venulosa</i> , Bl.	557
<i>venusta</i> , Kze	557
<i>Vespertilionis</i> , Lab.	549
<i>Victoriae</i> , hort.	560
<i>Walkeri</i> , Bk.	568
<i>Wallichiana</i> , Ag.	572
<i>Warburgii</i> , Christ	575
<i>Whitfordi</i> , Copel.	566
<i>Zollingeri</i> , Mett.	569

R.

Reniformiae (<i>Adiantum</i>)	519
---	-----

S.

<i>Saccoloma</i>	281
<i>moluccanum</i> , Mett.	282
<i>sorbifolium</i> , Christ	282
<i>Sadleria</i>	588
<i>cyatheoides</i> , Klf.	588
<i>Sagenia</i>	254
<i>alata</i> , Bedd.	241
<i>calcareae</i> , J. Sm.	282

Sagema

<i>coadunata</i> , J. Sm.	249
<i>decurrens</i> , Houlst.	247
<i>ferruginea</i> , Christ	516
<i>gigantea</i> , Bedd.	811
<i>var. minor</i>	811
<i>Griffithii</i> , Bedd.	250
<i>Labrusca</i> , Christ	517
<i>Lobbii</i> , Christ	244
<i>mamillosa</i> , Moore	247
<i>melanocaulon</i> , Moore	245
<i>pachyphylla</i> , Moore	252
<i>palmata</i> , Christ	257
<i>polymorpha</i> , Christ	245
<i>pteropoda</i> , Christ	242
<i>repanda</i> , Moore	238
<i>rufipubis</i> , Christ	516
<i>semibipinnata</i> , Moore	825
<i>subdigitata</i> , Christ	245
<i>subtriphylla</i> , Bedd.	246
<i>vasta</i> , Moore	241
<i>Vitis</i> , Christ	516
<i>Salpichlaena</i>	585
<i>Schaffneria</i>	477
<i>Schellolepis</i> (<i>Schellopsis</i>)	568
<i>Schizaea</i>	115, 803
<i>dichotoma</i> , Sm.	116, 803
<i>digitata</i> , Sw,	116, 803
<i>fistulosa</i> , Lab.	116
<i>var. malaccana</i>	116
<i>Forsteri</i> , Spr.	116
<i>malaccana</i> , Bk.	116
<i>Schizaeaceae</i>	109
<i>Schizocaena</i>	
<i>Brunonis</i> , J. Sm.	15
<i>Schizoloma</i>	275, 824
<i>angustum</i> , Copel.	277
<i>cordatum</i> , Gaud.	279

Schizoloma

<i>coriaceum</i> , v. A. v. R.	280, 824
<i>davallioides</i> , Moore	274
<i>divergens</i> , Kuhn	278
<i>ensifolium</i> , J. Sm.	280
<i>fuligineum</i> , Copel	277
<i>Guerinianum</i> , Gaud.	288, 824
<i>var. ovatum</i>	277
<i>heterophyllum</i> , J. Sm.	281
<i>induratum</i> , C. Chr.	277
<i>jamesonioides</i> , Copel.	276
<i>lobata</i> , Bedd.	274
<i>ovatum</i> , Copel.	277
<i>pellaeiforme</i> , C. Chr.	278
<i>recurvatum</i> , Moore	274
<i>trilobatum</i> , v. A. v. R.	280
<i>Walkerae</i> , Kuhn	278
<i>Schizostege</i>	548
<i>calocarpa</i> , Copel.	548
<i>pachysora</i> , Copel.	548
<i>Scolopendrium</i>	477
<i>longifolium</i> , Pr.	480
<i>mambara</i> , Bail.	478
<i>pinnatum</i> , J. Sm.	480
<i>schizocarpum</i> , Copel	479
<i>Scutigera</i>	707
<i>Scyphularia</i>	296
<i>Selliguea</i>	569
<i>campyloneuroides</i> , Bedd.	674
<i>caudiformis</i> , J. Sm.	676
<i>decurrens</i> , Pr.	677
<i>elliptica</i> , Bedd.	677
<i>Feei</i> , Bory	675
<i>Hamiltoniana</i> , Bedd.	675
<i>heterocarpa</i> , Bl.	676
<i>Maingayi</i> , Bedd.	677
<i>macrophylla</i> , Bl.	674
<i>membranacea</i> , Bl.	675

- Selliguea*
Selliguea, Christ 675
- Sitolobium*
cuneatum, J. Sm. 805
samoense, Brack. 804
- Soromanes* 725
- Sphaerostephanos*
asplenioides, J. Sm. 232
- Stegania*
Patersoni, R. Br. 579
- Stegnogramme* 486
aspidioides, Bl. 508
- Stenochlaena* 718, 855
aculeata, Kze 722
areolaris, Copel. 855
arthropteroides, Christ 719
dubia, v. A. v. R. 721
gracilis, Kze 722
juglandifolia, Pr. 720
laurifolia, Pr. 719
palustris, Bedd. 720, 722
Raciborskii, C. Chr. 720
scandens, J. Sm. 720
Smithii, v. A. v. R. 720
sorbifolia, J. Sm. 720, 722
var. dimorphophylla ... 480
subtrifoliata, Copel. 721
- Stenoloma* 258
chinensis, Bedd. 259
- Stenosemia* 726
aurita, J. Sm. 521
aurita, Pr. 726
pinnata, Copel. 727
Teysmanniana, Diels 727
- Synammia* 568
- Synaphlebium* 260
recurvatum, J. Sm. 274
- Syngramma(e)* 539, 851
- Syngramma(e)*
alismifolia, J. Sm. 545
Boerlageana, v. A. v. R. ... 545
borneensis, J. Sm. ... 541
cartilagidens, Diels 542
Dayi, Bedd. 541
fraxinea, Bedd. 527
Hookeri, C. Chr. 545
Hosei, Diels 545
lanceolata, Diels 545
Lobbiana, J. Sm. 541
obtusifolia, J. Sm. 541
pinnata, J. Sm. 546
quinata, Carr. 545
valleculata, C. Chr. 540
vittaeformis, J. Sm. 545
Wallichii, Bedd. 545
Zollingeri, Diels 851
- T.**
- Taeniopsis* 555
amboinensis, Bedd. 561
falcata, Bedd. 560
lineata, J. Sm. 561
scolopendrina, J. Sm. 561
- Taenitis* 562
blechnoides, Sw. 565
var. obtusa 562
interrupta, Hk. & Grev. 565
obtusa, Hk. 562
pusilla, Mett. 560
simplicivenia, Ces. 560
stenophylla, Christ 563
- Tapeinidium* 315
amboinense, C. Chr. 315
Denhami, C. Chr. 316
gracile, v. A. v. R. 315
oligophlebium, C. Chr. 314

- Tapeinidium**
pinnatum, C. Chr. 514
var. gracile 515
- Tectaria** 254
ambigua, Copel. 521, 812
Barberi, Copel. 515
Bryanti, Copel. 514, 519
calcarea, Copel. 256
Christii, Copel. 249
crenata, Cav. 258, 251, 252
decurrens, Copel. 246, 247
devexa, Copel. 810
grandifolia, Copel. 249
irregularis, Copel. 519, 520
var. Brongniartii 515
var. macrodon 516
irrigua, Copel. 245, 518
Labrusca, Copel. 517
latifolia, Copel. 249
leuzeana, Copel. 810
malayensis, Copel. 249
melanocaulon, Copel. 245, 249
Menyanthidis, Copel. 244
pentaphylla, v. A. v. R. 519
polymorpha, Copel. 243
süfolia, Copel. 249, 251
subcaudata, v. A. v. R. 825
subtriphylla, Copel. 246
iernatensis, v. A. v. R. 825
vasta, Copel. 241
- Teratophyllum** 718
aculeatum, Mett. 722
- Thamnopteris** 456
Nidus, Pr. 459
Phyllioidis, Pr. 850
Simonsiana, Moore 458
stipitata, Pr. 445
- Thayeria** 697
- Thayeria**
cornucopia, Copel. 700
nectarifera, Copel. 700
- Todea**
alpina, Bk. 757
Fraseri, Hk. & Grev. 756
Wilkesiana, Brack. 756
- Toxopteris** 559
- Trichomanes** 82, 799
aculeatum, J. Sm. 798
adiantoides, L. 464
anceps, Hk., var. B. 799
aphlebioides, Christ 102
apifolium, Pr. 104
asplenioides, Pr. 101
Aswijkii (*Asnykii*), Rac. 88
auriculatum, Bl. 96
Bauerianum, Endl. 104
Beccarianum, Ces. 86
bilabiatum, Nees & Bl. 94
var. bilingue 102
bilingue, Hk. 102
bimarginatum, V. D. B. 86
bipunctatum, Poir. 95
var. capillatum 95
var. nana-subflabellata 95
brevipes, Bk. 94, 102
capillatum, Taschn. 95
cellulosum, Hk. 105
chinense, L. 259
Christii, Copel. 800
contiguum, Forst. 616
corticola, Bedd. 89
Cumingii, C. Chr. 99
cupressoides, Desv. 105
var. selaceum 801
demissum, Forst. 797
denticulatum, Bk. 79

Trichomanes

<i>dichotomum</i> , Kze	96
<i>diffusum</i> , Bl.	90, 799
<i>digitatum</i> , Sw.	89
<i>var. palmatifidum</i>	90
<i>dissectum</i> , J. Sm.	96
<i>ericoides</i> , Hedw.	105
<i>Filicula</i> , Bedd.	98
<i>Filicula</i> , Bory	94, 95
<i>filiculoides</i> , Christ	95
<i>flabellatum</i> , V. D. B.	89
<i>flaccidum</i> , Forst.	805
<i>foeniculaceum</i> , Bory	105
<i>gemmatum</i> , J. Sm.	105
<i>giganteum</i> , Bory	105
<i>glauco-fuscum</i> , Hk.	95
<i>gracile</i> , V. D. B.	98
<i>Ilenzaiense</i> , Bedd.	85
<i>hispidulum</i> , Mett.	97
<i>Hosei</i> , Bk.	98
<i>humile</i> , Forst.	94
<i>ignobile</i> , Ces.	97
<i>japonicum</i> , Thb.	551
<i>javanicum</i> , Bl. ...	100, 799, 800
<i>var. asplenioides</i>	101
<i>Kurzii</i> , Bedd.	89
<i>Lauterbachii</i> , Christ	97
<i>longisetum</i> , Bory	105
<i>maximum</i> , Bl.	99, 799
<i>meifolium</i> , Bl.	104
<i>meifolium</i> , Bory	105
<i>melanorhizon</i> , Hk.	102
<i>Merrilli</i> , Copel.	801
<i>microchilum</i> , Bk.	95
<i>millefolium</i> , Pr.	799
<i>minutum</i> , Bl.	90
<i>Motleyi</i> , V. D. B.	85, 86
<i>var. pannosum</i>	86

Trichomanes

<i>multifidum</i> , Forst.	798
<i>muscoides</i> , Bk.	86
<i>nanum</i> , V. D. B.	89
<i>neilgherrense</i> , Bedd.	86
<i>nitidulum</i> , V. D. B.	88, 799
<i>Nymani</i> , Christ	94
<i>obscurum</i> , Bl.	105
<i>omphalodes</i> , C. Chr.	85
<i>pallidum</i> , Bl.	95
<i>palmatifidum</i> , K. Müll.	90
<i>pannosum</i> , Ces,	86
<i>parviflorum</i> , Poir.	105
<i>parvulum</i> , Poir.	88
<i>peltatum</i> , Bk.	85
<i>pluma</i> , Hk.	106
<i>polyanthos</i> , Hk.	104
<i>proliferum</i> , Bl.	90, 91
<i>pyxidiferum</i> , L.	98
<i>racemosum</i> , V. D. B.	102
<i>radicans</i> , Sw.	99
<i>rhomboideum</i> , J. Sm.	800
<i>rigidum</i> , Sw.	102, 105
<i>rupiculum</i> , Rac.	89
<i>saxatile</i> , Moore	105
<i>saxifragoides</i> , Pr.	88
<i>serrulatum</i> , Bk.	95
<i>setaceum</i> , V. D. B.	801
<i>Smithii</i> , Hk.	99
<i>speciosum</i> , Willd.	99
<i>strigosum</i> , Thb.	825
<i>sublimbatum</i> , K. Müll.	86
<i>sumatranum</i> , v. A. v. R. ..	800
<i>tenuifolium</i> , Burm.	559
<i>tenuissimum</i> , Christ	102
<i>Teysmanni</i> , V. D. B.	91
<i>trichophyllum</i> , Moore	106
<i>tunbridgense</i> , Sm.	797

Trichomanes	
<i>vestitum</i> , <i>Bk.</i>	96
<i>Warburgii</i> , <i>Christ</i>	97
<i>Wernerii</i> , <i>Rst.</i>	799
Trichomanopsis	84
Triphlebia	479
<i>dimorphophylla</i> , <i>Bk.</i>	480
<i>linza</i> , <i>Bk.</i>	479
<i>longifolia</i> , <i>Bk.</i>	480
<i>pinnata</i> , <i>Bk.</i>	480

V.

Vittaria	553
<i>acrostichoides</i> , <i>Hk. & Grev.</i> .	712
<i>alternans</i> , <i>Copel.</i>	559
<i>amboinensis</i> , <i>Fée</i>	561
<i>angustifolia</i> , <i>Bl.</i>	555
<i>Bensei</i> , <i>v. A. v. R.</i>	557
<i>caudiformis</i> , <i>Bl.</i>	554
<i>crassifolia</i> , <i>Bk.</i>	555
<i>debilis</i> , <i>Kuhn</i>	559
<i>divergens</i> , <i>Rb.</i>	278
<i>elongata</i> , <i>Sw.</i>	556
<i>var. ensiformis</i>	555
<i>var. genuina</i>	556
<i>var. zosterifolia</i>	556
<i>ensifolia</i> , <i>Bl.</i>	555
<i>ensiformis</i> , <i>Sw.</i>	555
<i>falcata</i> , <i>Kze</i>	560
<i>hirta</i> , <i>Fée</i>	558
<i>intermedia</i> , <i>Bl.</i>	555
<i>lineata</i> , <i>Sm.</i>	561

Vittaria	
<i>lloydiiifolia</i> , <i>Rac.</i>	554
<i>longicoma</i> , <i>Christ</i>	559
<i>Merrillii</i> , <i>Christ</i>	557
<i>minor</i> , <i>Fée</i>	560
<i>pachystemma</i> , <i>Christ</i>	556
<i>parvula</i> , <i>Bory</i>	559
<i>philippinensis</i> , <i>Christ</i>	562
<i>plantaginea</i> , <i>Hk. & Grev.</i> ...	555
<i>pumila</i> , <i>Mett.</i>	559
<i>pusilla</i> , <i>Bl.</i>	553, 560
<i>rigida</i> , <i>Klf.</i>	555
<i>scolopendrina</i> , <i>Thw.</i>	561
<i>scolopendrioides</i> , <i>Mett.</i>	561
<i>subcoriacea</i> , <i>Christ</i>	558
<i>sulcata</i> , <i>Kuhn</i>	560
<i>taeniophylla</i> , <i>Copel.</i>	561
<i>zosterifolia</i> , <i>Willd.</i>	556

W.

Wibelia	314
<i>amboynensis</i> , <i>Cuhn.</i>	315
<i>Denhami</i> , <i>Kuhn</i>	316
<i>gracilis</i> , <i>Christ</i>	315
<i>oligophlebia</i> , <i>Christ</i>	314
<i>pinnata</i> , <i>Fée</i>	315
Woodsieae	121
Woodwardia	590
<i>auriculata</i> , <i>Bl.</i>	591
<i>prolifera</i> , <i>Hk. & Arn.</i>	591
<i>radicans</i> , <i>Sm.</i>	591

ADDITIONS, MODIFICATIONS AND CORRECTIONS.

This list principally contains additions and modifications that could not be inserted in the Appendix; most of them may be cut out and pasted on the places indicated.

Typographical and grammatical errors of *no* importance with regard to the purpose of the work are *not* considered.

(P. = Page; l. = line; fr. b. = from bottom).

P. XXXI, l. 1—7 *should be modified as follows:*

GENUS 66. SYNGRAMMA. Veins mostly uniting 1—3 \times at or near the margin, sometimes anastomosing copiously, but then distinctly visible like the costa; areolae without free included veinlets. No main veins. Sori with paraphyses.

GENUS 67. HEMIONITIS. Veins anastomosing copiously, distinctly visible like the costa; areolae without free included veinlets. No main veins. Sori without paraphyses.

GENUS 67a. HEMIGRAMMA. Differs from genera 66 and 67 by the distinct main veins, and the areolae provided with free veinlets.

P. XXXVII, l. 4 *should be modified as follows:*

86a *Leptochilus (Gymnopteris)*, 67a *Hemigramma*.

P. 3, l. 2 fr. b.; cuneate, *should be*: clavate,

P. 14, l. 14 fr. b.; *rupturing irregularly.* *should be*:

rupturing irregularly or breaking down by a slit.

l. 5—1 fr. b. *should be modified as follows*:

§. Veins \pm 4—5 on each side of the costulae.

(21) C. Havilandii.

P. 15, l. 1—2 *should be modified as follows*:

§.§. Veins \pm 7—10 on each side of the costulae.

l. 5—4 fr. b.; *brunonis*, *should be*: *Brunonis*,

P. 17, l. 4; (not *Hk.*), Christ, Farnkr. d. Erde, 327; *should be*: Bedd., Handb. Ind. Ferns, Suppl., 2;

P. 21, l. 5; serrate *should be*: or serrate

l. 7 fr. b. (et passim); lower 1—3 veins *should be*: 1—5 lower veins

P. 23, l. 4; to $1\frac{1}{2}$ c.M. *should be*: to 2 c.M.

l. 2 fr. b.; soon deciduous *should be*: caducous.

P. 27, l. 7 fr. b.; near the base. *should be*: occupying the lower veins.

P. 28, l. 2 fr. b.; fimbriated *should be*: ferruginous, fimbriate

P. 32, l. 11—12; not divaricating. *should be*: whether or not divaricating.

l. 15; *Omit*: (not *Hk.*),

P. 37, l. 13; **A. caudata**, **J. Sm.** — *Omit this, it being* **Cyathea caudata**, **Copel.** — See the Appendix.

l. 16 fr. b. (et passim); semicoriaceous *should be*: subcoriaceous

P. 39, l. 10; broadly ovate *should be*: broadly oval,

l. 14 & 5 fr. b.; var *congesta*. — *Omit this, it being a variety of* *Al-sophila calocoma*, *Christ.* — See the Appendix.

P. 40, l. 17 fr. b.; **A. ornata**, **Scott** & var. — *Insert*: See the Appendix.

P. 41, l. 4 fr. b.; less glaucous *should be*: under surface less glaucous.

P. 47, l. 10; * **D. Blumei**, **Moore** *should be*: * (1) **D. Blumei**, **Moore.**

P. 48, l. 5; **C. barometz**, **J. Sm.** — *Insert*: See the Appendix.

P. 55, l. 13; *Omit the comma after*: sporangia

l. 11 fr. b.; *after* recognized by, *insert*: a distinct articulation or

P. 58, l. 10; **G. glauca**, **Hk.** — *Insert*: See the Appendix.

l. 7, fr. b.; bifurcate *should be*: \pm 2 \times furcate

P. 59, l. 16; *after long, insert: entire or emarginate at the apex.*

P. 65, l. 11; *for: climates—read: zones.*

P. 67, l. 1; P. 68, l. 17; P. 69, l. 16 fr. b.; *for: obovate—read: ovate*

P. 70, l. 5; **H. subdemissum**, *Christ.* — *Insert: See the Appendix.*

l. 5 fr. b.; *obovate should be: ovate,*

P. 71, l. 8 fr. b.; **H. Blumeanum**, *Spr.* — *Insert: See the Appendix.*

P. 71, l. 3 & 17 fr. b.; P. 72, l. 4; P. 75, l. 3 & 10, & 7 fr. b.; P. 74, l. 1 & 7; P. 77, l. 17 fr. b.; P. 78, l. 7 & 13 fr. b.; P. 80, l. 12 fr. b.; P. 95, l. 12 fr. b.; P. 94, l. 16; P. 95, l. 7; P. 96, l. 13 fr. b.; P. 104, l. 1; P. 143, l. 8; *for: obovate — read: ovate*

P. 75, l. 13—10 fr. b. should be modified as follows:

*. *Indusium without crest-like ridges at the base.*

(29) *H. tunbridgense.*

.. *Indusium with some (1 or more) indistinct, hardly elevated, longitudinal, crest-like ridges at the base.*

(30) *H. holochilum.*

P. 76, l. 13; *on the back. should be: on the back throughout.*

l. 17—18 should be modified as follows:

1.1.1. *Indusium provided with irregularly spinoso-fimbriate appendages on the back at the base.*

l. 15 fr. b.; *after down, insert: valves retuso-denticulate.*

l. 13 fr. b.; *after down, insert: valves sharply toothed.*

l. 11—10 fr. b. should be modified as follows:

1.1.1.1. *Indusium provided with rather close, longitudinal crests on the back at the base.* (44) *H. brachyglossum.*

P. 78, l. 14; *climates should be: zones*

l. 13—12 fr. b.; *surrounded by a row of longitudinal crests, should be: provided with some (1 or more) indistinct, hardly elevated, longitudinal, crest-like ridges,*

P. 81, l. 15; *on te back, should be: on the back throughout,*

P. 82, l. 4—5 *should be modified as follows:*

of the way down, the base provided with deeply and irregularly spinoso-fimbriate appendages, the valves slightly retuso-denticulate towards the apex.

- P. 82, l. 6 fr. b.; *for*: surrounded by a row of — *read*: provided with
P. 84, l. 1, & 8 fr. b.; ovate *should be*: oblong,
P. 91, l. 11 fr. b.; *for*: *appearantly* — *read*: *apparently*
P. 93, l. 10 (et passim); *for*: 2^d — *read*: 2nd
P. 99, l. 17; **T. maximum, Bl.** — *Insert*: See the Appendix.
P. 100, l. 15 fr. b.; **T. javanicum, Bl.** — *Insert*: See the Appendix.
P. 109, l. 4 fr. b.; *Omit*: (Not in Malaya). — See the Appendix.
P. 121, l. 16 fr. b.; fastened all round the receptacle, *should be*: fastened all round the receptacle, but apparently so in *Diplaziopsis*,
P. 122, l. 15; *for*: not articulate &c. — *read*: rarely articulate to the rhizome.
P. 123, l. 4—6 *should be modified as follows*:

2. Indusium cupuliform, globose or sausage-shaped, fastened beneath the sorus on all sides, but apparently so in genus N^o. 13, open at the top or bursting down irregularly. § 1. **Woodsiae**.

P. 126—127.

§ 2. **DAVALLIEAE.** — *Modify the key to the genera*:

1. Sori solitary and terminal on the veins; indusium attached by a very narrow base *to the apex of the vein only*, otherwise free, the outer edge lacerato-lobate. No. 31b. **Leptolepia**.
- 1.1. Sori solitary and dorsal or terminal on the veins, or uniting the apices of 2 or more veins; indusium attached by a broad base or by the base and sides *to the vein and the parenchyma*, the outer edge rather entire.
2. Fronds divided; leaflets dimidiate, i. e. the half on the lower side of the *midrib* cut away. No. 29. **Lindsaya**.
- 2.2. Fronds simple or divided, if divided the leaflets not dimidiate.
3. Stipes articulate to the rhizome; rhizome scaly. Sori roundish or oblong, solitary on the vein tips.
4. Indusium fastened by a broad base, the sides and outer edge free. No. 32. **Humata**.
- 4.4. Indusium fastened by the base and sides, the outer edge free. No. 34. **Davallia**.
- 3.3. Stipes not articulate to the rhizome; rhizome hairy, scaly or fibrillose.
- a. Sori occupying the broad apex of the ± cuneate ultimate segments. No. 28. **Odontosoria**.
- a.a. Sori not as in a. Ultimate segments not ± cuneate.

THE UNIVERSITY OF CHICAGO

PHYSICS DEPARTMENT

PHYSICS 311

LECTURE 10: ELECTROSTATICS

1. Electric field of a point charge

2. Gauss's law

3. Electric potential

4. Capacitance

5. Energy of a charge distribution

6. Summary

- b.** Sori uniting the tips of 2 or more veins or veinlets.
- *. Fronds pinnate, pinnae articulate to the rachis.
(No. 17. *Nephrolepis*, § *Lindsayopsis*).
- *.*. Fronds simple or divided, if divided the leaflets not articulate to the rachis.

No. 50. **Schizoloma.**

- b.b.** Sori dorsal or terminal, solitary on the veins or veinlets, sometimes confluent in pairs.
- c.** Sori dorsal, or terminal on very short, rather inconspicuous lateral veinlets.
- ×. Indusium attached at the base, often cucullate, at length reflexed.

No. 27. **Cystopteris.**

- ×.×. Indusium attached at the base and sides, half-cupshaped, never reflexed.
- (No. 55. *Asplenium*, § *Loxoscapha*).

- c.c.** Sori always terminal on the veins or rather long veinlets.
- d.** Sori placed in conspicuously cucullate marginal outgrowths, which are closed by the ± flattened, transversally oblong indusia.

No. 31a. **Saccoloma** (*sorbifolium*).

- d.d.** Sori not as in **d.**

- e.** Rhizome erect. Stipes tufted.
- +. Indusium cuneate, open at the outer edge. (*moluccanum*).

No. 31a. **Saccoloma**

- +.+. Indusium roundish, attached by a broad base, otherwise free.

No. 33. **Acrophorus.**

- e.e.** Rhizome creeping. Stipes remote or approximate.
- f.** Segments rhomboidal or linear-triangular, unequal-sided at the base. Receptacle distinct.

No. 35. **Microlepia.**

f.f. Segments oblong or linear, not or hardly unequal-sided at the base. Receptacle indistinct.

No. 56. **Tapeinidium.**

P. 128, l. 14; bearing \pm 5—6 *should be*: bearing generally \pm 2—12

l. 11 fr. b.; No. 42. *should be*: No. 42a.

l. 10—9 fr. b. *should be modified as follows*:

3.3. Sori roundish, on terminal, \perp -shaped expansions of the veins or veinlets, or uniting the apices of the branches of forked veins. No. 42b. **Schizostege.**

3.3.3. Sori linear or linear-oblong, the receptacle uniting the vein tips, sometimes running the veins transversally.

a. &c.

P. 129, l. 16; No. 45. *should be*: No. 45a.

l. 19—17 fr. b. *should be modified as follows*:

*. Stipes black. Fronds simple or divided, if divided the segments adnate, connected at the base. § *Eudoryopteris.*

No. 42a. **Doryopteris,**

.. Stipes not black. Fronds divided, at least the primary segments not connected, not adnate at the base.

×. Sori occupying both anterior and posterior edges of the ultimate segments. No. 45a. **Pteris.**

×.×. Sori occupying only the posterior edge of the ultimate segments, the anterior side destitute of sori.

No. 45b. **Hemipteris.**

P. 132, l. 11; **Syngramme** *should be*: **Syngamma,**

l. 8—4 fr. b. *should be modified as follows*:

3.3. Fronds thinner; veins distinctly visible when seen towards the light. Sori free or reticulate, running along the veins.

1914

1915

1916

1917

1918

1919

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

1932

1933

1934

1935

1936

1937

1938

1939

1940

1941

1942

1943

1944

1945

1946

1947

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

4. Main veins present. Areolae with free included veinlets. No. 67a. **Hemigramma**.
- 4.4. Main veins wanting. Areolae without free veinlets.
5. Sori with paraphyses. No. 66. **Syngamma**, § *Dictyogramme*.
- 5.5. Sori without paraphyses. No. 67. **Hemionitis**.

P. 133, l. 2; **Monogramme** should be: **Monogramma**.

P. 134, l. 3—9 should be modified as follows:

- §. Sori more or less oblique to the costa and margin, many on each side of the costa. No. 71. **Polypodium**, § *Selliguea*.
- §.§. Sori parallel to the costa or margin, 1 on each side of the costa.
- *. Sori in marginal or submarginal, 2-lipped grooves. Grass-like plants. (No. 69. *Vittaria*).
- *.*. Sori not in marginal or submarginal grooves. No grass-like plants.
- +. Fertile fronds simple, as are the barren ones. No. 75. **Drymoglossum**.
- +.+. Fertile fronds 3-partite, barren sagittate. No. 76. **Christopteris**.

P. 139, l. 4 fr. b.; **13. DIPLAZIOPSIS**, *Christensen*. — *Insert*: See the Appendix.

P. 141, l. 11; for: *creeping*. — read: *creeping or erect, but not arborescent*.

l. 10 fr. b.; for: *Stipes ebeneous, muricate*. — read: *Stipes ebeneous or castaneous, muricate or aculeolate*.

P. 152, l. 15 fr. b.; P. 153, l. 1; obovate — should be: lanceolate,

l. 17 fr. b.; after var. *brachypus*, *insert*: See the Appendix.

P. 155, l. 7; scaly should be: sparingly scaly.

l. 14; after scaly, *insert*: not rarely glabrous in age.

P. 157, l. 11; *lanuginosa*, *Hk.*, should be:

lanuginosa, *Hk.* (not *Wall.*),


P. 160, l. 10 fr. b.; *cordifolia*, should be: *cordifolia*,

P. 162, l. 12; C. Chr., *Ind. Fil.*; should be: *Gen. Fil.*, tab. III.;

l. 9 fr. b.; on should be: or on

P. 164, l. 9 fr. b.; *scaly or fibrillose*. should be:

scaly, fibrillose or hairy.



- P. 164, l. 7 fr. b.; *rather coriaceous.* should be: *papyraceous to coriaceous.*
l. 5 fr. b.: *ferrugineous.* should be: *ferrugineous or fulvous.*
- P. 165, l. 2; *scaly or fibrillose.* should be: *scaly, fibrillose or hairy.*
- P. 166, l. 4; *for: 165* — *read: 165*;
l. 7; *for: 926* — *read: 996*;
- P. 168, l. 4 fr. b.; *for: 19* — *read: 18*;
- P. 171, l. 18—19° should be modified as follows:
- d.* *Surfaces naked.* (7) P. Trimeni.
d.d. *Upper surface ± hairy towards the margin, lower slightly puberulous.* (8) P. devexa.
- l. 14 fr. b.; *Mett.*, should be: *Hk.* (not *Mett.*),
- P. 175, l. 6; **P. gigantea**, *Pr.* — *Insert: See the Appendix.*
l. 17 fr. b.; *for: Farns, 255* — *read: Ferns, 225*,
l. 5 fr. b.; *for: P. Bakeri, v. A. v. R.*—*read: P. Trimeni, Bedd., Handb. Ind. Ferns, 225.* — See the Appendix.
- P. 174, l. 12; **P. devexa**, *v. A. v. R.* — See the Appendix.
- P. 180, l. 10 fr. b.; *C. Chr.*, should be: *O. Ktze*, C. Chr.,
- P. 186, l. 3; *Schott*, should be: *Scott*,
- P. 188, l. 14 fr. b.; P. 189, l. 14; *obovate should be: ovate*
- P. 197, l. 16 fr. b.; *for: 178* — *read: 179.*
- P. 199, l. 10 & 8 fr. b.; *obovate should be: ovate*
- P. 202, l. 7; *for: XCX* — *read: XCV*;
l. 3 fr. b.; *for: tab. CCLXIX* — *read: 142* (not tab. CCLXIX);
l. 2 » ».; *after var. pallida,* — *insert: See the Appendix.*
- P. 205, l. 15 fr. b.; *for: pinnae* — *read: ones*
- P. 209, l. 2 fr. b.; P. 210, l. 1; *for: otaria* — *read: Otaria*,
- P. 210, l. 10 fr. b.; *for: Bedd.*, — *read: Par.*, Bedd.,
- P. 215, l. 16 fr. b.; *for: caute* — *read: acute*,
- P. 214, l. 16 fr. b.; *reduced pinnae. should be: reduced lower pinnae.*
- P. 225, l. 6; **D. caudiculata**, *v. A. v. R.* — *Insert: See the Appendix.*
- P. 228, l. 6; 2—5 *should be: 2—5*
- P. 230, l. 11 fr. b.; *for: Nephrodium* — *read: Aspidium*
- P. 252, l. 3; *for: 84* — *read: 94*;
- P. 256, l. 5 fr. b.; **Kunstleri**, *should be: Künstleri*,
- P. 241, l. 16; *for: Everettii*, & Everetti, — *read: Everettii*, & Everettii,

- P. 244, l. 14; *For*: 49 — *read*: 59,
P. 251, l. 14 fr. b.; *Mett.*, should be: *Mett.* (not *Bl.*),
P. 252, l. 11 fr. b.; *impressum* should be: *immersum*,
P. 253, l. 8; *for*: 206 — *read*: 106.
P. 257, l. 15 fr. b.; *tristis* should be: *triste*,
P. 259, l. 3; *Burck*, should be: *Mett.*,
P. 260, l. 14; *for*: *tendence* — *read*: *tendency*
P. 262, l. 18 fr. b.; *for*: 79 — *read*: 97,
P. 267, l. 5 fr. b.; *for*: 148 — *read*: 149
P. 268, l. 15; *for*: 298 — *read*: 208,
P. 269, l. 9 fr. b.; *for*: *Syn. Fil.* — *read*: *Syn. Fil.*, 106;
P. 270, l. 11; *flbellulata* should be: *flabellulata*,
l. 14; *for*: I, tab. LXIII — *read*: I, 211, tab. LXIII, A;
l. 15; *for*: *Ferns S.I.*, tab. XXIV; *L. tenera*, *Dry.*, l. c., tab. X;
— *read*: l. c.; *L. tenera*, *Dry.*, l. c., tab. X; *Bedd.*, *Ferns S.I.*,
tab XXIV;
P. 274, l. 9; *for*: *S. recurvata*, *Bedd.* — *read*: *S. recurvatum*, *Moore*, *Bedd.*,
P. 281, l. 12; *for*: 148 — *read*: 149
P. 282, l. 7, & 2 fr. b.; *for*: *inequalis* — *read*: *inaequalis*,
l. 3 fr. b.; *for*: *Diels*, etc. — *read*: *Christ*, *Farnkr. d. Erde*, 310;
P. 285, l. 3 & 9; *for*: *higher ones* — *read*: *following*.
P. 286, l. 5; *for*: **J. Sm.**, *Copel.*, — *read*: **J. Sm.** (not *Wall.*), *Copel.*,
P. 287, l. 2; *for*: *intramarginalis*, *Bl.*, — *read*: *intermarginalis*, *Moore* (not *Bl.*),
l. 13; *for*: (not *Sm.*) — *read*: (not *J. Sm.*),
l. 4 fr. b.; CXXXVII should be: CXXXVIII,
P. 288, l. 4 fr. b.; *pedeta* should be: *pedata*,
l. 3 fr. b.; *L.*, should be: *L.f.*,
P. 289, l. 13 fr. b.; *XVL* should be: *XLV*,
P. 300, l. 8; *for*: **Mett.**, — *read*: **Kuhn**,
P. 307, l. 12; *for*: (*fejensis*) — *read*: (*fejeensis*),
P. 509, l. 3 fr. b.; *Hk.* should be: *Goldm.* (not *Hk.*),
l. 16 fr. b.; XXVI should be: XXXVI,
P. 311, l. 16; *for* **Copel.**, — *read*: **Copel.** (not *Harr.*),
P. 312, l. 8 & 18 fr. b.; *Davalia* should be: *Davallia*
P. 313, l. 4 fr. b.; **Chistensen.** should be: **Christensen.**
P. 315, l. 1; *for*: 7 & 8 — *read*: 5 & 7;
l. 18; *for*: *Christ*, — *read*: *Christ* (not *Bl.*),
l. 3 & 5 fr. b.; *amboinensis* should be: *amboynesis*,



- P. 319, l. 3 fr. b.; *for*: With view on — *read*: In view of
P. 323, l. 16; *for*: *Hk.*, — *read*: *Hk.* (not *Willd.*),
P. 336, l. 1 & 4 fr. b.; *for*: 116 — *read*: 146
P. 345, l. 2; *for*: *Burm.* — *read*: *R. Br.*,
P. 347, l. 4; *for*: 146 — *read*: 151;
P. 359, l. 14 fr. b.; *for*: *spinulose serrate* — *read*: *spinulose or serrate*
P. 367, l. 8; *for*: in and size — *read*: in size and
P. 380, l. 17 fr. b.; **L. areolaris**, *Harr.* — Omit this. See the Appendix.
P. 381, l. 1 fr. b.; *Cunn.* should be: **A. Cunn.**,
P. 412, l. 12; major *should be*: majus,
l. 17 fr. b.; **major** *should be*: **majus**:
P. 435, l. 15; **A. umbrosum**, *Pr.* — Omit this. See the Appendix.
l. 4 fr. b.; **A. silvaticum**, *Milde.* — Omit this. See the
Appendix.
P. 450, l. 2 fr. b.; *for*: group — *read*: group **a.a.**
l. 1 fr. b.; *for*: *Fors* — *read*: *Forst.*
P. 510, l. 8; *for*: *sumbensi* — *read*: *sumbensis*,
P. 535, l. 13; *A. semicostatum* should be: *A. semicostatum*,
P. 540, l. 1; P. 544, l. 11; § 4. **HEMIGRAMMA.** — Omit this. See the Appendix.
P. 544, l. 12; **S. Zollingeri**, *Diels.* — Omit this. See the Appendix.
P. 547, l. 1; **H. gymnopteroidea**, *Copel.* — Omit this. See the Ap-
pendix.
P. 551, l. 5; *for*: *wit* — *read*: *with*
P. 570, l. 12; *for*: 1. — *read*: **1.**
P. 575, l. 2 & 1 fr. b. *should be modified as follows*:

(5) **P. Friderici et Pauli**, *Christ*, Ann. Btz., XV, 148, tab. XVI,
fig. 21, a & b (not XIX, 37).


- P. 585, l. 3 fr. b.; *for*: 2—3 × forked — *read*: 2—3-furcate,
P. 598, l. 2 fr. b.; *for*: in — *read*: is
P. 617, l. 14 fr. b.; *for* *integriore.* — *read*: *integrius.*
l. 11—10 fr. b. *should be modified as follows*:

*. *Sori slightly immersed; paraphyses yellow-brown.*

(116a) *P. Koningsbergeri.*

.. *Sori deeply immersed; paraphyses black-brown.* (116b) *P. verrucosum.*

- P. 618, l. 12 fr. b.; *for*: **integriore**, — *read*: **integrius**,



P. 619, l. 19; one *should be*: one).

l. 8—7 fr. b. *should be modified as follows*:

the Agricultural Department in Netherlands India. The description is taken from 3—4 years old plants, and I am inclined to suppose that the adult plants have the fronds large and more copiously pinnate.

Java (Preanger Regencies, v. A. v. R.).

P. 622, l. 9 fr. b.; *for*: 8—50 — *read*: 850

l. 7 fr. b.; *for*: $\frac{1}{2}$ — $2\frac{1}{2}$ — *read*: $\frac{1}{2}$ —5

P. 625, l. 17; *for*: *var. pumilum*; — *read*: *var. pumilum*;

P. 677, l. 14; *decurrens*, Hk. — *should be*: *decurrens*, Hk.,

P. 682, l. 6 fr. b.; **spaerostichus** *should be*: **sphaerostichus**,

P. 684, l. 15; Mett., *should be*: Mett.,

P. 688, l. 9 fr. b.; Kze, *should be*: Kze,

P. 778, l. 11 fr. b.; *for*: *subcarnosmm*, — *read*: *subcarnosum*,

P. 824, l. 2 fr. b.; **Wall.** *should be*: **Mett.**

P. 848, l. 10; *devexum*, Kze *should be*: *devexum*, Kze



587.
A355
F.C.

MALAYAN FERNS.

HANDBOOK TO THE DETERMINATION

OF

THE FERNS

OF

THE MALAYAN ISLANDS

(INCL. THOSE OF THE MALAY PENINSULA, THE PHILIPPINES
AND NEW GUINEA).

BY


CAPT. C. R. W. K. VAN ALDERWERELT VAN ROSENBURGH.

(CORRECTING SHEET).

PUBLISHED BY

THE DEPARTMENT OF AGRICULTURE
NETHERLANDS INDIA.

BATAVIA
LANDSDRUKKERIJ
1909.



MALAYAN FERNS.

ADDITIONS, MODIFICATIONS

AND

CORRECTIONS.

II.

List **A** contains some additions and modifications to my handbook on MALAYAN FERNS, which were omitted in the Appendix and in the former list of additions and modifications.

List **A** is an enumeration of grammatical and typographical errors, the correction of which is advisable, though not positively necessary to the purpose of the work.

Most of these corrections may be cut out and pasted over the places indicated.

(P. = Page; l. = line; fr. b. = from bottom).

TOKYO, June 1909.

v. A. v. R.

A.

P. XXXII, l. 10 fr. b.; P. 122, l. 2 fr. b. — *For*: fertile segments.

— *read*: fertile fronds or segments.

P. XL, l. 1 fr. b. — *For*: *fork branch, &c.* — *read*:

fork branch, undulate (wavy) for sinuated with rounded lobes and sinuses, joining (joined) for connected, united or confluent, curved sometimes for flexuose, &c.

P. 3, l. 7 fr. b. — *Strike out the comma after sporangia*

P. 4, l. 3 fr. b. — *Strike out the comma after segments*



- P. 15, l. 13 fr. b. — Strike out the comma after *vault*
- P. 16. **C. philippinensis**. — Strike out the description of this species. — *Insert*: See the Appendix.
- P. 21, l. 5—6. — *Strike out*: Sori close to the costae, 1 on each side of the costulae. — *Insert*: See the Appendix.
- P. 55, l. 10, & l. 5 fr. b. — *For*: Tropical—*read*: Tropical,
— *after*: hemisphere, — *insert*: rarely northern.
- P. 59, l. 19 fr. b. — **Var. normalis**:
— *insert*: See the Appendix.
- P. 77. **H. Kurzii**. — *Insert*: See the Appendix.
- P. 79. **H. serrulatum**. — *Insert*: See the Appendix.
- P. 85, l. 18 fr. b. — near its centre — *should be*:
near the centre
- P. 97, l. 7—8 (et passim). — *For*: Texture (surfaces) pale-coloured.
— *read*: Texture (surfaces) pale (*eventually* pale-green).
- P. 112, l. 9. **Var. cristatum**: — *Insert*: See the Appendix.
L. trifurcatum. — *Insert*: See the Appendix.
- P. 125, l. 2—1 fr. b. *should be modified as follows*:

or sometimes extending on both surfaces in the plants with much contracted fertile fronds or segments. § 11. **Acrosticheae**.

- P. 124, l. 14; P. 132, l. 19. — *For*: free or reticulat'e.
— *read*: free, joining or reticulate.
- P. 142, l. 11 fr. b. (et passim).
— *For*: surfaces, paler coloured
— *read*: surfaces, paler
- P. 152. **O. cuspidata**. — *Insert*: See the Appendix.
- P. 153, l. 2 fr. b. — *For*: Sibaldii, Sibaldii,
— *read*: Sibbaldii, Sibbaldii,
- P. 154, l. 6. — *For*: **Sibaldii**: — *read*: **Sibbaldii**:
l. 8. — *For*: Sibaldii: — *read*: Sibbaldii:
- P. 158, l. 1 & 5 should be modified as follows:

b. *Sori at the tips of teeth or lobes.*

b.b. *Sori not at the tips of teeth or lobes.*

Strike out the note at the base of page 158.

P. 159, l. 17. — *For*: Sori submarginal, 1 to each crenation;

— *read*: Sori placed at the tips of the teeth;

P. 175. — § 3. **CYCLOSORUS**. &c should be modified as follows:

§ 3: **CYCLOSORUS**. Like § 2, but the indusium peltate, and round or cordate. Sori not confluent. Fertile segments not or hardly contracted.

§ 4. **CYCLODIUM**. Like § 3, but the sori confluent and the fertile segments distinctly contracted. Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 101, C—E. (Not in Malaya).

P. 176, l. 6 fr. b. — *For*: membranaceous or herbaceous.

— *read*: membranaceous, herbaceous or somewhat fleshy.

P. 177, l. 5—11 should be modified as follows:

!. Under surface villous, the hairs not rigid. (17) *D. calcarata*.

!.!. Under surface naked or glandular, or setose with rigid hairs.

✧. Naked or slightly glandular.

(19) *D. gracilescens*.

✧.✧. Densely glandular or setose.

(18) *D. viscosa*.

P. 183, l. 9 & 1 fr. b. — *N. Webbianum*, *Hk.* — Strike out this synonym and its description. — *Insert*: See the Appendix.

P. 186. **D. viscosa**. — Strike out the description of this species. — *Insert*: See the Appendix.

P. 188, l. 8 fr. b. — *For*: only those on the upper base

— *read*: only the lowest anterior segment

P. 197. **D. Raciborskii**. — *Insert*: See the Appendix.


P. 220. **D. brachyodus**. — *Insert*: See the Appendix.

P. 221. **D. Loheriana**. — *Insert*: See the Appendix.

P. 280, l. 5 fr. b — *For*: Australia and Polynesia.

— *read*: Australia, Polynesia, and West and South Africa.

P. 294, l. 13 fr. b. — *For*: Davalia — *read*: Davallia



P. 297, l. 17. — *For*: 2—3 pairs — *read*: 2—4 pairs

P. 319, l. 6. — *For*: the sori on its inner surface.

— *read*: the sporangia on its inner side.

P. 376, l. 7. — *For*: **tripartia**, — *read*: **tripartita**,

P. 517, l. 17. — *After* v. A. v. R., l. c., — *insert*: tab. I.

P. 539, l. 8—7 fr. b. *should be modified as follows*:

nately arranged, free or anastomosing more or less copiously; no main veins; areolae without free included veinlets.

Malaya to Polynesia.

P. 540, l. 4—5 *should be modified as follows*:

§ 4. **DICTYOGRAMME**. Veins anastomosing more copiously. Sori free or reticulate.

P. 544, l. 4 fr. b. — *For*: § 5. — *read*: § 4.

P. 546, l. 7—5 fr. b. *should be modified as follows*:

tinct, anastomosing copiously, bearing the sori throughout; no main veins, but the fronds provided with 3 or more radiating ribs originating from the base, 1 to each lobe or segment; areolae without free veinlets. Hk. Bk., Syn. Fil., tab. VI, fig. 59, a; Diels, in Engl. & Prantl, Nat. Pfl.Fam., I⁴, fig. 138, C—D.

A small genus, almost confined to the tropics.

P. 546, l. 4—1 fr. b. *should be modified as follows*:

✧. *Barren fronds cordate, fertile ones sagittate.*

(2) H. arifolia.

✧✧. *Both barren and fertile fronds palmately 5-lobed.*

(3) H. palmata.


P. 571, l. 11. — *After round.* — *insert*: (*Cyclocarpium*).

l. 11 fr. b. — *After oblong.* — *insert*: (*Grammitis*).

P. 655, l. 11. *For*: *integrius*, Copel., l. c. — *read*: *integrius* (*integriore*), Copel., l. c.

P. 737, l. 18. — *For*: *latifolia* — *read*: *latifolia*

P. 762. **A. evecata**. — *Insert*: See the Appendix.



- P. 796, l. 10 fr. b. — *For: loheri*, — *read: Loheri*,
P. 800, l. 17 fr. b. — *For: asplenoides*, — *read: asplenoides*,
P. 811, l. 11. — *For: Bedd.*, — *read: Moore, Bedd.*,
 l. 11 fr. b. — *For: from* — *read: form*
 l. 2 fr. b. — *For: Bedd.*, — *read: Moore*,
P. 843, — After *Acrosorus*. — *insert: 613*,
P. 857. *Cyclopeltis Presliana*. — *For: J. Sm.* — *read: Berk.*
 Above Cyclopeltis — *insert:* *Cyclocarpium*..... 571
 Cyclodium..... 175
P. 858. *Davallia dissecta*. — *For: Moore* — *read: J. Sm.*
P. 863. *Drynaria sparsisora*. — *For: Moore* — *read: Moore*
P. 867. — *Above Eupteris*. — *insert: Euprosaptia*..... 613
P. 868. — After *Grammitis*. — *insert: 571*,
P. 881, l. 19. — *For: Sibaldii*, — *read: Sibaldii*,
P. 885, l. 8. — *For: affine*, — *read: affine*,
P. 888. *Polypodium macrophyllum*. — *For: Bk.* — *read: Rw.*
 l. 12. — *For: integrus, Copel.*
 — *read: integrus (integriore), Copel.*
P. 895. *Sagenia gigantea*. — *For: Bedd.* — *read: Moore.*

B.

For: surfaces....., upper (the upper, the upper one, upper one) dark-
coloured, lower (the lower, the under one, under one) paler coloured
(pale-coloured, paler in colour) —


read: surfaces.....: upper surface dark (*eventually* dark-green), lower
paler (pale or *eventually* pale-green)

P. XXII, l. 3 & 6 (et passim). — *Strike out the comma after many and few
and wherever it is erroneously placed after a numeral, when this (the
numeral) precedes the object defined.*

P. XXIV, l. 17; P. 258, l. 17. —

For: Indusium ± of the same shape, attached at

read: Indusium ± of the same shape as and attached at



P. XXVI, l. 5—6 (et passim) —

For: Indusium \pm of the same shape,

read: Indusium \pm the shape of the sorus,

P. XXX, l. 14 & 13 fr. b. *should be modified as follows:*

costa or the margin, free, joining or reticulate.

Fronds not articulate to the rhizome.

P. XXXI. **Tribe IX. Grammitideae.** (et passim). —

For: each side of the costa, parallel to the latter

read: each side of and parallel to the costa

P. 4, l. 11 & 5 fr. b.; P. 5, l. 7 fr. b. —

For: without ring. — *read:* without a ring.

P. 24, l. 5 fr. b. — *For:* oblique, — *read:* oblique,

P. 28, l. 3 fr. b. — *For:* densly — *read:* densely

P. 31, l. 9. — *For:* *worl* — *read:* *whorl*

P. 33, l. 13 fr. b. (et passim). — *For:* sinusses — *read:* sinuses

P. 43, l. 8 fr. b.; P. 58, l. 10 fr. b.; P. 93, l. 4 fr. b.; P. 203, l. 12.

For: between — *read:* among

P. 58, l. 1 fr. b. — *For:* semi- — *read:* sub-

P. 59, l. 15 fr. b. (et passim). — *For:* troughout (troughtout, throughtout) —
read: throughout

P. 70, l. 4 fr. b. — *For:* schort — *read:* short

P. 75, l. 18; P. 359, l. 19. — *For:* *troughout* — *read:* *throughout*

P. 79, l. 8. — *For:* exerted, — *read:* exserted,

P. 85, l. 5 fr. b. — *For:* without costa;

— *read:* without a costa;

P. 89, l. 2. — *For:* channeled), — *read:* channelled),

P. 92, l. 12. —

For: *semicoriaceous*. — *read:* *subcoriaceous*.

P. 96, l. 16; P. 103, l. 15 fr. b.; P. 181, l. 2; P. 183, l. 19; P. 201, l. 1
fr. b.; P. 215, l. 13 fr. b.; P. 227, l. 13 fr. b.; P. 228, l. 17; P.
236, l. 18; P. 246, l. 15 fr. b.; P. 146, l. 14 fr. b. —

For: *semicoriaceous* — *read:* *subcoriaceous*

P. 102, l. 5. — *Strike out the comma after* marginate

P. 103, l. 15 fr. b. — *For:* schining; — *read:* shining;

P. 104, l. 14 fr. b. — *For:* dillated, — *read:* dilated,

l. 7 & 6 fr. b. — *For:* *into* — *read:* *in*

l. 10 fr. b. — *For:* *var. polyanthos*: — *read:* *var. polyanthos*:

and various other things, and the
... ..

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

- P. 105, l. 14 fr. b. — *For*: narrowly — *read*: narrowly
- P. 110, l. 12. — *For*: *strand*; — *read*: *strands*;
- P. 115, l. 5 fr. b.; P. 124, l. 1 fr. b.; P. 294, l. 10; P. 296, l. 15.—
For: appearantly — *read*: apparently
- P. 122, l. 20. — *For*: covered by — *read*: covered by the
- P. 122, l. 15—13 fr. b. (et passim). —
For: Sori on each side of the costa(e),
oblique to the latter
read: Sori on each side of the costa(e)
oblique to it (*eventually* them)
- P. 126, l. 10. — *For*: into — *read*: in
- P. 129, l. 14—15 fr. b.; P. 131, l. 16; P. 268, l. 17; P. 368, l. 9 fr. b. —
For: at utmost — *read*: at the utmost
- P. 154, l. 3 fr. b. — *For*: this — *read*: it
- P. 155, l. 12 fr. b. — *For*: beyond — *read*: above
- P. 157. — *For*: * **N. lindsayae**, — *read*: * (3) **N. lindsayae**,
- P. 166, l. 9. — *For*: ressembling — *read*: resembling
- P. 167, l. 16 fr. b. — *For*: cilate, — *read*: ciliate,
- P. 170, l. 7. — *For*: of the sorus, — *read*: as the sorus,
l. 9. — *For*: basiscope — *read*: basipetal
l. 12. — *For*: Pantropical. — *read*: Pantropical.
- P. 177, l. 19 fr. b.; P. 178, l. 13 fr. b. —
For: *without tooth* — *read*: *without a tooth*
- P. 179, l. 10 fr. b.; P. 285, l. 11. —
For: *at utmost* — *read*: *at the utmost*
- P. 189, l. 5; P. 253, l. 1. — *For*: exept — *read*: except
- P. 192. — *For*: * **D. filix mas**, — *read*: * (54) **D. filix mas**,
- P. 207, l. 13; P. 208, l. 20. —
For: *without gland* — *read*: *without a gland*
- P. 211, l. 17. — *For*: paranchyma — *read*: parenchyma
- P. 214, l. 17 fr. b. — *Strike out the comma after the*
- P. 216. — *For*: * **D. urophylla**, — *read*: * (83) **D. urophylla**,
- P. 218, l. 8 & 9. — *For*: puberous — *read*: puberulous
- P. 226, l. 10. — *Strike out the comma after largest*
- P. 229, l. 6 fr. b. — *For*: wether — *read*: whether
- P. 236, l. 16; P. 238, l. 1; P. 521, l. 8 fr. b.; P. 522, l. 14; P. 523, l. 8 fr. b.
For: in — *read*: into
- P. 236, l. 7 fr. b. — *For*: fugaceous. — *read*: fugacious.

120
Section 10 of the Act

Section 11 of the Act

Section 12

Section 12 of the Act

Section 13 of the Act

Section 14 of the Act

Section 15 of the Act

Section 16 of the Act

Section 17 of the Act


Section 18 of the Act

Section 19 of the Act

- P. 238, l. 15 (et passim). —
For: sori to a row; — *read*: sori in a row;
l. 15 fr. b. — *For*: paleaceous — *read*: paleaceous
- P. 247, l. 16. — *For*: papilliae — *read*: papillae
- P. 251, l. 2. — *For*: surface sfinely — *read*: surfaces finely
- P. 269, l. 17 fr. b. — *For*: midial — *read*: medial
- P. 270, l. 7, & l. 13 — 12 fr. b. —
For: upper and outer edges — *read*: upper and outer edge
- P. 271. — *For*: *(25) **L. borneensis**, — *read*: *(24) **L. borneensis**,
l. 16 fr. b. — *For*: Leaflets — *read*: Leaflets
- P. 276, l. 3. — *For*: anastomising — *read*: anastomosing.
- P. 277, l. 12 fr. b. — *Strike out the comma after scales*
- P. 292, l. 4 fr. b.; P. 326, l. 3; P. 650, l. 12. —
For: wich — *read*: which
- P. 293, l. 8. — *Strike out the comma after base*
- P. 303, l. 6. — *For*: elleptical, — *read*: elliptical,
- P. 304, l. 2. — *For*: rhey — *read*: they
- P. 314, l. 2. — *For*: ot — *read*: to
- P. 321, l. 10 fr. b.; P. 561, l. 10 fr. b.; P. 774, l. 7 fr. b.; P. 28, l. 8.
For: Brasilia — *read*: Brazil
- P. 327, l. 1 & 13. — *For*: straight (straihgt) — *read*: straight
- P. 329, l. 4 fr. b. — *For*: to the latter. — *read*: to it.
- P. 339, l. 10. — *For*: linear, — *read*: linear-
- P. 337, l. 2. — *For*: a white or yellow powder.
read: white or yellow powder.
- P. 345, l. 7 fr. b. — *For*: pinnaty, — *read*: pinnate,
- P. 377, l. 17. — *For*: to te — *read*: to the
- P. 380, l. 8. — *For*: New-Zeeland. — *read*: New Zealand.
- P. 381, l. 3 fr. b.; P. 464, l. 8.
For: lenght, — *read*: length,
- P. 386, l. 10 fr. b.; P. 387, l. 12. — *For*: side — *read*: side,
- P. 398, l. 12. — *For*: (inner side), — *read*: (inner) side,
- P. 410, l. 7 fr. b. — *For*: distand — *read*: distant
- P. 412, l. 12. *should be modified as follows*:

Fil., 236; — var. majus (*major*), Bedd., Handb. Ind. Ferns, Suppl., 40.

l. 13 fr. b. — *For*: major: — *read*: majus:



- P. 426, l. 4 fr. b. — *For: redu ed.* — *read: reduced.*
P. 428, l. 8 fr. b. — *For: without sinus*
— *read: without a sinus*
l. 6 fr. b. — *For: Forbes* — *read: Forbes'*
P. 439, l. 14. — *For: sessille* — *read: sessile*
P. 451, l. 20. — *For: decoumpound,* — *read: decompound*
l. 11 fr. b. — *For: l nes.* — *read: lines.*
P. 465, l. 15 fr. b.; P. 490, l. 8; P. 542, l. 8 fr. b. —
For: the latter — *read: it (occasionally them)*
P. 489, l. 12. — *For: a the base,* — *read: at the base,*
P. 508, l. 17 fr. b. — *For: olong* — *read: along*
l. 16 fr. b. — *For: contracted.* — *read: contracted.*
P. 533, l. 14 fr. b. — *For: linear-* — *read: linear-*
P. 534, l. 13. — *For: Polinesia.* — *read: Polynesia.*
P. 542, l. 13. — *For: ferlile* — *read: fertile*
P. 546, l. 1. — *For: su1-* — *read: sur-*
P. 560, l. 6, & l. 14—13 fr. b.; P. 561, l. 2. —
For: groove in — *read: groove is in*
P. 575, l. 3 fr. b. — *For: New Gainea.* — *read: New Guinea.*
P. 587, l. 17. — *For: tbe* — *read: the*
P. 608, l. 18. — *For: Polynesiu.* — *read: Polynesia.*
P. 618, l. 12 fr. b. *should be modified as follows:*

* (115a) **P. integrus** (*integriore*), *Copel.*, Philipp. Journ., II, 159.

- P. 626, l. 9—8 fr. b. — *For: iudusia* — *read: indusia*
P. 635, l. 1 fr. b. — *For: ereso-ciliate,* — *read: eroso-ciliate,*
P. 639, l. 7 fr. b. — *For: arreolae* — *read: areolae*
P. 690, l. 11. — *For: stalkek;* — *read: stalked;*
P. 709, l. 18. — *For: spicimen* — *read: specimen*
P. 756, l. 4 fr. b. — *For: an* — *read: and*
P. 775, l. 14. — *For: Americ.* — *read: America.*
P. 785, l. 1 fr. b. — *For: th elower* — *read: the lower*
P. 788, l. 15. — *For: trunc,* — *read: trunk,*
l. 11 fr. b. — *For: some times* — *read: sometimes*
P. 793, l. 2 fr. b. — *For: (furfuraceous?,* — *read: (furfuraceous?),*
P. 805, l. 7 fr. b. — *For: Midanao.* — *read: Mindanao.*
P. 818, l. 18. — *For: capsulus.* — *read: capsules.*

①



SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00613 8473